

## Using geodetic measurements to improve estimates of Antarctica's GIA and present-day mass balance

Gunter, B.C.; Engels, Olga; Riva, Riccardo; Meister, R.; Muir, Alan; King, M.A.; van den Broeke, M.R.; Hashemi Farahani, Hassan; Ditmar, Pavel; More Authors

### Publication date

2015

### Document Version

Final published version

### Citation (APA)

Gunter, B. C., Engels, O., Riva, R., Meister, R., Muir, A., King, M. A., van den Broeke, M. R., Hashemi Farahani, H., Ditmar, P., & More Authors (2015). *Using geodetic measurements to improve estimates of Antarctica's GIA and present-day mass balance*. Abstract from 26th IUGG general assembly, Prague, Czech Republic.

### Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

### Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

### Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

**JG1 Dynamics of the Cryosphere from Geometric and Gravimetric Observations (IAG, IACS)**

24-Jun-2015, 16:30 - 18:00

**Abstract content:****Using geodetic measurements to improve estimates of Antarctica's GIA and present-day mass balance**

This presentation will provide an overview and discussion of a methodology in which satellite altimetry, satellite gravimetry, and climate data sets are used to generate empirical estimates of present-day Antarctic glacial isostatic adjustment (GIA), as well as corresponding ice mass change estimates. One of the benefits of this approach is its ability to provide more reliable uncertainties of these estimates based on the error characteristics of the input observations. The resulting empirical GIA uplift rates show both similarities and differences from traditional GIA models based on ice history reconstruction. This presentation will explore some of these differences, in particular for regions such as the Amundsen Sea Sector in West Antarctica and the Phillipi/Denman sectors of East Antarctica. Comparisons will also be made between empirical estimates made with a range of satellite gravimetry (GRACE) and satellite altimetry (ICESat, Envisat) data sets. The combinations suggest that the approach has the potential to reduce the uncertainty surrounding both Antarctic GIA and ice mass change estimates and provide new insights into the impact that recent ice load changes may have on present-day uplift rates.

**Author(s):**

B. Gunter<sup>1,2</sup>, O. Didova<sup>2</sup>, R.E.M. Riva<sup>2</sup>, V.R. Barletta<sup>3</sup>, R. Meister<sup>4</sup>, A. Shepherd<sup>5</sup>, A. Muir<sup>6</sup>, M.A. King<sup>7</sup>, S.R.M. Ligtenberg<sup>8</sup>, J.T.M. Lenaerts<sup>8</sup>, M.R. van den Broeke<sup>8</sup>, H. Hashemi Farahani<sup>2</sup>, P. Ditmar<sup>2</sup>.

<sup>1</sup>Georgia Institute of Technology, School of Aerospace Engineering, Atlanta, USA.

<sup>2</sup>Delft University of Technology, Geoscience and Remote Sensing, Delft, Netherlands.

<sup>3</sup>Ohio State University, School of Earth Sciences, Columbus, USA.

<sup>4</sup>Technical University of Denmark, National Space Institute, Copenhagen, Denmark.

<sup>5</sup>University of Leeds, School of Earth and Environment, Leeds, United Kingdom.

<sup>6</sup>University College London, Earth Sciences, London, United Kingdom.

<sup>7</sup>University of Tasmania, School of Land and Food, Hobart, Australia.

<sup>8</sup>Utrecht University, Institute for Marine and Atmospheric research Utrecht IMAU, Utrecht, Netherlands.

**Keywords:**

Antarctica glacial isostatic adjustment mass balance satellite altimetry gravimetry climate data GPS

[<< Back to session](#)