

Complete genome sequence of the Escherichia coli phage Ayreon

Vlot, Marnix; Nobrega, Franklin L.; Wong, Che F.A.; Liu, Yue; Brouns, Stan J.J.

DOI

[10.1128/genomeA.01354-17](https://doi.org/10.1128/genomeA.01354-17)

Publication date

2018

Document Version

Final published version

Published in

Genome Announcements

Citation (APA)

Vlot, M., Nobrega, F. L., Wong, C. F. A., Liu, Y., & Brouns, S. J. J. (2018). Complete genome sequence of the Escherichia coli phage Ayreon. *Genome Announcements*, 6(2), Article e01354-17. <https://doi.org/10.1128/genomeA.01354-17>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.



Complete Genome Sequence of the *Escherichia coli* Phage Ayreon

Marnix Vlot,^a Franklin L. Nobrega,^b Che F. A. Wong,^{a*} Yue Liu,^{a*} Stan J. J. Brouns^{a,b}

^aLaboratory of Microbiology, Department of Agrotechnology and Food Sciences, Wageningen University, Wageningen, The Netherlands

^bDepartment of Bionanoscience, Kavli Institute of Nanoscience, Delft University of Technology, Delft, The Netherlands

ABSTRACT We report the whole-genome sequence of a new *Escherichia coli* temperate phage, Ayreon, comprising a linear double-stranded DNA (dsDNA) genome of 44,708 bp.

Phage Ayreon was isolated from pond water samples in Wageningen, The Netherlands (51°58'9.2"N, 5°40'43.1"E) using *Escherichia coli* strain KD471 as the host, a derivative of *E. coli* K-12 derivative strain KD263 (1) lacking clustered regularly interspaced short palindromic repeat (CRISPR)-associated genes *Cas1* and *Cas2*. Infected cultures were given time to allow for lysogeny. Formed lysogens were induced by using both UV exposure and mitomycin C, demonstrating that phage Ayreon is a temperate phage. Transmission electron microscopy revealed icosahedral capsids (± 57 nm) and long flexible tails (± 120 nm), which are characteristic of *Siphoviridae* (2).

Phage DNA was extracted using the SDS-proteinase K protocol previously described (3). Library preparation and sequencing were performed by BaseClear (The Netherlands) using the Illumina HiSeq 2500 platform. About 1,282,954 short reads were generated with a mean 3,478-fold coverage of the genome. The resulting sequences were *de novo* assembled using the CLC Genomics Workbench version 8.5.1. The contig sequences were corrected with Pilon version 1.11 (4) and linked using the SSPACE Premium scaffold version 2.3 (5). Gapped regions within the scaffolds were partially closed in an automated manner using GapFiller version 1.10 (6). Annotation and identification of open reading frames (ORFs) were performed using the Rapid Annotations using Subsystems Technology (RAST) annotation server (7) followed by manual curation of all predicted proteins against the NCBI protein database using BLASTp (8) and Pfam domain searches (9). tRNAs were predicted with tRNAscan-SE version 1.21 (10), and promoters and terminators were identified using motif searches for TTGACAN(15,18)TATAAT with a maximum of two mismatches and ARNold (11), respectively. The genome packaging strategy was predicted by phylogenetic analysis of the large terminase subunit (12).

Phage Ayreon has a linear double-stranded DNA with a genome size of 44,708 bp and a G+C content of 50.1%, which is very similar to the G+C content of its bacterial host (50.8%). The phage has 59 predicted ORFs, of which 25 could not be assigned to a function. Three predicted promoters and 13 predicted Rho-independent terminators were identified; no tRNAs were identified. Phylogenetic analysis of the large terminase subunit suggested that phage Ayreon uses cohesive end site (*cos*) packaging; *cos* sites are expected to be located within ~ 1 kbp upstream of the small terminase subunit (13). Considering this, the genome of phage Ayreon was opened so that it would begin with the small terminase subunit and end with the expected location of the *cos* site.

The highest degree of similarity was observed with phage cdt1 (GenBank accession number AB285204), a cyclomodulin producing prophage (14), with 78% coverage and

Received 2 November 2017 Accepted 21 November 2017 Published 11 January 2018

Citation Vlot M, Nobrega FL, Wong CFA, Liu Y, Brouns SJJ. 2018. Complete genome sequence of the *Escherichia coli* phage Ayreon. Genome Announc 6:e01354-17. <https://doi.org/10.1128/genomeA.01354-17>.

Copyright © 2018 Vlot et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Stan J. J. Brouns, stanbrouns@gmail.com.

* Present address: Che F. A. Wong, Department of Biotechnology and Food Science, Norwegian University of Science and Technology, Trondheim, Norway; Yue Liu, Laboratory of Food Microbiology, Wageningen University, Wageningen, The Netherlands.

