



Delft University of Technology

## The climate impact of hypersonic transport

Emmerig, J.; Jockel, P.; Grewe, V.

### Publication date

2020

### Document Version

Final published version

### Citation (APA)

Emmerig, J., Jockel, P., & Grewe, V. (2020). *The climate impact of hypersonic transport*. 186-186. Abstract from 3rd ECATS conference.

### Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

### Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

### Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

## THE CLIMATE IMPACT OF HYPERSONIC TRANSPORT

J. Emmerig<sup>1,2</sup>, P. Jöckel<sup>1</sup> & V. Grewe<sup>1,2</sup>

<sup>1</sup> *Institut für Physik der Atmosphäre, DLR-Oberpfaffenhofen, Germany, johannes.emmerig@dlr.de*

<sup>2</sup> *Delft University of Technology, Aircraft Noise and Climate Effects, The Netherlands*

**Abstract.** Supersonic transport was the subject of intense debate in the 1970s and commercial operation was eventually abandoned until recently due to economic and environmental concerns. Flight emissions at stratospheric altitude differ from tropospheric emissions mainly in terms of longevity. Long lifetimes of chemically reactive emissions, especially in the presence of the stratospheric ozone layer, require a detailed investigation of the long-term impact of emissions at this altitude.

Recent studies show a faster degradation of stratospheric water vapor with increasing altitude, driven by photolysis and chemical reaction with O<sup>1</sup>D. This is seen as an opportunity for civil hypersonic transport. However, the climate impact of hypersonic flight has not yet been investigated.

This is why our study focuses on the emissions of hydrogen-powered hypersonic aircraft fleets (H<sub>2</sub>O, NO<sub>x</sub>, H<sub>2</sub>) in the middle and upper stratosphere (27 and 36 km). Three different scenarios based on the HIKARI emission data allow an altitude dependent comparison of hypersonic emissions. The scenarios were simulated with ECHAM5/MESy (v2.54.0), including a newly developed submodel H2OEMIS to integrate external water vapor emissions into the cycle of specific humidity.

Additional simulations using different models for comparison are planned with Didier Hauglustaine (LSCE) in the context of project 'Stratofly' funded by EU-Horizon 2020.

### ACKNOWLEDGEMENTS

The authors received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 769246 (project STRATOFly).