

A primary cutaneous nocardiosis of the hand

Camozzota, Camilla; Goldman, Alberto; Tchernev, Georgi; Lotti, Torello; Wollina, Uwe

DOI

[10.3889/oamjms.2017.106](https://doi.org/10.3889/oamjms.2017.106)

Publication date

2017

Document Version

Final published version

Published in

Open Access Macedonian Journal of Medical Sciences

Citation (APA)

Camozzota, C., Goldman, A., Tchernev, G., Lotti, T., & Wollina, U. (2017). A primary cutaneous nocardiosis of the hand. *Open Access Macedonian Journal of Medical Sciences*, 5(4 Special Issue GlobalDermatology), 470-472. <https://doi.org/10.3889/oamjms.2017.106>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

A Primary Cutaneous Nocardiosis of the Hand

Camilla Camozzota¹, Alberto Goldman², Georgi Tchernev³, Torello Lotti⁴, Uwe Wollina⁵

¹Santa Casa de Misericórdia Hospital, Porto Alegre, Porto Alegre, RS, Brazil; ²Clinica Goldman, Porto Alegre, RS, Brazil; ³Department of Dermatology, Venereology and Dermatologic Surgery, Medical Institute of Ministry of Interior, and Onkoderma Polyclinic for Dermatology and Dermatologic Surgery, Sofia, Bulgaria; ⁴Department of Dermatology, University of Rome "G. Marconi", Rome, Italy; ⁵Department of Biotechnology, Delft University of Technology, 2628 BC, Delft, The Netherlands; ⁵Department of Dermatology and Allergology, Academic Teaching Hospital Dresden-Friedrichstadt, Friedrichstrasse 41, 01067, Dresden, Germany

Abstract

Citation: Camozzota C, Goldman A, Tchernev G, Lotti T, Wollina U. A Primary Cutaneous Nocardiosis of the Hand. Open Access Maced J Med Sci. 2017 Jul 25; 5(4):470-472. https://doi.org/10.3889/oamjms.2017.106

Keywords: Actinomycetes; Nocardiosis; Hand; Antibiosis; Sabouraud agar; Primary cutaneous infection.

***Correspondence:** Uwe Wollina, Department of Dermatology and Allergology, Academic Teaching Hospital Dresden-Friedrichstadt, Friedrichstrasse 41, 01067, Dresden, Germany. E-mail: uwollina@googlemail.com

Received: 26-Mar-2017; Revised: 10-Apr-2017; Accepted: 11-Apr-2017; Online first: 20-Jul-2017

Copyright: © 2017 Camilla Camozzota, Alberto Goldman, Georgi Tchernev, Torello Lotti, Uwe Wollina. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

Funding: This research did not receive any financial support.

Competing Interests: The authors have declared that no competing interests exist.

BACKGROUND: Nocardiosis is caused by an aerobic actinomycete, most commonly introduced through the respiratory tract. The *Nocardiae* are gram-positive, partially acid-fast bacteria. Primary cutaneous nocardiosis infections are rare and caused by the traumatic introduction of organisms percutaneously. The manifestation is usually an opportunistic infection. Cutaneous involvement may develop as one of four types: mycetoma, lymphocutaneous infection, superficial skin infection, or systemic disease with cutaneous involvement. Diagnosis and evaluation of appropriate specimens are principally by culture.

CASE PRESENTATION: A 55-year-old female patient with diabetes type II presented with chronic skin lesions on the hand. Otherwise, her medical history was unremarkable. There were no signs of systemic disease. Direct examination of swabs demonstrated gramme bacteria and culture on Sabouraud agar was positive for *Nocardia* spp. The specimen of nocardiae was not identified. The patient was treated during nine months with sulfamethoxazole plus trimethoprim. There was an important clinical improvement of the cutaneous aspect of the lesions in hand. Some scars and fibrosis remained after nocardiosis.

CONCLUSIONS: Primary cutaneous nocardiosis of the hand is a rare condition. The clinical diagnosis is difficult, and culture is mandatory.

Introduction

The genus *Nocardia* has widely distributed a group of bacteria found in soil, organic matter, fresh and salt water. *Nocardia* and *Rhodococcus* belong to the family *Nocardiaceae* of the suborder of so-called aerobic actinomycetes that includes *Mycobacterium*, *Corynebacterium*, *Gordona*, and *Tsukamurella*. *Nocardia* includes more than 50 different species, but *N. asteroides complex* is responsible for most of the human infections [1]

Nocardia is aerobic, filamentous gram-positive, atypical acid-fast bacteria that can cause localised or systemic infections mostly in immunocompromised patients, i.e. post-transplantation, in renal insufficiency, chronic lung disease, human immunodeficiency, cancer and lymphoma or HIV/AIDS. Infections occur either by inhalation or direct skin inoculation [1]

Primary cutaneous nocardiosis, however, is a rare disease characterised by nodules, subcutaneous abscess formation, ulcerations, pyoderma or cellulitis. In contrast to systemic nocardiosis, mostly immunocompetent patients get affected. The most frequently isolated species is *N. brasiliensis*. *N. asteroides* and *Nocardia otitidiscaviarum* and some other species have only occasionally been isolated [1, 2]

The major differential diagnoses of primary cutaneous nocardiosis are bacterial soft tissue infections caused by *Staphylococcus aureus* or *Streptococci spp.*, but nocardiosis tends to be more indolent. An untreated infection can develop into lymphocutaneous nocardiosis with sporotrichosis and ulceroglandular tularemia as important differential diagnoses. *Nocardia* bacteremia is uncommon [3, 4]

Case Report

A 55-year-old female patient from Brazil presented with a chronic skin lesion on the hand. She suffered from diabetes mellitus type II but had no other risk factors in her medical history.

On examination, we observed an erythematous lesion on her right hand with plaque-like thickening and superimposed partly ulcerated nodules (Fig. 1 a, b). The lesion was moderately painful. There was no lymphadenopathy. Ultrasound investigations remained unremarkable.



Figure 1: Nocardiosis of the hand (Mycetoma). Initial presentation with inflammatory plaques and nodules of the hand partially ulcerated (a, b). Complete remission with scarring after nine months of antibiotic (c, d)

Direct microbiological examination demonstrated gram-positive bacteria. After ten day-culture on Sabouraud dextrose agar at 35 degrees, Celsius colonies with a chalky appearance and purple or white colour could be identified. The texture was smooth or heaped. Microscopy of the colonies revealed gram-positive branching filaments of 0.8 μm diameter characteristic for *Nocardia spp* (Fig. 2). Unfortunately, molecular techniques for *Nocardiae*

spp were not available.

A biopsy was taken for histology which found pustules and fragments of granulation tissue were seen with neutrophilic infiltration. Granulomatous alterations were absent. Intense neutrophilic infiltrates with a pustule and abscess formation is characteristic for cutaneous infection by *Nocardia spp*. The diagnosis of primary cutaneous nocardiosis was confirmed.

The patient was treated during nine months with 160 mg trimethoprim- 800 mg sulfamethoxazole twice daily. There was an important clinical improvement of the cutaneous aspect of the lesions in hand. Some scars and fibrosis remained after nocardiosis (Fig. 1 c, d).



Figure 2: *Nocardia spp.* on Sabouraud agar. Upper part – view from above, lower part – view from below

Discussion

Nocardia species are Gram-positive, weakly acid-fast with Kinyoun stain, and non-acid-fast with the Ziehl-Neelsen stain, and develop branching filaments only in aerobic culture. *Nocardia spp.* can be grown on Sabouraud agar, which is a selective medium for these bacteria. Colony morphology and

smell are other characteristics used for their identification [1, 5].

In a specialized laboratory analysis, 16S rDNA, multilocus sequence typing (MLST) using housekeeping genes for genotyping, or matrix-assisted laser desorption ionization-time of flight mass spectrometry (MALDI-TOF MS) is employed for species identification [6-9]. Unfortunately, molecular assays were not available in our case.

In Brazil, pulmonary disease is the most common nocardiosis [7]. *N. farcinica* and *N. asiatica* have been isolated from rare cases of primary cutaneous nocardiosis in Brazil [8]. Cutaneous involvement may develop as one of four types: superficial skin infection, mycetoma, lymphocutaneous infection, or systemic disease with cutaneous involvement [1]. Our patient can be classified into the first group. We treated the patient successfully with trimethoprim-sulfamethoxazole what is the most commonly used sulfonamide preparation. Some species are often resistant to this combined drug [8, 10]. Therefore, we can exclude *N. otitidiscaviarum*, *N. nova* and *N. farcinica* as responsible species for the primary cutaneous nocardiosis we observed.

Primary cutaneous nocardiosis of the hand, as in our patient, raises possible differential diagnoses, like staphylococcal or streptococcal soft tissue infections, ulceroglandular tularemia, or sporotrichosis [1]. Epithelioid cell sarcoma is another important differential diagnosis confirmed by histopathology [11]. In the case of mycetoma of the hand, the differential diagnosis confirmed by culture could be either eumycetoma or actinomycetoma. *Nocardia spp.*, *Actinomadura spp.*, and *Streptomyces spp.* cause actinomycetoma, while eumycetoma is due to infection with *Madurella mycetomatis*, *Leptosphaeria spp.*, and related species [12]. Mycetoma of the hand can lead to severe soft tissue and bone destruction [12]. In patients with the immunocompromising disease, primary cutaneous nocardiosis of the hand may lead to severe complications such as cellulitis-like nocardiosis [13], sporotrichoid nocardiosis [14] or necrotizing nocardiosis – an emergency [15].

In conclusion, nocardiosis should be considered even in immunocompetent patients with rather indolent infections of the hands. Rapid identification *Nocardia spp.*, early and sufficient antibiotic treatment ensure a good prognosis.

References

1. Brown-Elliott BA, Brown JM, Conville PS, Wallace RJ Jr. Clinical and laboratory features of the *Nocardia* spp. Based on current molecular taxonomy. *Clin Microbiol Rev.* 2006;19:259-282.

<https://doi.org/10.1128/CMR.19.2.259-282.2006> PMID:16614249
PMCID:PMC1471991

2. Smego RA Jr, Gallis HA. The clinical spectrum of *Nocardia brasiliensis* infections in the United States. *Rev Infect Dis.* 1984;6:164-180. <https://doi.org/10.1093/clinids/6.2.164>

3. Kim YK, Sung H, Jung J, Yu SN, Lee JY, Kim SH, et al. Impact of immune status on the clinical characteristics and treatment outcomes of nocardiosis. *Diagn Microbiol Infect Dis.* 2016;85:482-487. <https://doi.org/10.1016/j.diagmicrobio.2016.05.004> PMID:27241370

4. Chen B, Tang J, Lu Z, Wang N, Gao X, Wang F. Primary cutaneous nocardiosis in a patient with nephrotic syndrome: A case report and review of the literature. *Medicine (Baltimore).* 2016;95:e2490. <https://doi.org/10.1097/MD.0000000000002490> PMID:26817885 PMCID:PMC4998259

5. Magalhães GM, Oliveira SC, Soares AC, Machado-Pinto J, de Resende MA. Mycetoma caused by *Nocardia caviae* in the first Brazilian patient. *Int J Dermatol.* 2010;49:56-58. <https://doi.org/10.1111/j.1365-4632.2009.04263.x> PMID:20465613

6. Du P, Hou X, Xie Y, Xu S, Li L, Zhang J, et al. Genotyping of *Nocardia farcinica* with multilocus sequence typing. *Eur J Clin Microbiol Infect Dis.* 2016;35:771-778. <https://doi.org/10.1007/s10096-016-2596-x> PMID:26972429

7. Baio PV, Ramos JN, dos Santos LS, Soriano MF, Ladeira EM, Souza MC, et al. Molecular identification of nocardia isolates from clinical samples and an overview of human nocardiosis in Brazil. *PLoS Negl Trop Dis.* 2013;7:e2573. <https://doi.org/10.1371/journal.pntd.0002573> PMID:24340116 PMCID:PMC3854972

8. Condas LA, Ribeiro MG, Muro MD, de Vargas AP, Matsuzawa T, Yazawa K, et al. Molecular identification and antimicrobial resistance pattern of seven clinical isolates of *Nocardia* spp. In Brazil. *Rev Inst Med Trop Sao Paulo.* 2015;57:251-256. <https://doi.org/10.1590/S0036-46652015000300012> PMID:26200967 PMCID:PMC4544251

9. Girard V, Mailler S, Polsinelli S, Jacob D, Saccomani MC, Celliere B, et al. Routine identification of *Nocardia* species by MALDI-TOF mass spectrometry. *Diagn Microbiol Infect Dis.* 2017;87:7-10. <https://doi.org/10.1016/j.diagmicrobio.2016.09.024> PMID:27802877

10. Hashemi-Shahraki A, Heidarieh P, Bostanabad SZ, Hashemzadeh M, Feizabadi MM, Schraufnagel D, et al. Genetic diversity and antimicrobial susceptibility of *Nocardia* species among patients with nocardiosis. *Sci Rep.* 2015;5:17862. <https://doi.org/10.1038/srep17862> PMID:26638771 PMCID:PMC4671095

11. Wollina U, Schönlebe J, Haroske G, Unger L, Kittner T, Tchernev G, Chokoeva AA, Lotti T. Nodular epithelioid sarcoma of the upper limb. A case report and review of the literature. *J Biol Regul Homeost Agents.* 2015;29(Suppl):23-26. PMID:26016962

12. Omer RF, Seif El Din N, Abdel Rahim FA, Fahal AH. Hand mycetoma: The Mycetoma Research Centre experience and literature review. *PLoS Negl Trop Dis.* 2016;10:e0004886. <https://doi.org/10.1371/journal.pntd.0004886> PMID:27483367 PMCID:PMC4970814

13. Asgeirsson H, Sigurdardottir B. Nocardiosis in immunocompromised host presenting as cellulitis. *Laeknabladid.* 2010;96:423-425. PMID:20519773

14. Wang AW, D'Cruz M, Leung M. Primary cutaneous nocardiosis of the hand: a case report and literature review. *Hand Surg.* 2002;7:155-157. <https://doi.org/10.1142/S0218810402000868> PMID:12365059

15. Ricci JA, Weil AA, Eberlin KR. Necrotizing cutaneous nocardiosis of the hand: A case report and review of the literature. *J Hand Microsurg.* 2015;7:224-227. <https://doi.org/10.1007/s12593-015-0173-7> PMID:26078549 PMCID:PMC4461627