

Differences in mouthbar complex geometry and grain size trends across systems with different sediment supply compositions

van der Vegt, Helena; Storms, Joep; Walstra, Dirk-Jan; Li, Liang; Nordahl, Kjetil; Martinius, Allard; Howes, Nick

Publication date

2017

Document Version

Final published version

Citation (APA)

van der Vegt, H., Storms, J., Walstra, D.-J., Li, L., Nordahl, K., Martinius, A., & Howes, N. (2017). *Differences in mouthbar complex geometry and grain size trends across systems with different sediment supply compositions*. Abstract from EGU General Assembly 2017, Vienna, Austria.

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.



Differences in mouthbar complex geometry and grain size trends across systems with different sediment supply compositions

Helena van der Vegt (1), Joep Storms (1), Dirk-Jan Walstra (1,2), Liang Li (1), Kjetil Nordahl (3), Allard Martinius (3,1), and Nick Howes (4)

(1) Delft University of Technology, Delft, The Netherlands, (2) Deltares, Delft, The Netherlands, (3) Statoil Research Centre, Trondheim, Norway, (4) Shell Projects and Technology, Houston, USA

In deltaic environments, preserved mouth bar deposits are the sand-rich building blocks of the delta front. While different sediment supply composition will necessarily lead to different grain size trends in the mouth bars, this relationship may not always be linear. We investigate this relationship through a set of process based models of delta formation, using the well-established modelling software Delft3D. We show how the different sediment supply composition generates deltas with different channel morphologies and -dynamics. These channel networks act as the conduits through which sediment is transported to the delta front. In addition to the sediment supply composition, the different morphologies and dynamics of the channel networks represent an additional control on the depositional trends and preservation potential of the mouth bar complexes.