

**Upper and lower bounds for the optimal constant in the extended Sobolev inequality.  
Derivation and numerical results**

Nasibov, Sh M.; Veling, E. J.M.

**DOI**

[10.7153/jmi-2019-13-52](https://doi.org/10.7153/jmi-2019-13-52)

**Publication date**

2019

**Document Version**

Final published version

**Published in**

Journal of Mathematical Inequalities

**Citation (APA)**

Nasibov, S. M., & Veling, E. J. M. (2019). Upper and lower bounds for the optimal constant in the extended Sobolev inequality. Derivation and numerical results. *Journal of Mathematical Inequalities*, 13(3), 753-778. <https://doi.org/10.7153/jmi-2019-13-52>

**Important note**

To cite this publication, please use the final published version (if applicable).  
Please check the document version above.

**Copyright**

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

**Takedown policy**

Please contact us and provide details if you believe this document breaches copyrights.  
We will remove access to the work immediately and investigate your claim.

## UPPER AND LOWER BOUNDS FOR THE OPTIMAL CONSTANT IN THE EXTENDED SOBOLEV INEQUALITY. DERIVATION AND NUMERICAL RESULTS

SH. M. NASIBOV AND E. J. M. VELING\*

(Communicated by J. Pečarić)

*Abstract.* We prove and give numerical results for two lower bounds and eleven upper bounds to the optimal constant  $k_0 = k_0(n, \alpha)$  in the inequality

$$\|u\|_{2n/(n-2\alpha)} \leq k_0 \|\nabla u\|_2^\alpha \|u\|_2^{1-\alpha}, \quad u \in H^1(\mathbb{R}^n),$$

for  $n = 1$ ,  $0 < \alpha \leq 1/2$ , and  $n \geq 2$ ,  $0 < \alpha < 1$ .

This constant  $k_0$  is the reciprocal of the infimum  $\lambda_{n,\alpha}$  for  $u \in H^1(\mathbb{R}^n)$  of the functional

$$\Lambda_{n,\alpha} = \frac{\|\nabla u\|_2^\alpha \|u\|_2^{1-\alpha}}{\|u\|_{2n/(n-2\alpha)}}, \quad u \in H^1(\mathbb{R}^n),$$

where for  $n = 1$ ,  $0 < \alpha \leq 1/2$ , and for  $n \geq 2$ ,  $0 < \alpha < 1$ .

The lowest point in the point spectrum of the Schrödinger operator  $\tau = -\Delta + q$  on  $\mathbb{R}^n$  with the real-valued potential  $q$  can be expressed in  $\lambda_{n,\alpha}$  for all  $q_- = \max(0, -q) \in L^p(\mathbb{R}^n)$ , for  $n = 1$ ,  $1 \leq p < \infty$ , and  $n \geq 2$ ,  $n/2 < p < \infty$ , and the norm  $\|q_-\|_p$ .

### 1. Introduction

Here, we present the derivations and the results of some numerical evaluations for the optimal constant  $k_0 = k_0(n, \alpha)$  in the estimate

$$\|u\|_{2n/(n-2\alpha)} \leq k_0 \|\nabla u\|_2^\alpha \|u\|_2^{1-\alpha}, \quad u \in H^1(\mathbb{R}^n), \quad (1)$$

for  $n = 1$ ,  $0 < \alpha \leq 1/2$ , and  $n \geq 2$ ,  $0 < \alpha < 1$ .

For  $n = 1$ ,  $k_0$  is known explicitly (see [1], [2], [3] and [4, Lemma 2.1, (2.4)])

$$k_0(1, \alpha) = 2^\alpha \alpha^{\alpha/2} (1 - \alpha)^{-(1-\alpha)/2} (1 - 2\alpha)^{(1-2\alpha)/2} B\left(\frac{1}{2}, \frac{1}{2\alpha}\right)^{-\alpha}, \quad (2)$$

for  $0 < \alpha < 1/2$ , and  $k_0(1, 1/2) = 1$ ,

*Mathematics subject classification* (2010): 26D15, 41A44.

*Keywords and phrases:* Sobolev inequality, optimal constant, lower bound, upper bound.

\* Corresponding author.

where  $B(p, q)$  is the Beta Function

$$B(p, q) = \int_0^1 x^{p-1}(1-x)^{q-1} dx = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}, \quad \Re p > 0, \quad \Re q > 0. \quad (3)$$

For  $n \geq 2$ , a number of authors has dealt with estimates for  $k_0(n, \alpha)$  for some specific values or in a general sense: [5], [6], [7], [8], [9], [10], [11], [4], [12], [13], [14], [15].

The value  $k_0$  equals the reciprocal value of the infimum  $\lambda_{n,\alpha}$  of the functional  $\Lambda_{n,\alpha}$ :

$$\lambda_{n,\alpha} = \inf_{u \in H^1(\mathbb{R}^n)} \Lambda_{n,\alpha}, \quad \text{with} \quad (4)$$

$$\Lambda_{n,\alpha} = \frac{\|\nabla u\|_2^\alpha \|u\|_2^{1-\alpha}}{\|u\|_{2n/(n-2\alpha)}}, \quad u \in H^1(\mathbb{R}^n), \quad (5)$$

where  $0 < \alpha \leq 1/2$  if  $n = 1$ , and  $0 < \alpha < 1$  if  $n \geq 2$ .

One of the motivations to study this functional comes from the fact that the lowest point in the point spectrum of the Schrödinger operator can be expressed by the infimum  $\lambda_{n,\alpha}$  of this functional  $\Lambda_{n,\alpha}$ . So, for the Schrödinger operator  $\tau = -\Delta + q$  on  $\mathbb{R}^n$  with the real-valued potential  $q$  such that  $q = q_+ - q_-$ , where

$$q_+ = \max(0, q) \in L_{loc}^2(\mathbb{R}^n), \quad (6)$$

$$q_- = \max(0, -q) \in L^p(\mathbb{R}^n), \quad \begin{aligned} n = 1: & \quad 1 \leq p < \infty, \\ n \geq 2: & \quad n/2 < p < \infty. \end{aligned} \quad (7)$$

the lowest point in the point spectrum for all such  $q$  expressed as

$$l(n, \alpha) = \inf_{q_- \in L^p(\mathbb{R}^n)} \inf_{u \in H^1(\mathbb{R}^n)} \frac{\|\nabla u\|_2^2 + \int_{\mathbb{R}^n} q |u|^2 dx}{\|u\|_2^2} \|q_-\|_p^{-1/(1-\alpha)}, \quad (8)$$

with  $\alpha = n/(2p)$ ,

will be

$$l(n, \alpha) = -(1-\alpha)\alpha^{\alpha/(1-\alpha)}\lambda_{n,\alpha}^{-2/(1-\alpha)}, \quad \begin{aligned} 0 < \alpha \leq 1/2 \text{ if } n = 1, \\ 0 < \alpha < 1 \text{ if } n \geq 2, \end{aligned} \quad (9)$$

see among others [10], [4].

The corresponding Euler equation belonging to the infimum  $\lambda_{n,\alpha}$  of the functional  $\Lambda_{n,\alpha}(u)$  reads

$$-\alpha \frac{\Delta u}{\|\nabla u\|_2^2} + (1-\alpha) \frac{u}{\|u\|_2^2} - \frac{u|u|^\rho}{\|u\|_{\rho+2}^{\rho+2}} = 0, \quad (10)$$

$$\text{with } \rho = \frac{4\alpha}{(n-2\alpha)}, \quad \alpha = \frac{\rho n}{2(\rho+2)},$$

which can be scaled in the form (see [10], [4])

$$\begin{aligned}
 &-\frac{d^2}{dr^2}u - \frac{(n-1)}{r} \frac{d}{dr}u - u|u|^p + u = 0, \quad r = |x| > 0, \\
 &\frac{d}{dr}u(0) = 0, \quad \lim_{r \rightarrow \infty} u(r) = 0.
 \end{aligned} \tag{11}$$

We have used a scaling such that

$$\alpha \|u\|_2^2 = (1 - \alpha) \|\nabla u\|_2^2 = \alpha(1 - \alpha) \|u\|_{\rho+2}^{\rho+2}, \tag{12}$$

which is always possible by scaling the function and the argument. And the infimum  $\lambda_{n,\alpha}$  will then be found as (with  $\bar{u}_{n,\alpha}$  the unique positive (see [16]) solution of (11))

$$\frac{1}{k_0(n, \alpha)} = \lambda_{n,\alpha} = \alpha^{\alpha/2} (1 - \alpha)^{(n(1-\alpha)-2\alpha)/(2n)} \left[ \|\bar{u}_{n,\alpha}\|_2^2 \right]^{\alpha/n} = \chi(\alpha) \left( \frac{\|\bar{u}_{n,\alpha}\|_2^2}{1 - \alpha} \right)^{\alpha/n},$$

for  $0 < \alpha < 1, n \geq 2,$  (13)

with  $\chi(\alpha) = \sqrt{\alpha^\alpha (1 - \alpha)^{1-\alpha}}.$  (14)

The values  $k_0(n, \alpha)$  for  $\alpha = 1$  is covered by the special form of the Sobolev embedding

$$\|w\|_t \leq \frac{1}{C_T(n, s)} \|\nabla w\|_s, \quad t = sn/(n - s), \quad 1 \leq s < n, \quad w \in H^{1,s}(\mathbb{R}^n), \tag{15}$$

where  $C_T(n, s)$  is the optimal constant and

$$\begin{aligned}
 H^{1,s}(\mathbb{R}^n) &= \text{completion of } \{w \mid w \in C^1(\mathbb{R}^n), \|u\|_{1,s}^s = \|u\|_s^s + \|\nabla u\|_s^s < \infty\} \\
 &\text{with respect to the norm } \|\cdot\|_{1,s}.
 \end{aligned} \tag{16}$$

If we take  $\alpha = 1$  and  $s = 2$  in (1), we have  $k_0(n, 1) = 1/\lambda_{n,1} = 1/C_T(n, 2), n \geq 3$ . Since  $H^1(\mathbb{R}^2) \not\hookrightarrow L^\infty(\mathbb{R}^2)$ , it follows that  $\lambda_{2,1} = C_T(2, 2) = 0$ , and so  $k_0(2, 1)$  is not defined. The numbers  $C_T(n, s)$  are known explicitly by the work of [17] and [18], see also [19]

$$C_T(n, s) = n^{1/s} \left( \frac{n-s}{s-1} \right)^{(s-1)/s} \left[ \sigma_n B \left( \frac{n}{s}, n + 1 - \frac{n}{s} \right) \right]^{1/n}, \quad 1 < s < n, \tag{17}$$

$$C_T(n, 1) = n\omega_n^{1/n}, \quad n \geq 2, \tag{18}$$

where  $\sigma_n$  the surface area of the unit ball in  $\mathbb{R}^n$ ,  $\omega_n$  the volume of the unit ball in  $\mathbb{R}^n$

$$\omega_n = \pi^{n/2} / \Gamma(1 + n/2), \tag{19}$$

$$\sigma_n = n\omega_n = 2\pi^{n/2} / \Gamma(n/2), \tag{20}$$

$$B(a, b) = \Gamma(a)\Gamma(b) / \Gamma(a + b), \quad a, b > 0, \tag{21}$$

and there is equality in (15) for functions of the form

$$w_{n,s}(x_1, \dots, x_n) = \left\{ a + b|x|^{s/(s-1)} \right\}^{1-n/s}, \quad a, b > 0, \quad 1 < s < n. \quad (22)$$

From now on, we concentrate on the optimal constant  $k_0(n, \alpha)$ . Firstly, we list a number of estimates, two lower bounds and eleven different upper bounds for  $k_0(n, \alpha)$  with references if published. Thereafter, we proof the estimates also for the published bounds.

## 2. Lower bounds

### 2.1. Lower bound 1

$$k_0 > \underline{k}_0(\alpha) = \left[ \frac{\alpha^\alpha}{\pi^\alpha e^\alpha (1-\alpha)^\alpha \left[ \ln \left( \frac{1}{1-\alpha} \right) \right]^\alpha} \right]^{1/2}, \quad n = 2, \quad 0 < \alpha < 1. \quad (23)$$

### 2.2. Lower bound 2

$$k_0 > \underline{\underline{k}}_0(n, \alpha) = \left[ \frac{1}{n^n} \left( \frac{2}{\pi} \right)^{2\alpha} (n - 2\alpha)^{n-2\alpha} \right]^{1/4}, \quad n \geq 2, \quad 0 < \alpha < 1. \quad (24)$$

## 3. Upper bounds

### 3.1. Upper bound 1

$$k_0 < \overline{k}_0(n, \alpha) = \frac{1}{\chi(\alpha)} \left[ \frac{\sigma_n}{2} B \left( \frac{n}{2}, \frac{n(1-\alpha)}{2\alpha} \right) \right]^{\alpha/n} k_B \left( \frac{2n}{n+2\alpha} \right), \quad (25)$$

for  $n \geq 2$ ,  $0 < \alpha < 1$ ,

with  $\chi(\alpha)$  defined in (14),  $\sigma_n$  defined in (20),

with  $B(p, q)$  defined in (3),

$$\text{and with } k_B(p) = \left[ \left( \frac{p}{2\pi} \right)^{1/p} \left( \frac{p'}{2\pi} \right)^{-1/p'} \right]^{n/2}, \quad \frac{1}{p} + \frac{1}{p'} = 1. \quad (26)$$

See [10, Theorem 1], [12, Proposition 1] and [15, Theorem 1]. Remark that

$$n = 2, \quad B \left( 1, \frac{1-\alpha}{\alpha} \right) = \frac{\alpha}{1-\alpha}.$$

### 3.2. Upper bound 2

$$k_0 < \overline{\overline{k_0}}(n, \alpha) = \frac{1}{\chi(\alpha)} \left[ k_B \left( \frac{n}{n-2\alpha} \right) k_B^2 \left( \frac{2n}{n+2\alpha} \right) \|G(x)\|_{n/(n-2\alpha)} \right]^{1/2}, \quad (27)$$

for  $n \geq 2$ ,  $0 < \alpha < 1$ ,

with  $\chi(\alpha)$  defined in (14),  $k_B(p)$  defined in (26),

$$\text{and with } G(x) = \frac{K_{(n-2)/2}(|x|)}{|x|^{(n-2)/2}}, \quad K_\nu \text{ is the modified Bessel function.} \quad (28)$$

See [10, Theorem 2] and [15].

Remark that for  $n = 2$ ,  $\alpha = 1/2$

$$\|G(x)\|_2 = \left( 2\pi \int_0^\infty K_0^2(r) r dr \right)^{1/2} = \pi^{1/2},$$

and for  $n = 3$ , and general  $\alpha$

$$\|G(x)\|_{3/(3-2\alpha)} = \sqrt{\frac{\pi}{2}} (4\pi)^{(3-2\alpha)/3} \left( \frac{3-2\alpha}{3} \right)^{2-2\alpha} \left[ \Gamma \left( \frac{6-6\alpha}{3-2\alpha} \right) \right]^{(3-2\alpha)/3},$$

because  $K_{1/2}(x) = \sqrt{\frac{\pi}{2x}} \exp(-x)$ .

### 3.3. Upper bound 3

$$k_0 < \overline{\overline{\overline{k_0}}}(n, \alpha) = \frac{1}{\chi(\alpha)} \frac{1}{\sqrt{(1-\alpha)}} k_B \left( \frac{n}{n-\alpha} \right) k_B \left( \frac{2n}{n+2\alpha} \right) \times \|G(x)\|_{n/(n-\alpha)}, \quad \text{for } n \geq 2, \quad 0 < \alpha < 1, \quad (29)$$

with  $\chi(\alpha)$  defined in (14), with  $k_B(p)$  defined in (26),

and with  $G(x)$  defined in (28).

### 3.4. Upper bound 4

$$k_0 < \overline{\overline{\overline{\overline{k_D,1}}}}(n, \alpha) = A(n, \alpha)^\gamma, \quad n \geq 2, \quad 0 < \alpha < 1, \quad (30)$$

$$\text{with } A(n, \alpha) = \left[ \frac{2\alpha(n-\alpha)}{\pi n(n-2\alpha)^2} \right]^{\theta/2} \left[ 1 - \frac{n\alpha}{2(n-\alpha)} \right]^{(n-2\alpha)/(2n)} \quad (31)$$

$$\times \left[ \frac{\Gamma\left(\frac{n}{\alpha} - 1\right)}{\Gamma\left(\frac{n}{\alpha} - 1 - \frac{n}{2}\right)} \right]^{\theta/n},$$

and with  $\theta = \frac{\alpha(n - 2\alpha)}{2n - 2\alpha - \alpha n}$ ,  $\gamma = \frac{2n - 2\alpha - \alpha n}{n - 2\alpha}$ . (32)

### 3.5. Upper bound 5

$$k_0 < \overline{k_{D,2}}(n, \alpha) = A(n, \alpha)^\alpha \overline{k_0}(n, \alpha)^{1-\theta}, \quad n \geq 2, \quad 0 < \alpha < 1, \quad (33)$$

with  $A(n, \alpha)$  defined in (31),  $\overline{k_0}(n, \alpha)$  defined in (25),

and with  $\theta = \frac{\alpha(n - 2\alpha)}{2n - 2\alpha - \alpha n}$ , defined in (32).

Compare [4, Theorem 1.7 (1.30)].

### 3.6. Upper bound 6

$$k_0 < \overline{k_{D,3}}(n, \alpha) = A(n, \alpha)^\alpha \overline{k_0}(n, \alpha)^{1-\theta}, \quad n \geq 2, \quad 0 < \alpha < 1, \quad (34)$$

with  $A(n, \alpha)$  defined in (31),  $\theta$  defined in (32),

and with  $\overline{k_0}(n, \alpha)$  defined in (27).

Compare [4, Theorem 1.7 (1.30)].

### 3.7. Upper bound 7

$$k_0 < \overline{k_{I,1}}(n, \alpha) = 1/k_{V,1}(n, \alpha), \quad n \geq 3, \quad 1/2 < \alpha < 1, \quad (35)$$

$$\text{with } k_{V,1}(n, \alpha) = \overline{k_0} \left( n, \frac{1}{2} \right)^{-\alpha_1} k_T(n)^{-(1-\alpha_1)}, \quad \alpha_1 = 2(1 - \alpha), \quad (36)$$

with  $\overline{k_0}(n, \alpha)$  defined in (25),

$$\text{and with } k_T(n) = \frac{1}{C_T(n, 2)} = \frac{1}{\sqrt{\pi n(n-2)}} \left[ \frac{\Gamma(n)}{\Gamma\left(\frac{n}{2}\right)} \right]^{1/n}, \quad (37)$$

where  $C_T(n, 2)$  is defined in (17).

See [4, Theorem 1.7, (1.30),  $\theta' = 1/2$ ,  $\theta'' = 1$ , with the restriction  $n \geq 3$ ].

### 3.8. Upper bound 8

$$k_0 < \overline{k_{I,2}}(n, \alpha) = 1/k_{V,2}(n, \alpha), \quad n \geq 3, \quad \alpha_V < \alpha < 1, \quad (38)$$

$$\text{with } k_{V,2}(n, \alpha) = \overline{k_0}(n, \alpha_V)^{-\alpha_2} k_T(n)^{-(1-\alpha_2)}, \quad (39)$$

with  $\overline{k_0}(n, \alpha)$  defined in (25),  $k_T(n)$  defined in (37),

$$\text{and with } \alpha_2 = \frac{1-\alpha}{1-\alpha_V}, \quad (40)$$

where  $\alpha_V$  follows from

$$\alpha_V = \alpha_V(n) = \frac{n}{2p_V}, \quad \text{where } p_V \text{ is the solution of} \quad (41)$$

$$\ln\left(\frac{n-p}{p-1}\right) + \frac{n-p}{p(p-1)} + \psi(p) - \psi(n+1-p) = 0, \quad (42)$$

$$\psi(x) = \frac{\frac{d}{dx}\Gamma(x)}{\Gamma(x)}, \quad x > 0, \quad 1 < p < n, \quad n \geq 2.$$

See [4, Theorem 1.7 (1.30),  $\theta' = \theta_N (= \alpha_V)$ ,  $\theta'' = 1$ , with the restriction  $n \geq 3$ ]. See Section 5.3 for numerical values of  $\alpha_V(n)$ ,  $n = 2, \dots, 10$ .

### 3.9. Upper bound 9

$$k_0 < \overline{k_{I,3}}(n, \alpha) = 1/k_{V,3}(n, \alpha), \quad n \geq 3, \quad \alpha_V < \alpha < 1, \quad (43)$$

with  $\alpha_V$  defined in (41),

$$\text{with } k_{V,3}(n, \alpha) = k_{L,V}(n, \alpha_V)^{\alpha_2} k_T(n)^{-(1-\alpha_2)}, \quad \alpha_2 \text{ defined in (40),} \quad (44)$$

$$\text{with } k_{L,V}(n, \alpha) = [\alpha C_T(n, 2\alpha)]^\alpha, \quad (45)$$

with  $C_T(n, s)$  defined in (17), that is

$$C_T(n, s) = n^{1/s} \left(\frac{n-s}{s-1}\right)^{(s-1)/s} \left[\sigma_n B\left(\frac{n}{s}, n+1-\frac{n}{s}\right)\right]^{1/n}, \quad 1 < s < n,$$

and with  $k_T(n)$  defined in (37),  $k_T(n) = 1/C_T(n, 2)$ .

Compare [4, Theorem 1.7 (1.30) and (1.32),  $\theta' = \theta_N (= \alpha_V)$ ,  $\theta'' = 1$ , with the restriction  $n \geq 3$ ].

### 3.10. Upper bound 10

$$k_0 < \overline{k_{L,V}}(n, \alpha) = [\alpha_V C_T(n, 2\alpha_V)]^{-\alpha}, \quad (46)$$



$$\begin{aligned}
 n &\geq 2, \quad 0 < \alpha \leq \alpha_V, \\
 k_0 &< \overline{k_{L,V}}(n, \alpha) = 1/k_{L,V}(n, \alpha) = [\alpha C_T(n, 2\alpha)]^{-\alpha}, \\
 n &\geq 2, \quad \alpha_V \leq \alpha < 1,
 \end{aligned} \tag{47}$$

with  $\alpha_V$  defined in (41),  $C_T(n, s)$  defined in (17).

See [4, Theorem 1.7, (1.32)].

### 3.11. Upper bound 11

$$k_0 < \overline{k_B}(n, \alpha) = k_T(n)^\alpha, \quad n \geq 3, \quad 0 < \alpha < 1, \tag{48}$$

with  $k_T(n)$  defined in (37).

See [4, Theorem 1.7 (1.33),  $\theta' = 0$ ,  $\theta'' = 1$ , with the restriction  $n \geq 3$ ].

## 4. Proofs

### 4.1. Lower bounds

We take as trial function in (5) the function

$$u_{n,\alpha} = a \exp(-br^\mu), \quad a, b, \mu > 0. \tag{49}$$

We need the following general integral (see [20, (5.9.1)])

$$\int_0^\infty \exp(-mr^\mu) r^{\nu-1} dr = \frac{1}{\mu} \left(\frac{1}{m}\right)^{\nu/\mu} \Gamma\left(\frac{\nu}{\mu}\right). \tag{50}$$

For this trial function the following three integrals become ( $\sigma_n = 2\pi^{n/2}/\Gamma(n/2)$ , the surface area of the unit ball in  $\mathbb{R}^n$ , see (20))

$$\int_{\mathbb{R}^n} u_{n,\alpha}^2(x) dx = \sigma_n \int_0^\infty a^2 e^{-2br^\mu} r^{n-1} dr = \sigma_n a^2 \frac{1}{\mu} \left(\frac{1}{2b}\right)^{n/\mu} \Gamma\left(\frac{n}{\mu}\right), \tag{51}$$

$$\begin{aligned}
 \int_{\mathbb{R}^n} (\nabla u_{n,\alpha}(x))^2 dx &= \sigma_n \int_0^\infty a^2 b^2 \mu^2 r^{2(\mu-1)} e^{-2br^\mu} r^{n-1} dr \\
 &= \sigma_n a^2 \frac{\mu}{4} \left(\frac{1}{2b}\right)^{(n-2)/\mu} \Gamma\left(2 + \frac{n-2}{\mu}\right),
 \end{aligned} \tag{52}$$

$$\begin{aligned}
 \int_{\mathbb{R}^n} u_{n,\alpha}^{\rho+2}(x) dx &= \sigma_n \int_0^\infty a^{\rho+2} e^{-(\rho+2)br^\mu} r^{n-1} dr \\
 &= \sigma_n a^{\rho+2} \frac{1}{\mu} \left(\frac{1}{(\rho+2)b}\right)^{n/\mu} \Gamma\left(\frac{n}{\mu}\right).
 \end{aligned} \tag{53}$$

**4.2. Lower bound 1**

For  $n = 2$ , and general  $\mu$  the three integrals (51), (52) and (53) become

$$\int_{\mathbb{R}^2} u_{2,\alpha}^2(x) dx = 2\pi \int_0^\infty a^2 e^{-2br^\mu} r dr = \sigma_2 a^2 \frac{1}{\mu} \left(\frac{1}{2b}\right)^{2/\mu} \Gamma\left(\frac{2}{\mu}\right), \tag{54}$$

$$\int_{\mathbb{R}^2} (\nabla u_{2,\alpha}(x))^2 dx = 2\pi \int_0^\infty a^2 b^2 \mu^2 r^{2(\mu-1)} e^{-2br^\mu} r dr = \sigma_2 a^2 \frac{\mu}{4} \Gamma(2), \tag{55}$$

$$\begin{aligned} \int_{\mathbb{R}^2} u_{2,\alpha}^{\rho+2}(x) dx &= 2\pi \int_0^\infty a^{\rho+2} e^{-(\rho+2)br^\mu} r dr \\ &= \sigma_2 a^{\rho+2} \frac{1}{\mu} \left(\frac{1}{(\rho+2)b}\right)^{2/\mu} \Gamma\left(\frac{2}{\mu}\right). \end{aligned} \tag{56}$$

Let  $a, b$  be variable and  $\mu$  fixed, we use the two scaling relations (12)

$$\alpha \sigma_2 a^2 \frac{1}{\mu} \left(\frac{1}{2b}\right)^{2/\mu} \Gamma\left(\frac{2}{\mu}\right) = (1 - \alpha) \sigma_2 a^2 \frac{\mu}{4} \Gamma(2), \tag{57}$$

$$\sigma_2 a^2 \frac{1}{\mu} \left(\frac{1}{2b}\right)^{2/\mu} \Gamma\left(\frac{2}{\mu}\right) = (1 - \alpha) \sigma_2 a^{\rho+2} \frac{1}{\mu} \left(\frac{1}{(\rho+2)b}\right)^{2/\mu} \Gamma\left(\frac{2}{\mu}\right). \tag{58}$$

This gives for the optimal values for  $(a, b) = (a_0, b_0)$

$$a^\rho = a_0^\rho = \left(\frac{\rho+2}{2}\right)^{\frac{\mu+2}{\mu}}, \quad b^{2/\mu} = b_0^{2/\mu} = \frac{2\rho\Gamma\left(\frac{2}{\mu}\right)}{\mu^2 2^{2/\mu}}.$$

$$\begin{aligned} k_0(2, \alpha) &= \frac{1}{\chi(\alpha)} \left(\frac{1 - \alpha}{\|\bar{u}_{2,\alpha}\|_2^2}\right)^{\alpha/2} \\ &> \underline{k}_0(\alpha) = \frac{1}{\chi(\alpha)} \left\{ \frac{(1 - \alpha)2\rho}{2\pi \left[\mu^{\rho/2} \left(\frac{\rho}{2} + 1\right)^{1+2/\mu}\right]^{2/\rho}} \right\}^{\alpha/2}. \end{aligned} \tag{59}$$

Consider now  $\mu$  as variable to minimize  $\underline{k}_0(\alpha)$  by maximizing the denominator

$$\begin{aligned} \max_{0 < \mu < \infty} \left[\mu^{\rho/2} \left(\frac{\rho}{2} + 1\right)^{1+2/\mu}\right] &= \left[\frac{2e \ln(1 + \rho/2)}{\rho/2}\right]^{\rho/2} (1 + \rho/2), \\ \text{for } \mu_0 &= \frac{2 \ln(1 + \rho/2)}{\rho/2}. \end{aligned}$$

This gives for (59)

$$\underline{k}_0(\alpha) = \frac{1}{\chi(\alpha)} \left\{ \frac{2(1 - \alpha)(\rho/2)^2}{2\pi e \ln(1 + \rho/2) (1 + \rho/2)^{2/\rho}} \right\}^{\alpha/2}$$

$$= \left[ \frac{\alpha^\alpha}{\pi^\alpha e^\alpha (1-\alpha)^\alpha \left[ \ln \left( \frac{1}{1-\alpha} \right) \right]^\alpha} \right]^{1/2}, \quad (60)$$

which equals (23).

### 4.3. Lower bound 2

For general  $n$  and  $\mu = 2$  the three integrals (51), (52) and (53) become

$$\int_{\mathbb{R}^n} u_{n,\alpha}^2(x) dx = \sigma_n \int_0^\infty a^2 \exp(-2br^2) r^{n-1} dr = \sigma_n a^2 \frac{1}{2} \left( \frac{1}{2b} \right)^{n/2} \Gamma\left(\frac{n}{2}\right), \quad (61)$$

$$\begin{aligned} \int_{\mathbb{R}^n} (\nabla u_{n,\alpha}(x))^2 dx &= \sigma_n \int_0^\infty a^2 b^2 4r^2 \exp(-2br^2) r^{n-1} dr \\ &= \sigma_n a^2 \frac{1}{2} \left( \frac{1}{2b} \right)^{(n-2)/2} \Gamma\left(1 + \frac{n}{2}\right), \end{aligned} \quad (62)$$

$$\begin{aligned} \int_{\mathbb{R}^n} u_{n,\alpha}^{\rho+2}(x) dx &= \sigma_n \int_0^\infty a^2 \exp(-(\rho+2)r^2) r^{n-1} dr \\ &= \sigma_n a^{\rho+2} \frac{1}{2} \left( \frac{1}{(\rho+2)b} \right)^{n/2} \Gamma\left(\frac{n}{2}\right). \end{aligned} \quad (63)$$

Using the two scaling relations (12)

$$\alpha \sigma_n a^2 \frac{1}{2} \left( \frac{1}{2b} \right)^{n/2} \Gamma\left(\frac{n}{2}\right) = (1-\alpha) \sigma_n a^2 \frac{1}{2} \left( \frac{1}{2b} \right)^{(n-2)/2} \Gamma\left(1 + \frac{n}{2}\right), \quad (64)$$

$$\sigma_n a^2 \frac{1}{2} \left( \frac{1}{2b} \right)^{n/2} \Gamma\left(\frac{n}{2}\right) = (1-\alpha) \sigma_n a^{\rho+2} \frac{1}{2} \left( \frac{1}{(\rho+2)b} \right)^{n/2} \Gamma\left(\frac{n}{2}\right), \quad (65)$$

we get  $(a, b) = (a_0, b_0)$

$$a^\rho = a_0^\rho = \frac{1}{1-\alpha} \left( \frac{n}{n-2\alpha} \right)^{n/2}, \quad b = b_0 = \frac{\alpha}{n(1-\alpha)},$$

where we use all the time the relation  $\rho = \frac{4\alpha}{n-2\alpha}$ . Using (61) and (13) we find lower bound 2 (24)

$$\underline{k}_0(n, \alpha) = \left[ \frac{1}{n^n} \left( \frac{2}{\pi} \right)^{2\alpha} (n-2\alpha)^{n-2\alpha} \right]^{1/4}, \quad n \geq 2, \quad 0 < \alpha < 1. \quad (66)$$

### 4.4. Upper bounds

We introduce the standard notations

$$r = \frac{2n}{n-2\alpha}, \quad \rho = r-2 = \frac{4\alpha}{n-2\alpha}, \quad (67)$$

and so

$$\alpha = \frac{\rho n}{2(\rho + 2)} = \frac{n}{2} \left( \frac{r-2}{r} \right). \quad (68)$$

For the proof of upper bound 1 we need a less well-known inequality which we present here as Lemma.

LEMMA 1. See [21] and [13, Lemma 1]. For  $u \in L^2(\mathbb{R}^n)$ ,  $|x|u \in L^2(\mathbb{R}^n)$ ,  $x \in \mathbb{R}^n$ ,  $0 < \alpha < 1$ ,

$$\|u\|_{\frac{2n}{n+2\alpha}} \leq \frac{1}{\chi(\alpha)} \left[ \frac{\sigma_n}{2} B \left( \frac{n}{2}, \frac{n(1-\alpha)}{2\alpha} \right) \right]^{\alpha/n} \| |x|u \|_2^\alpha \|u\|_2^{1-\alpha}. \quad (69)$$

Equality will be reached for functions

$$u(x) = \frac{A}{\left( B + C|x|^2 \right)^{\frac{n+2\alpha}{4\alpha}}}, \quad \text{with } A, B, C \text{ arbitrary.}$$

*Proof.* We start with the inequality

$$\int_{\mathbb{R}^n} f^s g^t dx \leq \left( \int_{\mathbb{R}^n} f dx \right)^s \left( \int_{\mathbb{R}^n} g dx \right)^t, \quad s+t=1, \quad (70)$$

and we make the choices

$$s = p/2, \quad t = 1 - p/2. \quad f^s = \left( |u|^2 (a + b|x|^2) \right)^{p/2}, \quad g^t = \left( a + b|x|^2 \right)^{-p/2}.$$

This makes for (70)

$$\int_{\mathbb{R}^n} |u|^p dx \leq \left( \int_{\mathbb{R}^n} \left( |u|^2 (a + b|x|^2) \right) dx \right)^{p/2} \left( \int_{\mathbb{R}^n} \left( a + b|x|^2 \right)^{-\frac{p/2}{1-p/2}} dx \right)^{(1-p/2)},$$

or for  $p = (\rho + 2) / (\rho + 1) = 2n / (n + 2\alpha)$  and so  $\rho = 4\alpha / (n - 2\alpha)$

$$\begin{aligned} \int_{\mathbb{R}^n} |u|^p dx &= \|u\|_{\frac{\rho+2}{\rho+1}}^{\frac{\rho+2}{\rho+1}} \leq \left( \int_{\mathbb{R}^n} \left( |u|^2 (a + b|x|^2) \right) dx \right)^{\frac{\rho+2}{2(\rho+1)}} \\ &\quad \times \left( \int_{\mathbb{R}^n} \left( a + b|x|^2 \right)^{-\frac{\rho+2}{\rho}} dx \right)^{\frac{\rho}{2(\rho+1)}}. \end{aligned} \quad (71)$$

We define

$$I_0 = \left( \int_{\mathbb{R}^n} \left( a + b|x|^2 \right)^{-\frac{\rho+2}{\rho}} dx \right).$$

In a standard way this integral can be calculated as

$$I_0 = a^{-\frac{4-(n-2)\rho}{2\rho}} b^{-\frac{n}{2}} \left[ \frac{\sigma_n}{2} B \left( \frac{n}{2}, \frac{\rho+2}{\rho} - \frac{n}{2} \right) \right].$$

We make now the choice

$$b = \|u\|_2^2 / \| |x|u \|_2^2,$$

such that (71) transforms into

$$\begin{aligned} \|u\|_{\frac{\rho+2}{\rho+1}}^2 &\leq \left( \int_{\mathbb{R}^n} (|u|^2 (a + b|x|^2)) dx \right) \\ &\times \left( a^{-\frac{(4-(n-2)\rho)}{2\rho}} b^{-\frac{n}{2}} \left[ \frac{\sigma_n}{2} B \left( \frac{n}{2}, \frac{\rho+2}{\rho} - \frac{n}{2} \right) \right] \right)^{\frac{\rho}{\rho+2}}, \end{aligned}$$

or

$$\|u\|_{\frac{\rho+2}{\rho+1}}^2 \leq (a+1) a^{-(1-\alpha)} \|u\|_2^{2-n\frac{2\alpha}{n}} \| |x|u \|_2^{2(-\frac{n}{2})\frac{2\alpha}{n}} \left[ \frac{\sigma_n}{2} B \left( \frac{n}{2}, \frac{n(1-\alpha)}{2\alpha} \right) \right]^{\frac{2\alpha}{n}}.$$

We still have the free parameter  $a$ . We minimize the function  $h(a) = (a+1) a^{-(1-\alpha)}$ . By standard means this minimum will be found for  $a_0 = (1-\alpha)/\alpha$  and  $h(a_0) = \alpha^{-\alpha} (1-\alpha)^{-1+\alpha} = \chi^{-2}(\alpha)$ , by (14). Finally, we arrive at

$$\|u\|_{\frac{\rho+2}{\rho+1}} = \|u\|_{\frac{2n}{n+2\alpha}} \leq \frac{1}{\chi(\alpha)} \left[ \frac{\sigma_n}{2} B \left( \frac{n}{2}, \frac{n(1-\alpha)}{2\alpha} \right) \right]^{\frac{\alpha}{n}} \|u\|_2^{1-\alpha} \| |x|u \|_2^\alpha.$$

Equality in (70) will be reached if  $f = Cg$ ,  $C$  arbitrary, so

$$\left( |u|^2 (a + b|x|^2) \right) = C (a + b|x|^2)^{-\frac{\rho/2}{1-\rho/2}}, \quad a, b \text{ arbitrary,}$$

or

$$u(x) = C (a + b|x|^2)^{-\frac{\rho+1}{\rho}} = \frac{C}{(A + B|x|^2)^{\frac{n+2\alpha}{4\alpha}}}, \quad a, A, b, B \text{ arbitrary. } \square$$

LEMMA 2. See [4, Theorem 1.7, Case i), formula (1.30)]. For  $0 < \alpha < 1$ ,  $n \geq 2$  there holds the logconvexity of  $k_0(n, \alpha)$

$$\begin{aligned} k_0(n, \alpha) &< (k_0(n, \alpha'))^\theta (k_0(n, \alpha''))^{1-\theta}, \quad 0 < \theta < 1, \\ &\text{with } \alpha = \theta \alpha' + (1-\theta) \alpha'', \quad \alpha' \neq \alpha''. \end{aligned} \tag{72}$$

*Proof.* By the Hölder inequality

$$\|v\|_r < \|v\|_{r'}^\theta \|v\|_{r''}^{1-\theta}, \quad 0 < \theta < 1, \quad 1/r = \theta/r' + (1-\theta)/r'', \quad r' \neq r'', \tag{73}$$

which inequality is strict, since  $r' \neq r''$ . For the choice  $r = 2n/(n-2\alpha)$ , the condition for application of (73) implies  $\alpha = \theta \alpha' + (1-\theta) \alpha''$ , and so

$$\Lambda_{N,\alpha}(v) = \frac{\|\nabla v\|_2^\alpha \|v\|_2^{1-\alpha}}{\|v\|_r} > \left( \frac{\|\nabla v\|_2^{\alpha'} \|v\|_2^{1-\alpha'}}{\|v\|_{r'}} \right)^\theta \left( \frac{\|\nabla v\|_2^{\alpha''} \|v\|_2^{1-\alpha''}}{\|v\|_{r''}} \right)^{1-\theta}$$

$$= \Lambda_{N,\alpha'}^\theta(v) \Lambda_{N,\alpha''}^{1-\theta}(v), \quad (74)$$

and this implies the assertion of Lemma 2, since (see (4))

$$\frac{1}{k_0(n, \alpha)} = \lambda_{n, \alpha} = \inf_{u \in H^1(\mathbb{R}^n)} \Lambda_{n, \alpha}. \quad \square$$

#### 4.5. Upper bound 1

See the proof in [12, Proposition 1] or [15, Theorem 1]. For completeness we sketch the proof. We use the following sharp form of the Hausdorff-Young inequality due to Babenko (see [22, Section II. Babenko's inequality])

$$\|u\|_{\frac{2n}{n-2\alpha}} \leq k_b \left( \frac{2n}{n+2\alpha} \right) \|\widehat{u}\|_{\frac{2n}{n+2\alpha}}, \quad (75)$$

with  $\widehat{u} = \left( \frac{1}{2\pi} \right)^{n/2} \int_{\mathbb{R}^n} \exp(-i(x, \xi)) u(x) dx$ .

Application of Lemma 1 (69) for the Fourier Transform of  $u$ , the function  $\widehat{u}$ , gives (combined with (75))

$$\begin{aligned} \|u\|_{\frac{2n}{n-2\alpha}} &\leq k_b \left( \frac{2n}{n+2\alpha} \right) \|\widehat{u}\|_{\frac{2n}{n+2\alpha}} \\ &\leq k_b \left( \frac{2n}{n+2\alpha} \right) \frac{1}{\chi(\alpha)} \left[ \frac{\sigma_n}{2} B \left( \frac{n}{2}, \frac{n(1-\alpha)}{2\alpha} \right) \right]^{\alpha/n} \|\xi\|_2^\alpha \|\widehat{u}\|_2^{1-\alpha}. \end{aligned}$$

Due to the Parseval-Steklov relations for Fourier transforms  $\|\widehat{u}\|_2 = \|u\|_2$  and  $\|\xi\|_2 \|\widehat{u}\|_2 = \|\nabla u\|_2$ , we arrive at formula (25), the first upper bound, so

$$\overline{k_0}(n, \alpha) = k_b \left( \frac{2n}{n+2\alpha} \right) \frac{1}{\chi(\alpha)} \left[ \frac{\sigma_n}{2} B \left( \frac{n}{2}, \frac{n(1-\alpha)}{2\alpha} \right) \right]^{\alpha/n}. \quad (76)$$

#### 4.6. Upper bound 2

See the proof in [15, Theorem 1]. For completeness we sketch the proof. We apply the Beckner-Young's Inequality, see [22, Section III. Young's inequality], for  $f \in L^p(\mathbb{R}^n)$ ,  $g \in L^q(\mathbb{R}^n)$ ,

$$\|f * g\|_r \leq (A_p A_q A_r)^n \|f\|_p \|g\|_q, \quad 1 \leq p, q, r < \infty, \quad 1 + \frac{1}{r} = \frac{1}{p} + \frac{1}{q}, \quad (77)$$

$$\text{where } A_p = \left[ p^{1/p} / p^{(1/p')} \right]^{1/2}, \quad \text{with } \frac{1}{p} + \frac{1}{p'} = 1.$$

Note that  $k_b(p) = (2\pi)^{(-1/p+1/p')n/2} A_p^n$ .

We apply this inequality (77) for the solution of (11)  $\bar{u}_{n,\alpha}(r)$  written as  $\psi_0(x)$ ,  $x \in \mathbb{R}^n$ , in convolution form.  $\psi_0$  satisfies

$$\Delta\psi_0 - \psi_0 = -\psi_0^{\rho+1}. \tag{78}$$

By application of the Fourier Transform on the equation

$$\Delta\psi_{0,\delta} - \psi_{0,\delta} = \delta, \quad x \in \mathbb{R}^n,$$

with  $\delta$  the Dirac delta function, we find for the Fourier Transform  $\widehat{\psi_{0,\delta}}$

$$\widehat{\psi_{0,\delta}} = -\left(\frac{1}{2\pi}\right)^{n/2} \frac{1}{(1+\xi^2)}, \quad \text{because } \widehat{\delta} = \left(\frac{1}{2\pi}\right)^{n/2},$$

which gives for  $\psi_{0,\delta}$

$$\psi_{0,\delta} = -\left(\frac{1}{2\pi}\right)^{n/2} G(x), \quad \text{with } G(x) = \frac{K_{(n-2)/2}(|x|)}{|x|^{\frac{n-2}{2}}},$$

see [23, Chapter 8, p. 289]. And so we find for  $\psi_0$  the integral equation

$$\psi_0 = -\left(\frac{1}{2\pi}\right)^{n/2} G * (-\psi_0^{\rho+1}) = \left(\frac{1}{2\pi}\right)^{n/2} G * \psi_0^{\rho+1}. \tag{79}$$

Now, we apply (77) with  $f = G$ ,  $g = \psi_0^{\rho+1}$ ,  $r = \rho + 2$ ,  $p = (\rho + 2) / 2$ ,  $q = (\rho + 2) / (\rho + 1)$ , so  $r' = q$ , and we have

$$\begin{aligned} \|\psi_0\|_{\rho+2} &= \left(\frac{1}{2\pi}\right)^{n/2} \left\| G * \psi_0^{\rho+1} \right\|_{\rho+2} \\ &\leq \left(\frac{1}{2\pi}\right)^{n/2} \left( A_{(\rho+2)/2} A_{(\rho+2)/(\rho+1)}^2 \right)^n \|G\|_{(\rho+2)/2} \left\| \psi_0^{\rho+1} \right\|_{(\rho+2)/(\rho+1)} \\ &= k_b \left(\frac{\rho+2}{2}\right) k_b^2 \left(\frac{\rho+2}{\rho+1}\right) \|G\|_{(\rho+2)/2} \|\psi_0\|_{\rho+2}^{\rho+1}. \end{aligned} \tag{80}$$

From (80) we get

$$\|\psi_0\|_{\rho+2}^{\rho+2} \geq \left[ k_b \left(\frac{\rho+2}{2}\right) k_b^2 \left(\frac{\rho+2}{\rho+1}\right) \|G\|_{(\rho+2)/2} \right]^{-\left(\frac{\rho+2}{\rho}\right)}. \tag{81}$$

By (12) this becomes

$$\|\psi_0\|_2^2 \geq (1-\alpha) \left[ k_b \left(\frac{\rho+2}{2}\right) k_b^2 \left(\frac{\rho+2}{\rho+1}\right) \|G\|_{(\rho+2)/2} \right]^{-\left(\frac{\rho+2}{\rho}\right)},$$

and by (13) we have

$$\chi(\alpha) \left( \frac{\|\bar{u}_{n,\alpha}\|_2^2}{1-\alpha} \right)^{\alpha/n} = \frac{1}{k_0(n,\alpha)}.$$

Since  $\|\bar{u}_{n,\alpha}\|_2^2 = \|\psi_0\|_2^2$  (by definition) and  $\alpha/n = \rho/(2(\rho + 2))$

$$k_0(n, \alpha) \leq \frac{1}{\chi(\alpha)} \left[ k_b \left( \frac{\rho + 2}{2} \right) k_b^2 \left( \frac{\rho + 2}{\rho + 1} \right) \|G\|_{(\rho+2)/2} \right]^{1/2}.$$

This equals the announced upper bound 2 (27), because  $(\rho + 2)/2 = n/(n - 2\alpha)$  and  $(\rho + 2)/(\rho + 1) = 2n/(n + 2\alpha)$ :

$$\begin{aligned} k_0(n, \alpha) &\leq \frac{1}{\chi(\alpha)} \left[ k_B \left( \frac{n}{n - 2\alpha} \right) k_B^2 \left( \frac{2n}{n + 2\alpha} \right) \|G(x)\|_{n/(n-2\alpha)} \right]^{1/2} \\ &= \bar{k}_0(n, \alpha). \end{aligned} \tag{82}$$

### 4.7. Upper bound 3

We follow the same strategy as for the upper bound 2. We apply (77) with  $f = G$ ,  $g = \psi_0^{\rho+1}$ ,  $p = 2(\rho + 2)/(\rho + 4)$ ,  $q = (\rho + 2)/(\rho + 1)$ ,  $r = 2$ , so  $r' = 2$ , and we have

$$\begin{aligned} \|\psi_0\|_2 &= \left( \frac{1}{2\pi} \right)^{n/2} \|G * \psi_0^{\rho+1}\|_2 \\ &\leq \left( \frac{1}{2\pi} \right)^{n/2} (A_{2(\rho+2)/(\rho+4)} A_{(\rho+2)/(\rho+1)})^n \times \|G\|_{2(\rho+2)/(\rho+4)} \|\psi_0^{\rho+1}\|_{(\rho+2)/(\rho+1)} \\ &= k_b \left( \frac{2(\rho + 2)}{\rho + 4} \right) k_b \left( \frac{\rho + 2}{\rho + 1} \right) \|G\|_{2(\rho+2)/(\rho+4)} \|\psi_0\|_{\rho+2}^{\rho+1}. \end{aligned} \tag{83}$$

By (12) this becomes

$$(1 - \alpha) \|\psi_0\|_{\rho+2}^{\rho+2} \leq \left[ k_b \left( \frac{2(\rho + 2)}{\rho + 4} \right) k_b \left( \frac{\rho + 2}{\rho + 1} \right) \|G\|_{2(\rho+2)/(\rho+4)} \right]^2 \|\psi_0\|_{\rho+2}^{2(\rho+1)}.$$

This can be rewritten as

$$\|\psi_0\|_{\rho+2}^\rho \geq (1 - \alpha) \left[ k_b \left( \frac{2(\rho + 2)}{\rho + 4} \right) k_b \left( \frac{\rho + 2}{\rho + 1} \right) \|G\|_{2(\rho+2)/(\rho+4)} \right]^{-2}, \tag{84}$$

and by (13) we have

$$\chi(\alpha) \left( \frac{\|\bar{u}_{n,\alpha}\|_2^2}{1 - \alpha} \right)^{\alpha/n} = \chi(\alpha) \left( \|\bar{u}_{n,\alpha}\|_{\rho+2}^{\rho+2} \right)^{\alpha/n} = \frac{1}{k_0(n, \alpha)}.$$

Since  $\|\bar{u}_{n,\alpha}\|_{\rho+2}^{\rho+2} = \|\psi_0\|_{\rho+2}^{\rho+2}$  (by definition) and  $\alpha/n = \rho/(2(\rho + 2))$  there follows

$$k_0(n, \alpha) \leq \frac{1}{\chi(\alpha)} \frac{1}{\sqrt{1 - \alpha}} \left[ k_b \left( \frac{2(\rho + 2)}{\rho + 4} \right) k_b \left( \frac{\rho + 2}{\rho + 1} \right) \|G\|_{2(\rho+2)/(\rho+4)} \right].$$

This equals the announced upper bound 3 (29), because  $2(\rho + 2)/(\rho + 4) = n/(n - \alpha)$  and  $(\rho + 2)/(\rho + 1) = 2n/(n + 2\alpha)$ :



$$k_0(n, \alpha) \leq \frac{1}{\chi(\alpha)} \frac{1}{\sqrt{(1-\alpha)}} \left[ k_b \left( \frac{n}{(n-\alpha)} \right) k_b \left( \frac{2n}{(n+2\alpha)} \right) \|G\|_{n/(n-\alpha)} \right] \tag{85}$$

$$= \overline{\overline{\overline{k_0}}}(n, \alpha).$$

**4.8. Upper bound 4**

We start with the inequality

$$\|u\|_{2p} \leq A \|\nabla u\|_2^\theta \|u\|_{p+1}^{1-\theta}, \quad u \in L^{p+1}(\mathbb{R}^n), \nabla u \in L^2(\mathbb{R}^n), |u|^{2p} \in L^1(\mathbb{R}^n), \tag{86}$$

for  $n = 2, p > 1$ , and for  $n \geq 3, 1 < p \leq n/(n-2)$ ,

$$\theta = \frac{n(p-1)}{p(n+2-(n-2)p)}, \tag{87}$$

with the optimal constant

$$A = \left( \frac{y(p-1)^2}{2\pi n} \right)^{\frac{\theta}{2}} \left( \frac{2y-n}{2y} \right)^{\frac{1}{2p}} \left( \frac{\Gamma(y)}{\Gamma(y-\frac{n}{2})} \right)^{\frac{\theta}{n}}, \quad y = \frac{p+1}{p-1}, \tag{88}$$

see [24, Theorem 1].

Next, we apply the Cauch-Schwarz’s Inequality in the form

$$\|u\|_{p+1} \leq \|u\|_{2p}^\eta \|u\|_2^{1-\eta}, \quad \text{for } \eta = \frac{p}{p+1}, \tag{89}$$

and insert this inequality in the right-hand side of (86) to obtain

$$\|u\|_{2p} \leq A \|\nabla u\|_2^\theta \|u\|_{2p}^{\eta(1-\theta)} \|u\|_2^{(1-\eta)(1-\theta)},$$

or

$$\|u\|_{2p}^{1-\eta(1-\theta)} \leq A \|\nabla u\|_2^\theta \|u\|_2^{(1-\eta)(1-\theta)},$$

or

$$\|u\|_{2p} \leq A^{\frac{1}{1-\eta(1-\theta)}} \|\nabla u\|_2^{\frac{\theta}{1-\eta(1-\theta)}} \|u\|_2^{\frac{(1-\eta)(1-\theta)}{1-\eta(1-\theta)}}. \tag{90}$$

For the choice of  $p = n/(n-2\alpha)$  as in (1) we find after some calculations, using (87)

$$\theta = \frac{\alpha(n-2\alpha)}{2n-2\alpha-\alpha n}, \quad \frac{\theta}{1-\eta(1-\theta)} = \alpha, \tag{91}$$

$$\frac{(1-\eta)(1-\theta)}{1-\eta(1-\theta)} = 1-\alpha, \quad y = \frac{n-\alpha}{\alpha},$$

and

$$\frac{1}{1-\eta(1-\theta)} = \frac{2n-2\alpha-\alpha n}{n-2\alpha} \equiv \gamma. \tag{92}$$

Using the identities (91) and (92) we arrive at

$$\|u\|_{2n/(n-2\alpha)} \leq A^\gamma \|\nabla u\|_2^\alpha \|u\|_2^{1-\alpha}, \tag{93}$$

which is inequality (1) and where  $A^\gamma$  equals, using  $y = n/\alpha - 1$ ,  $p - 1 = 2\alpha/(n - 2\alpha)$

$$A^\gamma = \left( \frac{2\alpha(n - \alpha)}{\pi n(n - 2\alpha)^2} \right)^{\frac{\alpha}{2}} \left( 1 - \frac{n\alpha}{2(n - \alpha)} \right)^{(2n-2\alpha-\alpha n)/(2n)} \times \left( \frac{\Gamma(\frac{n}{\alpha} - 1)}{\Gamma(\frac{n}{\alpha} - 1 - \frac{n}{2})} \right)^{\frac{\alpha}{n}}, \tag{94}$$

so we found the announced upper bound 4 (30)

$$\overline{k_{D,1}}(n, \alpha) = A^\gamma, \quad \text{with } A = A(n, \alpha) \text{ defined in (31)}. \tag{95}$$

### 4.9. Upper bound 5

We observe that there holds trivially

$$k_0(n, \alpha) = k_0(n, \alpha)^\theta k_0(n, \alpha)^{1-\theta}. \tag{96}$$

Make now the choice  $\theta = \alpha(n - 2\alpha)/(2n - 2\alpha - \alpha n)$  see (32), then

$$k_0(n, \alpha)^\theta < \overline{k_{D,1}}(n, \alpha)^\theta = (A(n, \alpha)^\gamma)^\theta = A(n, \alpha)^\alpha, \tag{97}$$

since  $\gamma\theta = \alpha$  (see (92)) and further

$$k_0(n, \alpha)^{1-\theta} < \overline{k_0}(n, \alpha)^{1-\theta}. \tag{98}$$

Insertation of (97) and (98) into (96) gives upper bound 5:

$$k_0 < \overline{k_{D,2}}(n, \alpha) = A(n, \alpha)^\alpha \overline{k_0}(n, \alpha)^{1-\theta}, \quad n \geq 2, \quad 0 < \alpha < 1. \tag{99}$$

### 4.10. Upper bound 6

There holds trivially

$$k_0(n, \alpha) = k_0(n, \alpha)^\theta k_0(n, \alpha)^{1-\theta}. \tag{100}$$

Make now the choice  $\theta = \alpha(n - 2\alpha)/(2n - 2\alpha - \alpha n)$  see (32), then

$$k_0(n, \alpha)^\theta < \overline{k_{D,1}}(n, \alpha)^\theta = (A(n, \alpha)^\gamma)^\theta = A(n, \alpha)^\alpha, \tag{101}$$

since  $\gamma\theta = \alpha$  (see (92)) and further

$$k_0(n, \alpha)^{1-\theta} < \overline{\overline{k_0}}(n, \alpha)^{1-\theta}. \tag{102}$$

Insertation of (101) and (102) into (100) gives upper bound 6:

$$k_0 < \overline{k_{D,3}}(n, \alpha) = A(n, \alpha)^\alpha \overline{\overline{k_0}}(n, \alpha)^{1-\theta}, \quad n \geq 2, \quad 0 < \alpha < 1. \tag{103}$$

By the way, it is clear that in this way more upper bounds can be constructed.

**4.11. Upper bound 7**

This inequality is an application of [4, Theorem 1.7, (1.30),  $\theta' = 1/2$ ,  $\theta'' = 1$ , with the restriction  $n \geq 3$ ], as follows. Apply Lemma 2 with the choices  $\alpha' = 1/2$ ,  $\alpha'' = 1$  and  $\theta = 2(1 - \alpha)$ . See the results for the case  $\alpha = 1$  in the Introduction, equation (15). Application of (72) for  $n \geq 3$ :

$$\begin{aligned} k_0(n, \alpha) &< \overline{k_0} \left( n, \frac{1}{2} \right)^{2(1-\alpha)} k_0(n, 1)^{2\alpha-1} = \overline{k_0} \left( n, \frac{1}{2} \right)^{2(1-\alpha)} (C_T(n, 2))^{-2\alpha+1} \\ &= \overline{k_0} \left( n, \frac{1}{2} \right)^{2(1-\alpha)} (k_T(n))^{2\alpha-1}, \quad n \geq 3, \quad 1/2 < \alpha < 1. \end{aligned} \tag{104}$$

The last restriction comes from the requirement that  $\theta < 1$ . We made the choice to bound  $k_0 \left( n, \frac{1}{2} \right)$  by  $\overline{k_0} \left( n, \frac{1}{2} \right)$ . Equation (104) represents the announced upper bound 7

$$\overline{k_{I,1}}(n, \alpha) = \overline{k_0} \left( n, \frac{1}{2} \right)^{\alpha_1} k_T(n)^{(1-\alpha_1)}, \alpha_1 = 2(1 - \alpha), n \geq 3, 1/2 < \alpha < 1. \tag{105}$$

**4.12. Upper bound 8**

This inequality is an application of [4, Theorem 1.7, (1.30),  $\theta' = \theta_N (= \alpha_V)$ ,  $\theta'' = 1$ , with the restriction  $n \geq 3$ ], as follows. Apply Lemma 2 with the choices  $\alpha' = \alpha_V$ ,  $\alpha'' = 1$  and  $\theta = \alpha_2 = (1 - \alpha)/(1 - \alpha_V)$ . See the results for the case  $\alpha = 1$  in the Introduction, equation (15). Application of (72) for  $n \geq 3$  and for  $\alpha_V < \alpha < 1$ :

$$\begin{aligned} k_0(n, \alpha) &< \overline{k_0}(n, \alpha_V)^{\alpha_2} k_0(n, 1)^{1-\alpha_2} = \overline{k_0}(n, \alpha_V)^{\alpha_2} (C_T(n, 2))^{-(1-\alpha_2)} \\ &= \overline{k_0}(n, \alpha_V)^{\alpha_2} (k_T(n))^{(1-\alpha_2)}, \quad n \geq 3, \quad \alpha_V < \alpha < 1. \end{aligned} \tag{106}$$

We again made the choice to bound  $k_0(n, \alpha_V)$  by  $\overline{k_0}(n, \alpha_V)$ . The value  $\alpha_V$  can be chosen freely and has been chosen here as the argument value for the optimum of the expression  $\alpha C_T(n, 2\alpha)$ , see further at the proof for upper bound 10. Equation (106) represents the announced upper bound 8

$$\overline{k_{I,2}}(n, \alpha) = \overline{k_0}(n, \alpha_V)^{\alpha_2} k_T(n)^{(1-\alpha_2)}, \alpha_2 = (1 - \alpha)/(1 - \alpha_V), n \geq 3, \alpha_V < \alpha < 1. \tag{107}$$

**4.13. Upper bound 9**

This inequality is an application of [4, Theorem 1.7, (1.30),  $\theta' = \theta_N (= \alpha_V)$ ,  $\theta'' = 1$ , with the restriction  $n \geq 3$ ], as follows. Apply Lemma 2 with the choices  $\alpha' = \alpha_V$ ,  $\alpha'' = 1$  and  $\theta = \alpha_2 = (1 - \alpha)/(1 - \alpha_V)$ . See the results for the case  $\alpha = 1$  in the Introduction, equation (15). Application of (72) for  $n \geq 3$  and for  $\alpha_V < \alpha < 1$ :

$$\begin{aligned} k_0(n, \alpha) &< \overline{k_{L,V}}(n, \alpha_V)^{\alpha_2} k_0(n, 1)^{1-\alpha_2} = (\alpha_V C_T(n, 2\alpha_V))^{-\alpha_V \alpha_2} (C_T(n, 2))^{-(1-\alpha_2)} \\ &= (\alpha_V C_T(n, 2\alpha_V))^{-\alpha_V \alpha_2} (k_T(n))^{(1-\alpha_2)}, \quad n \geq 3, \quad \alpha_V < \alpha < 1. \end{aligned} \tag{108}$$

Here, we bounded  $k_0(n, \alpha_V)$  by  $\overline{k_{L,V}}(n, \alpha_V)$ , i.e. the upper bound 10 (46). The value  $\alpha_V$  can be chosen freely and has been chosen here as the argument value for the optimum of the expression  $\alpha C_T(n, 2\alpha)$ , see further at the proof for upper bound 10. Equation (108) represents the announced upper bound 9

$$\begin{aligned} \overline{k_{1,3}}(n, \alpha) &= (\alpha_V C_T(n, 2\alpha_V))^{-\alpha_V \alpha_2} k_T(n)^{(1-\alpha_2)}, \\ \alpha_2 &= (1-\alpha)/(1-\alpha_V), \quad n \geq 3, \quad \alpha_V < \alpha < 1. \end{aligned} \tag{109}$$

**4.14. Upper bound 10**

Firstly, we prove

$$k_0(n, \alpha) < (\alpha C_T(n, 2\alpha))^{-\alpha}, \quad n \geq 2, \quad 1/2 < \alpha < 1. \tag{110}$$

This result has been given in [4, Theorem 1.7, (1.31)] and was inspired by [6, (1.5)], by making the transformation  $w = u^{1/\alpha}$  for  $v > 0$  in (15) as follows

$$\begin{aligned} C_T(n, s) &\leq \frac{\|\nabla w\|_s}{\|w\|_t} = \frac{\|\nabla u^{1/\alpha}\|_s}{\|u^{1/\alpha}\|_t} = \frac{1/\alpha \|u^{(1-\alpha)/\alpha} \nabla u\|_s}{\|u^{1/\alpha}\|_t} \quad [t = sn/(n-s)] \\ &= \frac{1}{\alpha} \frac{\left(\int (\nabla u)^s u^{s(1-\alpha)/\alpha} dx\right)^{1/s}}{\left(\int u^{t/\alpha} dx\right)^{1/t}} \quad \begin{array}{l} \text{[apply Hölder inequality,} \\ 1/P + 1/Q = 1] \end{array} \\ &\leq \frac{1}{\alpha} \frac{\left(\int (\nabla u)^{sP} dx\right)^{1/(sP)} \left(\int u^{2s(1-\alpha)/\alpha} dx\right)^{1/(sQ)}}{\left(\int u^{t/\alpha} dx\right)^{1/t}} \quad \begin{array}{l} \text{[take } P = 2/s, \\ Q = 2/(2-s)] \end{array} \\ &= \frac{1}{\alpha} \frac{\left(\int (\nabla u)^2 dx\right)^{1/2} \left(\int u^{2s(1-\alpha)/\alpha} dx\right)^{(2-s)/(2s)}}{\left(\int u^{t/\alpha} dx\right)^{1/t}} \quad \begin{array}{l} \text{[take } s = 2\alpha, \text{ and} \\ r = t/\alpha = 2n/(n-2\alpha)] \end{array} \\ &= \frac{1}{\alpha} \frac{\|\nabla u\|_2 \|u\|_2^{(1-\alpha)/\alpha}}{\|u\|_r^{1/\alpha}} = \frac{1}{\alpha} (\Lambda_{n,\alpha}(u))^{1/\alpha}, \end{aligned} \tag{111}$$

for the choice  $s = 2\alpha$ . We have to restrict  $\alpha$  to the interval  $1/2 \leq \alpha \leq 1$  to give  $C_T(n, 2\alpha)$  a meaning. Again, the inequality is strict since  $w = \overline{w}_{n,\alpha}^\alpha$  does not equal a function  $w_{n,s}$  (see (22)), with  $s = 2\alpha$ . So (111) implies

$$\lambda_{n,\alpha} = \inf_{u \in H^1(\mathbb{R}^n)} \Lambda_{n,\alpha}(u) > (\alpha C_T(n, 2\alpha))^\alpha,$$

and this equivalent with

$$k_0(n, \alpha) = 1/\lambda_{n,\alpha} < (\alpha C_T(n, 2\alpha))^{-\alpha}, \quad n \geq 2, \quad 1/2 < \alpha < 1.$$

Application of Lemma 2 with  $\alpha'' = 0$ ,  $\theta = \alpha/\alpha'$ , and  $k_0(n, 0) = 1$  gives

$$k_0(n, \alpha) < \left( (\alpha' C_T(n, 2\alpha'))^{-\alpha'} \right)^{\alpha/\alpha'} = (\alpha' C_T(n, 2\alpha'))^{-\alpha}.$$

Since  $\alpha'$  can still be chosen freely, we can improve this inequality by maximizing the  $(\alpha' C_T(n, 2\alpha'))$ . In a standard way we find that there is a unique value  $\alpha_V \in (1/2, 1)$  which optimizes this expression, see [4, Proof Theorem 1.7, (1.32)] for details. Finally we find the announced upper bound 10

$$k_0 < \overline{k_{L,V}}(n, \alpha) = [\alpha_V C_T(n, 2\alpha_V)]^{-\alpha}, \quad n \geq 2, 0 < \alpha \leq \alpha_V, \tag{112}$$

$$k_0 < \overline{k_{L,V}}(n, \alpha) = 1/k_{L,V}(n, \alpha) = [\alpha C_T(n, 2\alpha)]^{-\alpha}, \quad n \geq 2, \alpha_V \leq \alpha < 1, \tag{113}$$

where the value for  $\alpha_V$  follows from

$$\alpha_V = \alpha_V(n) = \frac{n}{2p_V}, \text{ where } p_V \text{ is the solution of} \tag{114}$$

$$\ln\left(\frac{n-p}{p-1}\right) + \frac{n-p}{p(p-1)} + \psi(p) - \psi(n+1-p) = 0, \tag{115}$$

$$\psi(x) = \frac{d}{dx} \frac{\Gamma(x)}{\Gamma(x)}, \quad x > 0, \quad 1 < p < n, \quad n \geq 2.$$

In both expressions (112) and (113) the second argument in  $C_T$  is larger than 1, as required. The value  $\alpha_V$  has also been used in the upper bounds 8 and 9.

**4.15. Upper bound 11**

This inequality is a combination of the Hölder inequality (73)

$$\|u\|_r < \|u\|_{r'}^\theta \|u\|_{r''}^{1-\theta}, \quad 0 < \theta < 1, \quad 1/r = \theta/r' + (1-\theta)/r'', \quad r' \neq r'', \tag{116}$$

and the Sobolev embedding (15)

$$\|u\|_t \leq \frac{1}{C_T(n, 2)} \|\nabla u\|_2, \quad t = 2n/(n-2), \quad n \geq 3. \tag{117}$$

For the choice  $r = 2n/(n-2\alpha)$ ,  $\theta = \alpha$ ,  $r'' = 2$  in (116), we find  $r' = 2n/(n-2)$ , which is just the value applicable for the Sobolev embedding (117). These two estimates combined gives

$$\|u\|_{2n/(n-2\alpha)} < \left(\frac{1}{C_T(n, 2)}\right)^\alpha \|\nabla u\|_2^\alpha \|u\|_2^{1-\alpha} = k_T(n)^\alpha \|\nabla u\|_2^\alpha \|u\|_2^{1-\alpha}, \quad n \geq 3. \tag{118}$$

So, we found the announced upper bound 11

$$k_0 < \overline{k_B}(n, \alpha) = k_T(n)^\alpha, \quad n \geq 3, \quad 0 < \alpha < 1. \tag{119}$$

**5. Numerical evaluations lower and upper bounds**

In order to assess the quality of the estimates we have calculated the numbers  $\lambda_{n,\alpha}$  for  $n = 2, 3, 4, 5, 10$  and  $\alpha = 0.05 + (i-1)0.005$ ,  $i = 1, 2, 3, \dots, 176$  up till  $\theta = 0.925$ . The method is the same as used in the paper [4]. This method to find  $\lambda_{n,\alpha}$  consists of a

shooting technique to find that value  $\bar{u}(0) = u_0$  such that  $\bar{u}(r)$  is a positive solution of (11) with  $\lim_{r \rightarrow \infty} \bar{u}(r) = 0$ . Therefore, we transformed the interval  $r \in (0, \infty)$  into  $s = r/(1+r) \in (0, 1)$ . The transformed differential equation becomes, with  $w(s) = u(r)$ ,  $0 < s < 1$ ,

$$(1-s)^4 \frac{d^2}{ds^2} w + \left\{ \left( \frac{(n-1)}{s} - 2 \right) (1-s)^3 \right\} \frac{d}{ds} w - w |w|^{(n+2\alpha)/(n-2\alpha)-1} - w = 0,$$

$$w(0) = v_0, \quad \frac{d}{ds} w(0) = 0. \tag{120}$$

The aim now is to find a value  $v_0$  such that for  $w(0) = v_0$ ,  $\frac{d}{ds} w(0) = 0$ , we find  $w(1) = 0$ . We solved the transformed differential equation (120) by means of a numerical integration method (Runge-Kutta of the fourth order) with a self-adapting step-size routine such that a prescribed maximal relative error ( $\epsilon_{rel}$ ) in each component ( $w(s), \frac{d}{ds} w(s)$ ) has been satisfied. We made the choice  $\epsilon_{rel} = 10^{-15}$ . For every value of  $v_0$  the numerical integrator will find some point  $s = s(v_0) \in (0, 1)$  where either  $w(s) < 0$ , or  $\frac{d}{ds} w(s) > 0$ . At that point  $s$  the integration will be stopped. This integrator is coupled to a numerical zero-finding routine (see ([25])), which can also be applied for finding a discontinuity. The function  $f$  for which such a discontinuity has to be found is specified by if  $w(s(v_0)) < 0$ ,  $f(v_0) = -(1-s(v_0))$  else (that means thus  $\frac{d}{ds} w(s(v_0)) > 0$ )  $f(v_0) = (1-s(v_0))$ . The sought value  $v_0$  has been found if this numerical routine has come up with two values  $v_0$  and  $v_0^1$  such that  $|v_0 - v_0^1| < r_p |v_0| + a_p$ , (with  $r_p = a_p = 10^{-15}$  relative and absolute precisions, respectively) and  $|f(v_0)| \leq |f(v_0^1)|$ , while  $sign(f(v_0)) = -sign(f(v_0^1))$ . During the integration processes the norms in (12) will be calculated. As a check upon this procedure the following expressions

$$\|\bar{u}_{n,\alpha}\|_2^2 / (1-\alpha), \quad \|\nabla \bar{u}_{n,\alpha}\|_2^2 / \alpha, \quad \|\bar{u}_{n,\alpha}\|_{2n/(n-2\alpha)}^{2n/(n-2\alpha)}, \tag{121}$$

are compared. They should be all equal, see (12). The eigenvalue  $\lambda_{n,\alpha}$  is found then by (13).

**5.1. Some numerical results for values for  $\alpha = 1/3, 2/3$  and  $n = 2$**

Here, we give for  $n = 2$  and for particular values of  $\alpha$  ( $\alpha = 1/3$  and  $2/3$ ) the upper and lower bounds which are applicable. Compare these with [10,  $\alpha = 1/3$ ] and [6,  $\alpha = 2/3$ ].

$\alpha$	$k_0$	$\underline{k}_0$	$\underline{\underline{k}}_0$
$n = 2$			
1/3	7.2493833e-001	7.2431703e-001	7.2184608e-001
2/3	6.0129905e-001	5.9737503e-001	5.6854280e-001

Table 1: Functional,  $n = 2$ , Lower bounds 1 - 2.

$\alpha$	$k_0$	$\overline{k_0}$	$\overline{\overline{k_0}}$	$\overline{\overline{\overline{k_0}}}$
$n = 2$				
1/3	7.2493833e-001	7.2978972e-001	7.3987840e-001	7.8567080e-001
2/3	6.0129905e-001	6.4335375e-001	6.1742806e-001	7.2152108e-001

Table 2: Functional,  $n = 2$ , Upper bounds 1 - 3.

$\alpha$	$k_{D,1}$	$k_{D,2}$	$k_{D,3}$	$k_{L,V}$
$n = 2$				
1/3	7.3907188e-001	7.3132861e-001	7.3974392e-001	7.7547470e-001
2/3	6.8278406e-001	6.5623746e-001	6.3848696e-001	6.1088706e-001

Table 3:  $n = 2$ , Upper bounds 4 - 6 and 10.

**5.2. Numerical results for  $\alpha = 0.05, \dots, 0.925$  ( $\Delta = 0.005$ ) and  $n = 2, 3, 4, 5, 10$**

In the Supplementary Material to this paper we present tables which give the results of the numerical calculations of the functional  $k_0(n, \alpha)$  and the lower and upper bounds, based on the technique described above (see also [4]).

Values "0.0000000e+000" has to be interpreted as "Not Applicable". The lower and upper bounds have been calculated using the software package Matlab™.

**5.3. Results for the zeros  $p_V$  and  $\alpha_V = n/(2p_V)$**

The zeros  $p_V$  as defined in (42) are given below in the Table 4;  $\alpha_V(n) = n/(2p_V)$ . The asymptotic expressions are

$$p_V(n) = 2n/3 + 5/18 + O(1/n), \quad n \rightarrow \infty, \tag{122}$$

$$\alpha_V(n) = 3/4 - 5/(16n) + O(1/n^2), \quad n \rightarrow \infty, \tag{123}$$

$n$	$p_V$	$p_{V,asympt}$	$p_V - p_{V,asympt}$
		$= 2n/3 + 5/18$	
2	1.6474176e+000	1.6111111e+000	3.6306497e-002
3	2.3044430e+000	2.2777778e+000	2.6665194e-002
4	2.9654018e+000	2.9444444e+000	2.0957401e-002
5	3.6283253e+000	3.6111111e+000	1.7214200e-002
6	4.2923606e+000	4.2777778e+000	1.4582787e-002
7	4.9570820e+000	4.9444444e+000	1.2637555e-002
8	5.6222549e+000	5.6111111e+000	1.1143822e-002
9	6.2877400e+000	6.2777778e+000	9.9621751e-003
10	6.9534493e+000	6.9444444e+000	9.0048448e-003

Table 4: The zeros  $p_V$  for  $n = 2, \dots, 10$  and their asymptotic approximations.

$n$	$\alpha_V$	$\alpha_{V,asympt}$	$\alpha_V - \alpha_{V,asympt}$
		$= 3/4 - 5/(16n)$	
2	6.0701063e-001	5.9375000e-001	1.3260630e-002
3	6.5091652e-001	6.4583333e-001	5.0831867e-003
4	6.7444485e-001	6.7187500e-001	2.5698490e-003
5	6.8902311e-001	6.8750000e-001	1.5231128e-003
6	6.9891612e-001	6.9791667e-001	9.9945530e-004
7	7.0606054e-001	7.0535714e-001	7.0339854e-004
8	7.1145831e-001	7.1093750e-001	5.2081118e-004
9	7.1567845e-001	7.1527778e-001	4.0067485e-004
10	7.1906759e-001	7.1875000e-001	3.1758674e-004

Table 5: The zeros  $\alpha_V = n/(2p_V)$  for  $n = 2, \dots, 10$  and their asymptotic approximations.

### 6. Discussion

With respect to the lower bounds it is clear based on the numerical results in the Supplementary Material to this paper (Tables 4-8 and Fig. 3 in "Comparison Functional with Lower bounds for Functional" therein) that the lower bound for  $n = 2$ ,  $\underline{k}_0(\alpha)$ , is superior to the lower bound  $\underline{k}_0(2, \alpha)$ .

With respect to the upper bounds the situation is more complicated. For the range of  $n$  ( $n = 2, 3, 4, 5$  and  $n = 10$ ) and  $\alpha$  ( $0.05 \leq \alpha \leq 0.925$  with steps  $\Delta\alpha = 0.005$ ) we have examined there are just four upper bounds which are superior, see the Table 6 and the Figures 1, 2, 3, 4 and 5.




Figure 1: Best bounds for  $n = 2$ .




Figure 2: Best bounds for  $n = 3$ .






Figure 3: Best bounds for  $n = 4$ .




Figure 4: Best bounds for  $n = 5$ .




Figure 5: Best bounds for  $n = 10$ .

$n$	Range $\alpha$	Upper bound #	Expression Upper bound
2	(0.050, 0.495)	1	$\overline{k_0}(2, \alpha)$
2	0.500	1 = 2	$\overline{k_0}(2, 1/2) = \overline{k_0}(2, 1/2)$
2	[0.505, 0.615)	2	$\overline{k_0}(2, \alpha)$
2	(0.620, 0.745)	10	$\overline{k_{L,V}}(2, \alpha)$
2	(0.750, 0.925)	2	$\overline{k_0}(2, \alpha)$
3	(0.050, 0.590)	1	$\overline{k_0}(3, \alpha)$
3	(0.595, 0.925)	10	$\overline{k_{L,V}}(3, \alpha)$
4	(0.050, 0.590)	1	$\overline{k_0}(4, \alpha)$
4	(0.595, 0.605)	4	$\overline{k_{D,1}}(4, \alpha)$
4	(0.610, 0.925)	10	$\overline{k_{L,V}}(4, \alpha)$
5	(0.050, 0.565)	1	$\overline{k_0}(5, \alpha)$
5	(0.570, 0.630)	4	$\overline{k_{D,1}}(5, \alpha)$
5	(0.635, 0.925)	10	$\overline{k_{L,V}}(5, \alpha)$
10	(0.050, 0.535)	1	$\overline{k_0}(10, \alpha)$
10	(0.540, 0.675)	4	$\overline{k_{D,1}}(10, \alpha)$
10	(0.680, 0.925)	10	$\overline{k_{L,V}}(10, \alpha)$

Table 6: Optimal upper bounds for  $n = 2, 3, 4, 5, 10$ .

We remark that  $\overline{k_0}(2, 1/2) = \overline{\overline{k_0}}(2, 1/2) = 2^1 3^{-3/4} \pi^{-1/4}$ , and  $\overline{k_0}(3, 3/4) = \overline{\overline{k_0}}(3, 3/4) = 2^{7/4} 3^{-3/2} \pi^{-1/4}$  see [15, equation (12) and (17)].

As can be seen from the figures in the Supplementary Material to this paper, for larger values of  $n$  almost all bounds come close to the actual value for  $k_0(n, \alpha)$ ; see the Figures 7, 12, 28, 32, 37, 42, 46 and 51 therein, for  $n = 10$ .

## REFERENCES

- [1] B. v. SZ. NAGY, *Über Integralgleichungen zwischen einer Funktion und ihrer Ableitung*, Acta Sci. Math. (Szeged), 10 (1941), 64–74.
- [2] E. H. LIEB AND W. E. THIRRING, *Inequalities for the moments of the eigenvalues of the Schrödinger Hamiltonian and their relation to Sobolev inequalities*, Studies in Mathematical Physics, Essays in Honor of Valentine Bargmann (E. H. Lieb and B. Simon and A. S. Wightman, Eds.), 1976, Princeton University Press, 269–303.
- [3] E. J. M. VELING, *Optimal lower bounds for the spectrum of a second order linear differential equation with a  $p$ -integrable coefficient*, Proc. Roy. Soc. Edinburgh Sect. A, 92 (1982), 95–101.
- [4] E. J. M. VELING, *Lower bounds for the infimum of the spectrum of the Schrödinger operator in  $\mathbf{R}^N$  and the Sobolev inequalities*, Journal of Inequalities in Pure and Applied Mathematics, 3, no. 4 (2002), Article 63, Corrigendum in JIPAM, 4, no. 5, (2003), Article 109.
- [5] W. A. STRAUSS, *Existence of solitary waves in higher dimensions*, Comm. Math. Phys., 55 (1977), 149–162.
- [6] H. A. LEVINE, *An estimate for the best constant in a Sobolev inequality involving three integral norms*, Ann. Mat. Pura Appl. (4), 124 (1980), 181–197.
- [7] W. THIRRING, *A course in mathematical physics III. Quantum mechanics of atoms and molecules*, Springer, New York, 1981.
- [8] G. ROSEN, *Necessary conditions on potential functions for nonrelativistic bound states*, Phys. Rev. Lett., 49 (1982), 1885–1887.
- [9] M. I. WEINSTEIN, *Nonlinear Schrödinger equations and sharp interpolation estimates*, Comm. Math. Phys., 87 (1983), 567–576.
- [10] SH. M. NASIBOV, *On optimal constants in some Sobolev inequalities and their application to a nonlinear Schrödinger equation*, Soviet. Math. Dokl., 40, no. 1 (1990), 110–115, Translation of Dokl. Akad. Nauk SSSR 307(3) (1989), 538–542.
- [11] JACK GUNSON, *Inequalities in mathematical physics*, Inequalities, fifty years on from Hardy, Littlewood and Pólya (W. Norrie Everitt, Eds.), Lecture notes in pure and applied mathematics series, no. 129 (1991), 53–79, Proceedings of the international conference, July 13–17, 1987, University of Birmingham, U.K., London Mathematical Society, Marcel Dekker, Inc., New York, Basel, Hong Kong.
- [12] SH. M. NASIBOV, *An upper bound for the sharp constant in the Sobolev inequality and its application to an interior Dirichlet problem for the homogeneous stationary Schrödinger equation  $\Delta u + q(x)u = 0$* , Dokl. Math., 72, no. 1 (2005), 547–550, Translation of Dokl. Akad. Nauk 403, no. 4, (2005), 443–447.
- [13] SH. M. NASIBOV, *On an integral inequality and its application to the proof of the entropy inequality*, Math. Notes, 84, no. 2 (2008), 218–223, Translation of Mat. Zametki 84, no. 2, (2008), 231–237.
- [14] SH. M. NASIBOV, *A sharp constant in a Sobolev-Nirenberg inequality and its application to the nonlinear Schrödinger equation*, Izv. Math. 73, no. 3 (2009), 555–577, Translation of Izv. Ross. Akad. Nauk Ser. Mat. 73, no. 3, (2009), 127–150.
- [15] SH. M. NASIBOV AND M. A. NAMAZOV, *On a Sobolev inequality [in Russian]*, Proceedings of the Institute of Applied Mathematics [Baku, Azerbaijan] 2, no. 2 (2013), 187–195.
- [16] MAN KAM KWONG, *Uniqueness of positive solutions of  $\Delta u - u + u^p = 0$  in  $\mathbf{R}^N$* , Arch. Ration. Mech. Anal. 105, no. 3 (1989), 243–266.
- [17] T. AUBIN, *Problèmes isopérimétriques et espaces de Sobolev*, J. Differential Geom. 11 (1976), 573–598.
- [18] G. TALENTI, *Best constant in Sobolev inequality*, Ann. Mat. Pura Appl. (4) 110 (1976), 353–372.
- [19] E. H. LIEB, *Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities*, Ann. of Math. (2) 118 (1983), 349–374.

- [20] FRANK W. J. OLVER, DANIEL W. LOZIER, RONALD F. BOISVERT AND CHARLES W. CLARK, *NIST Handbook of mathematical functions*, Cambridge University Press & National Institute of Standards and Technology (2010), New York, N.Y., USA, <http://dlmf.nist.gov>.
- [21] V. S. VLADIMOROV, *Equations of mathematical physics*, Marcel Dekker, New York (1971).
- [22] WILLIAM BECKNER, *Inequalities in Fourier analysis*, Annals of Mathematics, Second Series 102, no. 1 (1975), 159–182.
- [23] S. M. NIKOL'SKII, *Approximation of functions of several variables and imbedding theorems*, Springer-Verlag, Berlin, Heidelberg, New York, 1975, Translated from the Russian by J. M. Dankin.
- [24] MANUEL DEL PINO AND JEAN DOLBEAULT, *Best constants for Gagliardo–Nirenberg inequalities and applications to nonlinear diffusions*, J. Math. Pures Appl. 81 (2002), 847–875.
- [25] J. C. P. BUS AND T. J. DEKKER, *Two efficient algorithms with guaranteed convergence for finding a zero of a function*, ACM Trans. Math. Software 1, no. 4 (1975), 330–345.

(Received March 25, 2017)

*Sh. M. Nasibov*  
*Baku State University*  
*e-mail: nasibov\_sharif@hotmail.com*

*E. J. M. Veling*  
*Delft University of Technology*  
*Faculty of Civil Engineering and Geosciences*  
*Water Resources Section, P.O. Box 5048, NL-2600 GA*  
*Delft, The Netherlands*  
*e-mail: E.J.M.Veling@TUDelft.nl & ed.veling@gmail.com*