

## **Fibroepithelioma of Pinkus (FeP) located in the left lower quadrant of the abdomen - Case Report and review of the literature**

Mihai, Mara Madalina; Voicu, Cristiana; Lupu, Mihai; Koleva, Nely; Patterson, James W.; Lotti, Torello; Lotti, Jacopo; França, Katlein; Batashki, Atanas; Bakardzhiev, Ilko

**DOI**

[10.3889/oamjms.2017.079](https://doi.org/10.3889/oamjms.2017.079)

**Publication date**

2017

**Document Version**

Final published version

**Published in**

Open Access Macedonian Journal of Medical Sciences

**Citation (APA)**

Mihai, M. M., Voicu, C., Lupu, M., Koleva, N., Patterson, J. W., Lotti, T., Lotti, J., França, K., Batashki, A., Bakardzhiev, I., Wollina, U., & Tchernev, G. (2017). Fibroepithelioma of Pinkus (FeP) located in the left lower quadrant of the abdomen - Case Report and review of the literature. *Open Access Macedonian Journal of Medical Sciences*, 5(4 Special Issue GlobalDermatology), 439-444. <https://doi.org/10.3889/oamjms.2017.079>

**Important note**

To cite this publication, please use the final published version (if applicable). Please check the document version above.

**Copyright**

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

**Takedown policy**

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

# Fibroepithelioma of Pinkus (FeP) Located in the Left Lower Quadrant of the Abdomen - Case Report and Review of the Literature

Mara Madalina Mihai<sup>1</sup>, Cristiana Voicu<sup>2</sup>, Mihai Lupu<sup>3</sup>, Nely Koleva<sup>4</sup>, James W. Patterson<sup>5</sup>, Torello Lotti<sup>6</sup>, Jacopo Lotti<sup>7</sup>, Katlein Franca<sup>8</sup>, Atanas Batashki<sup>9</sup>, Ilko Bakardzhiev<sup>10</sup>, Uwe Wollina<sup>11</sup>, Georgi Tchernev<sup>12\*</sup>

<sup>1</sup>Oncologic Dermatology and Allergology Department, "Carol Davila" University of Medicine and Pharmacy; 17 Marasti Blvd, Bucharest, Romania; <sup>2</sup>Dermatology Department, Poliso Clinic, 26Z Timisoara Blvd, Bucharest, Romania; <sup>3</sup>Dermatology Department, Medas Medical Centre, 41 Nicolae Grigorescu Blvd, Bucharest, Romania; <sup>4</sup>Medical Institute of Ministry of Interior (MVR), Department of Dermatology and Dermatologic Surgery, General Skobelev 79, 1606 Sofia, Bulgaria; <sup>5</sup>University of Virginia Health System, 1215 Lee Street, Charlottesville, VA 22908, USA; <sup>6</sup>University of Rome "G. Marconi", Rome, Italy; <sup>7</sup>Department of Biotechnology, Delft University of Technology, 2628 BC, Delft, The Netherlands; <sup>8</sup>Department of Nuclear, Subnuclear and Radiation Physics, University of Rome "G. Marconi", Rome, Italy; <sup>9</sup>Institute for Bioethics & Health Policy; <sup>10</sup>Department of Dermatology & Cutaneous Surgery; <sup>11</sup>Department of Psychiatry & Behavioral Sciences, University of Miami Miller School of Medicine - Miami, FL, USA; <sup>12</sup>Specialist in Abdominal and Thoracic Surgery, Department of special Surgery, Medical University of Plovdiv, bul. "Peshtersko shose" Nr 66, 4000, Plovdiv, Bulgaria; <sup>10</sup>Medical College, Medical University of Varna, Varna 9000, Bulgaria; <sup>11</sup>Department of Dermatology and Allergology, Academic Friedrichstrasse 41, 01067, Dresden, Germany; <sup>12</sup>Medical Institute of Ministry of Interior (MVR), Department of Dermatology and Dermatologic Surgery, General Skobelev 79, 1606 Sofia, Bulgaria; Onkoderma - Policlinic for Dermatology, Venereology and Dermatologic Surgery, 26 General Skobelev blvd., Sofia, Bulgaria

## Abstract

**Citation:** Mihai MM, Voicu C, Lupu L, Koleva N, Patterson JW, Lotti T, Lotti J, Franca K, Batashki A, Bakardzhiev I, Wollina U, Tchernev G. Fibroepithelioma of Pinkus (FeP) Located in the Left Lower Quadrant of the Abdomen - Case Report and Review of the Literature. Open Access Maced J Med Sci. 2017 Jul 25; 5(4):439-444. https://doi.org/10.3889/oamjms.2017.079

**Keywords:** fibroepithelioma of Pinkus; basal cell carcinoma; trichoblastoma; trichoepithelioma.

**\*Correspondence:** Professor Dr Georgi Tchernev PhD, Chief of 1) Department of Dermatology, Venereology and Dermatologic Surgery, Medical Institute of Ministry of Interior (MVR), General Skobelev 79, 1606 Sofia; 2) Onkoderma- Policlinic for Dermatology and Dermatologic Surgery, General Skobelev 26, Sofia, Bulgaria. GSM: 00359885588424. E-mail: georgi\_tchernev@yahoo.de

**Received:** 01-Apr-2017; **Revised:** 22-Apr-2017; **Accepted:** 23-Apr-2017; **Online first:** 19-Jul-2017

**Copyright:** © Mara Madalina Mihai, Cristiana Voicu, Mihai Lupu, Nely Koleva, James W. Patterson, Torello Lotti, Jacopo Lotti, Katlein Franca, Atanas Batashki, Ilko Bakardzhiev, Uwe Wollina, Georgi Tchernev. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

**Funding:** This research did not receive any financial support.

**Competing Interests:** The authors have declared that no competing interests exist.

**BACKGROUND:** Fibroepithelioma of Pinkus (FeP) is an uncommon and controversial skin lesion, sharing features of both basal cell carcinoma (BCC) and trichoepithelioma. In this article, we present a case of FeP and synthesise current concepts on the etiopathogenesis, diagnosis and treatment of this uncommon tumour.

**CASE REPORT:** We report the case of an 88-year-old male patient presenting to the dermatology clinic for a sharply demarcated, pink, exophytic cutaneous tumour situated in the left inguinal region. The histopathological examination performed after complete surgical excision of the lesion revealed a diagnosis of FeP. A systematic review of the literature was conducted. The terms 'fibroepithelioma' and 'Pinkus' have been searched in bibliographical databases, including PubMed and Google Scholar, without time limitation up to February 15th, 2017. Seventy-nine articles that fulfilled all the required conditions were identified. Relevant citations and additional articles identified from references have been assessed. The systematic review included a total number of 452 cases of FeP.

**CONCLUSION:** Even though FeP is considered a relatively rare tumour, its true incidence rate might be higher than previously believed. The clinical aspects of the lesion described in this paper and its location in the left lower quadrant of the abdomen are classic features of FeP. Histopathologic examination revealed features of both BCC and trichoepithelioma. Further epidemiological studies are required to clarify whether patients with FEP should be screened for the occurrence of other malignancies.

## Introduction

Fibroepithelioma of Pinkus (FeP) is an uncommon and controversial skin lesion, sharing features of both basal cell carcinoma (BCC) and trichoepithelioma. While some authors differentiate trichoblastoma from trichoepithelioma, the World Health Organization classifies the terms synonymously [1]. In the present paper, we will use the term trichoepithelioma.

FeP was described for the first time by Hermann Pinkus in 1953, who regarded this skin lesion as a variant of BCC [2], a theory that continued for many years [3, 4]. Recent research has highlighted the fact that BCC and trichoepithelioma have the same origin; that is, the epithelial stem cells of the hair follicle [5, 6]. Such a concept has led to the proposal that FeP might be a trichoblastic tumour intermediate between trichoepithelioma and BCC [7]. The clinical presentation of FeP is that of a flesh-coloured, well-demarcated, sessile, dome-shaped papule or

pedunculated tumour frequently often located in the lumbosacral area. The clinical differential diagnosis includes benign tumours such as intradermal nevus, fibroma, acrochordon, and seborrheic keratosis [8], as well as a variety of cutaneous neoplasms, including basal cell carcinoma and amelanotic melanoma. The pathogenesis of FeP is still a matter of debate. As is the case with conventional BCC, theories include mutations in the tumour suppressor gene P53 and the PATCHED gene, inducing inhibitory signals in the Hedgehog pathway [7]. There are several reports on the dermatoscopic, reflectance confocal microscopic, and histopathologic features of FeP. The treatment of choice is complete surgical excision. Other surgical techniques, such as cryosurgery, electrodesiccation and Mohs micrographic surgery have also been performed successfully [7].

In this article, we present a case of FeP and summarise current concepts on the etiopathogenesis, diagnosis and treatment of this uncommon tumour.

## Case Report

An 88-year-old male presented to the dermatology clinic for evaluation, diagnosis and treatment of a skin tumour that had been slowly growing for the preceding five years in the left inguinal region. There were no complaints regarding pain or itching of the skin lesion. The patient's medical history revealed several cardiovascular risk factors: an elevated blood pressure and non-insulin dependent diabetes mellitus type 2 but was otherwise unremarkable. Clinical examination revealed a sharply demarcated, pink, exophytic cutaneous tumour with a lobulated, cerebriform surface, measuring approximately 5 x 3.6 cm situated in the left lower quadrant of the abdomen (Fig. 1a). Paraclinical Diagnostic tests were unremarkable. Abdominal ultrasound, chest x-ray and ultrasound of the axillary and inguinal lymph node groups showed no evidence of disease progression or other abnormal findings. In the light of the history of tumour growth and clinical differential diagnosis, the lesion was surgically excised with wide margins under local anaesthesia, thus creating an elliptical defect (Fig. 1b), followed by the primary closure (Fig. 1c) and by the application of antiseptic dressings. Postoperatively, the patient was well and was discharged with a set of follow-up instructions. Histopathologic examination of the excised cutaneous tumour showed a polypoid tumoral proliferation with superficial ulcerated areas (Fig. 2b). Focal solid cribriform areas (Fig. 2a), tumour islands and long anastomosing strands and columns of basaloid cells with scant cytoplasm, hyperchromatic nuclei and moderate cytologic atypia could be seen projecting downwards from the epidermis into the

papillary dermis. Relatively frequent mitoses could be seen along with abundant fibro hyaline tumoral stroma. Based on the clinical and microscopic features, a final diagnosis of fibroepithelioma of Pinkus was made. The pathology report confirmed its complete excision, with tumour-free surgical resection margins.



Figure 1: 1a) Cutaneous tumour in the left inguinal region; 1b) Elliptic surgical defect after haemostasis, ready for reconstruction; 1c) Primary closure of the defect with interrupted non-absorbable sutures

## Discussion

FeP was first described as a distinct clinicopathologic entity by Hermann Pinkus in 1953 [2]. In a retrospective study that included over 900 epitheliomas, he described four cases with unique clinical and microscopic features, emphasising their importance in enhancing the understanding of epitheliomas in general [2].

While FeP is considered a relatively rare tumour, it is probably underreported. It can be easily confused with other benign skin tumours that may not be treated or biopsied [9]. After a thorough literature review, we identified 79 articles reporting a total number of 452 FeP cases. FeP usually appears in the fourth to fifth decades of life [10], although two cases of FeP in children have been reported [11, 12]. In the largest clinical study, Bowen et al. (2005) observed a slightly higher prevalence rate in women: 54% of 114 patients [13].




Figure 2: Several histopathologic images from the tumour, showing branching cords of basaloid epithelium, some narrow (2a - lower left, 2c and 2d - centre, and figure 2e - upper left) and some more broad (2b and 2f), within a cellular fibrotic stroma

After almost 65 years since the first description of FeP, its classification and pathogenesis are still matters of debate. The following alternatives were taken into consideration: a premalignant tumour, a fenestrated trichoepithelioma subtype, or an indolent BCC subtype. The first reports described FeP as a BCC subtype. Due to its indolent, long-term course, and histological particularities it was further classified as a premalignant tumour - a precursor of BCC. In 2005, Bowen et al. argued that FeP is a trichoepithelioma rather than a BCC subtype, based upon their histopathological and immunohistochemical studies of 75 FePs [13]. The same year Ackerman et al. provided criteria for the histopathologic differentiation of FeP from trichoblastoma (trichoepithelioma) and classified the fibroepithelial tumour of Pinkus as a trichoblastic (Basal-cell) carcinoma [4], reflecting the controversial nature of the problem.

It is well known that BCC can spread directly into the dermis, by perifollicular or perineural extension. However, BCC might also spread via eccrine ducts. In 1994, Stern et al. hypothesised that eccrine ducts might serve as the starting point of FeP development, based on the immunohistochemistry of 9 tumours that stained positively for carcinoembryonic antigen (CEA) [14]. This glycoprotein is usually found in eccrine sweat glands as well as gastrointestinal tumours and fetal tissues. In 2007 Stern and Haupt provided another argument that supports this theory: FeP lesions have been found in glabrous skin areas that lack hair follicles [15]. In 2007, Kurokawa et al. also supported the hypothesis, proposing that FeP may originate from intraepidermal eccrine ducts and afterwards proliferate into the dermis [16]. An important observation is that BCC might replace a FeP or develop independently by the eccrine ductal spread.[14]

Further contributing to the debate regarding the true origin of this tumour, it has been shown that both Merkel cells and androgen receptors are found in FeP.

In a study of 76 adnexal tumours of skin Wollina and Schrepel (2006) reported that trichoepitheliomas had an increased proliferative activity but not an increased number of MC (17). In basal cell carcinomas and trichofolliculomas, however, MC was found near proliferating tumour cells. Androgen receptors are expressed in BCCs, but only exceptionally in trichoepitheliomas [7]. Katona et al. (2007) observed that 10 out of 13 FeP, as well as 8 out of 11 BCCs, stained positively for androgen receptors [7]; only 2 out of 15 trichoepitheliomas expressed androgen receptors [7]. These results support the theory that FeP is an indolent BCC subtype. Merkel cells are a feature of benign follicular tumours, and it is well known that moderate hyperplasia of Merkel cells is present in chronic sun-damaged skin and hypertrophic actinic keratoses [18]. Merkel cells are found in trichoepitheliomas but are

absent in BCCs [7]. This supports the theory that FeP may be a fenestrated trichoepithelioma subtype.

To provide more clarity in this debate, Sellheyer et al. (2012) used a stem cell marker, PHLDA1 (TDAG51), which is expressed in the basal cell layer during embryogenesis and is present in trichoepitheliomas, but not BCCs. The results were consistent with a mixed histological pattern, showing an anastomosing network of thin cellular strands positive for PHLDA1 and basaloid nests with negative results for this marker [3]. The authors concluded that FeP is a premalignant lesion with a specific type of epidermal hyperplasia that stains positively for PHLDA-1 and has the capability of developing multifocal BCCs [3].

Previous literature reports have linked FeP with several risk factors:

**Genetic factors:** Some authors theorise that, like BCC, mutations in p53 and PATCHED-1 genes may also be responsible for the development of FeP [7, 19] and that both tumours originate in the follicular germinative cells. Others suggest that FeP might be a premalignant lesion that progresses to BCC by acquiring additional genetic mutations [20] or that it might develop from seborrheic keratosis, based on some histopathologic similarities [4]. Also, FePs have been identified in continuity with both BCC and seborrheic keratosis [4].

**Radiotherapy:** There is a frequent occurrence of FeP in patients with a history of radiotherapy. Hartschuch et al. observed hyperplasia of Merkel cells in chronic radiation dermatitis, as well as their presence in FePs and trichoepitheliomas, though they are absent in BCCs. The researchers further suggested that Merkel cells might be responsible for the benign biological behaviour of the tumours that have them [18]. Although previous irradiation might constitute the initial carcinogenic factor in some cases, this is not invariably the case, and the evolution and prognosis of the tumours are the same when compared to those due to other triggering factors [21]. **Sun-exposure:** It was observed that unlike BCCs, FeP has a predilection for sun-protected skin, is often located in the dorsal, lumbar and sacral regions. Bowen et al. showed that only 5% of tumours develop in anatomic sites that receive significant amounts of solar elastosis [13].

**Association with other neoplasms:** Some authors have suggested that FeP might be a reactive process (22) associated with other neoplasms, including breast cancer [23, 24], extramammary Paget's disease [25], gastrointestinal neuroendocrine tumours [26] or BCCs and Gorlin-Goltz syndrome [27]. Longo et al. (2016) investigated whether FeP is an expression of a more complex gastrointestinal syndrome and observed that in 9 of the 49 cases it was associated with gastrointestinal tumours [26]. The expression of CEA in both FeP and gastrointestinal tumours could suggest a common pathogenesis [14].

Further epidemiological studies are required to clarify whether or not these associations are coincidental and whether patients with FeP should be screened for the occurrence of other malignancies [26]

**Chronic inflammation:** There has been a case report of malignant degeneration in a chronic lower limb ulcer with the histological image of a FeP [28]. Clinically, most FePs appear in individuals between 40 and 60 years old, with a history of BCC [19], of solitary or multiple slow-growing papules or plaques. Typically, the lesions appear as flesh-coloured, pink, red, grey or even brown [29, 30], firm, sessile or pedunculated papules with a broad base. As previously noted, and in contrast to most BCCs, FeP develops mainly in sun-protected areas such as the lower back, inguinal, and crural areas, or the extremities. Atypical clinical forms of FeP may present as multiple lesions with different presentations [30]; these include ulcerated lesions or tumours arising in atypical locations such as the head, axillae, torso, umbilicus [31], plantar region [14, 32], or even on mucocutaneous junctions [33].

FeP is considered to be either a rare tumour or an underdiagnosed one because of its non-specific clinical appearance. Therefore, it can easily be mistaken for a wide variety of skin lesions such as acrochordons, intradermal or compound nevi, sebaceous nevi, seborrheic keratoses, fibromas, lipomas, cysts, and even amelanotic melanomas [10, 30, 34, 35].

Regarding diagnostic methods, the dermoscopy of FeP is not specific, as there are mixed features of BCC and trichoepithelioma, consisting of polymorphous vessels - mainly short arborizing and dotted types, gray-brown and gray-blue areas and dots, and shiny white streaks (also known as crystalline structures) [8, 36], secondary to fibrosis. Some structures similar to BCC or seborrheic keratosis may be present, such as ulceration, large ovoid nests, and milia-like cysts. Recently, Kornreich et al. reported an additional dermoscopic feature of FeP; namely, a white network, which could be more specific for FeP [37]. The white network is the manifestation of elongated hyperplastic, anastomosing epithelial strands [37]. This finding, together with additional BCC-related dermoscopic features, may facilitate the diagnosis of FeP [37].

The main reflectance confocal microscopy findings of FeP include a fenestrated pattern corresponding to the fibrous stroma and the presence of bright cells in pigmented lesions [38].

The histopathological features of FeP are distinctive, characterised by long, anastomosing strands of basaloid cells, embedded in a fibrous stroma, that project downwards from the epidermis and extend into the papillary dermis, giving the tumour a honeycomb or sponge-like appearance [22]. The cells from the edge of the strands are columnar and

arranged in a palisade. Sometimes, follicular germ-like structures can be identified within the tumour [4]. FeP usually has a distinct interface with the normal, underlying dermis [13], but sometimes tumour cells may extend into the reticular dermis [4].

FeP is typically treated by complete surgical excision, with generally excellent results [10]. Other possible treatment options include cryosurgery, electrodesiccation, or Mohs micrographic surgery [7]. In contrast to the treatment of some BCCs, topical Imiquimod 5% has been proven to be ineffective in the treatment of FeP [36]. Overall, FeP is a not-aggressive tumour with no metastatic potential and good prognosis after complete surgical excision [39].

In conclusion, the tumour presented in our case report appeared in a male in his eighties, which stands in contrast to epidemiological studies in which the majority of FePs appear to occur in women in the fourth to fifth decades of life. Nevertheless, the clinical features of the lesion and its location in the left lower quadrant of the abdomen are classical presenting features of FeP. The systematic review performed in this paper includes a total number of 452 FePs that have been reported in the medical literature. Even though FeP is considered a relatively rare tumour, its true incidence rate might be higher than suggested in published studies. Outside of dermatology, FeP is a relatively unknown tumour in the medical field, and, as indicated by our review, it can be easily confused with other benign or even malignant tumours.

The histopathologic findings appear to support Ackerman's theory that FeP is a 'trichoblastic (basal-cell) carcinoma' [4], sharing both features of BCC and trichoepithelioma. To further sustain this argument, the history of our patient's tumour evolution raises the hypothesis that a lesion originating as a trichoblastoma may have acquired additional genetic mutations over time, progressing to the premalignant lesion described in our histopathology report.

There is no known history of radiotherapy in our patient, and the abdominal ultrasound, chest x-ray and ultrasound studies of the axillary and inguinal lymph node groups showed no evidence of disease progression or other malignancies. Further epidemiological studies will be required to clarify whether or not these associations are merely hypothetical or fortuitous, or if patients with FeP should be carefully screened for the occurrence of other malignancies.

## References

1. Hurt MA, Cribier B, Kaddu S, Schulz T, Kutzner H, Hartschuh W. Chapter 3—appendageal tumors. In: LeBoit PE, Burg G, Weedon D, Sarasin A, editors. World Health Organization classification of tumors: pathology and genetics of skin tumors. Lyon: IARC, 2006: p. 152–153.



2. Pinkus H. Premalignant fibroepithelial tumors of skin. *AMA Arch Derm Syphilol.* 1953; 67(6):598-615. <https://doi.org/10.1001/archderm.1953.01540060060009> PMID:13050184
3. Sellheyer K, Nelson P, Kutzner H. Fibroepithelioma of Pinkus is a true basal cell carcinoma developing in association with a newly identified tumour-specific type of epidermal hyperplasia. *Br J Dermatol.* 2012; 166(1):88-97. <https://doi.org/10.1111/j.1365-2133.2011.10612.x> PMID:21910710
4. Ackerman AB, Gottlieb GJ. Fibroepithelial tumor of pinkus is trichoblastic (Basal-cell) carcinoma. *Am J Dermatopathol.* 2005; 27(2):155-9. <https://doi.org/10.1097/01.dad.0000154395.14971.71>
5. Cotsarelis G. Epithelial stem cells: a folliculocentric view. *J Invest Dermatol.* 2006; 126(7):1459-68. <https://doi.org/10.1038/sj.jid.5700376> PMID:16778814
6. Peterson SC, Eberl M, Vagnozzi AN, Belkadi A, Veniaminova NA, Verhaegen ME, Bichakjian CK, Ward NL, Dlugosz AA, Wong SY. Basal cell carcinoma preferentially arises from stem cells within hair follicle and mechanosensory niches. *Cell Stem Cell.* 2015; 16(4):400-12. <https://doi.org/10.1016/j.stem.2015.02.006> PMID:25842978 PMCID:PMC4387376
7. Katona TM, Ravis SM, Perkins SM, Moores WB, Billings SD. Expression of androgen receptor by fibroepithelioma of Pinkus: evidence supporting classification as a basal cell carcinoma variant? *Am J Dermatopathol.* 2007; 29(1):7-12. <https://doi.org/10.1097/01.dad.0000245201.59762.d2> PMID:17284955
8. Reggiani C, Zalaudek I, Piana S, Longo C, Argenziano G, Lallas A, Pellacani G, Moscarella E. Fibroepithelioma of Pinkus: case reports and review of the literature. *Dermatology.* 2013; 226(3):207-11. <https://doi.org/10.1159/000348707> PMID:23711617
9. Zalaudek I, Ferrara G, Broganelli P, Moscarella E, Mordente I, Giacomel J, Argenziano G. Dermoscopy patterns of fibroepithelioma of pinkus. *Arch Dermatol.* 2006; 142(10):1318-22. <https://doi.org/10.1001/archderm.142.10.1318> PMID:17043187
10. Cohen PR, Tschen JA. Fibroepithelioma of Pinkus presenting as a sessile thigh nodule. *Skinmed.* 2003; 2(6):385-7. <https://doi.org/10.1111/j.1540-9740.2003.02491.x>
11. Pan Z, Huynh N, Sarma DP. Fibroepithelioma of pinkus in a 9-year-old boy: a case report. *Cases J.* 2008; 1(1):21. <https://doi.org/10.1186/1757-1626-1-21> PMID:18588684 PMCID:PMC2474574
12. Scalvenzi M, Francia MG, Falletti J, Balato A. Basal cell carcinoma with fibroepithelioma-like histology in a healthy child: report and review of the literature. *Pediatr Dermatol.* 2008; 25(3):359-63. <https://doi.org/10.1111/j.1525-1470.2008.00683.x> PMID:18577044
13. Bowen AR, LeBoit PE. Fibroepithelioma of pinkus is a fenestrated trichoblastoma. *Am J Dermatopathol.* 2005; 27(2):149-54. <https://doi.org/10.1097/01.dad.0000138051.71415.fe>
14. Stern JB, Haupt HM, Smith RR. Fibroepithelioma of Pinkus. Eccrine duct spread of basal cell carcinoma. *Am J Dermatopathol.* 1994; 16(6):585-7. <https://doi.org/10.1097/00000372-199412000-00002> PMID:7864295
15. Stern JB, Haupt HM. Basal cell carcinoma of the sole: implications for fibroepithelioma of pinkus. *Am J Dermatopathol.* 2007; 29(5):494. <https://doi.org/10.1097/DAD.0b013e3181462889> PMID:17890926
16. Kurokawa I, Yokoyama T, Nishimura K, Hakamada A, Isoda K, Yamanaka K, Tsubura A, Mizutani H. Pinkus tumor may originate from intraepidermal eccrine ducts and proliferate in the dermis. *Oncol Rep.* 2007; 17(1):49-54. <https://doi.org/10.3892/or.17.1.49>
17. Wollina U, Schrepel U. Merkel cells in adnexal neoplasias of skin. *Čes-slov Derm.* 2006;81(4):206-9.
18. Hartschuh W, Schulz T. Merkel cell hyperplasia in chronic radiation-damaged skin: its possible relationship to fibroepithelioma of Pinkus. *J Cutan Pathol.* 1997; 24(8):477-83. <https://doi.org/10.1111/j.1600-0560.1997.tb01321.x> PMID:9331893
19. Tarallo M, Cigna E, Fino P, Lo Torto F, Corrias F, Scuderi N. Fibroepithelioma of Pinkus: variant of basal cell carcinoma or trichoblastoma? Case report. *G Chir.* 2011; 32(6-7):326-8. PMID:21771402
20. Misago N, Suzuki Y, Miura Y, Narisawa Y. Giant polypoid basal cell carcinoma with features of fibroepithelioma of Pinkus and extensive cornification. *Eur J Dermatol.* 2004; 14(4):272-5. PMID:15319165
21. Cahuzac P, Hermier C, Thivolet J. [Premalignant fibroepithelial tumors of Pinkus and previous irradiation: report of a survey of 20 patients]. *Ann Dermatol Venereol.* 1982; 109(4):355-7. PMID:7125488
22. Haddock ES, Cohen PR. Fibroepithelioma of Pinkus Revisited. *Dermatol Ther (Heidelb).* 2016; 6(3):347-62. <https://doi.org/10.1007/s13555-016-0123-8> PMID:27329375 PMCID:PMC4972729
23. Bryant J. Fibroepithelioma of Pinkus overlying breast cancer. *Arch Dermatol.* 1985; 121(3):310. <https://doi.org/10.1001/archderm.1985.01660030030013> PMID:2983617
24. Sandbank M, Bashan D. Multiple trichoepithelioma and breast carcinoma. Simultaneous appearance of epithelioma adenoides cysticum and infiltrating lobular carcinoma of the breast. *Arch Dermatol.* 1978; 114(8):1230. <https://doi.org/10.1001/archderm.1978.01640200082026> PMID:677927
25. Warner TF, Burgess H, Mohs FE. Extramammary Paget's disease in fibroepithelioma of Pinkus. *J Cutan Pathol.* 1982; 9(5):340-4. <https://doi.org/10.1111/j.1600-0560.1982.tb01070.x> PMID:6292274
26. Longo C, Pellacani G, Tomasi A, Mandel VD, Ponti G. Fibroepithelioma of Pinkus: Solitary tumor or sign of a complex gastrointestinal syndrome. *Mol Clin Oncol.* 2016; 4(5):797-800. <https://doi.org/10.3892/mco.2016.794>
27. Gorlin RJ, Goltz RW. Multiple nevoid basal-cell epithelioma, jaw cysts and bifid rib. A syndrome. *N Engl J Med.* 1960; 262:908-12. <https://doi.org/10.1056/NEJM196005052621803> PMID:13851319
28. Ebner H, Niebauer G. Malignant degeneration of an ulcer cruris under the histological image of a fibroepithelioma Pinkus. *Z Haut Geschlechtskr.* 1967; 42(12):417-21. PMID:5590296
29. Strauss RM, Edwards S, Stables GI. Pigmented fibroepithelioma of Pinkus. *Br J Dermatol.* 2004; 150(6):1208-9. <https://doi.org/10.1111/j.1365-2133.2004.06018.x> PMID:15214913
30. Sehli Attafi S, Jones M, Faza'a B, Zermani R, Rommany SR. Pinkus tumour: an unusual case. *Pathologica.* 2013; 105(4):140-1. PMID:24466765
31. Inskip M, Longo C, Haddad A. Two adjacent individual fibroepithelioma of Pinkus of the umbilicus-one pink, one pigmented-a case report and review of the literature. *Dermatol Pract Concept.* 2016; 6(2):17-20. <https://doi.org/10.5826/dpc.0602a04> PMID:27222767 PMCID:PMC4866622
32. Lin TC, Lee TL, Lin TY, Wu PY. An infrequent case of neoplasm with fibroepithelioma of pinkus and hidradenomatous features arising at the umbilicus: a rare finding. *Am J Dermatopathol.* 2011; 33(7):750-2. <https://doi.org/10.1097/DAD.0b013e31820978cf> PMID:21885941
33. Karatas D, Erten S. Pinkus tumor at the nostril floor. *J Craniofac Surg.* 2014; 25(4):e317-8. <https://doi.org/10.1097/SCS.0000000000000582> PMID:24978450
34. Gellin GA, Bender B. Giant premalignant fibroepithelioma. *Arch Dermatol.* 1966; 94(1):70-3. <https://doi.org/10.1001/archderm.1966.01600250076015>
35. Jones CC, Ansari SJ, Tschen JA. Cystic fibroepithelioma of pinkus. *J Cutan Pathol.* 1991; 18(3):220-2. <https://doi.org/10.1111/j.1600-0560.1991.tb00157.x> PMID:1918510
36. Zalaudek I, Leinweber B, Ferrara G, Soyer HP, Ruocco E,

Argenziano G. Dermoscopy of fibroepithelioma of pinkus. J Am Acad Dermatol. 2005; 52(1):168-9.

<https://doi.org/10.1016/j.jaad.2004.06.033> PMID:15627105

37. Kornreich DA, Lee JB. White network in fibroepithelioma of Pinkus. JAAD Case Rep. 2016; 2(5):400-02.

<https://doi.org/10.1016/j.jidcr.2016.07.011> PMID:27766301

PMCID:PMC5065638

38. Longo C, Soyer HP, Pepe P, Casari A, Wurm EM, Guitera P, Pellacani G. In vivo confocal microscopic pattern of fibroepithelioma of pinkus. Arch Dermatol. 2012; 148(4):556.

<https://doi.org/10.1001/archdermatol.2011.945> PMID:22508888

39. Su MW, Fromer E, Fung MA. Fibroepithelioma of pinkus. Dermatol Online J. 2006; 12(5):2. PMID:16962017