

Analytical treatments of time-fractional seventh-order nonlinear equations via Elzaki transform

Ali, Liaqat; Zou, Guang; Li, Na; Mehmood, Kashif; Fang, Pan; Khan, Adnan

DOI

[10.1007/s10665-023-10326-y](https://doi.org/10.1007/s10665-023-10326-y)

Publication date

2024

Document Version

Final published version

Published in

Journal of Engineering Mathematics

Citation (APA)

Ali, L., Zou, G., Li, N., Mehmood, K., Fang, P., & Khan, A. (2024). Analytical treatments of time-fractional seventh-order nonlinear equations via Elzaki transform. *Journal of Engineering Mathematics*, 145(1), Article 1. <https://doi.org/10.1007/s10665-023-10326-y>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

Green Open Access added to TU Delft Institutional Repository

'You share, we take care!' - Taverne project

<https://www.openaccess.nl/en/you-share-we-take-care>

Otherwise as indicated in the copyright section: the publisher is the copyright holder of this work and the author uses the Dutch legislation to make this work public.



Analytical treatments of time-fractional seventh-order nonlinear equations via Elzaki transform

Liaquat Ali^{1,2} · Guang Zou¹ · Na Li¹ · Kashif Mehmood² · Pan Fang³ · Adnan Khan⁴

Received: 19 October 2023 / Accepted: 14 December 2023
© The Author(s), under exclusive licence to Springer Nature B.V. 2024

Abstract

In this article, we'll show how to solve the time-fractional seventh-order Lax's Korteweg–de Vries and Kaup–Kupershmidt equations analytically using the homotopy perturbation approach, the Adomian decomposition method, and the Elzaki transformation. The KdV equation is a general integrable equation with an inverse scattering transform-based solution that arises in a variety of physical applications, including surface water waves, internal waves in a density stratified fluid, plasma waves, Rossby waves, and magma flow. Fractional derivative is described in the Caputo sense. The solutions to fractional partial differential equation is computed using convergent series. The numerical computations and graphical representations of the analytical results obtained using the homotopy perturbation and decomposition techniques. Moreover, plots that are simple to grasp are used to compare the integer order and fractional-order solutions. After only a few iterations, we may easily obtain numerical results that provide us better approximations. The exact solutions and the derived solutions were observed to be very similar. The suggested methods have also acquired the highest level of accuracy. The most prevalent and convergent techniques for resolving nonlinear fractional-order partial differential issues are the applied techniques.

Keywords Analytical techniques · Caputo operator · Elzaki Transform · Kaup–Kupershmidt (KK) equation · Lax's Korteweg–de Vries equation

Mathematics Subject Classification 35A20 · 26A33 · 34A25 · 35Q53

✉ Guang Zou
zoug@sustech.edu.cn

¹ Department of Ocean Science and Engineering, Southern University of Science and Technology, Shenzhen, People's Republic of China

² College of Civil Engineering and Architecture, Zhejiang University, Hangzhou 310058, China

³ Department of Maritime and Transport Technology, Delft University of Technology, 2600 AA Delft, The Netherlands

⁴ Department of Mathematics, Abdul Wali Khan University Mardan, Mardan 23200, Pakistan

1 Introduction

The derivative of a function can be extended in any order using the branch of calculus known as fractional calculus. The use of fractional calculus across numerous disciplines of applied science and engineering to describe the properties of various real physical phenomena has captured the attention of many scholars in particular fields of applied science and engineering in recent years. The concept of the fractional derivative has been established in response to the problems brought on by heterogeneity. The fast development of mathematical techniques with computer packages led to many researchers working on FC to illustrate their belvederes while studying complex models. Recently, senior scholars provided a number of novel ideas for the FC, and those provided the framework. FC is a commonly used idea and theory that is connected to real-world initiatives [1–5]. Integrals and derivatives are addressed by FC to an arbitrary real or complex order. Recently, a number of fractional operators, including Caputo, Caputo Fabrizio, Atangana–Baleanu, Katugampola, Hilfer, etc. have been proposed and implemented for dealing with real-world applications. The fractional calculus has demonstrated that it is the best tool for studying problems in the actual world. The Caputo fractional derivative is often used in practical applications, as it enables one to include the traditional initial and boundary conditions in formulating mathematical models. Moreover, as in the integer-order derivative, the Caputo fractional derivative of a constant is zero [6].

A detailed association among an unknown function and its partial derivatives is expressed using partial differential equations (PDEs). Nearly every area of engineering and study uses PDEs. PDEs are now being used more frequently in disciplines like biology, economics, image processing, graphics and social sciences. As a result, relevant functions in these variables can be formed when some independent variables relate with one another in every of the areas listed before. This enables the modeling of diverse processes through equations for the corresponding functions. There are many aspects to the study of PDEs. Developing techniques for identifying explicit solutions was the conventional approach that predominated the eighteenth century. It's vital to note that certain highly complicated problems were not solved via computers. In most cases, it is also preferred for the solution to be distinct and robust to minor data interruptions. If these requirements are satisfied, the equation can be understood theoretically [7–9].

Modern calculus tools like fractional partial differential equations (FPDEs) can be used to model a number of events in the applied sciences and engineering. Researchers began to be interested in fractional calculus since it was difficult to treat nonlinear real-world processes in conventional calculus [10–16]. The proper description of the behavior of significant physical processes in this context depends on the approximation of FPDEs and analytical solutions. In light of the aforementioned assertion, mathematicians have developed and taken into practice a number of computations and analytical techniques to determine the solutions for a variety of significant mathematical models that represent challenges. Mathematicians continue to make the best efforts possible in this area despite the fact that computing the analytical and sometimes even approximative solutions of some nonlinear FPDEs and systems of FPDEs is very difficult. Several studies have been done throughout the years in the domains

of science and engineering all across the world, and many approaches have been created to offer the best solutions attainable. The world is always being faced with new, tough, and complex difficulties and problems; examples include [17–21]. This is an unstoppable process, and new techniques are being established on a daily basis. The literature has suggested a variety of approaches to addressing FDEs, which include the Fractional complex transform [22], Finite difference methods [23], Adomian Decomposition method [24], Residual power series method [25], Homotopy Analysis method [26], Differential transform method [27], the Variational Iteration method [28], and the predictor-corrector approach [29].

This article develops the solution of time-fractional seventh-order Lax's Korteweg–de Vries equation and Kaup–Kupershmidt equation using the Elzaki transform decomposition technique (ETDM) and the Homotopy Perturbation Transform Method (HPTM). The Elzaki transform was created by Tarig Elzaki to help resolving ordinary and PDEs in the time domain simpler. On the other hand, the Adomian decomposition approach [30, 31], which is well known, provides precise solutions in the form of a convergent series for the solution of linear and nonlinear, homogeneous and nonhomogeneous differential equations. In 1998, he introduced HPM [32, 33]. According to this method, the accuracy is assumed to be an in series solution with a large number of terms that quickly converges to the actual derived solution. Nonlinear PDEs can be successfully solved using this method. A higher degree of accuracy was demonstrated when the HPTM findings were contrasted with the actual solutions to the problems. The newly developed technique is a mix form of HPM and the Elzaki transform. The time-fractional seventh-order nonlinear equations are solved analytically using the existing methods, which are demonstrated to be highly efficient. The outcomes of the suggested techniques are reliable and offer precise solutions to the desired issues. Our methods produced infinite series as the results in the numerical examples. When we write the series in closed form, it gives precise solutions to the relevant equations. Researchers can use this study as a fundamental reference to examine these strategies and employ it in many applications to get accurate and approximative results in a few easy steps. The results of fractional problem analysis using the suggested methodologies are also used to examine the issues from a fractional aspect.

The Korteweg–de Vries (KdV) equation is an example of a partial differential equation. It has been used as a model for the evolution and interaction of nonlinear waves to describe a wide range of physical phenomena. This equation was derived as an evolution that controlled the propagation of long, low-amplitude, one-dimensional surface gravity waves in a shallow water channel [34]. Nowadays, the KdV equation is applied in many areas of physics, such as lattice dynamics, collision-free hydro-magnetic waves, stratified internal waves, ion-acoustic waves, and plasma physics [35]. Some possible physical phenomena in the framework of quantum mechanics have been represented using a KdV model. It serves as a model for the propagation of shock wave, turbulence, solitons, mass transport in fluid dynamics, boundary layer behavior, continuum mechanics and aerodynamics [36]. In this study, we aim to solve a fractional-order nonlinear Lax's Korteweg–de Vries equation using two analytical methods.

$$\begin{aligned}
D_1^\mu \mathcal{Y}(\omega, 1) = & -140\mathcal{Y}^3(\omega, 1)\mathcal{Y}_\omega(\omega, 1) - 70\mathcal{Y}_\omega^3(\omega, 1) \\
& - 280\mathcal{Y}(\omega, 1)\mathcal{Y}_\omega(\omega, 1)\mathcal{Y}_{\omega\omega}(\omega, 1) - 70\mathcal{Y}^2(\omega, 1)\mathcal{Y}_{\omega\omega\omega}(\omega, 1) \\
& - 70\mathcal{Y}_{\omega\omega}(\omega, 1)\mathcal{Y}_{\omega\omega\omega}(\omega, 1) - 42\mathcal{Y}_\omega(\omega, 1)\mathcal{Y}_{\omega\omega\omega\omega}(\omega, 1) \\
& - 14\mathcal{Y}(\omega, 1)\mathcal{Y}_{\omega\omega\omega\omega\omega}(\omega, 1) - \mathcal{Y}_{\omega\omega\omega\omega\omega\omega}(\omega, 1), \\
& 0 < \mu \leq 1,
\end{aligned} \tag{1}$$

and Kaup–Kupershmidt equation

$$\begin{aligned}
D_1^\mu \mathcal{Y}(\omega, 1) = & -2016\mathcal{Y}^3(\omega, 1)\mathcal{Y}_\omega(\omega, 1) - 630\mathcal{Y}_\omega^3(\omega, 1) \\
& - 2268\mathcal{Y}(\omega, 1)\mathcal{Y}_\omega(\omega, 1)\mathcal{Y}_{\omega\omega}(\omega, 1) - 504\mathcal{Y}^2(\omega, 1)\mathcal{Y}_{\omega\omega\omega}(\omega, 1) \\
& - 252\mathcal{Y}_{\omega\omega}(\omega, 1)\mathcal{Y}_{\omega\omega\omega}(\omega, 1) - 147\mathcal{Y}_\omega(\omega, 1)\mathcal{Y}_{\omega\omega\omega\omega}(\omega, 1) \\
& - 42\mathcal{Y}(\omega, 1)\mathcal{Y}_{\omega\omega\omega\omega\omega}(\omega, 1) - \mathcal{Y}_{\omega\omega\omega\omega\omega\omega}(\omega, 1), \\
& 0 < \mu \leq 1,
\end{aligned} \tag{2}$$

The parameter μ here denotes the order of the fractional derivative. These equations serve as a mathematical representation of the complex physical processes that develop in biology, physics, chemistry and engineering. Examples include nonlinear optics, quantum mechanics, plasma physics, long wave propagation in shallow water under gravity, and fluid mechanics. Several researchers have used the modified Cole–Hopf transformation method [37], variational iteration method [38], and pseudospectral method [39] to solve the seventh-order Lax’s Korteweg–de Vries. In [40], time-fractional Rosenau–Hyman equation is solved numerically using the residual power series technique and perturbation-iteration algorithm which is a model that is comparable to KdV. Pomeau et al. [41] investigated the stability of the KdV equation in terms of singular perturbation and came up with the classical model of Eq. 2.

The format of the present article is as follows: In Sect. 2, we begin with the fundamental concept of fractional calculus. In Sects. 3 and 4, we go over the core ideas behind the suggested methods. In Sects. 5 we give the convergence analysis of the suggested techniques. These methods are used in Sect. 6 to solve the time-fractional Lax’s Korteweg–de Vries and Kaup–Kupershmidt (KK) issues with the given initial condition. The conclusion is presented in Sect. 6.

2 Preliminaries

We presented some fundamental concept of fractional calculus.

2.1 Definition

The fractional derivative in Abel–Riemann manner is taken as [42–44]

$$D^\mu \mathcal{Y}(\omega) = \begin{cases} \frac{d^\zeta}{d\omega^\zeta} \mathcal{Y}(\omega), & \mu = \zeta, \\ \frac{1}{\Gamma(\zeta - \mu)} \frac{d}{d\omega^\zeta} \int_0^\omega \frac{\mathcal{Y}(\phi)}{(\omega - \phi)^{\mu - \zeta + 1}} d\phi, & \zeta - 1 < \mu < \zeta, \end{cases}$$

where $\varsigma \in Z^+$, $\mu \in R^+$ and

$$D^{-\mu} \mathcal{Y}(\omega) = \frac{1}{\Gamma(\mu)} \int_0^\omega (\omega - \phi)^{\mu-1} \mathcal{Y}(\phi) d\phi, \quad 0 < \mu \leq 1.$$

2.2 Definition

The fractional integral in Abel–Riemann manner is taken as [42–44]

$$J^\mu \mathcal{Y}(\omega) = \frac{1}{\Gamma(\mu)} \int_0^\omega (\omega - \phi)^{\mu-1} \mathcal{Y}(\omega) d\omega, \quad \omega > 0, \quad \mu > 0.$$

with below properties

$$J^\mu \omega^\varsigma = \frac{\Gamma(\varsigma + 1)}{\Gamma(\varsigma + \mu + 1)} \omega^{\varsigma+\mu},$$

$$D^\mu \omega^\varsigma = \frac{\Gamma(\varsigma + 1)}{\Gamma(\varsigma - \mu + 1)} \omega^{\varsigma-\mu}.$$

2.3 Definition

The fractional derivative in Caputo manner is taken as [42–44]

$$D^\mu \mathcal{Y}(\omega) = \begin{cases} \frac{1}{\Gamma(\varsigma-\mu)} \int_0^\omega \frac{\mathcal{Y}^\varsigma(\phi)}{(\omega-\phi)^{\mu-\varsigma+1}} d\phi, & \varsigma - 1 < \mu < \varsigma, \\ \frac{d^\varsigma}{d\omega^\varsigma} \mathcal{Y}(\omega), & \varsigma = \mu. \end{cases} \tag{3}$$

with below properties

$$J_\omega^\mu D_\omega^\mu \mathcal{Y}(\omega) = g(\omega) - \sum_{k=0}^m g^k(0^+) \frac{\omega^k}{k!}, \quad \text{for } \omega > 0, \text{ and } \varsigma - 1 < \mu \leq \varsigma, \quad \varsigma \in N. \tag{4}$$

$$D_\omega^\mu J_\omega^\mu \mathcal{Y}(\omega) = g(\omega).$$

2.4 Definition

The ET of Caputo operator is taken as

$$E[D_\omega^\mu \mathcal{Y}(\omega)] = s^{-\mu} E[\mathcal{Y}(\omega)] - \sum_{k=0}^{\varsigma-1} s^{2-\mu+k} \mathcal{Y}^{(k)}(0), \quad \text{where } \varsigma - 1 < \mu < \varsigma.$$

3 Analysis of the HPTM

To present the concept of HPTM, we examine the FPDE of the form

$$D_1^\mu \mathcal{Y}(\omega, \mathfrak{t}) = \mathcal{F}_1[\omega] \mathcal{Y}(\omega, \mathfrak{t}) + \mathcal{G}_1[\omega] \mathcal{Y}(\omega, \mathfrak{t}), \quad 0 < \mu \leq 1, \quad (5)$$

with initial condition

$$\mathcal{Y}(\omega, 0) = \kappa(\omega).$$

Here, $D_1^\mu = \frac{\partial^\mu}{\partial \mathfrak{t}^\mu}$ denotes the fractional Caputo operator of order μ , and $\mathcal{F}_1[\omega]$, $\mathcal{G}_1[\omega]$ are linear and nonlinear functions.

Apply the ET, we get

$$\mathbb{E}[D_1^\mu \mathcal{Y}(\omega, \mathfrak{t})] = \mathbb{E}[\mathcal{Y}_1[\omega] \mathcal{Y}(\omega, \mathfrak{t}) + \mathcal{R}_1[\omega] \mathcal{Y}(\omega, \mathfrak{t})], \quad (6)$$

$$\frac{1}{u^\mu} \{M(u) - u^2 \mathcal{Y}(\omega, 0)\} = \mathbb{E}[\mathcal{F}_1[\omega] \mathcal{Y}(\omega, \mathfrak{t}) + \mathcal{G}_1[\omega] \mathcal{Y}(\omega, \mathfrak{t})]. \quad (7)$$

On simplification we have

$$M(u) = u^2 \mathcal{Y}(\omega, 0) + u^\mu \mathbb{E}[\mathcal{F}_1[\omega] \mathcal{Y}(\omega, \mathfrak{t}) + \mathcal{G}_1[\omega] \mathcal{Y}(\omega, \mathfrak{t})]. \quad (8)$$

Apply the inverse ET, we get

$$\mathcal{Y}(\omega, \mathfrak{t}) = \mathcal{Y}(\omega, 0) + \mathbb{E}^{-1}[u^\mu \mathbb{E}[\mathcal{F}_1[\omega] \mathcal{Y}(\omega, \mathfrak{t}) + \mathcal{G}_1[\omega] \mathcal{Y}(\omega, \mathfrak{t})]]. \quad (9)$$

By applying homotopy perturbation method (HPM) to (9), we have

$$\mathcal{Y}(\omega, \mathfrak{t}) = \mathcal{Y}(\omega, 0) + \epsilon [\mathbb{E}^{-1}[u^\mu \mathbb{E}[\mathcal{F}_1[\omega] \mathcal{Y}(\omega, \mathfrak{t}) + \mathcal{G}_1[\omega] \mathcal{Y}(\omega, \mathfrak{t})]]]. \quad (10)$$

The basic series form solution is as

$$\mathcal{Y}(\omega, \mathfrak{t}) = \sum_{k=0}^{\infty} \epsilon^k \mathcal{Y}_k(\omega, \mathfrak{t}), \quad (11)$$

having homotopy parameter $\epsilon \in [0, 1]$.

The nonlinear term is taken as

$$\mathcal{G}_1[\omega] \mathcal{Y}(\omega, \mathfrak{t}) = \sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{Y}). \quad (12)$$

The homotopy polynomial H_k is determined as

$$H_k(\mathcal{Y}_0, \mathcal{Y}_1, \dots, \mathcal{Y}_n) = \frac{1}{\Gamma(n+1)} D_\epsilon^k \left[\mathcal{G}_1 \left(\sum_{i=0}^{\infty} \epsilon^i \mathcal{Y}_i \right) \right]_{\epsilon=0}, \quad (13)$$

with $D_\epsilon^k = \frac{\partial^k}{\partial \epsilon^k}$.

By inserting (11) and (12) in (10), we have

$$\sum_{k=0}^{\infty} \epsilon^k \mathcal{Y}_k(\omega, 1) = \mathcal{Y}(\omega, 0) + \epsilon \times \left(\mathbb{E}^{-1} \left[u^\mu \mathbb{E}\{\mathcal{F}_1 \sum_{k=0}^{\infty} \epsilon^k \mathcal{Y}_k(\omega, 1) + \sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{Y})\} \right] \right). \tag{14}$$

Equating the ϵ coefficient, we have

$$\begin{aligned} \epsilon^0 : \mathcal{Y}_0(\omega, 1) &= \mathcal{Y}(\omega, 0), \\ \epsilon^1 : \mathcal{Y}_1(\omega, 1) &= \mathbb{E}^{-1} [u^\mu \mathbb{E}(\mathcal{F}_1[\omega] \mathcal{Y}_0(\omega, 1) + H_0(\mathcal{Y}))], \\ \epsilon^2 : \mathcal{Y}_2(\omega, 1) &= \mathbb{E}^{-1} [u^\mu \mathbb{E}(\mathcal{F}_1[\omega] \mathcal{Y}_1(\omega, 1) + H_1(\mathcal{Y}))], \\ &\vdots \\ &\vdots \\ &\vdots \\ \epsilon^k : \mathcal{Y}_k(\omega, 1) &= \mathbb{E}^{-1} [u^\mu \mathbb{E}(\mathcal{F}_1[\omega] \mathcal{Y}_{k-1}(\omega, 1) + H_{k-1}(\mathcal{Y}))], \quad k > 0, k \in N. \end{aligned} \tag{15}$$

Hence, the series form solution of the proposed method is as

$$\mathcal{Y}(\omega, 1) = \lim_{M \rightarrow \infty} \sum_{k=1}^M \mathcal{Y}_k(\omega, 1). \tag{16}$$

4 Analysis of the ETDM

To present the concept of ETDM, we examine the FPDE of the form

$$D_1^\mu \mathcal{Y}(\omega, 1) = \mathcal{F}_1(\omega, 1) + \mathcal{G}_1(\omega, 1), \quad 0 < \mu \leq 1, \tag{17}$$

with initial condition

$$\mathcal{Y}(\omega, 0) = \kappa(\omega).$$

Here, $D_1^\mu = \frac{\partial^\mu}{\partial 1^\mu}$ denotes the fractional Caputo operator of order μ , and \mathcal{F}_1 and \mathcal{G}_1 are linear and non-linear functions.

Apply the ET, we get

$$\begin{aligned} \mathbb{E}[D_1^\mu \mathcal{Y}(\omega, 1)] &= \mathbb{E}[\mathcal{F}_1(\omega, 1) + \mathcal{G}_1(\omega, 1)], \\ \frac{1}{u^\mu} \{M(u) - u^2 \mathcal{Y}(\omega, 0)\} &= \mathbb{E}[\mathcal{F}_1(\omega, 1) + \mathcal{G}_1(\omega, 1)]. \end{aligned} \tag{18}$$

On simplification we have

$$M(u) = u \mathcal{Y}(\omega, 0) + u^\mu \mathbb{E}[\mathcal{F}_1(\omega, 1) + \mathcal{G}_1(\omega, 1)]. \tag{19}$$

Apply the inverse ET, we get

$$\mathcal{Y}(\omega, 1) = \mathcal{Y}(\omega, 0) + \mathbb{E}^{-1}[u^\mu \mathbb{E}[\mathcal{F}_1(\omega, 1) + \mathcal{G}_1(\omega, 1)]]. \tag{20}$$

In terms of ADM, the basic series form solution is as

$$\mathcal{Y}(\omega, 1) = \sum_{m=0}^{\infty} \mathcal{Y}_m(\omega, 1). \tag{21}$$

The nonlinear term is taken as

$$\mathcal{G}_1(\omega, 1) = \sum_{m=0}^{\infty} \mathbb{A}_m(\mathcal{Y}), \tag{22}$$

with

$$\mathbb{A}_m(\mathcal{Y}_0, \mathcal{Y}_1, \mathcal{Y}_2, \dots, \mathcal{Y}_m) = \frac{1}{m!} \left[\frac{\partial^m}{\partial \ell^m} \left\{ \mathcal{G}_1 \left(\sum_{m=0}^{\infty} \ell^m \mathcal{Y}_m \right) \right\} \right]_{\ell=0}, \quad m = 0, 1, 2, \dots \tag{23}$$

By inserting (21) and (22) in (20), we have

$$\sum_{m=0}^{\infty} \mathcal{Y}_m(\omega, 1) = \mathcal{Y}(\omega, 0) + \mathbb{E}^{-1} u^\mu \left[\mathbb{E} \left\{ \mathcal{F}_1 \left(\sum_{m=0}^{\infty} \mathcal{Y}_m(\omega, 1) \right) + \sum_{m=0}^{\infty} \mathbb{A}_m(\mathcal{Y}) \right\} \right]. \tag{24}$$

By equating both sides, we have

$$\begin{aligned} \mathcal{Y}_0(\omega, 1) &= \mathcal{Y}(\omega, 0), \\ \mathcal{Y}_1(\omega, 1) &= \mathbb{E}^{-1} [u^\mu \mathbb{E}\{\mathcal{F}_1(\mathcal{Y}_0) + \mathbb{A}_0\}]. \end{aligned} \tag{25}$$

Hence, the general solution of the proposed method for $m \geq 1$ is as

$$\mathcal{Y}_{m+1}(\omega, 1) = \mathbb{E}^{-1} [u^\mu \mathbb{E}\{\mathcal{F}_1(\mathcal{Y}_m) + \mathbb{A}_m\}].$$

5 Convergence analysis

In this section, the proposed approaches convergence are illustrated.

Theorem 5.1 *Assume that the accurate solution of (5) is $\Psi(\theta, 1)$ and let $\Psi(\theta, 1), \Psi_n(\theta, 1) \in H$ and $\wp \in (0, 1)$, where H illustrates the Hilbert space. The obtained solution $\sum_{q=0}^{\infty} \Psi_q(\theta, 1)$ converge to $\Psi(\theta, 1)$ if $\Psi_q(\theta, 1) \leq \Psi_{q-1}(\theta, 1) \quad \forall q > A$, i.e., for any $\theta > 0 \exists A > 0$, such that $\|\Psi_{q+n}(\theta, 1)\| \leq \beta, \forall m, n \in N$.*

Proof Consider a sequence of $\sum_{q=0}^{\infty} \Psi_q(\theta, 1)$.

$$\begin{aligned}
 \mathcal{J}_0(\theta, 1) &= \Psi_0(\theta, 1), \\
 \mathcal{J}_1(\theta, 1) &= \Psi_0(\theta, 1) + \Psi_1(\theta, 1), \\
 \mathcal{J}_2(\theta, 1) &= \Psi_0(\theta, 1) + \Psi_1(\theta, 1) + \Psi_2(\theta, 1), \\
 \mathcal{J}_3(\theta, 1) &= \Psi_0(\theta, 1) + \Psi_1(\theta, 1) + \Psi_2(\theta, 1) + \Psi_3(\theta, 1), \\
 &\vdots \\
 \mathcal{J}_q(\theta, 1) &= \Psi_0(\theta, 1) + \Psi_1(\theta, 1) + \Psi_2(\theta, 1) + \dots + \Psi_q(\theta, 1).
 \end{aligned}
 \tag{26}$$

We must illustrate that $\mathcal{J}_q(\theta, 1)$ forms a "Cauchy sequence" in order to attain the chosen result. Also, let's take

$$\begin{aligned}
 \|\mathcal{J}_{q+1}(\theta, 1) - \mathcal{J}_q(\theta, 1)\| &= \|\Psi_{q+1}(\theta, 1)\| \leq \wp \|\Psi_q(\theta, 1)\| \\
 &\leq \wp^2 \|\Psi_{q-1}(\theta, 1)\| \leq \wp^3 \|\Psi_{q-2}(\theta, 1)\| \dots \\
 &\leq \wp_{q+1} \|\Psi_0(\theta, 1)\|.
 \end{aligned}
 \tag{27}$$

For $q, n \in N$, we have

$$\begin{aligned}
 &\|\mathcal{J}_q(\theta, 1) - \mathcal{J}_n(\theta, 1)\| \\
 &= \|\Psi_{q+n}(\theta, 1)\| = \|\mathcal{J}_q(\theta, 1) - \mathcal{J}_{q-1}(\theta, 1) + (\mathcal{J}_{q-1}(\theta, 1) - \mathcal{J}_{q-2}(\theta, 1)) \\
 &\quad + (\mathcal{J}_{q-2}(\theta, 1) - \mathcal{J}_{q-3}(\theta, 1)) + \dots + (\mathcal{J}_{n+1}(\theta, 1) - \mathcal{J}_n(\theta, 1))\| \\
 &\leq \|\mathcal{J}_q(\theta, 1) - \mathcal{J}_{q-1}(\theta, 1)\| + \|(\mathcal{J}_{q-1}(\theta, 1) - \mathcal{J}_{q-2}(\theta, 1))\| \\
 &\quad + \|(\mathcal{J}_{q-2}(\theta, 1) - \mathcal{J}_{q-3}(\theta, 1))\| + \dots + \|(\mathcal{J}_{n+1}(\theta, 1) - \mathcal{J}_n(\theta, 1))\| \\
 &\leq \wp^q \|\Psi_0(\theta, 1)\| + \wp^{q-1} \|\Psi_0(\theta, 1)\| + \dots + \wp^{q+1} \|\Psi_0(\theta, 1)\| \\
 &= \|\Psi_0(\theta, 1)\| (\wp^q + \wp^{q-1} + \wp^{q+1}) \\
 &= \|\Psi_0(\theta, 1)\| \frac{1 - \wp^{q-n}}{1 - \wp^{q+1}} \wp^{n+1}.
 \end{aligned}
 \tag{28}$$

As $0 < \wp < 1$, and $\Psi_0(\theta, 1)$ are bound, so take $\beta = 1 - \wp / (1 - \wp_{q-n}) \wp^{n+1} \|\Psi_0(\theta, 1)\|$, and we get

$$\|\Psi_{q+n}(\theta, 1)\| \leq \beta, \quad \forall q, n \in N.
 \tag{29}$$

Hence, $\{\Psi_q(\theta, 1)\}_{q=0}^{\infty}$ makes a "Cauchy sequence" in H . It proves that the sequence $\{\Psi_q(\theta, 1)\}_{q=0}^{\infty}$ is a convergent sequence with the limit $\lim_{q \rightarrow \infty} \Psi_q(\theta, 1) = \Psi(\theta, 1)$ for $\exists \Psi(\theta, 1) \in \mathcal{H}$ which complete the proof. \square

Theorem 5.2 Assume that $\sum_{h=0}^k \Psi_h(\theta, 1)$ is finite and $\Psi(\theta, 1)$ reflect the series solution. Considering $\wp > 0$ with $\|\Psi_{h+1}(\theta, 1)\| \leq \|\Psi_h(\theta, 1)\|$, the maximum absolute error is determined as

$$\|\Psi(\theta, \mathfrak{1}) - \sum_{h=0}^k \Psi_h(\theta, \mathfrak{1})\| < \frac{\wp^{k+1}}{1 - \wp} \|\Psi_0(\theta, \mathfrak{1})\|. \tag{30}$$

Proof Suppose $\sum_{h=0}^k \Psi_h(\theta, \mathfrak{1})$ is finite which implies that $\sum_{h=0}^k \Psi_h(\theta, \mathfrak{1}) < \infty$.
 Let us consider

$$\begin{aligned} \|\Psi(\theta, \mathfrak{1}) - \sum_{h=0}^k \Psi_h(\theta, \mathfrak{1})\| &= \left\| \sum_{h=k+1}^{\infty} \Psi_h(\theta, \mathfrak{1}) \right\| \\ &\leq \sum_{h=k+1}^{\infty} \|\Psi_h(\theta, \mathfrak{1})\| \\ &\leq \sum_{h=k+1}^{\infty} \wp^h \|\Psi_0(\theta, \mathfrak{1})\| \\ &\leq \wp^{k+1} (1 + \wp + \wp^2 + \dots) \|\Psi_0(\theta, \mathfrak{1})\| \\ &\leq \frac{\wp^{k+1}}{1 - \wp} \|\Psi_0(\theta, \mathfrak{1})\|. \end{aligned} \tag{31}$$

which complete the proof of theorem. □

Theorem 5.3 *The result of (17) is unique when $0 < (\varphi_1 + \varphi_2) \left(\frac{1^\wp}{\Gamma(\wp+1)}\right) < 1$.*

Proof Let $H = (C[J], \|\cdot\|)$ with the norm $\|\phi(\mathfrak{1})\| = \max_{\mathfrak{1} \in J} |\phi(\mathfrak{1})|$ is Banach space, \forall continuous function on J . Let $I : H \rightarrow H$ is a non-linear mapping, where

$$\mathcal{M}_{l+1}^C = \mathcal{M}_0^C + \mathbb{E}^{-1} [u^\wp \mathbb{E} [\mathcal{F}_1(\mathcal{M}_l(\theta, \mathfrak{1})) + \mathcal{G}_1(\mathcal{M}_l(\theta, \mathfrak{1}))]], \quad l \geq 0.$$

Suppose that $|\mathcal{F}_1(\mathcal{M}) - \mathcal{F}_1(\mathcal{M}^*)| < \varphi_1 |\mathcal{M} - \mathcal{M}^*|$ and $|\mathcal{G}_1(\mathcal{M}) - \mathcal{G}_1(\mathcal{M}^*)| < \varphi_2 |\mathcal{M} - \mathcal{M}^*|$, where $\mathcal{M} := \mathcal{M}(\theta, \mathfrak{1})$ and $\mathcal{M}^* := \mathcal{M}^*(\theta, \mathfrak{1})$ are two separate function values and φ_1, φ_2 are Lipschitz constants.

$$\begin{aligned} \|I\mathcal{M} - I\mathcal{M}^*\| &\leq \max_{\mathfrak{1} \in J} |\mathbb{E}^{-1} [u^\wp \mathbb{E} [\mathcal{F}_1(\mathcal{M}) - \mathcal{F}_1(\mathcal{M}^*)] \\ &\quad + u^\wp \mathbb{E} [\mathcal{G}_1(\mathcal{M}) - \mathcal{G}_1(\mathcal{M}^*)]]| \\ &\leq \max_{\mathfrak{1} \in J} \left[\varphi_1 \mathbb{E}^{-1} [u^\wp \mathbb{E} [|\mathcal{M} - \mathcal{M}^*|]] \right. \\ &\quad \left. + \varphi_2 \mathbb{E}^{-1} [u^\wp \mathbb{E} [|\mathcal{M} - \mathcal{M}^*|]] \right] \\ &\leq \max_{\mathfrak{1} \in J} (\varphi_1 + \varphi_2) \left[\mathbb{E}^{-1} [u^\wp \mathbb{E} [|\mathcal{M} - \mathcal{M}^*|]] \right] \\ &\leq (\varphi_1 + \varphi_2) \left[\mathbb{E}^{-1} [u^\wp \mathbb{E} [|\mathcal{M} - \mathcal{M}^*|]] \right] \\ &= (\varphi_1 + \varphi_2) \left(\frac{1^\wp}{\Gamma(\wp + 1)} \right) \|\mathcal{M} - \mathcal{M}^*\| \end{aligned} \tag{32}$$

It is contraction as $0 < (\varphi_1 + \varphi_2) \left(\frac{1^\wp}{\Gamma(\wp+1)}\right) < 1$. The result of (17) is unique by means of Banach fixed point theorem. \square

Theorem 5.4 *The result of (17) is convergent.*

Proof Let $\mathcal{M}_m = \sum_{r=0}^m \mathcal{M}_r(\theta, 1)$. To show that \mathcal{M}_m is a Cauchy sequence in H. Let

$$\begin{aligned} \|\mathcal{M}_m - \mathcal{M}_n\| &= \max_{i \in J} \left| \sum_{r=n+1}^m \mathcal{M}_r \right|, \quad n = 1, 2, 3, \dots \\ &\leq \max_{i \in J} \left| \mathbb{E}^{-1} \left[u^\wp \mathbb{E} \left[\sum_{r=n+1}^m (\mathcal{F}_1(\mathcal{M}_{r-1}) + \mathcal{G}_1(\mathcal{M}_{r-1})) \right] \right] \right| \\ &= \max_{i \in J} \left| \mathbb{E}^{-1} \left[u^\wp \mathbb{E} \left[\sum_{r=n+1}^{m-1} (\mathcal{F}_1(\mathcal{M}_r) + \mathcal{G}_1(\mathcal{M}_r)) \right] \right] \right| \tag{33} \\ &\leq \max_{i \in J} |\mathbb{E}^{-1}[u^\wp \mathbb{E}[(\mathcal{F}_1(\mathcal{M}_{m-1}) - \mathcal{F}_1(\mathcal{M}_{n-1}) + \mathcal{G}_1(\mathcal{M}_{m-1}) - \mathcal{G}_1(\mathcal{M}_{n-1}))]]| \\ &\leq \varphi_1 \max_{i \in J} |\mathbb{E}^{-1}[u^\wp \mathbb{E}[(\mathcal{F}_1(\mathcal{M}_{m-1}) - \mathcal{F}_1(\mathcal{M}_{n-1}))]]| \\ &\quad + \varphi_2 \max_{i \in J} |\mathbb{E}^{-1}[u^\wp \mathbb{E}[(\mathcal{G}_1(\mathcal{M}_{m-1}) - \mathcal{G}_1(\mathcal{M}_{n-1}))]]| \\ &= (\varphi_1 + \varphi_2) \left(\frac{1^\wp}{\Gamma(\wp + 1)}\right) \|\mathcal{M}_{m-1} - \mathcal{M}_{n-1}\| \end{aligned}$$

Let $m = n + 1$, then

$$\begin{aligned} \|\mathcal{M}_{n+1} - \mathcal{M}_n\| &\leq \varphi \|\mathcal{M}_n - \mathcal{M}_{n-1}\| \leq \varphi^2 \|\mathcal{M}_{n-1} - \mathcal{M}_{n-2}\| \leq \dots \leq \varphi^n \|\mathcal{M}_1 - \mathcal{M}_0\|, \tag{34} \end{aligned}$$

where $\varphi = (\varphi_1 + \varphi_2) \left(\frac{1^\wp}{\Gamma(\wp+1)}\right)$. Similarly, we have

$$\begin{aligned} \|\mathcal{M}_m - \mathcal{M}_n\| &\leq \|\mathcal{M}_{n+1} - \mathcal{M}_n\| + \|\mathcal{M}_{n+2} - \mathcal{M}_{n+1}\| + \dots + \|\mathcal{M}_m - \mathcal{M}_{m-1}\|, \\ &(\varphi^n + \varphi^{n+1} + \dots + \varphi^{m-1}) \|\mathcal{M}_1 - \mathcal{M}_0\| \leq \varphi^n \left(\frac{1 - \varphi^{m-n}}{1 - \varphi}\right) \|\mathcal{M}_1\|, \tag{35} \end{aligned}$$

As $0 < \varphi < 1$, we get $1 - \varphi^{m-n} < 1$. Hence,

$$\|\mathcal{M}_m - \mathcal{M}_n\| \leq \frac{\varphi^n}{1 - \varphi} \max_{i \in J} \|\mathcal{M}_1\|. \tag{36}$$

Since $\|\mathcal{M}_1\| < \infty$, $\|\mathcal{M}_m - \mathcal{M}_n\| \rightarrow 0$ when $n \rightarrow \infty$. Hence, \mathcal{M}_m is a Cauchy sequence in H, illustrating that the series \mathcal{M}_m is convergent. \square

6 Applications

Example 6.1 Assume the seventh-order TFLK-dV equation:

$$\begin{aligned}
 D_1^\wp \mathcal{M}(\theta, 1) = & -140\mathcal{M}^3(\theta, 1)\mathcal{M}_\theta(\theta, 1) - 70\mathcal{M}_\theta^3(\theta, 1) \\
 & - 280\mathcal{M}(\theta, 1)\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) - 70\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\
 & - 70\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 42\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \quad (37) \\
 & - 14\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1), \\
 & 0 < \wp \leq 1,
 \end{aligned}$$

with initial condition

$$\mathcal{M}(\theta, 0) = 2\rho^2 \operatorname{sech}^2(\rho\theta).$$

Apply the ET, we get

$$\begin{aligned}
 \mathbb{E} \left(\frac{\partial^\wp \mathcal{M}}{\partial 1^\wp} \right) = & \mathbb{E} \left[- 140\mathcal{M}^3(\theta, 1)\mathcal{M}_\theta(\theta, 1) - 70\mathcal{M}_\theta^3(\theta, 1) \right. \\
 & - 280\mathcal{M}(\theta, 1)\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) - 70\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\
 & - 70\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 42\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \quad (38) \\
 & \left. - 14\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right].
 \end{aligned}$$

After, we get

$$\begin{aligned}
 \frac{1}{u^\wp} \{M(u) - u^2\mathcal{M}(\theta, 0)\} = & \mathbb{E} \left[- 140\mathcal{M}^3(\theta, 1)\mathcal{M}_\theta(\theta, 1) - 70\mathcal{M}_\theta^3(\theta, 1) \right. \\
 & - 280\mathcal{M}(\theta, 1)\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) \\
 & - 70\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\
 & - 70\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 42\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\
 & - 14\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) \\
 & \left. - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right], \quad (39)
 \end{aligned}$$

$$\begin{aligned}
 M(u) = & u\mathcal{M}(\theta, 0) + u^\wp \mathbb{E} \left[- 140\mathcal{M}^3(\theta, 1)\mathcal{M}_\theta(\theta, 1) - 70\mathcal{M}_\theta^3(\theta, 1) \right. \\
 & - 280\mathcal{M}(\theta, 1)\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) - 70\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\
 & - 70\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 42\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\
 & - 14\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) \\
 & \left. - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right]. \quad (40)
 \end{aligned}$$

Apply the inverse ET, we get

$$\begin{aligned}
 \mathcal{M}(\theta, 1) &= \mathcal{M}(\theta, 0) + \mathbb{E}^{-1} \left[u^{\wp} \left\{ \mathbb{E} \left[-140\mathcal{M}^3(\theta, 1)\mathcal{M}_{\theta}(\theta, 1) - 70\mathcal{M}_{\theta}^3(\theta, 1) \right. \right. \right. \\
 &\quad - 280\mathcal{M}(\theta, 1)\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) \\
 &\quad - 70\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 70\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\
 &\quad - 42\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\
 &\quad - 14\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) \\
 &\quad \left. \left. \left. - \mathcal{M}_{\theta\theta\theta\theta\theta\theta\theta}(\theta, 1) \right] \right\} \right], \\
 \mathcal{M}(\theta, 1) &= 2\rho^2 \operatorname{sech}^2(\rho\theta) + \mathbb{E}^{-1} \left[u^{\wp} \left\{ \mathbb{E} \left[-140\mathcal{M}^3(\theta, 1)\mathcal{M}_{\theta}(\theta, 1) - 70\mathcal{M}_{\theta}^3(\theta, 1) \right. \right. \right. \\
 &\quad - 280\mathcal{M}(\theta, 1)\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) \\
 &\quad - 70\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 70\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\
 &\quad - 42\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\
 &\quad - 14\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) \\
 &\quad \left. \left. \left. - \mathcal{M}_{\theta\theta\theta\theta\theta\theta\theta}(\theta, 1) \right] \right\} \right]. \tag{41}
 \end{aligned}$$

In HPM manner, the basic series form solution is as:

$$\begin{aligned}
 \sum_{k=0}^{\infty} \epsilon^k \mathcal{M}_k(\theta, 1) &= \left(2\rho^2 \operatorname{sech}^2(\rho\theta) \right) + \left(\mathbb{E}^{-1} \left[u^{\wp} \mathbb{E} \left[-140 \left(\sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{M}) \right) \right. \right. \right. \\
 &\quad - 70 \left(\sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{M}) \right) - 280 \left(\sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{M}) \right) \\
 &\quad - 70 \left(\sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{M}) \right) - 70 \left(\sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{M}) \right) \\
 &\quad - 42 \left(\sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{M}) \right) - 14 \left(\sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{M}) \right) \\
 &\quad \left. \left. \left. - \left(\sum_{k=0}^{\infty} \epsilon^k \mathcal{M}_k(\theta, 1) \right)_{\theta\theta\theta\theta\theta\theta\theta} \right] \right] \right). \tag{42}
 \end{aligned}$$

Equating the ϵ coefficient, we have

$$\begin{aligned}
 \epsilon^0 : \mathcal{M}_0(\theta, 1) &= 2\rho^2 \operatorname{sech}^2(\rho\theta), \\
 \epsilon^1 : \mathcal{M}_1(\theta, 1) &= \frac{256\rho^9 1^{\wp} \tanh(\rho\theta) \operatorname{sech}^2(\rho\theta)}{\Gamma(\wp + 1)},
 \end{aligned}$$

$$\begin{aligned} \epsilon^2 : \mathcal{M}_2(\theta, 1) &= \frac{163841^{2\wp} \rho^{16} (\cosh(2\rho\theta) - 2) \operatorname{sech}^4(\rho\theta)}{\Gamma(2\wp + 1)}, \\ &\vdots \end{aligned}$$

Hence, the series form solution of the proposed method is as

$$\begin{aligned} \mathcal{M}(\theta, 1) &= \mathcal{M}_0(\theta, 1) + \mathcal{M}_1(\theta, 1) + \mathcal{M}_2(\theta, 1) + \dots \\ \mathcal{M}(\theta, 1) &= 2\rho^2 \operatorname{sech}^2(\rho\theta) + \frac{256\rho^9 1^\wp \tanh(\rho\theta) \operatorname{sech}^2(\rho\theta)}{\Gamma(\wp + 1)} \\ &\quad + \frac{163841^{2\wp} \rho^{16} (\cosh(2\rho\theta) - 2) \operatorname{sech}^4(\rho\theta)}{\Gamma(2\wp + 1)} + \dots \end{aligned}$$

Solution by means of ETDM

Apply the ET, we get

$$\begin{aligned} \mathbb{E} \left\{ \frac{\partial^\wp \mathcal{M}}{\partial 1^\wp} \right\} &= \mathbb{E} \left[-140\mathcal{M}^3(\theta, 1)\mathcal{M}_\theta(\theta, 1) - 70\mathcal{M}_\theta^3(\theta, 1) \right. \\ &\quad - 280\mathcal{M}(\theta, 1)\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) - 70\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\ &\quad - 70\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 42\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\ &\quad \left. - 14\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right]. \end{aligned} \tag{43}$$

After, we get

$$\begin{aligned} \frac{1}{u^\wp} \{M(u) - u^2 \mathcal{M}(\theta, 0)\} &= \mathbb{E} \left[-140\mathcal{M}^3(\theta, 1)\mathcal{M}_\theta(\theta, 1) - 70\mathcal{M}_\theta^3(\theta, 1) \right. \\ &\quad - 280\mathcal{M}(\theta, 1)\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) \\ &\quad - 70\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 70\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\ &\quad - 42\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\ &\quad \left. - 14\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right], \end{aligned} \tag{44}$$

$$\begin{aligned} M(u) &= u^2 \mathcal{M}(\theta, 0) + u^\wp \mathbb{E} \left[-140\mathcal{M}^3(\theta, 1)\mathcal{M}_\theta(\theta, 1) \right. \\ &\quad - 70\mathcal{M}_\theta^3(\theta, 1) - 280\mathcal{M}(\theta, 1)\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) \\ &\quad - 70\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 70\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\ &\quad - 42\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) - 14\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\ &\quad \left. - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right]. \end{aligned} \tag{45}$$

Apply the inverse ET, we get

$$\begin{aligned} \mathcal{M}(\theta, 1) = & \mathcal{M}(\theta, 0) + \mathbb{E}^{-1} \left[u^\rho \left\{ \mathbb{E} \left[-140\mathcal{M}^3(\theta, 1)\mathcal{M}_\theta(\theta, 1) - 70\mathcal{M}_\theta^3(\theta, 1) \right. \right. \right. \\ & - 280\mathcal{M}(\theta, 1)\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) \\ & - 70\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 70\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\ & - 42\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\ & \left. \left. \left. - 14\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right] \right\} \right], \end{aligned} \tag{46}$$

$$\begin{aligned} \mathcal{M}(\theta, 1) = & 2\rho^2 \operatorname{sech}^2(\rho\theta)) + \mathbb{E}^{-1} \left[u^\rho \left\{ \mathbb{E} \left[-140\mathcal{M}^3(\theta, 1)\mathcal{M}_\theta(\theta, 1) \right. \right. \right. \\ & - 70\mathcal{M}_\theta^3(\theta, 1) - 280\mathcal{M}(\theta, 1)\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) \\ & - 70\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 70\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\ & - 42\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\ & \left. \left. \left. - 14\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right] \right\} \right]. \end{aligned}$$

In ADM manner, the basic series form solution is as:

$$\mathcal{M}(\theta, 1) = \sum_{m=0}^{\infty} \mathcal{M}_m(\theta, 1). \tag{47}$$

The nonlinear terms are taken as $\mathcal{M}^3(\theta, 1)\mathcal{M}_\theta(\theta, 1) = \sum_{m=0}^{\infty} \mathbb{A}_m$, $\mathcal{M}_\theta^3(\theta, 1) = \sum_{m=0}^{\infty} \mathbb{B}_m$, $\mathcal{M}(\theta, 1)\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) = \sum_{m=0}^{\infty} \mathbb{C}_m$, $\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) = \sum_{m=0}^{\infty} \mathbb{D}_m$, $\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) = \sum_{m=0}^{\infty} \mathbb{E}_m$, $\mathcal{M}_\theta(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) = \sum_{m=0}^{\infty} \mathbb{F}_m$, $\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) = \sum_{m=0}^{\infty} \mathbb{G}_m$. Thus, we get

$$\begin{aligned} \sum_{m=0}^{\infty} \mathcal{M}_m(\theta, 1) = & \mathcal{M}(\theta, 0) - \mathbb{E}^{-1} \left[u^\rho \left\{ \mathbb{E} \left[-140 \sum_{m=0}^{\infty} \mathbb{A}_m - 70 \sum_{m=0}^{\infty} \mathbb{B}_m \right. \right. \right. \\ & - 280 \sum_{m=0}^{\infty} \mathbb{C}_m - 70 \sum_{m=0}^{\infty} \mathbb{D}_m - 70 \sum_{m=0}^{\infty} \mathbb{E}_m - 42 \sum_{m=0}^{\infty} \mathbb{F}_m \\ & \left. \left. \left. - 14 \sum_{m=0}^{\infty} \mathbb{G}_m - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right] \right\} \right], \end{aligned}$$

$$\begin{aligned} \sum_{m=0}^{\infty} \mathcal{M}_m(\theta, 1) = & 2\rho^2 \operatorname{sech}^2(\rho\theta)) - \mathbb{E}^{-1} \left[u^\rho \left\{ \mathbb{E} \left[-140 \sum_{m=0}^{\infty} \mathbb{A}_m \right. \right. \right. \\ & \left. \left. - 70 \sum_{m=0}^{\infty} \mathbb{B}_m - 280 \sum_{m=0}^{\infty} \mathbb{C}_m \right] \right] \end{aligned}$$

$$\begin{aligned}
 & - 70 \sum_{m=0}^{\infty} \mathbb{D}_m - 70 \sum_{m=0}^{\infty} \mathbb{E}_m - 42 \sum_{m=0}^{\infty} \mathbb{F}_m \\
 & - 14 \sum_{m=0}^{\infty} \left[\mathbb{G}_m - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right] \Bigg\}. \tag{48}
 \end{aligned}$$

By equating both sides, we have

$$\mathcal{M}_0(\theta, 1) = 2\rho^2 \operatorname{sech}^2(\rho\theta).$$

On $m = 0$,

$$\mathcal{M}_1(\theta, 1) = \frac{256\rho^9 1^{\wp} \tanh(\rho\theta) \operatorname{sech}^2(\rho\theta)}{\Gamma(\wp + 1)}.$$

On $m = 1$,

$$\mathcal{M}_2(\theta, 1) = \frac{16384 1^{2\wp} \rho^{16} (\cosh(2\rho\theta) - 2) \operatorname{sech}^4(\rho\theta)}{\Gamma(2\wp + 1)}.$$

Hence, the series form solution of the proposed method is as

$$\mathcal{M}(\theta, 1) = \sum_{m=0}^{\infty} \mathcal{M}_m(\theta, 1) = \mathcal{M}_0(\theta, 1) + \mathcal{M}_1(\theta, 1) + \mathcal{M}_2(\theta, 1) + \dots$$

$$\begin{aligned}
 \mathcal{M}(\theta, 1) &= 2\rho^2 \operatorname{sech}^2(\rho\theta) + \frac{256\rho^9 1^{\wp} \tanh(\rho\theta) \operatorname{sech}^2(\rho\theta)}{\Gamma(\wp + 1)} \\
 &+ \frac{16384 1^{2\wp} \rho^{16} (\cosh(2\rho\theta) - 2) \operatorname{sech}^4(\rho\theta)}{\Gamma(2\wp + 1)} + \dots
 \end{aligned}$$

By choosing $\wp = 1$ we get

$$\mathcal{M}(\theta, 1) = 2\rho^2 \operatorname{sech}^2(\rho(\theta - 64\rho^6 1)). \tag{49}$$

Example 6.2 Let's suppose the seventh-order TK-K equation:

$$\begin{aligned}
 D_1^{\wp} \mathcal{M}(\theta, 1) &= -2016\mathcal{M}^3(\theta, 1)\mathcal{M}_{\theta}(\theta, 1) - 630\mathcal{M}_{\theta}^3(\theta, 1) \\
 &- 2268\mathcal{M}(\theta, 1)\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) - 504\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\
 &- 252\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 147\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\
 &- 42\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1), \quad 0 < \wp \leq 1,
 \end{aligned} \tag{50}$$

having initial source

$$\mathcal{M}(\theta, 0) = \frac{\rho^2}{3} \left(1 - \frac{3}{2} \tanh^2(\rho\theta) \right).$$

After, we have

$$\begin{aligned} \mathbb{E} \left(\frac{\partial^{\wp} \mathcal{M}}{\partial 1^{\wp}} \right) = \mathbb{E} \left[\right. & - 2016\mathcal{M}^3(\theta, 1)\mathcal{M}_{\theta}(\theta, 1) - 630\mathcal{M}_{\theta}^3(\theta, 1) \\ & - 2268\mathcal{M}(\theta, 1)\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) - 504\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\ & - 252\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 147\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\ & \left. - 42\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right]. \end{aligned} \tag{51}$$

After, we obtain

$$\begin{aligned} \frac{1}{u^{\wp}} \{M(u) - u^2 \mathcal{M}(\theta, 0)\} = \mathbb{E} \left[\right. & - 2016\mathcal{M}^3(\theta, 1)\mathcal{M}_{\theta}(\theta, 1) - 630\mathcal{M}_{\theta}^3(\theta, 1) \\ & - 2268\mathcal{M}(\theta, 1)\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) \\ & - 504\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 252\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\ & - 147\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\ & \left. - 42\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right], \end{aligned} \tag{52}$$

$$\begin{aligned} M(u) = u\mathcal{M}(\theta, 0) + u^{\wp} \mathbb{E} \left[\right. & - 2016\mathcal{M}^3(\theta, 1)\mathcal{M}_{\theta}(\theta, 1) - 630\mathcal{M}_{\theta}^3(\theta, 1) \\ & - 2268\mathcal{M}(\theta, 1)\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) - 504\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\ & - 252\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 147\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\ & \left. - 42\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right]. \end{aligned} \tag{53}$$

Apply the inverse ET, we get

$$\begin{aligned} \mathcal{M}(\theta, 1) = \mathcal{M}(\theta, 0) + \mathbb{E}^{-1} \left[\right. & u^{\wp} \left\{ \mathbb{E} \left[\right. \right. \\ & - 2016\mathcal{M}^3(\theta, 1)\mathcal{M}_{\theta}(\theta, 1) - 630\mathcal{M}_{\theta}^3(\theta, 1) \\ & - 2268\mathcal{M}(\theta, 1)\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) - 504\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\ & - 252\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 147\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\ & \left. \left. - 42\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right] \right\} \left. \right], \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}(\theta, 1) = & \frac{\rho^2}{3} \left(1 - \frac{3}{2} \tanh^2(\rho\theta)\right) + \mathbb{E}^{-1} \left[u^\wp \left\{ \mathbb{E} \left[-2016 \mathcal{M}^3(\theta, 1) \mathcal{M}_\theta(\theta, 1) \right. \right. \right. \\
 & - 630 \mathcal{M}_\theta^3(\theta, 1) - 2268 \mathcal{M}(\theta, 1) \mathcal{M}_\theta(\theta, 1) \mathcal{M}_{\theta\theta}(\theta, 1) \\
 & - 504 \mathcal{M}^2(\theta, 1) \mathcal{M}_{\theta\theta\theta}(\theta, 1) - 252 \mathcal{M}_{\theta\theta}(\theta, 1) \mathcal{M}_{\theta\theta\theta}(\theta, 1) \\
 & - 147 \mathcal{M}_\theta(\theta, 1) \mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) - 42 \mathcal{M}(\theta, 1) \\
 & \left. \left. \left. \mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right] \right\} \right]. \tag{54}
 \end{aligned}$$

In terms of HPM, the basic series form solution is as:

$$\begin{aligned}
 \sum_{k=0}^{\infty} \epsilon^k \mathcal{M}_k(\theta, 1) = & \left(\frac{\rho^2}{3} \left(1 - \frac{3}{2} \tanh^2(\rho\theta)\right) \right) \\
 & + \left(\mathbb{E}^{-1} \left[u^\wp \mathbb{E} \left[-2016 \left(\sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{M}) \right) - 630 \left(\sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{M}) \right) \right. \right. \right. \\
 & - 2268 \left(\sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{M}) \right) - 504 \left(\sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{M}) \right) \\
 & - 252 \left(\sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{M}) \right) - 147 \left(\sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{M}) \right) \\
 & \left. \left. \left. - 42 \left(\sum_{k=0}^{\infty} \epsilon^k H_k(\mathcal{M}) \right) - \left(\sum_{k=0}^{\infty} \epsilon^k \mathcal{M}_k(\theta, 1) \right)_{\theta\theta\theta\theta\theta\theta} \right] \right] \right). \tag{55}
 \end{aligned}$$

Equating the ϵ coefficient, we have

$$\begin{aligned}
 \epsilon^0 : \mathcal{M}_0(\theta, 1) &= \frac{\rho^2}{3} \left(1 - \frac{3}{2} \tanh^2(\rho\theta)\right), \\
 \epsilon^1 : \mathcal{M}_1(\theta, 1) &= -\frac{4\rho^9 1^\wp \tanh(\rho\theta) \operatorname{sech}^2(\rho\theta)}{3\Gamma(\wp + 1)}, \\
 \epsilon^2 : \mathcal{M}_2(\theta, 1) &= \frac{16\rho^{16} 1^{2\wp} (\cosh(2\rho\theta) - 2) \operatorname{sech}^4(\rho\theta)}{9\Gamma(2\wp + 1)}, \\
 &\vdots
 \end{aligned}$$

Hence, the series form solution of the proposed method is as

$$\begin{aligned}
 \mathcal{M}(\theta, 1) &= \mathcal{M}_0(\theta, 1) + \mathcal{M}_1(\theta, 1) + \mathcal{M}_2(\theta, 1) + \dots \\
 \mathcal{M}(\theta, 1) &= \frac{\rho^2}{3} \left(1 - \frac{3}{2} \tanh^2(\rho\theta)\right) - \frac{4\rho^9 1^\wp \tanh(\rho\theta) \operatorname{sech}^2(\rho\theta)}{3\Gamma(\wp + 1)} \\
 &\quad + \frac{16\rho^{16} 1^{2\wp} (\cosh(2\rho\theta) - 2) \operatorname{sech}^4(\rho\theta)}{9\Gamma(2\wp + 1)} + \dots
 \end{aligned}$$

Solution by means of ETDM

Apply the ET, we get

$$\begin{aligned} \mathbb{E} \left\{ \frac{\partial^{\rho} \mathcal{M}}{\partial 1^{\rho}} \right\} = \mathbb{E} & \left[-2016\mathcal{M}^3(\theta, 1)\mathcal{M}_{\theta}(\theta, 1) - 630\mathcal{M}_{\theta}^3(\theta, 1) \right. \\ & - 2268\mathcal{M}(\theta, 1)\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) - 504\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\ & - 252\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 147\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\ & \left. - 42\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right]. \end{aligned} \tag{56}$$

After, we obtain

$$\begin{aligned} \frac{1}{u^{\rho}} \{M(u) - u^2\mathcal{M}(\theta, 0)\} = \mathbb{E} & \left[-2016\mathcal{M}^3(\theta, 1)\mathcal{M}_{\theta}(\theta, 1) - 630\mathcal{M}_{\theta}^3(\theta, 1) \right. \\ & - 2268\mathcal{M}(\theta, 1)\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) \\ & - 504\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\ & - 252\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 147\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\ & \left. - 42\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right], \end{aligned} \tag{57}$$

$$\begin{aligned} M(u) = u^2\mathcal{M}(\theta, 0) + u^{\rho}\mathbb{E} & \left[-2016\mathcal{M}^3(\theta, 1)\mathcal{M}_{\theta}(\theta, 1) - 630\mathcal{M}_{\theta}^3(\theta, 1) \right. \\ & - 2268\mathcal{M}(\theta, 1)\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) - 504\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\ & - 252\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 147\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\ & \left. - 42\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right]. \end{aligned} \tag{58}$$

Apply the inverse ET, we get

$$\begin{aligned} \mathcal{M}(\theta, 1) = \mathcal{M}(\theta, 0) + \mathbb{E}^{-1} & \left[u^{\rho} \left\{ \mathbb{E} \left[-2016\mathcal{M}^3(\theta, 1)\mathcal{M}_{\theta}(\theta, 1) \right. \right. \right. \\ & - 630\mathcal{M}_{\theta}^3(\theta, 1) - 2268\mathcal{M}(\theta, 1)\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) \\ & - 504\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 252\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\ & - 147\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) - 42\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) \\ & \left. \left. \left. - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right] \right\} \right], \\ \mathcal{M}(\theta, 1) = \frac{\rho^2}{3} \left(1 - \frac{3}{2} \tanh^2(\rho\theta) \right) + \mathbb{E}^{-1} & \left[u^{\rho} \left\{ \mathbb{E} \left[-2016\mathcal{M}^3(\theta, 1)\mathcal{M}_{\theta}(\theta, 1) \right. \right. \right. \\ & \left. \left. \left. - 630\mathcal{M}_{\theta}^3(\theta, 1) - 2268\mathcal{M}(\theta, 1)\mathcal{M}_{\theta}(\theta, 1) \right] \right\} \right], \end{aligned}$$

$$\begin{aligned}
 & \mathcal{M}_{\theta\theta}(\theta, 1) - 504\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 252\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) \\
 & - 147\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) - 42\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) \\
 & - \mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) \Big] \Big]. \tag{59}
 \end{aligned}$$

In terms of ADM, the basic series form solution is as:

$$\mathcal{M}(\theta, 1) = \sum_{m=0}^{\infty} \mathcal{M}_m(\theta, 1). \tag{60}$$

Let us assume nonlinear terms by adomian polynomial as $\mathcal{M}^3(\theta, 1)\mathcal{M}_{\theta}(\theta, 1) = \sum_{m=0}^{\infty} \mathbb{A}_m$, $\mathcal{M}_{\theta}^3(\theta, 1) = \sum_{m=0}^{\infty} \mathbb{B}_m$, $\mathcal{M}(\theta, 1)\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta}(\theta, 1) = \sum_{m=0}^{\infty} \mathbb{C}_m$, $\mathcal{M}^2(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) = \sum_{m=0}^{\infty} \mathbb{D}_m$, $\mathcal{M}_{\theta\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta}(\theta, 1) = \sum_{m=0}^{\infty} \mathbb{E}_m$, $\mathcal{M}_{\theta}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta}(\theta, 1) = \sum_{m=0}^{\infty} \mathbb{F}_m$, $\mathcal{M}(\theta, 1)\mathcal{M}_{\theta\theta\theta\theta\theta}(\theta, 1) = \sum_{m=0}^{\infty} \mathbb{G}_m$. So, we get

$$\begin{aligned}
 \sum_{m=0}^{\infty} \mathcal{M}_m(\theta, 1) &= \mathcal{M}(\theta, 0) - \mathbb{E}^{-1} \left[u^{\wp} \left\{ \mathbb{E} \left[-2016 \sum_{m=0}^{\infty} \mathbb{A}_m - 630 \sum_{m=0}^{\infty} \mathbb{B}_m \right. \right. \right. \\
 & - 2268 \sum_{m=0}^{\infty} \mathbb{C}_m - 504 \sum_{m=0}^{\infty} \mathbb{D}_m - 252 \sum_{m=0}^{\infty} \mathbb{E}_m \\
 & \left. \left. \left. - 147 \sum_{m=0}^{\infty} \mathbb{F}_m - 42 \sum_{m=0}^{\infty} \mathbb{G}_m - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right] \right\} \right], \tag{61} \\
 \sum_{m=0}^{\infty} \mathcal{M}_m(\theta, 1) &= \frac{\rho^2}{3} \left(1 - \frac{3}{2} \tanh^2(\rho\theta) \right) - \mathbb{E}^{-1} \left[u^{\wp} \left\{ \mathbb{E} \left[-2016 \sum_{m=0}^{\infty} \mathbb{A}_m \right. \right. \right. \\
 & - 630 \sum_{m=0}^{\infty} \mathbb{B}_m - 2268 \sum_{m=0}^{\infty} \mathbb{C}_m - 504 \sum_{m=0}^{\infty} \mathbb{D}_m - 252 \sum_{m=0}^{\infty} \mathbb{E}_m \\
 & \left. \left. \left. - 147 \sum_{m=0}^{\infty} \mathbb{F}_m - 42 \sum_{m=0}^{\infty} \mathbb{G}_m - \mathcal{M}_{\theta\theta\theta\theta\theta\theta}(\theta, 1) \right] \right\} \right].
 \end{aligned}$$

By equating both sides, we have

$$\mathcal{M}_0(\theta, 1) = \frac{\rho^2}{3} \left(1 - \frac{3}{2} \tanh^2(\rho\theta) \right).$$

On $m = 0$,

$$\mathcal{M}_1(\theta, 1) = - \frac{4\rho^9 1^{\wp} \tanh(\rho\theta) \operatorname{sech}^2(\rho\theta)}{3\Gamma(\wp + 1)}.$$




Fig. 1 The graphical view of accurate and our approaches solution




Fig. 2 The graphical view of our approaches solution at $\varphi = 0.8, 0.6$




Fig. 3 The graphical view of our approaches solution at different orders of φ

Fig. 4 The graphical view of our approaches solution by means of absolute error




Table 1 Nature of accurate and our approaches solution for TFLK-dV equation at various orders of \wp

β	θ	$\wp = 0.7$	$\wp = 0.8$	$\wp = 0.9$	$\wp = 1(\text{approx})$	$\wp = 1(\text{exact})$
0.01	0.2	0.3195902	0.3190435	0.3185029	0.3179660	0.3179660
	0.4	0.3151229	0.3140570	0.3130030	0.3119561	0.3119561
	0.6	0.3068229	0.3052898	0.3037739	0.3022682	0.3022682
	0.8	0.2950949	0.2931661	0.2912589	0.2893645	0.2893645
	1	0.2804830	0.2782424	0.2760268	0.2738262	0.2738262
0.02	0.2	0.3196105	0.3190570	0.3185115	0.3179713	0.3179713
	0.4	0.3151626	0.3140833	0.3130198	0.3119665	0.3119665
	0.6	0.3068799	0.3053277	0.3037981	0.3022832	0.3022832
	0.8	0.2951667	0.2932137	0.2912892	0.2893833	0.2893833
	1	0.2805664	0.2782977	0.2760621	0.2738480	0.2738480
0.03	0.2	0.3196279	0.3190691	0.3185197	0.3179766	0.3179766
	0.4	0.3151964	0.3141070	0.3130357	0.3119769	0.3119769
	0.6	0.3069286	0.3053617	0.3038210	0.3022981	0.3022981
	0.8	0.2952280	0.2932565	0.2913180	0.2894020	0.2894020
	1	0.2806375	0.2783474	0.2760956	0.2738698	0.2738698
0.04	0.2	0.3196436	0.3190805	0.3185276	0.3179819	0.3179819
	0.4	0.3152270	0.3141291	0.3130511	0.3119872	0.3119872
	0.6	0.3069726	0.3053935	0.3038431	0.3023130	0.3023130
	0.8	0.2952833	0.2932966	0.2913459	0.2894208	0.2894208
	1	0.2807018	0.2783939	0.2761279	0.2738916	0.2738916
0.05	0.2	0.3196581	0.3190912	0.3185353	0.3179872	0.3179872
	0.4	0.3152553	0.3141501	0.3130661	0.3119976	0.3119976
	0.6	0.3070133	0.3054238	0.3038647	0.3023279	0.3023279
	0.8	0.2953346	0.2933346	0.2913731	0.2894395	0.2894395
	1	0.2807614	0.2784381	0.2761595	0.2739134	0.2739134

Table 2 Comparison of our approaches solution by means of absolute error at different orders of ρ

β	θ	$\rho = 0.7$	$\rho = 0.8$	$\rho = 0.9$	$\rho = 1(HPTM)$	$\rho = 1(YTDM)$
0.01	0.2	1.624198000E-04	1.077511700E-03	5.369338000E-04	3.500000000E-09	3.500000000E-09
	0.4	3.166760700E-03	2.100864300E-03	1.046877700E-03	3.000000000E-09	3.000000000E-09
	0.6	4.554641200E-03	3.021599100E-03	1.505686500E-03	2.800000000E-09	2.800000000E-09
	0.8	5.730431700E-03	3.801631100E-03	1.894382000E-03	2.300000000E-09	2.300000000E-09
	1	6.656800000E-03	4.416193700E-03	2.206622700E-03	1.700000000E-09	1.700000000E-09
	0.2	1.639227900E-04	1.085688600E-03	5.402287000E-04	1.380000000E-08	1.380000000E-08
	0.4	3.196054800E-03	2.116796900E-03	1.053291600E-03	1.270000000E-08	1.270000000E-08
	0.6	4.596768200E-03	3.044508800E-03	1.514905700E-03	1.100000000E-08	1.100000000E-08
0.03	0.8	5.783430300E-03	3.830451400E-03	1.905977600E-03	9.100000000E-09	9.100000000E-09
	1	6.718363500E-03	4.449670200E-03	2.214090100E-03	6.900000000E-09	6.900000000E-09
	0.2	1.651289300E-04	1.092525900E-03	5.430906000E-04	3.090000000E-08	3.090000000E-08
	0.4	3.219553800E-03	2.130110100E-03	1.058854000E-03	2.840000000E-08	2.840000000E-08
	0.6	4.630557300E-03	3.063648000E-03	1.522897100E-03	2.490000000E-08	2.490000000E-08
	0.8	5.825936100E-03	3.854525400E-03	1.916025900E-03	2.050000000E-08	2.050000000E-08
	1	6.767735900E-03	4.477631200E-03	2.225758100E-03	1.540000000E-08	1.540000000E-08
	0.2	1.661669700E-04	1.098570600E-03	5.456862000E-04	5.490000000E-08	5.490000000E-08
0.04	0.4	3.239768300E-03	2.141871300E-03	1.063890200E-03	5.070000000E-08	5.070000000E-08
	0.6	4.659618500E-03	3.080551100E-03	1.530128000E-03	4.430000000E-08	4.430000000E-08
	0.8	5.862491000E-03	3.875783500E-03	1.925114900E-03	3.640000000E-08	3.640000000E-08
	1	6.810193700E-03	4.502319400E-03	2.236310000E-03	2.730000000E-08	2.730000000E-08
	0.2	1.670910600E-04	1.104062700E-03	5.480917000E-04	8.580000000E-08	8.580000000E-08
	0.4	3.257754000E-03	2.152547700E-03	1.068548600E-04	7.920000000E-08	7.920000000E-08
	0.6	4.685470600E-03	3.095890500E-03	1.536811900E-04	6.930000000E-08	6.930000000E-08
	0.8	5.895005800E-03	3.895071800E-03	1.935513200E-04	5.680000000E-08	5.680000000E-08
1	6.847956500E-03	4.524717500E-03	2.246057600E-04	4.270000000E-08	4.270000000E-08	




Fig. 5 The graphical view of accurate and our approaches solution

On $m = 1$,

$$\mathcal{M}_2(\theta, 1) = \frac{16\rho^{16}1^{2\wp}(\cosh(2\rho\theta) - 2) \operatorname{sech}^4(\rho\theta)}{9\Gamma(2\wp + 1)}.$$

Hence, the series form solution of the proposed method is as

$$\mathcal{M}(\theta, 1) = \sum_{m=0}^{\infty} \mathcal{M}_m(\theta, 1) = \mathcal{M}_0(\theta, 1) + \mathcal{M}_1(\theta, 1) + \mathcal{M}_2(\theta, 1) + \dots$$

$$\begin{aligned} \mathcal{M}(\theta, 1) = & \frac{\rho^2}{3} \left(1 - \frac{3}{2} \tanh^2(\rho\theta) \right) - \frac{4\rho^9 1^{\wp} \tanh(\rho\theta) \operatorname{sech}^2(\rho\theta)}{3\Gamma(\wp + 1)} \\ & + \frac{16\rho^{16} 1^{2\wp} (\cosh(2\rho\theta) - 2) \operatorname{sech}^4(\rho\theta)}{9\Gamma(2\wp + 1)} + \dots \end{aligned}$$

By choosing $\wp = 1$ we get

$$\mathcal{M}(\theta, 1) = \frac{\rho^2}{3} \left(1 - \frac{3}{2} \tanh^2 \left(\rho \left[\theta + \frac{4\rho^6}{3} 1 \right] \right) \right). \tag{62}$$

Numerical simulation studies

In this study, the exact approximate solution of time-fractional seventh-order nonlinear equations has been studied using two novel approaches. The Caputo fractional derivative operator at any order for variable values of space and time is presented as exact analytical solutions for the time-fractional seventh-order nonlinear equations via Maple. The numerical results demonstrate the technique’s applicability, and the




Fig. 6 The graphical view of our approaches solution at $\varphi = 0.8, 0.6$





Fig. 7 The graphical view of our approaches solution at different orders of φ

Fig. 8 The graphical view of our approaches solution by means of absolute error



precision of the approach is assessed in light of the precise results. The suggested approaches solution plot of $\mathcal{M}(\theta, 1)$ is shown in Fig. 1a, while Fig. 1b shows the actual solution plot. Figure 2a,b display the fractional-order behavior of $\mathcal{M}(\theta, 1)$ for $\varphi = 0.8$ and 0.6. Figure 3a, b show the plots of $\mathcal{M}(\theta, 1)$ for the range of $\varphi = 0.25, 0.50, 0.75,$ and 1, whereas Fig.4 displays the error evaluation for the same equation generated

Table 3 Nature of the accurate and proposed approaches solution at various orders of \wp

β	θ	$\wp = 0.85$	$\wp = 0.90$	$\wp = 0.95$	$\wp = 1(\text{approx})$	$\wp = 1(\text{exact})$
0.01	0.2	0.06080016	0.06080053	0.06080090	0.06080126	0.06080127
	0.4	0.05947033	0.05947106	0.05947178	0.05947249	0.05947250
	0.6	0.05731134	0.05731240	0.05731345	0.05731449	0.05731450
	0.8	0.05440336	0.05440470	0.05440604	0.05440736	0.05440738
	1	0.05084969	0.05085128	0.05085287	0.05085443	0.05085444
0.02	0.2	0.06080014	0.06080052	0.060800894	0.06080125	0.06080126
	0.4	0.05947030	0.05947103	0.059471766	0.05947247	0.05947249
	0.6	0.05731130	0.05731236	0.057313426	0.05731445	0.05731448
	0.8	0.05440330	0.05440466	0.054406011	0.05440732	0.05440735
	1	0.05084963	0.05085123	0.050852830	0.05085438	0.05085441
0.03	0.2	0.06080013	0.06080051	0.06080088	0.06080124	0.06080125
	0.4	0.05947027	0.05947101	0.05947174	0.05947245	0.05947247
	0.6	0.05731126	0.05731233	0.05731340	0.05731442	0.05731445
	0.8	0.05440325	0.05440462	0.05440597	0.05440728	0.05440732
	1	0.05084957	0.05085118	0.05085279	0.05085433	0.05085438
0.04	0.2	0.06080012	0.06080050	0.06080087	0.06080123	0.06080124
	0.4	0.05947025	0.05947099	0.05947173	0.05947243	0.05947246
	0.6	0.05731122	0.05731230	0.05731337	0.05731439	0.05731443
	0.8	0.05440320	0.05440458	0.05440594	0.05440724	0.05440730
	1	0.05084952	0.05085114	0.05085275	0.05085429	0.05085435
0.05	0.2	0.06080011	0.06080049	0.06080086	0.06080122	0.06080124
	0.4	0.05947022	0.05947097	0.05947171	0.05947241	0.05947244
	0.6	0.05731119	0.05731227	0.05731334	0.05731436	0.05731441
	0.8	0.05440316	0.05440454	0.05440591	0.05440720	0.05440727
	1	0.05084946	0.05085109	0.05085271	0.05085424	0.05085432

by both methods. For various values of θ and \mathfrak{I} , the analytical solution to the equation $\mathcal{M}(\theta, \mathfrak{I})$ is displayed in Table 1 while the error evaluation has been evaluated in Table 2 for various values of θ and \mathfrak{I} . The proposed techniques solution plot of $\mathcal{M}(\theta, \mathfrak{I})$ is shown in Fig. 5a, while Fig. 5b shows the actual solution plot. Figure 6a, b display the graphical representations of $\mathcal{M}(\theta, \mathfrak{I})$ for $\wp = 0.8$ and 0.6 . Figure 7a, b show the plots of $\mathcal{M}(\theta, \mathfrak{I})$ for the range of $\wp = 0.25, 0.50, 0.75$, and 1 , whereas Fig. 8 displays the error evaluation for the same equation generated by both methods. For various values of θ and \mathfrak{I} , the analytical solution to the equation $\mathcal{M}(\theta, \mathfrak{I})$ is displayed in Table 3 while the error evaluation has been evaluated in Table 4 for various values of θ and \mathfrak{I} . It need to be noted that throughout the calculations, we used second-order approximations and that using accurate results to the problem gave us a better estimate. We could have obtained more accurate approximation solutions by increasing the order of the approximation, which results in more terms in the solution.

Table 4 Numerical simulation in sense of absolute error at numerous values of ρ

β	θ	$\rho = 0.85$	$\rho = 0.90$	$\rho = 0.95$	$\rho = 1(HPTM)$	$\rho = 1(YTDM)$
0.01	0.2	1.1037900000E-06	7.3351000000E-07	3.6446000000E-07	3.6500000000E-09	3.6500000000E-09
	0.4	2.1649300000E-06	1.4386700000E-06	7.1483000000E-07	7.1700000000E-09	7.1700000000E-09
	0.6	3.1442000000E-06	2.0894300000E-06	1.0381800000E-06	1.0410000000E-08	1.0410000000E-08
	0.8	4.0088700000E-06	2.6640400000E-06	1.3236800000E-06	1.3280000000E-08	1.3280000000E-08
	1	4.7347700000E-06	3.1464200000E-06	1.5633600000E-06	1.5680000000E-08	1.5680000000E-08
	0.2	1.1076800000E-06	7.3436000000E-07	3.6282000000E-07	7.3100000000E-09	7.3100000000E-09
	0.4	2.1725700000E-06	1.4403500000E-06	7.1163000000E-07	1.4330000000E-08	1.4330000000E-08
	0.6	3.1553100000E-06	2.0918800000E-06	1.0335200000E-06	2.0810000000E-08	2.0810000000E-08
0.03	0.8	4.0230300000E-06	2.6671500000E-06	1.3177400000E-06	2.6540000000E-08	2.6540000000E-08
	1	4.7514900000E-06	3.1501000000E-06	1.5563500000E-06	3.1350000000E-08	3.1350000000E-08
	0.2	1.1104600000E-06	7.3461000000E-07	3.6094000000E-07	1.0970000000E-08	1.0970000000E-08
	0.4	2.1780000000E-06	1.4408500000E-06	7.0795000000E-07	2.1500000000E-08	2.1500000000E-08
	0.6	3.1631900000E-06	2.0925800000E-06	1.0281600000E-06	3.1240000000E-08	3.1240000000E-08
	0.8	4.0330900000E-06	2.6680700000E-06	1.3109300000E-06	3.9810000000E-08	3.9810000000E-08
	1	4.7633600000E-06	3.1511700000E-06	1.5482900000E-06	4.7030000000E-08	4.7030000000E-08
	0.2	1.1125400000E-06	7.3449000000E-07	3.5892000000E-07	1.4620000000E-08	1.4620000000E-08
0.04	0.4	2.1821000000E-06	1.4406100000E-06	7.0397000000E-07	2.8660000000E-08	2.8660000000E-08
	0.6	3.1691400000E-06	2.0922400000E-06	1.0223900000E-06	4.1640000000E-08	4.1640000000E-08
	0.8	4.0406800000E-06	2.6676200000E-06	1.3035600000E-06	5.3080000000E-08	5.3080000000E-08
	1	4.7723200000E-06	3.1506400000E-06	1.5395900000E-06	6.2700000000E-08	6.2700000000E-08
	0.2	1.1141500000E-06	7.3410000000E-07	3.5678000000E-07	1.8270000000E-08	1.8270000000E-08
	0.4	2.1852400000E-06	1.4398300000E-06	6.9977000000E-07	3.5830000000E-08	3.5830000000E-08
	0.6	3.1736900000E-06	2.0911100000E-06	1.0162800000E-06	5.2050000000E-08	5.2050000000E-08
	0.8	4.0464900000E-06	2.6661900000E-06	1.2957800000E-06	6.6360000000E-08	6.6360000000E-08
1	4.7791800000E-06	3.1489500000E-06	1.5303900000E-06	7.8380000000E-08	7.8380000000E-08	

7 Conclusion

The ETDM and the HPTM are two unique methodologies that have been thoroughly examined in this work for solving non-linear fractional seventh-order Lax's Korteweg–de Vries and Kaup–Kupershmidt equations. The methods that are proposed are the combined form of the Elzaki transformation with the homotopy perturbation method and the Adomian decomposition approach. The fractional-order solutions give different dynamics for different fractional orders of the derivative. In comparison to numerical studies, which require more complex computations, the task can be completed quite simply and effectively using analytical solutions. After all, the researchers can now choose the fractional-order issue whose solution is comparable and extremely close to the experimental results of any physical problem. The graphical analysis of the revealed solutions was executed. The study's findings showed that the precise solutions offered and those actually found were very congruent. As the problems fractional orders change, different dynamical patterns emerge in the solutions, which are generated for various fractional orders. The tables show the applicability of the suggested methods by offering a variety of fractional-order results. The existing approaches have shown to be an efficient and straightforward process when compared to the precise solution. Finally, this research leads us to the conclusion that the suggested approaches are strong and useful mathematical tool for examining a variety of real problems that arise in the natural sciences and engineering and that may be represented by fractional differential equations.

Funding This research received no external funding.

Data availability The numerical data used to support the findings of this study are included within the article.

Declarations

Conflict of interest The authors declare that they have no competing interests.

References

1. Ismael HF, Bulut H, Baskonus HM (2021) W shaped surfaces to the nematic liquid crystals with three nonlinearity laws. *Soft Comput* 25:4513–4524
2. Yavuz M, Sulaiman TA, Yusuf A, Abdeljawad T (2021) The Schrödinger–KdV equation of fractional order with Mittag–Leffler nonsingular kernel. *Alex Eng J* 60(2):2715–2724
3. Samko SG, Kilbas AA, Marichev OI (1993) *Fractional integrals and derivatives*, vol 1. Gordon and Breach Science Publishers, Yverdon
4. Akinyemi L, Veerasha P, Şenol M, Rezazadeh H (2022) An efficient technique for generalized conformable Pochhammer–Chree models of longitudinal wave propagation of elastic rod. *Indian J Phys* 96(14):4209–4218
5. Alyobi S, Shah R, Khan A, Shah NA, Nonlaopon K (2022) Fractional analysis of nonlinear Boussinesq equation under Atangana–Baleanu–Caputo operator. *Symmetry* 14(11):2417
6. Caputo M (1967) Linear models of dissipation whose Q is almost frequency independent, part II. *Geophys J Int* 13:529–539
7. Carrier GF, Pearson CE (1988) *Partial differential equations, theory and technique*, 2nd edn. Academic Press, Boston

8. Koksai ME, Senol M, Unver AK (2019) Numerical simulation of power transmission lines. *Chin J Phys* 59:507–524
9. Halmos PR (1998) Introduction to Hilbert space and the theory of spectral multiplicity. American Mathematical Society–Chelsea Publications, Providence
10. Areshi M, Khan A, Shah R, Nonlaopon K (2022) Analytical investigation of fractional-order Newell–Whitehead–Segel equations via a novel transform. *Aims Math* 7(4):6936–6958
11. Shah NA, Hamed YS, Abualnaja KM, Chung JD, Shah R, Khan A (2022) A comparative analysis of fractional-order kaup–kupersmidt equation within different operators. *Symmetry* 14(5):986
12. Sunthrayuth P, Alyousef HA, El-Tantawy SA, Khan A, Wyal N (2022) Solving fractional-order diffusion equations in a plasma and fluids via a novel transform. *J Funct Spaces*
13. Alderremy AA, Aly S, Fayyaz R, Khan A, Shah R, Wyal N (2022) The analysis of fractional-order nonlinear systems of third order KdV and Burgers equations via a novel transform. *Complexity*
14. Shah NA, El-Zahar ER, Akgül A, Khan A, Kafle J (2022) Analysis of fractional-order regularized long-wave models via a novel transform. *J Funct Spaces*
15. Botmart T, Agarwal RP, Naeem M, Khan A, Shah R (2022) On the solution of fractional modified Boussinesq and approximate long wave equations with non-singular kernel operators. *AIMS Math* 7:12483–12513
16. Senol M, Tasbozan O, Kurt A (2021) Comparison of two reliable methods to solve fractional Rosenau–Hyman equation. *Math Methods Appl Sci* 44(10):7904–7914
17. Cresson J (2007) Fractional embedding of differential operators and Lagrangian systems. *J Math Phys* 48(3):033504
18. Katugampola UN (2011) New approach to a generalized fractional integral. *Appl Math Comput* 218(3):860–865
19. Kilbas AA, Saigo M, Saxena RK (2004) Generalized Mittag–Leffler function and generalized fractional calculus operators. *Integral Transform Spec Funct* 15(1):31–49
20. Klimek M (2005) Lagrangian fractional mechanics a noncommutative approach. *Czechoslovak J Phys* 55(11):1447–1453
21. Klimek M (2009) On solutions of linear fractional differential equations of a variational type. Publishing Office of Czestochowa University of Technology, Czestochowa
22. Li ZB, He JH (2010) Fractional complex transform for fractional differential equations. *Math Comput Appl* 15(5):970–973
23. Li C, Zeng F (2012) Finite difference methods for fractional differential equations. *Int J Bifurc Chaos* 22(04):1230014
24. Momani S, Odibat Z (2006) Analytical solution of a time-fractional Navier–Stokes equation by Adomian decomposition method. *Appl Math Comput* 177(2):488–494
25. Syam MI (2017) A numerical solution of fractional Lienard’s equation by using the residual power series method. *Mathematics* 6(1):1
26. Odibat Z, Momani S, Xu H (2010) A reliable algorithm of homotopy analysis method for solving nonlinear fractional differential equations. *Appl Math Modell* 34(3):593–600
27. Arikoglu A, Ozkol I (2007) Solution of fractional differential equations by using differential transform method. *Chaos Solitons Fractals* 34(5):1473–1481
28. Wu GC (2011) A fractional variational iteration method for solving fractional nonlinear differential equations. *Comput Math Appl* 61(8):2186–2190
29. Diethelm K, Ford NJ, Freed AD (2002) A predictor-corrector approach for the numerical solution of fractional differential equations. *Nonlinear Dyn* 29:3–22
30. Adomian G (1994) Solution of physical problems by decomposition. *Comput Math Appl* 27(9–10):145–154
31. Adomian G (1988) A review of the decomposition method in applied mathematics. *J Math Anal Appl* 135:501544
32. He JH (1999) Homotopy perturbation technique. *Comput Methods Appl Mech Eng* 178:257–262
33. He JH (2000) A coupling method of a homotopy technique and a perturbation technique for non-linear problems. *Int J Non-Linear Mech* 35:37–43
34. Korteweg DJ, De Vries G (1895) XLI. On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves. *Lond Edinb Dublin Philos Mag J Sci* 39(240):422–443
35. Fung MK (1997) KdV equation as an Euler–Poincaré equation. *Chin J Phys* 35(6S):789–796

36. El-Wakil SA, Abulwafa EM, Zahran MA, Mahmoud AA (2011) Time-fractional KdV equation: formulation and solution using variational methods. *Nonlinear Dyn* 65:55–63
37. Salas AH, Gómez SCA (2010) Application of the Cole–Hopf transformation for finding exact solutions to several forms of the seventh-order KdV equation. *Math Prob Eng*
38. Soliman AA (2006) A numerical simulation and explicit solutions of KdV–Burgers and Lax’s seventh-order KdV equations. *Chaos Solitons Fractals* 29(2):294–302
39. Akinyemi L (2019) q-Homotopy analysis method for solving the seventh-order time-fractional Lax’s Korteweg–de Vries and Sawada–Kotera equations. *ComputdAppl Math* 38(4):191
40. Senol M, Tasbozan O, Kurt A (2021) Comparison of two reliable methods to solve fractional Rosenau–Hyman equation. *Math Methods Appl Sci* 44(10):7904–7914
41. Pomeau Y, Ramani A, Grammaticos B (1988) Structural stability of the Korteweg–de Vries solitons under a singular perturbation. *Physica D* 31(1):127–134
42. Elzaki TM (2011) The new integral transform ‘Elzaki transform’. *Glob J Pure Appl Math* 7:57–64
43. Alshikh AA, Mahgob MMA (2016) A comparative study between laplace transform and two new integrals “ELzaki” transform and “Aboodh” transform. *Pure Appl Math J* 5:145
44. Elzaki T, Alkhateeb S (2015) Modification of Sumudu transform “Elzaki transform” and adomian decomposition method. *Appl Math Sci* 9:603–611

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.