

O 051 – Does video game-based balance-training improve gait stability in children with cerebral palsy?

Meyns, P.; Bras, C.; Harlaar, J.; van de Pol, L.; Barkhof, F.; Buizer, A.

DOI

[10.1016/j.gaitpost.2018.06.078](https://doi.org/10.1016/j.gaitpost.2018.06.078)

Publication date

2018

Document Version

Final published version

Published in

Gait and Posture

Citation (APA)

Meyns, P., Bras, C., Harlaar, J., van de Pol, L., Barkhof, F., & Buizer, A. (2018). O 051 – Does video game-based balance-training improve gait stability in children with cerebral palsy? *Gait and Posture*, 65(Supplement 1), 105-106. <https://doi.org/10.1016/j.gaitpost.2018.06.078>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

Green Open Access added to TU Delft Institutional Repository

'You share, we take care!' - Taverne project

<https://www.openaccess.nl/en/you-share-we-take-care>

Otherwise as indicated in the copyright section: the publisher is the copyright holder of this work and the author uses the Dutch legislation to make this work public.



ELSEVIER

Contents lists available at ScienceDirect

Gait & Posture

journal homepage: www.elsevier.com/locate/gaitpost

Short communication

O 051 – Does video game-based balance-training improve gait stability in children with cerebral palsy?

P. Meyns^{a,*}, C. Bras^b, J. Harlaar^c, L. van de Pol^d, F. Barkhof^e, A. Buizer^b

^a Hasselt University, Rehabilitation Research Institute REVAL, Diepenbeek, Belgium

^b VU University Medical Center Amsterdam, Rehabilitation Medicine, Amsterdam, Netherlands

^c Delft University of Technology, Biomechanical Engineering, Delft, Netherlands

^d VU University Medical Center Amsterdam, Child Neurology, Amsterdam, Netherlands

^e VU University Medical Center Amsterdam, Radiology and Nuclear Medicine, Amsterdam, Netherlands

1. Introduction

Video games are increasingly used in balance rehabilitation since literature has indicated promising effects on motivation [1] and functional outcomes [2] in several patient populations. Several pilot studies investigated the effectiveness of video game-based balance-training (VGBT) in children with cerebral palsy (CpC) [3,4], as these children experience poor balance control during standing and walking due to sensorimotor disorders resulting from non-progressive brain lesions [5]. These studies, however, yielded inconsistent results. This could be caused by the clinical balance scales used, which might be insensitive to training-induced improvements, as these scales are often ordinal-scale and composite scores of walking and standing balance [6].

2. Research question

Is VGBT effective to improve three balance outcomes in CpC; Pediatric Balance Scale (PBS; frequently used clinical ordinal-scale measure of functional balance), margins of stability (MoS; ratio-scale gait stability measure based on whole-body biomechanics) and MoS variability (sdMoS)?

3. Methods

Preliminary data of our registered trial (NTR6034) is presented. Eight CpC (8–16 years old) were included (Table 1), if they had: bilateral spastic CP, GMFCS-level 2, no surgery < 12 months, and no Botulinum-Toxin A injections < 6 months. VGBT comprised 6 weeks home-based X-box One Kinect (Microsoft) training. Kinect sports games focused on balance (tennis, football, bowling) were used 5x/week, 30 min/session. Time played was monitored. Pre and post-VGBT, par-

ticipants performed unperturbed treadmill walking trials at self-selected speed. Total body (Plug-In-Gait) kinematics were collected via a 10-camera Vicon motion-capture system at 120 Hz. Medio-lateral MoS was calculated as the position of the extrapolated center of mass (XCoM) relative to the lateral malleolus of the leading foot [7]. XCoM was defined as CoM, plus its velocity times a factor: $\sqrt{(\text{CoM height}/\text{acceleration of gravity})}$. High MoS indicates a large distance between XCoM and lateral border of base of support, and represents stable gait. Variability of MoS (sdMoS) was determined (high variability indicates less stable gait). Functional balance was assessed with the PBS; a 14-item balance measure concerning everyday tasks (on 56 points; high PBS indicates good balance).

4. Results

Only sdMoS improved significantly post-VGBT (Table 1).

5. Discussion

Preliminary results indicate that sdMoS is most sensitive to assess VGBT-induced balance improvements in CpC. This finding suggests a transfer from standing balance-training towards gait stability. Contrarily, VGBT did not induce improvements on PBS, which may indicate that PBS is less sensitive to training-induced changes or that training-induced sdMoS improvements are not clinically relevant. Half of the children increased (decreased) MoS after VGBT; i.e. the distance increased (decreased) between XCoM and lateral border of the base of support. These findings need further investigation taking into account whether individual patients showed increased step widths as compensation to decrease gait instability.

* Corresponding author.

E-mail address: pieter.meyns@gmail.com (P. Meyns).

Table 1
Patient characteristics and comparison between the pre- and post-VGBT balance outcomes.

		CPi01	CPi02	CPi03	CPi04	CPi05	CPi06	CPi07	CPi08	Mean	SD	Wilcoxon Signed Rank Z test	p-value
gender		male	male	male	male	male	female	male	male				
age (yrs)		10.6	9.8	11.9	8.1	10.6	15.8	14.3	13.6	11.8	2.6		
N strides	pre-VGBT	109	69	72	43	58	42	48	44				
	post-VGBT	48	50	47	48	52	38	47	43				
MoS (cm)	Pre-VGBT	14.6	17.5	20.3	17.1	15.1	22.8	15.9	18.6	17.8	2.8		
	Post-VGBT	13.3	16.9	21.4	17.0	17.6	27.0	15.0	19.6	18.5	4.3	24.00	0.401
sdMoS (cm)	Pre-VGBT	2.7	4.2	1.4	2.5	1.8	3.0	1.0	4.2	2.6	1.2		
	Post-VGBT	2.6	2.9	1.1	2.0	1.2	2.6	1.3	2.2	2.0	0.7	2.50	0.030
PBS	Pre-VGBT	41	51	54	51	50	49	56	41	49.1	5.4		
	Post-VGBT	46	49	55	50	48	49	56	52	50.6	3.5	12.50	0.673

Acknowledgment

PM was Marie-Sklodowska Curie-fellow (EU; n°660458).

References

[1] Meyns, *Dev. Neurorehabilit.* 9 (2017) 1–20.

[2] Tatla, *Dev. Med. Child Neurol.* 55 (2013) 593–601.
 [3] Jelsma, *Dev. Neurorehabilit.* 16 (2013) 27–37.
 [4] Brien, *Pediatr. Phys. Ther.* 23 (2011) 258–266.
 [5] Bruijn, *Res. Dev. Disabil.* 34 (2013) 1689–1699.
 [6] Meyns, *Gait Posture* 57 (2017) 234–235.
 [7] Hak, *PloS One* 8 (2013) e82842.