

The Thue–Morse Sequence in Base $3/2$

Dekking, F. M.

Publication date

2023

Document Version

Final published version

Published in

Journal of Integer Sequences

Citation (APA)

Dekking, F. M. (2023). The Thue–Morse Sequence in Base $3/2$. *Journal of Integer Sequences*, 26(2), Article 23.2.3.

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.



The Thue–Morse Sequence in Base $3/2$

F. M. Dekking
CWI Amsterdam and Delft University of Technology
Faculty EEMCS
P.O. Box 5031
2600 GA Delft
The Netherlands

F.M.Dekking@math.tudelft.nl

Abstract

We discuss the base $3/2$ representation of the natural numbers. We prove that the sum-of-digits function of the representation is a fixed point of a 2-block substitution on an infinite alphabet, and that this implies that sum-of-digits function modulo 2 of the representation is a fixed point $x_{3/2}$ of a 2-block substitution on $\{0, 1\}$. We prove that $x_{3/2}$ is invariant for taking the binary complement, and present a list of conjectured properties of $x_{3/2}$, which we think will be hard to prove. Finally, we make a comparison with a variant of the base $3/2$ representation, and give a general result on p - q -block substitutions.

1 Introduction

A natural number N is written in base $3/2$ if N has the form

$$N = \sum_{i \geq 0} d_i \left(\frac{3}{2}\right)^i, \tag{1}$$

with digits $d_i = 0, 1$ or 2 .

Base $3/2$ representations are also known as sesquinary representations of the natural numbers; see Propp [6]. We write these expansions as

$$\text{SQ}(N) = d_R(N) \cdots d_1(N)d_0(N) = d_R \cdots d_1 d_0.$$

We have, for example, $\text{SQ}(7) = 211$, since $2 \cdot (9/4) + (3/2) + 1 = 7$. See [A024629](#) for the continuation of Table 1. Ignoring leading 0's, the base $3/2$ representation of a number N is unique (see Section 3).

N	0	1	2	3	4	5	6	7	8	9	10
$\text{SQ}(N)$	0	1	2	20	21	22	210	211	212	2100	2101

Table 1: Base 3/2 expansions for $N = 1, \dots, 10$.

For $N \geq 0$ let

$$s_{3/2}(N) := \sum_{i=0}^{i=R} d_i(N)$$

be the sum-of-digits function of the base 3/2 expansions. We have (see [A244040](#))

$$s_{3/2} = 0, 1, 2, 2, 3, 4, 3, 4, 5, 3, 4, 5, 5, 6, 7, 4, 5, 6, 5, 6, 7, 7, 8, 9, 5, 6, 7, 5, 6, 7, 7, 8, 9, 8, 9, 10, \dots$$

In this note we study the base 3/2 analogue of the Thue–Morse sequence [A010060](#) (where the base equals 2), i.e., the sequence (see [A357448](#))

$$(x_{3/2}(N)) := (s_{3/2}(N) \bmod 2) = 0, 1, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 0, 1, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, \dots$$

The Thue Morse sequence is the fixed point starting with 0 of the substitution $0 \rightarrow 01, 1 \rightarrow 10$. This might be called a 1-2-block substitution.

Let $p \leq q$ be two natural numbers. A p - q -block substitution κ on an alphabet A is a map $\kappa : A^p \rightarrow A^q$. A p - q -block substitution κ acts on $(A^p)^*$ by defining

$$\kappa(w_1 w_2 \cdots w_{pm-1} w_{pm}) = \kappa(w_1 \cdots w_p) \cdots \kappa(w_{pm-p+1} \cdots w_{pm})$$

for $w_1 w_2 \cdots w_{pm-1} w_{pm} \in (A^p)^*$ and $m = 1, 2, \dots$. Its action extends to infinite sequences $x = x_0 x_1 \cdots$ by defining $\kappa : x \mapsto y$ by $y_{qm} \cdots y_{qm+q-1} = \kappa(x_{pm} \cdots x_{pm+p-1})$ for $m = 0, 1, \dots$

Theorem 1. *The sequence $x_{3/2}$ is a fixed point of the 2-3-block substitution*

$$\kappa : \begin{cases} 00 & \rightarrow 010 \\ 01 & \rightarrow 010 \\ 10 & \rightarrow 101 \\ 11 & \rightarrow 101 \end{cases}$$

Theorem 1 will be proved in Section 2.2.

2 Sum of digits function and Thue–Morse in base 3/2

2.1 Sum of digits function in base 3/2

Let $s_{3/2} = (0, 1, 2, 2, 3, 4, 3, 4, 5, 3, 4, 5, 5, 6, 7, 4, 5, \dots)$ be the sum-of-digits function of the base 3/2 expansions. To describe this sequence, we extend the notion of a p - q -block substitution to alphabets of infinite cardinality.

Theorem 2. *The sequence $s_{3/2}$ is the fixed point starting with 0 of the 2-3-block substitution given by*

$$a, b \mapsto a, a + 1, a + 2 \quad \text{for } a = 0, 1, 2, \dots \text{ and } b = 0, 1, 2, \dots$$

Proof. We have $d(0) = 0, d(1) = 1$ and from the uniqueness of the base $3/2$ expansions it follows immediately that $d(3N + r) = d(2N) + r$ for $N \geq 0$ and $r = 0, 1, 2$.

Thus $s_{3/2}(3N) = s_{3/2}(2N), s_{3/2}(3N + 1) = s_{3/2}(2N) + 1$, and $s_{3/2}(3N + 2) = s_{3/2}(2N) + 2$. This gives the result. \square

Remark 3. The base-4/3 version of this sequence is [A244041](#); the base-2 version is [A000120](#); the base-3 version is [A053735](#); the base-10 version is [A007953](#).

2.2 Thue–Morse in base $3/2$

Proof of Theorem 1. This follows directly from Theorem 2 by taking a and b modulo 2. \square

Although iterates of $\kappa : 00 \rightarrow 010, 01 \rightarrow 010, 10 \rightarrow 101, 11 \rightarrow 101$ are undefined, we can generate the fixed point $x_{3/2}$ by iteration of a map κ' defined by $\kappa'(w) = \kappa(w)$ if w has even length, and $\kappa'(v) = \kappa(w)$ if $v = w0$ or $v = w1$ has odd length.

The fact that the iterates of κ are undefined causes difficulty in establishing properties of $x_{3/2}$. This is similar to the lack of progress in the last 25 years to prove the conjectures on the Kolakoski sequence, which is also a fixed point of a 2-block substitution (cf. the papers [2, 3]). Here is a property that is open for the Kolakoski sequence [A000002](#), but can be proved for $x_{3/2}$.

Proposition 4. *If a word w occurs in $x_{3/2}$, then its binary complement \bar{w} defined by $\bar{0} = 1, \bar{1} = 0$, also occurs in $x_{3/2}$.*

Proof. First one checks this for all 16 words of length 6 that occur in $x_{3/2}$. Note that then also \bar{w} occurs for all w with $|w| \leq 6$, where $|w|$ denotes the length of w . Let u be a word of length $m \geq 7$. By adding at most 3 letters at the beginning and/or end of u one can obtain a word v with $|v| = 3n$ that occurs in $x_{3/2}$ at a position 0 modulo 3. But then Theorem 1 gives that $v = \kappa(w)$ for at least one word w occurring in $x_{3/2}$. The length of w is $|w| = 2n$. Since $\overline{\kappa(w)} = \kappa(\bar{w})$ the result follows by induction on $m = |u|$. For example, for $|u| = m = 7$, one has $|v| = 9$, and so $|w| = 6$. \square

Here are some conjectured properties of $x_{3/2}$.

Conjecture 5. $x_{3/2}$ is reversal invariant, i.e., if the word $w = w_1 \cdots w_m$ occurs in $x_{3/2}$ then $\overleftarrow{w} = w_m \cdots w_1$ occurs in $x_{3/2}$.

Conjecture 6. $x_{3/2}$ is uniformly recurrent, i.e., each word that occurs in $x_{3/2}$ occurs infinitely often, with bounded gaps between consecutive occurrences.

Conjecture 7. The frequencies $\mu[w]$ of the words w occurring in $x_{3/2}$ exist. Two conjectured values: $\mu[00] = 1/10, \mu[01] = 4/10$.

Conjecture 8. μ is invariant for binary complements, i.e., $\mu[w] = \mu[\bar{w}]$ for all words w .

Conjecture 9. μ is reversal invariant, i.e., $\mu[w] = \mu[\overleftarrow{w}]$ for all words w .

Conjecture 10. (Shallit) The critical exponent (=largest number of repeated blocks) of $x_{3/2}$ is 5.

3 Base $3/2$ and base $1/2 \cdot 3/2$

Many authors refer to the paper [1] from Akiyama, Frougny, and Sakarovitch for the properties of base $3/2$ expansions (see, e.g., Propp [6] and Rigo and Stipulanti [7]). However, the q/p expansions studied in paper [1] are different from the $3/2$ expansions that are usually considered as in Equation (1). In the paper [1]:

$$N = \sum_{i \geq 0} d_i \frac{1}{p} \left(\frac{q}{p}\right)^i, \quad (2)$$

with digits $d_i = 0, 1$ or 2 . We write $\text{AFS}(N)$ for the expansion of N .

Remark 11. There is a small notational problem here: Akiyama, Frougny, and Sakarovitch write about p/q expansions with $p > q$, but in this note we consider q/p expansions with $p \leq q$. This fits better with the p - q -block substitutions, and with the order of p and q in the alphabet.

Here is the table given in the paper [1] for the case $3/2$:

N	0	1	2	3	4	5	6	7	8	9	10
$\text{AFS}(N)$	ε	2	21	210	212	2101	2120	2122	21011	21200	21202

Table 2: Base $1/2 \cdot 3/2$ expansions for $N = 1, \dots, 10$.

These expansions will not even be found in the OEIS (at the moment).

The situation is clarified in the paper [5] by Frougny and Klouda. They consider both representations, called, respectively, base p/q and base $1/q \cdot p/q$ representations. In the present note these are called respectively base q/p and base $1/p \cdot q/p$ representations.

A combination of the results in [1] and [5] yields a proof of the uniqueness of the base $3/2$ expansions ($\text{QS}(N)$). There is also a direct proof of uniqueness in the paper by Edgar et al. [4]; see Theorem 1.1.

Note that $\text{AFS}(N) = \text{QS}(2N)$ for $N > 0$. So uniqueness of the base $3/2$ representation implies immediately uniqueness of the $1/2 \cdot 3/2$ representation $\text{AFS}(N)$. This observation obviously extends to base q/p .

Next we consider the question whether also the sequence $y_{3/2}$, the sum-of-digits function modulo 2 of the base $1/2 \cdot 3/2$ representation, is a fixed point of a 2-block substitution. This is indeed the case, and this 2-block substitution is given by Rigo and Stipulanti in [7].

Theorem 12. ([7]) $y_{3/2}$ is the fixed point with prefix 00 of the 2-3-block substitution

$$\kappa' : \begin{cases} 00 & \rightarrow 001 \\ 01 & \rightarrow 000 \\ 10 & \rightarrow 111 \\ 11 & \rightarrow 110 \end{cases}$$

In the paper [7] the proof of Theorem 12 is based on a generalization of Cobham's theorem to what are called \mathcal{S} -automatic sequences built on tree languages with a periodic labeled signature. Here we consider a more direct route, based on a simple closure property of p - q -block substitutions. Recall that a coding is a letter to letter map from one alphabet to another.

Theorem 13. Let $x = (x(N))$ be a fixed point of a p - q -block substitution. Let r be a positive integer. Then the sequence $(x(rN))$ is the fixed point of a coding of a p - q -block substitution.

Proof. If x is a fixed point of a p - q -block substitution, then x is also a fixed point of a pr - qr -block substitution. As new alphabet, take the words of length r occurring in x . On this alphabet, the pr - qr -block substitution induces a p - q -block substitution in an obvious way. Mapping each word of length r to its first letter is a coding that gives the result. \square

Alternative proof for Theorem 12. Apply Theorem 13 with $r = 2$. The 4-6-block substitution is given by

$$\begin{aligned} 0010 &\rightarrow 010101, & 0100 &\rightarrow 010010, & 0101 &\rightarrow 010010, & 0110 &\rightarrow 010101, \\ 1001 &\rightarrow 101010, & 1010 &\rightarrow 101101, & 1011 &\rightarrow 101101, & 1101 &\rightarrow 101010. \end{aligned}$$

Coding $00 \mapsto a$, $01 \mapsto b$, $10 \mapsto c$, $11 \mapsto d$, this induces the 2-3-block substitution

$$ac \rightarrow bbb, ba \rightarrow bac, bb \rightarrow bac, bc \rightarrow bbb, cb \rightarrow ccc, cc \rightarrow cdb, cd \rightarrow cdb, db \rightarrow ccc.$$

If we code further $a, b \mapsto 0$, and $c, d \mapsto 1$, then we obtain κ' from Theorem 12. \square

4 Acknowledgment

I am grateful to Jean-Paul Allouche for several useful comments. I also thank an anonymous referee for useful remarks.

References

- [1] S. Akiyama, C. Frougny, and J. Sakarovitch, Powers of rationals modulo 1 and rational base number systems, *Israel J. Math.* **168** (2008), 53–91.

- [2] F. M. Dekking, What is the long range order in the Kolakoski sequence?, in *The Mathematics of Long-Range Aperiodic Order*, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., Vol. 489, Kluwer Acad. Publ., 1997, pp. 115–125.
- [3] F. M. Dekking and M. Keane, Two-block substitutions and morphic words, arxiv preprint arXiv:2202.13548 [math.CO], 2022. Available at <https://arxiv.org/abs/2202.13548>.
- [4] T. Edgar, H. Olafson, and J. Van Alstine, Some combinatorics of rational base representations, preprint, 2014. Available at <https://community.plu.edu/~edgartj/preprints/basepqarithmic.pdf>.
- [5] C. Frougny and K. Klouda, Rational base number systems for p -adic numbers, *RAIRO Theor. Inform. Appl.* **46** (2019), 87–106.
- [6] J. Propp, How do you write one hundred in base $3/2$? Available at <https://tinyurl.com/2p9auyz6>. Accessed January 2023.
- [7] M. Rigo and M. Stipulanti, Automatic sequences: from rational bases to trees, *Disc. Math. Theor. Comput. Sci.* **24** (2022), #25.
- [8] N. J. A. Sloane et al., *On-Line Encyclopedia of Integer Sequences*, electronically available at <https://oeis.org>, 2023.

2020 *Mathematics Subject Classification*: Primary 11B85, Secondary 68R15
Keywords: Base $3/2$, Thue–Morse sequence, sum of digits, two-block substitution.

(Concerned with sequences [A000002](#), [A000120](#), [A007953](#), [A010060](#), [A024629](#), [A053735](#), [A244040](#), [A244041](#), and [A357448](#).)

Received February 9 2023; revised version received February 12 2023. Published in *Journal of Integer Sequences*, February 23 2023.

Return to [Journal of Integer Sequences home page](#).