



Delft University of Technology

Total land water storage change over 2003-2013 estimated from a global mass budget approach

Dieng, Habib B.; Champollion, Nicolas; Cazenave, A.; Wada, Y.; Schrama, E.; Meyssignac, B

DOI

[10.1088/1748-9326/10/12/124010](https://doi.org/10.1088/1748-9326/10/12/124010)

Publication date

2015

Document Version

Final published version

Published in

Environmental Research Letters

Citation (APA)

Dieng, H. B., Champollion, N., Cazenave, A., Wada, Y., Schrama, E., & Meyssignac, B. (2015). Total land water storage change over 2003-2013 estimated from a global mass budget approach. *Environmental Research Letters*, *10*(12), Article 124010. <https://doi.org/10.1088/1748-9326/10/12/124010>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

LETTER • OPEN ACCESS

Total land water storage change over 2003–2013 estimated from a global mass budget approach

To cite this article: H B Dieng *et al* 2015 *Environ. Res. Lett.* **10** 124010

View the [article online](#) for updates and enhancements.

Related content

- [GRACE, time-varying gravity, Earth system dynamics and climate change](#)
B Wouters, J A Bonin, D P Chambers *et al.*
- [Physical applications of GPS geodesy: a review](#)
Yehuda Bock and Diego Melgar
- [Sustainability of global water use: past reconstruction and future projections](#)
Yoshihide Wada and Marc F P Bierkens

Recent citations

- [Acceleration in the Global Mean Sea Level Rise: 2005-2015](#)
Shuang Yi *et al*
- [Greenland and Antarctica Ice Sheet Mass Changes and Effects on Global Sea Level](#)
Rene Forsberg *et al*
- [Global wetland contribution to 2000–2012 atmospheric methane growth rate dynamics](#)
Benjamin Poulter *et al*

Environmental Research Letters



LETTER

Total land water storage change over 2003–2013 estimated from a global mass budget approach

OPEN ACCESS

RECEIVED

20 August 2015

REVISED

28 October 2015

ACCEPTED FOR PUBLICATION

10 November 2015

PUBLISHED

8 December 2015

H B Dieng¹, N Champollion², A Cazenave^{1,2}, Y Wada^{3,4,5}, E Schrama⁶ and B Meysignac¹¹ LEGOS, 18 avenue E. Belin, F-31400 Toulouse, France² ISSI, Hallerstrasse 6, CH-3012 Bern, Switzerland³ Department of Physical Geography, Utrecht University, Heidelberglaan 2, 3584 CS Utrecht, The Netherlands⁴ NASA Goddard Institute for Space Studies, 2880 Broadway, New York, NY 10025, USA⁵ Center for Climate Systems Research, Columbia University, 2880 Broadway, New York, NY 10025, USA⁶ Faculty of Aerospace Engineering, Kluyverweg 1, 2629HS Delft, The NetherlandsE-mail: habib.dieng@legos.obs-mip.fr**Keywords:** land waters, sea level rise, global water mass budgetSupplementary material for this article is available [online](#)

Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

**Abstract**

We estimate the total land water storage (LWS) change between 2003 and 2013 using a global water mass budget approach. Hereby we compare the ocean mass change (estimated from GRACE space gravimetry on the one hand, and from the satellite altimetry-based global mean sea level corrected for steric effects on the other hand) to the sum of the main water mass components of the climate system: glaciers, Greenland and Antarctica ice sheets, atmospheric water and LWS (the latter being the unknown quantity to be estimated). For glaciers and ice sheets, we use published estimates of ice mass trends based on various types of observations covering different time spans between 2003 and 2013. From the mass budget equation, we derive a net LWS trend over the study period. The mean trend amounts to $+0.30 \pm 0.18 \text{ mm yr}^{-1}$ in sea level equivalent. This corresponds to a net decrease of $-108 \pm 64 \text{ km}^3 \text{ yr}^{-1}$ in LWS over the 2003–2013 decade. We also estimate the rate of change in LWS and find no significant acceleration over the study period. The computed mean global LWS trend over the study period is shown to be explained mainly by direct anthropogenic effects on land hydrology, i.e. the net effect of groundwater depletion and impoundment of water in man-made reservoirs, and to a lesser extent the effect of naturally-forced land hydrology variability. Our results compare well with independent estimates of human-induced changes in global land hydrology.

1. Introduction

Liquid fresh water on land is stored in various reservoirs: rivers, lakes, man-made reservoirs, wetlands and inundated areas, root zone (upper few meters of the soil) and aquifers (groundwater reservoirs). Terrestrial reservoirs continuously exchange with the atmosphere, oceans and land, through vertical and horizontal mass fluxes (precipitation, evaporation, transpiration of the vegetation, surface runoff and underground flow). Land water storage (LWS) varies with change in mean climate and climate variability. Human activities also directly affect LWS through water extraction from aquifers, building of dams along rivers, urbanization, wetland drainage,

land use and land cover changes, and deforestation. All these effects modify the water budget in river basins, and because of water mass conservation in the climate system, cause sea level changes. Studies based on hydrological modeling have not reported any clear long-term trend in global LWS over the past 60 years but only interannual variability (e.g., Ngo-Duc *et al* 2005). This is unlike human-induced factors such as dam building (Chao *et al* 2008) and groundwater extraction (Konikow 2011, Pokhrel *et al* 2012, Wada *et al* 2012, Wada 2015). Although their contributions to the global mean sea level (GMSL) are of opposite sign (<0 for dams, >0 for groundwater pumping), their net effect is responsible for a significant long-term positive trend at least for the recent decades

(other human-induced factors have negligible contributions to the GMSL). Building on the results from Konikow (2011) and Wada *et al* (2012), Church *et al* (2013) estimated that the net effect of dams and groundwater depletion (i.e., groundwater abstraction minus recharge; e.g., Wada 2015) on the GMSL amounted $0.38 \pm 12 \text{ mm yr}^{-1}$ over 1993–2010. This represents 12% of the observed GMSL rate of rise over this time span, an amount of the same order of magnitude as the Antarctic ice mass loss (see table 13.1 in Church *et al* 2013). Because of such a significant contribution to sea level, it is worth to examine this component in more detail. In addition, uncertainty of this component has direct impact on our capability to close the sea level budget, thus constrain missing contributions (due to lack of data) such as the deep ocean thermal expansion (see discussions on that topic in Llovel *et al* 2014, Dieng *et al* 2015a).

The effect of dams and man-made reservoirs has been estimated by Chao *et al* (2008). They reconstructed the history of water impoundment in the nearly 30 000 reservoirs built during the twentieth century and estimated the contribution to sea level by dams and artificial reservoirs (including seepage) at $-0.55 \pm 0.08 \text{ mm yr}^{-1}$ in sea level equivalent (SLE) during the last half-century, with a stabilization in recent years. Estimates of groundwater depletion are based on three methods (see Wada 2015): (1) volume-based method, (2) flux-based method, and (3) satellite observations from the GRACE space gravimetry mission. Each method has strengths and weaknesses. For example, GRACE gives a vertically integrated estimate of the water mass change; thus surface waters and soil moisture must be known and removed to estimate the ground water contribution. In addition, GRACE-based estimates do not yet have full global coverage for the estimation of groundwater depletion (Famiglietti 2014, Wada 2015). The volume- and flux-based methods lack global information and suffer from model uncertainties. Thus estimating groundwater depletion remains very challenging, as is the global dam contribution.

In this study, we develop another approach based on the global water mass budget of the climate system to estimate the total LWS change. Focusing on the January 2003–December 2013 time span (for which GRACE data are available), we compare the GRACE-based ocean mass change to the sum of mass components (glaciers, Greenland and Antarctica ice sheets, atmospheric water vapor and LWS). We neglect other mass components such as permafrost because global data are lacking, as well as change in the snow pack, previously shown to give negligible contribution to the GMSL beyond time spans larger than 1 year (Biancamaria *et al* 2011). In this mass budget approach, we use estimates of each component from different observational data sets, except for the net total LWS, the unknown quantity to be estimated. A mean LWS trend is first estimated over 2003–2013. Then, accounting

for increasing rate of change (acceleration) of several components (ocean mass, ice sheet mass balances) over the study time span, we investigate whether the LWS rate varies with time. To validate our results, we perform a similar analysis but instead of using GRACE, we estimate the ocean mass term from the satellite altimetry-based GMSL corrected for steric effects (i.e., effects of ocean temperature and salinity).

All results are expressed in terms of SLE change. Units are given in mm yr^{-1} and mm yr^{-2} for trend and acceleration respectively.

2. Method

To estimate the contribution of LWS change to sea level, we can simply consider the conservation of water mass in the Earth's system (e.g., Llovel *et al* 2010). Of course, LWS change could be derived from GRACE data over the continents, as done previously in a number of studies. However, considering that the GRACE resolution ($\sim 300\text{--}500 \text{ km}$) may be problematic in separating nearby masses (e.g., river basins and glaciers), our objective here is to use a different approach.

On time scales of years to decades, water mass changes inside the solid Earth (e.g., in the crust) can be neglected, so that only changes in land reservoirs, ocean and atmosphere need to be considered, with the mass conservation equation written as follows:

$$\begin{aligned} \Delta M_{\text{Ocean}}(t) + \Delta M_{\text{Glaciers}}(t) + \Delta M_{\text{Greenland}}(t) \\ + \Delta M_{\text{Antarct.}}(t) + \Delta M_{\text{Atm}}(t) \\ + \Delta M_{\text{LWS}}(t) = 0, \end{aligned} \quad (1)$$

where $\Delta M(t)$ represents changes with time t of water mass in the different reservoirs: ocean, glaciers (including small ice caps), Greenland ice sheet, Antarctica ice sheet, atmosphere and land water stores. Note that $\Delta M(t)$ may be either positive or negative. Using equation (1), we deduce the LWS component by simple rewriting as:

$$\begin{aligned} \Delta M_{\text{LWS}}(t) = - \left[\Delta M_{\text{Ocean}}(t) + \Delta M_{\text{Glaciers}}(t) \right. \\ \left. + \Delta M_{\text{Greenland}}(t) + \Delta M_{\text{Antarct.}}(t) \right. \\ \left. + \Delta M_{\text{Atm}}(t) \right]. \end{aligned} \quad (2)$$

As mentioned above, all contributions are expressed in terms of SLE.

3. Data

3.1. Ocean mass

For estimating the ocean mass component, we apply two approaches: (1) use of GRACE space gravimetry data over the oceans, and (2) estimate of the GMSL corrected for steric effects.

3.1.1. GRACE-based ocean mass

Three different data sets of the GRACE Release 05 products have been considered:

- (1) from the Texas University (CSR RL05),
- (2) from the German GeoForschungsZentrum (GFZ RL05)
- (3) from the Jet Propulsion Laboratory (JPL RL05).

To study the ocean mass evolution, a specific processing has been carried out by D. Chambers (described in Johnson and Chambers 2013; geocenter terms included; data available at https://dl.dropboxusercontent.com/u/31563267/ocean_mass_orig.txt). The data are provided as global mean (averaged over the 90°S–90°N domain) time series at monthly interval with associated uncertainty. The GIA (Glacial Isostatic Adjustment) effect is corrected for using the GIA correction computed in Chambers *et al* (2010). In the following, we consider the mean of the three data sets.

3.1.2. Ocean mass estimated from the GMSL corrected for steric effects

Changes in the GMSL result from steric effects plus ocean mass changes. Thus, the ocean mass component can be also derived from the difference ‘GMSL minus steric effects’. For that purpose we used the mean of six different satellite altimetry-based GMSL data sets: (1) Validation and Interpretation of Satellite Oceanographic (AVISO; <http://aviso.altimetry.fr/en/data/products/ocean-indicators-products/actualitesindicateurs-des-oceansniveau-moyen-des-mersindexhtml.html>); (2) University of Colorado (CU Release 5; <http://sealevel.colorado.edu/>); (3) National Oceanographic and Atmospheric Administration (NOAA; http://star.nesdis.noaa.gov/sod/lsa/SeaLevelRise/LSA_SLR_timeseries_global.php); (4) Goddard Space Flight Center (GSFC version 2; http://podaac-ftp.jpl.nasa.gov/dataset/MERGED_TP_J1_OSTM_OST_GMSL_ASCII_V2); (5) Commonwealth Scientific and Industrial Research Organization (CSIRO; http://cmar.csiro.au/sealevel/sl_data_cmar.html); (6) The European Space Agency/ESA Climate Change Initiative/CCI sea level data (<http://esa-sealevel-cci.org/>). Details on these data sets can be found in Dieng *et al* (2015a, 2015b).

For the steric component, instead of using Argo that suffer from gaps in the data coverage (e.g., in the Indonesian region; Dieng *et al* 2015b), we make use of the ORAS4 reanalysis (Balmaseda *et al* 2013) that provides ocean temperature and salinity down to 5350 m and global coverage. Note that over their common geographical and depth coverage, Argo-based and ORAS4-based steric sea level are in good agreement (see Dieng *et al* 2015b for a discussion).

Using the mean of the six GMSL products, we compute the ocean mass component by subtracting the ORAS4 steric component. It is simply called below ‘GMSL minus ORAS4’.

3.2. Atmospheric water vapor mass

To estimate change in atmospheric water vapor mass, we used the vertically integrated water vapor grids from the ERA Interim reanalysis performed by the European Center for Medium Range Weather Forecast/ECMWF (Dee *et al* 2011). The data are provided as 1.5° × 1.5° grids at monthly interval. We compute a globally averaged water vapor time series and express it in terms of SLE (see Dieng *et al* 2014 for details).

3.3. Greenland and Antarctica mass

For the ice sheet mass balances, we used two approaches: (1) time series given by Velicogna *et al* (2014) and from the ESA CCI Ice Sheet project (<http://esa-icesheets-cci.org>; see also Forsberg *et al* 2014), (2) published estimates of mass balance trends from the literature. For Greenland, we considered 52 published trend values based on 30 articles. For Antarctica, we used 24 published trend values based on 13 articles. Corresponding list of the 43 articles used in this study, as well as associated trend values are given in the supporting information (SI).

3.4. Glaciers mass



For the glaciers, we considered 4 published estimates of mass balance trends from the literature (Gardner *et al* 2013, Schrama *et al* 2014, Yi *et al* 2015). The Gardner *et al* data are a compilation of a large number of glacier mass balance estimates from different methods (these estimates are in terms of trends over the 2003–2009 time span). Global glacier mass trends from Schrama *et al* and Yi *et al* are based on GRACE and are given over different time intervals: January 2003–December 2013 (Schrama *et al* 2014); January 2005–December 2009 and January 2010–March 2014 (Yi *et al* 2015).

Note that the four glaciers estimates considered in this study do not include Greenland and Antarctica peripheral glaciers. Trend values are given in the SI.

4. Data analysis

When time series are used to estimate trends and accelerations, the annual & semi annual signals are removed by fitting 12-month and 6-month sinusoids.

Figure 1 shows the GRACE-based ocean mass (called GOM) time series over 2003–2013. This time series is an update of that previously used by Dieng *et al* (2015a, 2015b) to examine the closure of the sea level budget. A mean trend of $1.85 \pm 0.1 \text{ mm yr}^{-1}$ is estimated over the study time span. We fitted a degree 2 polynomial to the data, from which we deduce an acceleration of $0.29 \pm 0.04 \text{ mm yr}^{-2}$. The





acceleration is defined as 2 times the adjusted coefficient of the polynomial t^2 term. The quoted uncertainties represent 1 sigma errors estimated from the least-squares fit and accounting for the time series errors. In figure 1 is superimposed the ‘GMSL minus ORAS4’ ocean mass time series and associated uncertainty (note that the GOM uncertainty is not shown because smaller than the latter). The mean ‘GMSL minus ORAS4’ trend over 2003–2013 amounts to $2.03 \pm 0.11 \text{ mm yr}^{-1}$. Besides, the acceleration is found to be almost zero over the study time span.

Similarly, figure 2 shows the global atmospheric water vapor time series. The mean trend estimated from the time series is slightly negative (equal to $-0.04 \pm 0.04 \text{ mm yr}^{-1}$ SLE), indicating a small but

not significant increase in atmospheric water vapor content. Dieng *et al* (2014) considered other water vapor datasets and found little differences in terms of interannual variability and trend. As for the ocean mass data, we fitted a degree 2 polynomial but found zero acceleration.



Figure 3 shows a plot of the 52 published trend values of the Greenland mass balance (in mm yr^{-1} SLE). Horizontal lines represent the time span covered by each analysis. Note that several values are superimposed and are not visible on the figure. Averaging all published values provides a mean trend of $0.76 \pm 0.1 \text{ mm yr}^{-1}$ over 2003–2013 (this 1 sigma error is that obtained from the regression, accounting for errors provided with each estimate). This value can



be compared with other mean trend estimates over the same time span (2003–2013): $0.77 \pm 0.16 \text{ mm yr}^{-1}$ (Velicogna *et al* 2014), $0.63 \pm 0.1 \text{ mm yr}^{-1}$ (CCI data set), and $0.77 \pm 0.05 \text{ mm yr}^{-1}$ (Schrama *et al* 2014). All trends agree well within their respective error bars. We next performed a linear regression of the trend data shown in figure 3 to estimate the acceleration term (identified in figure 3 by the black line). The computed acceleration of the Greenland ice sheet mass loss amounts to $0.071 \pm 0.007 \text{ mm yr}^{-2}$. The acceleration provided by Velicogna *et al* (2014) equals $0.071 \pm 0.004 \text{ mm yr}^{-2}$, in perfect agreement with our estimate based on the 52 published values. The acceleration estimated from the CCI data amounts to $0.060 \pm 0.007 \text{ mm yr}^{-2}$, slightly less than the other two values.

Figure 4 is a plot similar to figure 3 but for the Antarctica ice sheet. The black straight line has also the

same meaning. Unlike Greenland, results for the Antarctica mass trends are much more scattered, even though there is indication of an acceleration (as reported in the literature; see also Church *et al* 2013). The mean trend over 2003–2013 based on all individual trends amounts to $0.34 \pm 0.12 \text{ mm yr}^{-1}$, a value slightly larger than those from Velicogna *et al* (2014) ($0.19 \pm 0.12 \text{ mm yr}^{-1}$), Schrama *et al* (2014) ($0.26 \pm 0.08 \text{ mm yr}^{-1}$), and the CCI data ($0.27 \pm 0.10 \text{ mm yr}^{-1}$). Such differences may be partly attributed to the considered GIA correction, highly important for Antarctica (see Schrama *et al* 2014 for a discussion). The acceleration adjusted on the 24 published trends considered in our study amounts to $0.051 \pm 0.011 \text{ mm yr}^{-2}$. This value is slightly larger than the Velicogna *et al* (2014) and CCI estimates (of $0.031 \pm 0.010 \text{ mm yr}^{-2}$ and $0.042 \pm 0.010 \text{ mm yr}^{-1}$, respectively).



A treatment similar to that applied for the ice sheets was performed for the glaciers mean trend and acceleration (see figure 5, with the black straight line representing the acceleration). The mean trend and acceleration over 2003–2013 are estimated to $0.58 \pm 0.1 \text{ mm yr}^{-1}$ and $-0.004 \pm 0.005 \text{ mm yr}^{-2}$. The acceleration is not significantly different from zero. Using GRACE only, Schrama *et al* (2014) estimated the mean trend of the glacier contribution over 2003–2013 to $0.44 \pm 0.03 \text{ mm yr}^{-1}$. The difference cannot be attributed to peripheral glaciers of Greenland and Antarctica ice sheets, not considered in either case. However, the GRACE only glacier estimate may be contaminated by land hydrology because of the poor GRACE resolution.

We are also aware that the estimated acceleration needs to be used with caution due to the few available glacier observation-based data sets.

5. Results

5.1. LWS trend over 2003–2013

In figure 6, we present a chart of the mean trends over 2003–2013 for ΔM_{Ocean} (from GRACE and from ‘GMSL minus ORAS4’), the sum of ΔM (atmospheric water vapor plus glaciers plus ice sheets) and the residuals (ΔM_{Ocean} —sum of ΔM). For the sums (and residuals as well), 3 trend values are considered. Sums a, b and c correspond to: (a) the average trend estimated with the published results (i.e., sum of mean Greenland

plus mean Antarctica plus mean glacier plus water vapor trends—values given in section 4 above-), (b) the Velicogna *et al* (2014)'s trends for Greenland and Antarctica, and (c) the CCI trends also for Greenland and Antarctica. For sums b and c, we consider for glaciers and water vapor trends, the same values as in sum a. All residuals are interpreted in terms of LWS trends over 2003–2013, and are expressed in SLE.

Residuals 1 and 2 (called res1 and res2) in figure 6 are based on ΔM_{Ocean} from GRACE and from 'GMSL minus ORAS4', to which 'sum a' is subtracted. We also compute the mean of res1 and res2. Corresponding trends amount to $0.21 \pm 0.18 \text{ mm yr}^{-1}$ (res1), $0.39 \pm 0.19 \text{ mm yr}^{-1}$ (res2) and $0.30 \pm 0.18 \text{ mm yr}^{-1}$ (mean). Residuals 3 and 4 (called res3 and res4) are based on sums b and c respectively, using the mean value of ΔM_{Ocean} from GRACE and 'GMSL minus ORAS4'. These are also plotted in figure 6 as well as LWS trends estimated by Wada *et al* (2012)—based on the flux method- for 2 cases: (1) only dams and groundwater depletion are accounted for, and (2) in addition to dams and ground waters, account of deforestation and wetland drainage (called Wada1 and Wada2 hereinafter). Trends over 2003–2013 for res3, res4, Wada1 and Wada2 amount to $0.44 \pm 0.24 \text{ mm yr}^{-1}$, $0.50 \pm 0.20 \text{ mm yr}^{-1}$, $0.39 \pm 0.11 \text{ mm yr}^{-1}$ and $0.54 \pm 0.12 \text{ mm yr}^{-1}$, respectively.

All computed residuals appear rather consistent in spite of the quite different data sets used. They also compare rather well with Wada1 & 2. Trend values are gathered in the SI.

5.2. LWS acceleration over 2003–2013

As mentioned above, figures 3 and 4 show clear acceleration for the Greenland and Antarctica mass balances (unlike the glacier and water vapor components). The ΔM_{Ocean} based on GRACE also displays important acceleration over the study time span. Using the mass budget equation, we can deduce the acceleration of the LWS residuals for this case. For res1, it amounts to $+0.17 \pm 0.04 \text{ mm yr}^{-2}$. However if we consider ΔM_{Ocean} based on 'GMSL minus ORAS4', acceleration of the residual time series (res2) becomes negative (and equal to $-0.08 \pm 0.05 \text{ mm yr}^{-2}$). Besides accelerations of Wada 1 and 2 are very small and non significant (of $0.008 \pm 0.010 \text{ mm yr}^{-2}$). With the data currently available, it does not seem possible to estimate any reliable LWS acceleration, nor to identify which term of the mass budget equation compensates the ice sheet mass balance acceleration. Our results suggest nevertheless that the LWS acceleration is not significantly different from zero over the 2003–2013 time span.

5.3. Interannual variability in LWS trends over 2003–2013

We computed short-term trends of the ΔM_{Ocean} time series based on GRACE and 'GMSL minus ORAS4',


over successive 2-year time spans (with 1 year overlap). To these short-term trends, we removed the acceleration term of sum a (i.e., the combined acceleration of glaciers, ice sheets and water vapor). The corresponding curves are shown in figure 7 (labeled res1 and res2). In figure 7 are superimposed four additional LWS 2-year trend curves (also with 1-year overlap) using: (1) LWS determined by Yi *et al* (2015) using GRACE over continental river basins, (2) LWS based on the ISBA/TRIP global hydrological model (Alkama *et al* 2010), (3) LWS based on the Water Gap Hydrological Model (WGHM, Döll *et al* 2014a), and (4) LWS based on the Modern Era Retrospective-analysis for Research and Applications (MERRA) Reanalysis (Mantas *et al* 2015). The ISBA/TRIP and WGHM land surface schemes calculate time variations of surface energy and water budgets in different soil layers. ISBA/TRIP only considers upper soil layers while WGHM accounts for groundwater and man-made reservoirs (Döll *et al* 2014a, 2014b). The MERRA dataset is the version 5.2.0 of the GEOS-5 data service. We used the total water storage in land reservoirs product that includes the groundwater component. As for ISBA/TRIP and WGHM, it is available as gridded time series at monthly interval over the 2003–2013 time span. We computed geographical averages, applying a cosine latitude weighting.

All six curves (expressed in SLE trends) exhibit large interannual variability, mostly related to El Niño-Southern Oscillation (ENSO) events (note for example the minimum corresponding to the 2011 La Niña). On average, a good correlation at interannual time scale is noticed between these six curves. But we note closer agreement between 2-year trend curves from res1 and Yi *et al* LWS on the one hand, and ISBA/TRIP and MERRA LWS on the other hand. The Wada 1 and Wada 2 short-term trends are also shown in figure 7 (bottom curves; signal amplified by a factor of 10). It is interesting to note that these also display a minimum in 2011. Although the latter only represent the direct anthropogenic components, the 2011 minimum likely reflects increased groundwater recharge during this La Niña episode.

All trends and accelerations estimates presented above are gathered in the SI.

6. Discussion

The mean trend in LWS estimated by the global mass budget approach developed in this study is found to be positive in terms of SLE over the 2003–2013 time span. The mean of the two estimates based on two different values of ΔM_{Ocean} is $0.30 \pm 0.18 \text{ mm yr}^{-1}$ SLE. This corresponds to an annual decrease in net LWS of $-108 \pm 64 \text{ km}^3 \text{ yr}^{-1}$. This quantity represents the combined effects of natural climate variability, anthropogenic climate change and direct anthropogenic factors. The uncertainty of this estimate directly relies



on the ocean mass trend uncertainty. Here we used two independent methods to estimate ΔM_{Ocean} and associated uncertainty, with quite consistent results. Wada *et al* (2012)'s results for the direct anthropogenic LWS components are only slightly larger (of $+0.39 \pm 0.11 \text{ mm yr}^{-1}$ SLE or $-140 \pm 40 \text{ km}^3 \text{ yr}^{-1}$ LWS trend) for the net effect of dams and ground water depletion. While the rate of reservoir impoundment exceeded groundwater depletion over most of the 20th century, for the recent years, groundwater depletion exceeds impoundment, thus the net effect leads to a positive contribution to the GMSL. In addition to Wada *et al* (2012), other estimates of the groundwater component have been published in the recent years: Konikow (2011) estimated that human-induced groundwater depletion contributed $0.34 \pm 0.07 \text{ mm yr}^{-1}$ to the GMSL rise over 1993–2008 (based mostly on observational methods). Pokhrel *et al* (2012) estimated much larger groundwater depletion over 1981–2007, amounting $1.0 \pm 0.16 \text{ mm yr}^{-1}$ SLE (or $0.8 \pm 0.25 \text{ mm yr}^{-1}$ when accounting for dams and natural climate variability). A recent study by Döll *et al* (2014b) based on the WGHM model combined with GRACE data finds a groundwater contribution to GMSL rise of $0.31 \pm 0.06 \text{ mm yr}^{-1}$ for the 2003–2009 time span. Thus, these estimates (except for Pokhrel *et al* 2012) agree well with our results.

Other studies based on GRACE have provided estimates of the total (natural plus anthropogenic) LWS change over different time spans. Considering the 33 largest river basins, Llovel *et al* (2010) and Jensen *et al* (2013) estimated to $-0.22 \pm 0.05 \text{ mm yr}^{-1}$ and $-0.20 \pm 0.04 \text{ mm yr}^{-1}$ the LWS contribution to

GMSL over 2002–2009. Two LWS trend estimates by Yi *et al* (2015) for 2005–2009 and 2010–2014 give $-0.27 \pm 0.25 \text{ mm yr}^{-1}$ and $0.38 \pm 0.48 \text{ mm yr}^{-1}$ respectively. Note that over the 2003–2013 period, the Yi *et al* (2015) and Schrama *et al* (2014) LWS trends (both based on GRACE data processing on land) are respectively slightly positive and negative ($+0.07 \pm 0.04 \text{ mm yr}^{-1}$ and $-0.06 \pm 0.09 \text{ mm yr}^{-1}$ respectively). This dispersion of LWS values and large associated uncertainties based on GRACE is not totally surprising. The GRACE LWS rate estimates are much dependent on the study period, considering the importance of the interannual variability (see figure 7), as discussed in Jensen *et al* (2013). Besides, the GRACE resolution does not allow unambiguous separation between nearby sources (Longuevergne *et al* 2010). This is particularly true in the region of the Ganges basin and Himalayan glaciers. Finally, studies considering a limited number of river basins (e.g., Llovel *et al* 2010) miss part of the signal, and the GIA correction, important in high latitudes, adds significant uncertainty. All together, the GRACE LWS estimates remain uncertain, in particular if study time spans are short (Landerer and Swenson 2012).

In this study, we proposed a different approach to estimate the net LWS contribution to GMSL change over a 10 year time span. Using a large number of different data sets for the mass components, we came up to a positive value for the LWS trend over 2003–2013 (in terms of SLE; i.e., decrease of total water storage on land) that likely reflects the net anthropogenic component, i.e., the dominant contribution of groundwater depletion versus dams, in good agreement with the Wada *et al* (2012) estimates.




Figure 8. Global and regional trends in surface water and groundwater use (adapted from Wada *et al* 2014).

As far as the acceleration is concerned, results from this study remain inconclusive. But they suggest that there is no significant acceleration in LWS change over the 2003–2013 decade, in agreement with Wada *et al* (2012)'s results. Over the 2000s, Jung *et al* (2010) suggest higher soil moisture and large water availability, which reduce the amount of irrigation needs and thus groundwater pumping. Figure 8 shows the global and regional trends in surface water and groundwater use. For South Asia (India, Pakistan), North America (USA), and Western Europe (Spain, Greece, Italy) where the sum of regional groundwater depletion accounts for more than half of the global total, the increase in groundwater pumping is low over the period. This substantially slowed down the increase in groundwater contribution to sea level rise. At the same time, negative contribution due to reservoir impoundment is also rather constant over the same period, with a sharp increase due to the Three Gorges Dams only over a few years. Considering the two terms being almost constant, the net contribution also became rather constant with negligible acceleration over the 2000s. This explains the results obtained in this study.

Finally, our study confirms that total LWS rate is highly variable on the short-term (interannual time scale) and much impacted by ENSO events. The net LWS rate due to direct anthropogenic forcing (Wada *et al* 2012) also responds to ENSO events, likely via aquifer recharge.

Acknowledgments

H B Dieng is supported by an ESA (European Space Agency) grant in the context of the Climate Change Initiative (CCI) project. Y Wada is supported by Japan Society for the Promotion of Science (JSPS) Oversea Research Fellowship (grant no. JSPS-2014-878). We thank S Yi for providing us with his GRACE-based land water time series, as well as B Decharme for

making available to us the outputs of the ISBA/TRIP hydrological model.

References

- Alkama R *et al* 2010 Global evaluation of the ISBA-TRIP continental hydrological system I. Comparison to GRACE terrestrial water storage estimates and *in situ* river discharges *J. Hydromet.* **11** 583–600
- Balmaseda M A, Mogens K and Weaver A 2013 Evaluation of the ECMWF ocean reanalysis ORAS4 Q. *J. R. Meteorol. Soc.* **139** 1132–61
- Biancamaria S, Cazanave A, Mognard N, Llovel W and Frappart F 2011 Satellite-based high latitudes snow volume trend, variability and contribution to sea level over 1989/2006 *Glob. Planet. Change* **75** 99–107
- Chambers D P, Wahr J, Tamisiea M E and Nerem R S 2010 Ocean mass from GRACE and glacial isostatic adjustment *J. Geophys. Res.* **115** B11415
- Chao B F, Wu Y H and Li Y S 2008 Impact of artificial reservoir water impoundment on global sea level *Science* **320** 212–4
- Church J A *et al* 2013 Sea level change *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker *et al* (Cambridge: Cambridge University Press)
- Dee D P *et al* 2011 The ERA-Interim reanalysis: configuration and performance of the data assimilation system Q. *J. R. Meteorol. Soc.* **137** 553–97
- Dieng H B, Cazenave A, Messignac B, Henry O, von Schuckmann K and Lemoine J M 2014 Effect of La Niña on the global mean sea level and north Pacific ocean mass over 2005–2011 *J. Geodetic Sci.* **4** 19–27
- Dieng H B, Palanisamy H, Cazenave A, Meyssignac B and von Schuckmann K 2015a The sea level budget since 2003: inference on the deep ocean heat content *Surv. Geophys.* **36** 209–29
- Dieng H, Cazenave A, von Schuckmann K, Ablain M and Meyssignac B 2015b Sea level budget over 2005–2013: missing contributions and data errors *Ocean Sci.* **11** 1–14
- Döll P, Fritsche M, Eicker A and Mueller Schmied H 2014a Seasonal water storage variations as impacted by water abstractions: comparing the output of a global hydrological model with GRACE and GPS observations *Surv. Geophys.* **35** 1311–31
- Döll P, Mueller S H, Schuh C, Portmann F T and Eicker A 2014b Global-scale assessment of groundwater depletion and related groundwater abstractions: combining hydrological

- modeling with information from well observations and GRACE satellites *Water Resour. Res.* **50** 5698–720
- Famiglietti J S 2014 The global groundwater crisis *Nat. Clim. Change* **4** 945–8
- Forsberg R et al 2014 Mass balance of Greenland from combined GRACE and satellite altimetry inversion, 2014 AGU fall meeting
- Gardner A S et al 2013 A reconciled estimate of glacier contributions to sea level rise, 2003–2009 *Science* **340** 852–7
- Jensen L, Rietbroek R and Kusche J 2013 Land water contribution to sea level and Jason-1 measurements *J. Geophys. Res.: Oceans* **118** 212–26
- Johnson G C and Chambers D P 2013 Ocean bottom pressure seasonal cycles and decadal trends from GRACE Release-05: Ocean circulation implications *J. Geophys. Res.: Oceans* **118** 4228–40
- Jung M et al 2010 Recent decline in the global land evapotranspiration trend due to limited moisture supply *Nature* **467** 951–4
- Konikow L F 2011 Contribution of global groundwater depletion since 1900 to sea-level rise *Geophys. Res. Lett.* **38** L17401
- Landerer F W and Swenson S C 2012 Accuracy of scaled GRACE terrestrial water storage estimates *Water Resour. Res.* **48** W04531
- Llovel W, Becker M, Cazenave A and Crétaux J F 2010 Contribution of land water storage change to global mean sea level from GRACE and satellite altimetry *C.R. Geosci.* **342** 179–88
- Llovel W, Willis J K, Landerer F W and Fukumori I 2014 Deep-ocean contribution to sea level and energy budget not detectable over the past decade *Nat. Clim. Change* **4** 1031–5
- Longuevergne L, Scanlon B R and Wilson C R 2010 GRACE Hydrological estimates for small basins: evaluating processing approaches on the High Plains Aquifer, USA *Water Resour. Res.* **46** W11517
- Mantas V M, Liu Z and Pereira A J S C 2015 A Web service and android application for the distribution of rainfall estimates and Earth observation data *Comput. Geosci.* **77** 66–76
- Ngo-Duc T, Laval K, Polcher J, Lombard A and Cazenave A 2005 Effects of land water storage on global mean sea level over the past 50 years *Geophys. Res. Lett.* **32** L09704
- Pokhrel Y N, Hanasaki N, Yeh P J-F, Yamada T, Kanae S and Oki T 2012 Model estimates of sea level change due to anthropogenic impacts on terrestrial water storage *Nat. Geosci.* **5** 389–92
- Schrama E J O, Wouters B and Rietbroek R 2014 A mascon approach to assess ice sheet and glacier mass balance and their uncertainties from GRACE data *J. Geophys. Res.: Solid Earth* **119** 6048–66
- Velicogna I, Sutterley T C and van den Broeke M R 2014 Regional acceleration in ice mass loss from Greenland and Antarctica using grace time variable gravity data *Geophys. Res. Lett.* **41** 8130–7
- Wada Y 2015 Modelling groundwater depletion at regional and global scales: present state and future prospects *Surv. Geophys.* in press (doi:10.1007/s10712-015-9347-x)
- Wada Y, van Beek L P H, Weiland F C S, Chao B F, Wu Y-H and Bierkens M F P 2012 Past and future contribution of global groundwater depletion to sea-level rise *Geophys. Res. Lett.* **39** L09402
- Wada Y, Wisser D and Bierkens M F P 2014 Global modeling of withdrawal, allocation and consumptive use of surface water and groundwater resources *Earth Syst. Dyn.* **5** 15–40
- Yi S, Sun W, Heki K and Qian A 2015 An increase in the rate of global mean sea level rise since 2010 *Geophys. Res. Lett.* **42** 3998–4006