

Recruiting participants for ergonomic research using self-reported stature and body mass

Kılıç, H.; Vledder, G.; Yao, X.; Elkhuisen, W.S.; Song, Y.; Vink, P.

DOI

[10.3233/WOR-220565](https://doi.org/10.3233/WOR-220565)

Publication date

2023

Document Version

Final published version

Published in

Work: a journal of prevention, assessment & rehabilitation

Citation (APA)

Kılıç, H., Vledder, G., Yao, X., Elkhuisen, W. S., Song, Y., & Vink, P. (2023). Recruiting participants for ergonomic research using self-reported stature and body mass. *Work: a journal of prevention, assessment & rehabilitation*, 76(4), 1509-1517. <https://doi.org/10.3233/WOR-220565>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

Recruiting participants for ergonomic research using self-reported stature and body mass

Halil Kılıç^a, Gerbera Vledder^{b,*}, Xinhe Yao^b, Willemijn S. Elkhuizen^b, Yu Song^b and Peter Vink^b
^a*Department of Woodworking Industrial Engineering, Faculty of Technology, Muğla Sıtkı Koçman University, Muğla, Turkey*
^b*Faculty of Industrial Design Engineering, Delft University of Technology, Delft, The Netherlands*

Received 30 September 2022

Accepted 6 April 2023

Abstract.

BACKGROUND: A valid distribution of key anthropometric parameters among participants is often a prerequisite of ergonomics research.

OBJECTIVE: In this paper, we investigated the accuracy of self-reported stature and body mass of the population in the Netherlands.

METHODS: Data from 4 experiments was synthesized where in each experiment, participants self-reported their stature and body mass prior to being measured, of which they were not notified before.

RESULTS: Statistical analysis of 249 records indicated that on average, participants overreported their stature by 1.31 cm and underreported their mass by 1.45 kg. This is especially true for people with a BMI ≥ 25 .

CONCLUSION: Two models were proposed to adjust the self-reported stature and body mass for ergonomic researchers in a survey or recruitment. Limitations in using the models are highlighted as well.

Keywords: Anthropometrics, body mass index, participatory research

1. Introduction

In the context of ergonomics research, the anthropometric measurements of participants are often a prerequisite. Self-reported techniques are often used to collect information on the stature, the body mass, and the calculated body mass index (BMI, in kg/m²) during survey and recruiting subjects. In different (online) surveys, the height and body mass can be important factors in evaluating physical activities, perceptions, etc. [1, 2]. For instance, Liu et al. asked

the height and the body mass of 90 participants via questionnaires for studying comfort of seats in a staggered configuration [3]. In an online survey designed by Srinivasan et al. on the topic of college students' personality under the impact of COVID-19 online classes, self-reported height and body mass of 897 subjects were collected as well [4]. To recruit subjects for an experiment, following the requirements and within the practical constraints, researchers often try to ensure a sparse distribution of key measurements, e.g. the stature, the body mass, among the participants to reduce the specificity of the target group and ensure the quality of the research [5, 6]. For instance, in evaluating comfort of an economy seat, researchers tried to select 97 test subjects out of 125 applicants

*Address for correspondence: Gerbera Vledder, Faculty of Industrial Design Engineering, Delft University of Technology, Landbergstraat 15, 2628CE, Delft, The Netherlands. E-mail: g.vledder@tudelft.nl.

based on the self-reported stature and body mass [7]. Besides, a better estimation of the actual stature and body mass based on self-reported data might accelerate the process of ergonomics experiments and reduce the cost. Measuring anthropometrics consumes valuable time and manpower. Extensive measurements are seen as “tedious and time consuming” [8]. For improving the accuracy, researchers also need to conduct prior-training sessions and repeatable measurements are often taken [9], which both cost extra time and efforts.

Literature supports the theory that the body mass can be underreported and the stature can be overreported in different contexts [10]. For instance, women under-reported their mass by a mean of 0.91 kg in the Health Initiative Observational Study [11], and men overreported their height by 1.22 cm on average in National Health and Nutrition Examination Survey [12]. However, there are also reports indicating that such self-reported body mass and stature were accurate enough in different studies, e.g. Hodge et al. suggested that BMI computed from self-reported weight and height is a valid measure in men and women across different socio-demographic groups [13]. Kee et al. also reported the high correlation between self-report mass and stature with direct measurements in the adolescents group [14].

The accuracy of the self-reported stature and body mass is influenced by many factors, e.g., the types of survey, the social-demographic contexts, the sex. The existence of self-stigma regarding the body mass [15] could influence the subjective ratings in ergonomic studies as well. In research on comparing three different survey data, Flegal et al. found that self-reported height, weight, BMI, and obesity prevalence were not identical across the National Health and Nutrition Examination Survey, the National Health Interview Survey, and the Behavioral Risk Factor Surveillance System, particularly for women [16]. Gugushvili & Jarosz also found that women living in rural areas were likely to overestimate their height [17]. Maukonen et al. also showed that for overweight and obese participants, the bias was generally higher than those of average weight. Regarding cultural background, the bias was larger in North America, while in Asian studies the bias seemed to be lower [10]. For instance, Xie et al. found that the stature was overestimated at an average of 0.42 cm, but there were no significant differences regarding the body mass and blood pressures between self-reported and measured data in Hong Kong [18]. Meanwhile, the stature and the body mass of a subject are not static during the day.

Vuvor and Harrison found that the mean stature variation from 7am to 7pm was about 1.61 cm [19]. The average mass fluctuation of a person in a short term, e.g. a day or several days, can also be 1 to 2 kg [20].

While the accuracy, the reliability and the associated factors are still in debate, the self-reported measurements are the easiest, or sometimes the only way, for acquiring the basic anthropometric parameters in the survey and selecting participants in ergonomics research. In this paper, we aim at building models to estimate the actual stature and body mass based on of the self-reported data of adults recruited in the Netherlands in the context of ergonomic research. The outcomes might help ergonomics researchers in using self-reported stature and body mass for survey and recruitment in ergonomics studies.

2. Materials and methods

We collected data from 4 ergonomics experiments conducted in the Faculty of Industrial Design Engineering, Delft University of Technology. All experiments focused on evaluating comfort of participants, but in different contexts as: Exp.1) flying with a turboprop aircraft (November, 2021), Exp.2) experiencing jet and turboprops noise (February, 2022), Exp.3) sitting with different postures (March, 2022) and Exp.4) sitting in train and aircraft seats (June, 2022). In those experiments, participants self-reported their stature and body mass prior to being measured, of which they were not informed beforehand. For instance, in Exp.4 an ergonomics research context was created on the topic of evaluating the comfort experience of sitting in train and aircraft seats (see Fig. 1). A large room divider was placed in the middle and anthropometric measurement tools were placed at another side of the room divider. Participants were invited to the experiment in different timespans of a day across 3 weeks. They were instructed to approach from the ergonomics research area where the anthropometric measurement tools were invisible to them. After acquiring the informed consent, participants were asked to fill in a questionnaire about their sex, age, nationality, stature and body mass. Then a researcher led them to another side of the room divider where the basic anthropometric measurements were taken.

All study protocols were approved by the human research ethical committee at Delft University of Technology. Respectively under ID number 1823



Fig. 1. Setup of a typical experiment.

(Exp.1), 1953 (Exp.2), 1228 (Exp.3), and 2248 (Exp.4). It is worth mentioning that participants were free to withdraw at any moment of the experiments and his/her data (if any) was destroyed on-site. From the 4 experiments, we collected 76, 15, 28 and 130 valid samples, resulting a total of 249 records. In Exp.1 approximately half of the participants were measured between 08 : 30 and 09 : 30, the other half was measured between 12 : 30 and 13 : 30. During Exp.2 all measurements were made between 13 : 00 and 13 : 30. For Exp.3 measurements were made in the afternoon. Exp.4 started at 09 : 00 and lasted until 18 : 00. G*Power calculation indicated that for identifying small effects (0.3) of paired samples using Wilcoxon signed rank test in a two-tails setup, 154 samples were needed at the power of 0.95. The 249 valid records were analyzed by statistical tools using a self-developed Python program. All participants took off their shoes and jacket and no extra adjustments, e.g. weight of clothing [21], were introduced as interviews of the participants revealed mixed scenarios regarding self-weighting. Based on statistical analysis results, different groups were highlighted.

3. Results

Table 1 presents the descriptive statistics of each experiment result as well as a whole. All participants were adults and their mean age is about 28 years. There are high correlations between the self-reported and the measured stature (0.981), and between the self-reported and the measured body mass (0.967). However, people overreported their stature by 1.31 cm on average, and underreported their body mass by a mean of 1.45 kg. Subsequently, the difference of BMIs based on self-reported data and the measured data is about 0.82 kg/m^2 . Regarding individual experiments, Exp.2 and 3 were targeting at the international population and in Exp.1 and 4, the Dutch population was the majority. Though populations differ, Mann-Whitney U test indicated that there was no statistically significant difference ($p > 0.01$) regarding BMIs calculated based on the self-reported data and the measured data between Dutch and international populations. Therefore, all experiment data were combined together in the following analysis.

Bland-Altman plots (Fig. 2) on the stature and the body mass affirm the differences. The stature difference is 0.74% higher than the mean value while the body mass difference is about 1.93% lower than the mean. Normality test results indicated that the distribution of the differences does not follow the normal distribution. Wilcoxon signed rank test results showed that the differences of the self-reported and the measured stature ($p < 0.01$), body mass ($p < 0.01$) and BMI ($p < 0.01$) are statistically significant.

Figure 3 presents two regression models where the 95% prediction and 95% confidence intervals are highlighted as well. In the model, the measured stature can be calculated from the self-reported

Table 1
Descriptive statistics of the collected data

	Exp. 1	Exp. 2	Exp. 3	Exp. 4	All experiments
Total number	76	15	28	130	249
Gender (female/male)	49/27	9/6	13/15	66/64	137/112
Age	33.42 ± 14.02	26.33 ± 3.66	27.82 ± 2.89	25.15 ± 9.15	28.04 ± 10.87
Nationality (NL/INT)	32/44	4/11	1/27	93/37	130/119
Mass (kg, self-reported)	71.40 ± 12.40	69.87 ± 16.07	62.29 ± 9.98	71.54 ± 12.57	70.36 ± 12.75
Mass (kg, measured)	73.11 ± 13.33	71.73 ± 17.05	62.14 ± 10.66	73.13 ± 12.97	71.81 ± 13.49
Mass difference (kg)	1.71 ± 3.19	1.86 ± 4.45	-0.14 ± 3.33	1.59 ± 3.40	1.45 ± 3.43
Stature (cm, self-reported)	177.31 ± 10.63	171.00 ± 10.45	168.02 ± 6.60	177.19 ± 10.30	175.82 ± 10.50
Stature (cm, measured)	176.24 ± 10.49	169.76 ± 9.28	166.03 ± 6.94	175.87 ± 9.90	174.51 ± 10.28
Stature difference (cm)	-1.06 ± 2.29	-1.24 ± 3.40	-1.99 ± 1.53	-1.31 ± 1.78	-1.31 ± 2.05
BMI (kg/m^2 , based on self-reported data)	22.61 ± 2.74	23.70 ± 3.95	21.99 ± 2.78	22.66 ± 2.54	22.63 ± 2.73
BMI (kg/m^2 , based on measured data)	23.43 ± 3.11	24.64 ± 4.10	22.49 ± 3.24	23.53 ± 2.92	23.45 ± 3.10
BMI difference (kg/m^2)	0.83 ± 0.95	0.94 ± 1.66	0.50 ± 1.23	0.87 ± 1.17	-0.82 ± 1.15




Fig. 2. Bland-Altman plots of the self-reported stature (a) and body mass (b) and regarding the means of using the two methods.

stature as:

$$\begin{aligned} \text{Measured Stature} &= 0.96375 * \\ &\text{Self Reported Stature} + 5.36 \end{aligned} \quad (1)$$

with a root mean square error (RMSE) of 2.24 CM. For the body mass,

$$\begin{aligned} \text{Measured Mass} &= 1.04469 * \\ &\text{Self Reported Mass} - 1.48 \end{aligned} \quad (2)$$

and the RMSE is 3.12 Kg.

Though all experiments were conducted in the context of ergonomic research, different factors, e.g. sex, nationality and BMI, might have influenced the accuracy of the self-reported stature and the body mass. We categorized the data by sex, nationalities and BMI where for BMI, the Centers for Disease Control and Prevention (CDC) recommendation (BMI < 18.5 is underweight, between 18.5 and 24.9 is healthy, between 25.0 and 29.9 is overweight and larger than 30.0 is obesity) [22] was adopted. Mann-Whitney U test results indicated that there was not statistically significant difference in those groups except the mass difference of the BMI < 25 group (180 subjects, mean = -0.76 kg) was significantly smaller




Fig. 3. The regression model (red lines) regarding the data.

($p < 0.01$) than the BMI ≥ 25 group (69 subjects, mean = -3.24 kg) as Fig. 4.

There are statistically significant differences of the measured stature and the body mass between the Dutch and the international groups. The mean stature of the Dutch group is 177.79 ± 9.17 cm versus 170.93 ± 10.26 cm of the international group. Regarding the body mass, the values are 75.23 ± 13.31 kg and 68.03 ± 13.75 kg for Dutch and international groups, respectively. However, we did not find statistically significant differences between the self-reported data and the measured data for both groups as shown in Fig. 5.

Literature indicated that sex and BMI are two important factors of the accuracy of the self-reported stature and body mass, we categorized all data using these two criteria as Table 2. It can be found that for the self-reported stature, all groups are within the accuracy of 2 cm where women are slightly bet-




Fig. 4. Measured stature and body mass vs the self-reported body mass and stature regarding different BMI groups.




Fig. 5. Measured stature and body mass vs the self-reported body mass and stature regarding Dutch & International groups.

ter than men (1.02 cm vs 1.55 cm). For the body mass, in general men are slightly more accurate than women (-1.24 vs -1.70 kg), but for the BMI 18.5–24.9 groups, both men and women are quite accurate. The largest difference was observed for men and women with BMI ≥ 25 groups, where the mean body mass differences were found as -2.95 kg and -3.5 kg, respectively. Though the mean difference of women is higher than that of the men, it is not statistically significant ($p > 0.01$). Regarding nationality, Dutch men and women BMI ≥ 25 groups underreported the mass about 2.70 and 3.66 kg, and for the International BMI ≥ 25 groups, the values are 3.50 and 3.89 kg, respectively.

4. Discussion

In this paper, we investigated the accuracy of self-reported stature and body mass regarding the anthropometric measurements within the ergonomics research context in the Netherlands. Experiment results support the declaration that people slightly overreport their stature and underreport their body mass, which is in accordance with some studies in Table 3. This is especially true for people with

BMI ≥ 25 , where statistically significant difference regarding the self-reported and the measured body mass was identified with a mean difference of -3.24 kg. Compared to a previous study on the Dutch population [21], the difference of the stature and the body mass are slightly larger, which might be that we did not introduce extra adjustments. However, it is in accordance with the review conducted by Maukonen et al. [10], where large BMI population tends to report less on their body mass.

Although the time of measurements for all experiment is known, the time and date of self-reported measurements are unknown. Vuvor & Harrison found that the daily stature variation of adults aged ≥ 30 years is as high as 1.61 cm [19]. Reilly et al. indicated that the peak variation of the stature can be 1.93 cm or 1.1% of the overall stature [27]. 71% of the height gained during the night was achieved in the first half of the night's sleep, and over 50% of the height loss in a day happened in the first hour of rising, and 80% was lost within 3 hours of arising. Considering the time of the experiment, e.g. Exp.2 and 3 took place after 3 hours of arising, the main decrease in stature changes already happened. The identified mean difference of the stature is 1.31 cm, which is slightly larger than 50% of the peak variation.

Table 2
Differences of self-reported mass and height regarding measurement for different groups

BMI	All groups	<18.5	18.5–24.9	25–29.9	>30
Men					
No.	137	1	92	37	7
Mean Stature	180.37 ± 8.50	n/a	179.81 ± 8.71	182.51 ± 7.52	178.57 ± 8.75
Mean Mass	77.49 ± 13.26	n/a	71.43 ± 8.71	89.13 ± 8.32	101.24 ± 10.80
Mean BMI	23.74 ± 3.26	n/a	22.03 ± 1.61	26.73 ± 1.35	31.68 ± 1.36
Stature Diff.	1.55 ± 2.16	n/a	1.38 ± 2.24	1.92 ± 2.03	1.71 ± 1.90
Mass Diff.	-1.24 ± 3.85	n/a	-0.50 ± 3.94	-3.03 ± 2.79	-2.53 ± 4.03
BMI Diff.	-0.77 ± 1.19	n/a	-0.48 ± 1.07	-1.45 ± 1.05	-1.27 ± 1.71
Stature Diff. in %	0.87% ± 1.21%	n/a	0.78% ± 1.28%	1.05% ± 1.08%	0.93% ± 1.05%
Mass Diff. in %	-1.28% ± 5.11%	n/a	-0.56% ± 5.40%	-3.34% ± 2.98%	-2.28% ± 3.86%
BMI Diff. in %	-0.90% ± 1.56%	n/a	-0.66% ± 1.54%	-1.60% ± 1.13%	-1.15% ± 1.66%
Women					
No.	112	2	85	22	3
Mean Stature	167.34 ± 7.30	173.40 ± 9.90	166.93 ± 7.36	167.79 ± 6.37	171.60 ± 11.75
Mean Mass	64.85 ± 10.11	55.25 ± 6.43	61.32 ± 6.95	75.64 ± 6.58	92.20 ± 12.23
Mean BMI	23.11 ± 2.89	18.34 ± 0.05	21.97 ± 1.66	26.83 ± 1.25	31.22 ± 0.64
Stature Diff.	1.02 ± 1.88	-1.40 ± 1.41	1.02 ± 1.98	1.03 ± 1.47	2.40 ± 0.95
Mass Diff.	-1.70 ± 2.82	-0.25 ± 0.64	-1.13 ± 2.67	-3.50 ± 2.42	-5.53 ± 2.74
BMI Diff.	-0.89 ± 1.10	0.20 ± 0.51	-0.68 ± 1.00	-1.56 ± 1.01	-2.61 ± 0.95
Stature Diff. in %	0.61% ± 1.14%	-0.79% ± 0.77%	0.62% ± 1.21%	0.61% ± 0.88%	1.39% ± 0.49%
Mass Diff. in %	-2.43% ± 4.30%	-0.52% ± 1.21%	-1.77% ± 4.38%	-4.67% ± 3.36%	-5.87% ± 2.45%
BMI Diff. in %	-1.31% ± 1.67%	0.31% ± 0.90%	-1.09% ± 1.67%	-2.09% ± 1.42%	-2.82% ± 0.83%

Table 3
Comparison of this study and some previous studies

Reference	Stature difference (in cm/%)		Mass difference (in kg/%)		Context	Country
	Men	Women	Men	Women		
[13]	2.08	1.07	-0.726	-1.406	Health Insurance Study	USA
[4]	1.23	0.60	-1.85	-1.40	European Prospective Investigation into Cancer and Nutrition	UK
[14]	1.7	1.1	-0.4	-1.1	The CAESAR Project 1999–2000	Italy, NL and North America
[11]	0.8	0.8	-0.9	-0.8	Amsterdam Health Monitor 2004	NL
[15]	0.3	2.3	-1.00	-2.8	Survey in a music festival	Australia
[16]	0.7	-0.6	-1.3	-1.4	European Health Examination Survey	12 EU countries
[5]	0.48	0.16	-1.55	-0.88	Cancer Prevention Study	USA
This study	1.55/0.87%	1.02/0.61%	-1.24/-1.28%	-1.70/-2.43%	Ergonomics studies	NL

Regarding the body mass, Turicchi et al. found that there are weekly, seasonal and Christmas patterns where within a week, the mean body mass fluctuations are 0.35% [28]. Whigham et al. indicated that the average clothing weight throughout the year was significantly greater in men (1.2 ± 0.3 kg) than in women (0.8 ± 0.3 kg) [29]. Besides, the timing for defecation (median 0.128 kg/cap/day) and urination (median 1.42 L/cap/day) might also influence of the

measured body mass [30]. Considering the uncertainty in the self-measurement time and conditions, it can be assumed that for groups with BMI < 18.5, the self-reported body mass is within the fluctuation ranges. However, for BMI \geq 25 groups, the difference between the self-report body mass and the measurement is larger.

In the measurement results, the mean heights are 180.37 cm and 167.34 cm for men and women,

respectively. These are close to the P50 data in the Dutch anthropometric database [31]. The smaller standard deviations of women on the self-reported body mass indicated that women might know their actual body mass better than men, as the self-weight frequency of women is higher than that of men [32, 33].

With the underreported mass, 35% (24 of 69) of the participants in the $BMI \geq 25$ group (mean $BMI = 25.9$) lowered their reported BMI to the range of $BMI < 25$ (mean = 24.11). The reasons that the self-reported body mass of participants in this group is lower than the measurement might be complicated. Besides sex, nationality, Althubaiti summarized that self-reported bias can also arise from social desirability, recall period, sampling approach, or selective recall these aspects [34]. The accordance with literature indicates that the ergonomics research context has little influence on the bias. For recall period, sampling approach, or selective recall, previous research indicated that the self-weighing frequency of $BMI \geq 85$ th percentile is even higher than that of the $15 \leq BMI < 85$ th percentile group [33]. Social desirability might be the most influential factor on the bias. The body shape is associated with the self-image of a person, and a positive self-image is an important aspect for people to recognize the assets, the limitations and the potentials, and keep positive motivation on tasks and liabilities. For this, people tend to think towards the positive side on their self-image rather than generate it based on facts, e.g. people are more accurate while their focus is on the external observers [35]. This might be more obvious for women, as they are more critical of their body shape [36]. Therefore, participants with larger BMI might want to report less of their body mass to maintain a positive self-image on the body shape, as in most cases the stature is more a nature and more visible than the body mass.

In summary, participants slightly overreported their height and underreported their mass. The stature differences between the self-reported and measured data slightly exceed the daily fluctuation range of a person. The differences of the self-reported body mass and measured results of the $BMI < 24.9$ group is within the daily fluctuation range. But for the $BMI \geq 25$ group, the difference is larger. For ergonomics studies where the anthropometric measurements were not possible, e.g. in survey, in recruitment, it is suggested that the self-reported stature and the body mass can be adjusted using the linear regression models (Eq.1 and 2). In this case, the RMSEs of the differences in the stature and the

mass can be reduced from 2.43 cm to 2.02 cm and from 3.71 kg to 3.42 kg, respectively. This is especially useful for the body mass of the $BMI \geq 25$ group where the RMSE is reduced from 4.37 to 3.06 kg. In the experiment, when there are constraints regarding both/either manpower and/or time, the proposed model can be used as a backup tool. However, ergonomics researchers shall be aware of the large RMSEs and risks of peak variations in using this technique, as the peak variations of the self-reported statures are -5.5% and 7% and for the self-report body mass, they are -12.8% and 22.9%.

This study has several limitations. Most of the International group participants were studying/working in the Netherlands. We did not find significant differences of self-reported data between Dutch and International groups. Maukonen et al. reported a difference between Asian and American studies, which means a comparison study in other countries might find more social-cultural factors that influence the accuracy of self-reported stature and body mass [10]. Though researchers paid extensive efforts in recruiting, the mean age (28.04 years) of participants was young and the number of participants, especially in the $BMI < 18.5$ and $BMI > 30$ groups, was low compared to other studies that were mainly based on national surveys. This prevented the use of advanced modeling tools to make a more accurate estimation based on more parameters, e.g. gender, measurement time in a day, time duration from the last self-measurement. However, the context of the study is ergonomics research, which casts a new view toward the bias of participants in self-reporting their stature and body mass.

5. Conclusion

In this paper, we compared the self-reported stature and body mass regarding the measurements in the context of ergonomics research, and try to provide a tool for ergonomic researchers and practitioners for estimating the actual stature and body mass based on self-reported data. Experiment results supported the findings reported in the literature that people slightly overreport their stature and underreport their body mass, especially in the $BMI \geq 25$ group. It is suggested that in ergonomic research especially in the survey and recruitment where conducting anthropometric measurement is not possible, or too expensive regarding both manpower and time, researchers can use the proposed models, with the understanding of

the limitations, to estimate the actual stature and body mass based on self-reported data, especially for the BMI ≥ 25 group.

Ethical considerations

All study protocols were approved by the human research ethical committee at Delft University of Technology (ID numbers 1823 (Exp.1), 1953 (Exp.2), 1228 (Exp.3), and 2248 (Exp.4).

Informed consent

All participants involved in this study signed an informed consent form before participating in this study. The informed consent followed the guidelines of the human research ethical committee at Delft University of Technology and was approved by this same committee.

Acknowledgments

The authors want to express their appreciation to all participants who joined the experiment at the Faculty of Industrial Design Engineering, Delft University of Technology.

Conflict of interest

The authors declare that they have no conflict of interest.

Funding

Experiments 1 and 2 received funding from the Clean Sky 2 Joint Undertaking (JU) under grant agreement 945583 (ComfDemo). Dr. Halil Kılıç is supported by the Scientific and Technological Research Institution of Turkey (TÜBİTAK) funding programme: 2219 International Postdoctoral Research Fellowship. Ms. Xinhe Yao is supported by the China Scholarship Council (201907720095).

References

- [1] Turrell G, Nathan A, Burton NW, Brown WJ, McElwee P, Barnett AG, et al. Cohort Profile: HABITAT—a longitudinal multilevel study of physical activity, sedentary behaviour and health and functioning in mid-to-late adulthood. *Int J Epidemiol.* 2021;50(3):730-1h.
- [2] Watson N, Wooden MP. The HILDA Survey: a case study in the design and development of a successful Household Panel Survey. *Longit Life Course Stud.* 2012;3(3): 369-81.
- [3] Liu Z, Rotte T, Anjani S, Vink P. Seat pitch and comfort of a staggered seat configuration. *Work.* 2021;68(s1):S151-9.
- [4] Srinivasan V, Vishnuram S, Suganthirababu P, Kumaresan A, Divyalaxmi P, Saranya Devi M, et al. Impact of COVID-19 online classes on morningness-eveningness personality among adult college students: A survey study. *Work.* 2021;70(3):695-700.
- [5] Roebuck JA, Kroemer KHE, Thomson GW. *Engineering anthropometry methods.* John Wiley & Sons, Inc.; 1975.
- [6] Anjani S, Song Y, Hou T, Ruiter IA, Vink P. The effect of 17-inch-wide and 18-inch-wide airplane passenger seats on comfort. *Int J Ind Ergon.* 2021;82:103097.
- [7] Vink P, Vledder G, Song Y, Herbig B, Reichherzer AS, Mansfield N. Aircraft interior and seat design: priorities based on passengers' opinions. *Int J Aviat Aeronaut Aerosp [Internet].* 2022 Jan 1 [cited 2022 Apr 28];9(1):3. Available from: <https://commons.erau.edu/ijaaa/vol9/iss1/3>
- [8] Dianat I, Molenbroek J, Castellucci HI. A review of the methodology and applications of anthropometry in ergonomics and product design. <https://doi-org.tudelft.idm.oclc.org/101080/0014013920181502817> [Internet]. 2018 Dec 2 [cited 2023 Jan 18];61(12):1696-720. Available from: <https://www.tandfonline-com.tudelft.idm.oclc.org/doi/abs/10.1080/00140139.2018.1502817>
- [9] Bravo G, Bragança S, Arezes PM, Molenbroek JFM, Castellucci HI. A literature review of anthropometric studies of school students for ergonomics purposes: Are accuracy, precision and reliability being considered? *Work.* 2018;60(1):3-17.
- [10] Maukonen M, Männistö S, Tolonen H. A comparison of measured versus self-reported anthropometrics for assessing obesity in adults: a literature review. *Scand J Public Health.* 2018;46(5):565-79.
- [11] Luo J, Thomson CA, Hendryx M, Tinker LF, Manson JE, Li Y, et al. Accuracy of self-reported weight in the Women's Health Initiative. *Public Health Nutr.* 2019;22(06):1019-28.
- [12] Spencer EA, Appleby PN, Davey GK, Key TJ. Validity of self-reported height and weight in 4808 EPIC-Oxford participants. *Public Health Nutr.* 2002;5(4):561-5.
- [13] Hodge JM, Shah R, McCullough ML, Gapstur SM, Patel AV. Validation of self-reported height and weight in a large, nationwide cohort of U.S. adults. Siegel R, editor. *PLoS One.* 2020;15(4):e0231229.
- [14] Kee CC, Lim KH, Sumarni MG, Teh CH, Chan YY, Nuur Hafizah MI, et al. Validity of self-reported weight and height: a cross-sectional study among Malaysian adolescents. *BMC Med Res Methodol.* 2017;17(1):85.
- [15] Pakpour AH, Tsai MC, Lin YC, Strong C, Latner JD, Fung XCC, et al. Psychometric properties and measurement invariance of the Weight Self-Stigma Questionnaire and Weight Bias Internalization Scale in children and adolescents. *Int J Clin Heal Psychol.* 2019;19(2): 150-9.

- [16] Flegal KM, Ogden CL, Fryar C, Afful J, Klein R, Huang DT. Comparisons of Self-Reported and Measured Height and Weight, BMI, and Obesity Prevalence from National Surveys: 1999–2016. *Obesity*. 2019;27(10):1711-9.
- [17] Gugushvili A, Jarosz E. Inequality, validity of self-reported height, and its implications for BMI estimates: An analysis of randomly selected primary sampling units' data. *Prev Med Reports*. 2019;16:100974.
- [18] Xie YJ, Ho SC, Liu ZM, Hui SSC. Comparisons of Measured and Self-Reported Anthropometric Variables and Blood Pressure in a Sample of Hong Kong Female Nurses. Wang J, editor. *PLoS One*. 2014;9(9):e107233.
- [19] Vuvor F, Harrison O. A study of the diurnal height changes among sample of adults aged 30 years and above in Ghana. *Biomed Biotechnol Res J*. 2017;1(2):113.
- [20] Bhutani S, Kahn E, Tasali E, Schoeller DA. Composition of two-week change in body weight under unrestricted free-living conditions. *Physiol Rep*. 2017;5(13):e13336.
- [21] Dijkshoorn H, Ujcic-Voortman JK, Viet L, Verhoeff AP, Uitenbroek DG. Ethnic variation in validity of the estimated obesity prevalence using self-reported weight and height measurements. *BMC Public Health*. 2011;11(1):408.
- [22] CDC. About Adult BMI. 2022.
- [23] Stewart AL. The reliability and validity of self-reported weight and height. *J Chronic Dis*. 1982;35(4):295-309.
- [24] Krul AJ, Daanen HAM, Choi H. Self-reported and measured weight, height and body mass index (BMI) in Italy, the Netherlands and North America. *Eur J Public Health*. 2011;21(4):414-9.
- [25] Bowring AL, Peeters A, Freak-Poli R, Lim MS, Gouillou M, Hellard M. Measuring the accuracy of self-reported height and weight in a community-based sample of young people. *BMC Med Res Methodol*. 2012;12(1):175.
- [26] Tolonen H, Koponen P, Mindell JS, Mannisto S, Giampaoli S, Dias CM, et al. Under-estimation of obesity, hypertension and high cholesterol by self-reported data: comparison of self-reported information and objective measures from health examination surveys. *Eur J Public Health*. 2014;24(6):941-8.
- [27] Reilly T, Tyrell A, Troup JDG. Circadian variation in human stature. *Chronobiol Int*. 1984;1(2):121-6.
- [28] Turicchi J, O'Driscoll R, Horgan G, Duarte C, Palmeira AL, Larsen SC, et al. Weekly, seasonal and holiday body weight fluctuation patterns among individuals engaged in a European multi-centre behavioural weight loss maintenance intervention. Ferrer R, editor. *PLoS One*. 2020;15(4):e0232152.
- [29] Whigham LD, Schoeller DA, Johnson LK, Atkinson RL. Effect of clothing weight on body weight. *Int J Obes*. 2013;37(1):160-1.
- [30] Rose C, Parker A, Jefferson B, Cartmell E. The Characterization of Feces and Urine: A Review of the Literature to Inform Advanced Treatment Technology. *Crit Rev Environ Sci Technol*. 2015;45(17):1827-79.
- [31] DINED. Anthropometric database TU Delft Ergonomics. Dined.NL. 2011.
- [32] Gavin KL, Linde JA, Pacanowski CR, French SA, Jeffery RW, Ho YY. Weighing frequency among working adults: Cross-sectional analysis of two community samples. *Prev Med Reports*. 2015;2:44-6.
- [33] Hahn SL, Pacanowski CR, Loth KA, Miller J, Eisenberg ME, Neumark-Sztainer D. Self-weighing among young adults: who weighs themselves and for whom does weighing affect mood? A cross-sectional study of a population-based sample. *J Eat Disord*. 2021;9(1):37.
- [34] Althubaiti A. Information bias in health research: definition, pitfalls, and adjustment methods. *J Multidiscip Healthc*. 2016;211.
- [35] Bašić Z, Quercia S. The influence of self and social image concerns on lying. *Games Econ Behav*. 2022;133:162-9.
- [36] Buote VM, Wilson AE, Strahan EJ, Gazzola SB, Papps F. Setting the bar: Divergent sociocultural norms for women's and men's ideal appearance in real-world contexts. *Body Image*. 2011;8(4):322-34.