

Anaerobic carboxydotrophy in sulfur-respiring haloarchaea from hypersaline lakes

Sorokin, Dmitry Y.; Merkel, Alexander Y.; Messina, Enzo; Tugui, Claudia; Pabst, Martin; Golyshin, Peter N.; Yakimov, Michail M.

DOI

[10.1038/s41396-022-01206-x](https://doi.org/10.1038/s41396-022-01206-x)

Publication date

2022

Document Version

Final published version

Published in

ISME Journal

Citation (APA)

Sorokin, D. Y., Merkel, A. Y., Messina, E., Tugui, C., Pabst, M., Golyshin, P. N., & Yakimov, M. M. (2022). Anaerobic carboxydotrophy in sulfur-respiring haloarchaea from hypersaline lakes. *ISME Journal*, *16*(6), 1534-1546. <https://doi.org/10.1038/s41396-022-01206-x>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Green Open Access added to TU Delft Institutional Repository

'You share, we take care!' - Taverne project




<https://www.openaccess.nl/en/you-share-we-take-care>

Otherwise as indicated in the copyright section: the publisher is the copyright holder of this work and the author uses the Dutch legislation to make this work public.

ARTICLE



Anaerobic carboxydrotrophy in sulfur-respiring haloarchaea from hypersaline lakes

Dimitry Y. Sorokin^{1,2}, Alexander Y. Merkel¹, Enzo Messina³, Claudia Tugui², Martin Pabst², Peter N. Golyshin⁴ and Michail M. Yakimov⁵

© The Author(s), under exclusive licence to International Society for Microbial Ecology 2022

Anaerobic carboxydrotrophy is a widespread catabolic trait in bacteria, with two dominant pathways: hydrogenogenic and acetogenic. The marginal mode by direct oxidation to CO₂ using an external *e*-acceptor has only a few examples. Use of sulfidic sediments from two types of hypersaline lakes in anaerobic enrichments with CO as an *e*-donor and elemental sulfur as an *e*-acceptor led to isolation of two pure cultures of anaerobic carboxydrotrophs belonging to two genera of sulfur-reducing haloarchaea: *Halanaeroarchaeum* sp. HSR-CO from salt lakes and *Halalkaliarchaeum* sp. AArc-CO from soda lakes. Anaerobic growth of extremely halophilic archaea with CO was obligatory depended on the presence of elemental sulfur as the electron acceptor and yeast extract as the carbon source. CO served as a direct electron donor and H₂ was not generated from CO when cells were incubated with or without sulfur. The genomes of the isolates encode a catalytic Ni₂Fe-CODH subunit *CooS* (distantly related to bacterial homologs) and its Ni-incorporating chaperone *CooC* (related to methanogenic homologs) within a single genomic locus. Similar loci were also present in a genome of the type species of *Halalkaliarchaeum* closely related to AArc-CO, and the ability for anaerobic sulfur-dependent carboxydrotrophy was confirmed for three different strains of this genus. Moreover, similar proteins are encoded in three of the four genomes of recently described carboxydrate-utilizing sulfur-reducing haloarchaea belonging to the genus *Halapricum* and in two yet undescribed haloarchaeal species. Overall, this work demonstrated for the first time the potential for anaerobic sulfur-dependent carboxydrotrophy in extremely halophilic archaea.

The ISME Journal (2022) 16:1534–1546; <https://doi.org/10.1038/s41396-022-01206-x>

INTRODUCTION

Prokaryotic carboxydrotrophy is based on two different types of unrelated CO-dehydrogenase enzymes (CODH), namely (Cu,Mo,FeS,flavin)-CODH (CoxLMS), which is present in aerobic or denitrifying prokaryotes, and (Ni,FeS)-CODH (*CooS*) present in anaerobic carboxydrotrophs [1–5]. Aerobic carboxydrotrophs are predominantly proteobacteria [6], although members of other phyla have recently been found to aerobically oxidize CO, including *Chloroflexi* (growth and oxidation) [7] and the extremely halophilic archaea of the class *Halobacteria* (oxidation without growth) [8]. Anaerobic carboxydrotrophy is mainly present in three groups of secondary anaerobic prokaryotes: (i) acetogenic bacteria and archaea via the Wood-Ljungdahl pathway, resulting in the production of CO₂, acetate and/or ethanol; (ii) methanogenic archaea via the archaeal variety of the Wood-Ljungdahl pathway, leading to formation of methane or acetate; and (iii) hydrogenogenic carboxydrotrophic bacteria and archaea, producing H₂ and CO₂ [7, 9, 10].

The microbial carboxydrotrophy is poorly investigated in hypersaline habitats. Until now, halophilic carboxydrotrophy has only been observed aerobically by organisms classified within two closely related haloalkaliphilic *Gammaproteobacteria* genera: *Alkalispirillum* and *Alkalilimnicola* [11, 12]. The potential for

oxidation of CO at low concentrations using O₂, nitrate or perchlorate (albeit without growth) was shown for several members of extremely halophilic archaea belonging to genera *Halorubrum*, *Natronorubrum*, *Haloarcula* and *Halobaculum* [8, 13]. These bacteria and archaea have aerobic type of (Mo,Cu)-CODH. However, the utilization of CO as the energy source for aerobic carboxydrotrophic growth was proven only for the *Alkalispirillum/Alkalilimnicola* group [12].

Since protons are the final electron acceptors of hydrogenogenic pathway, anaerobic carboxydrotrophy would be extremely unlikely in soda lake environment, while both acetogenic and methanogenic CO utilization can be feasible. Indeed, for extremely haloalkaliphilic microorganisms, hydrogenogenic anaerobic carboxydrotrophy has never been reported.

Although for some hydrogenogenic carboxydrotrophs the addition of an external electron acceptor such as sulfur, thiosulfate, or sulfate led to a partial redirection of CO-derived electrons from protons to a more favorable acceptor, this did not completely stop the H₂ formation [14–17]. The same seems to be true for those carboxydrotrophic acetogens that can utilize electron acceptors other than CO₂. In this case, some CO-derived electrons is diverted to reduction of the external acceptor, as was shown for the euryarchaeon *Archaeoglobus fulgidus* in the

¹Winogradsky Institute of Microbiology, Federal Research Centre of Biotechnology, Russian Academy of Sciences, Moscow, Russia. ²Department of Biotechnology, Delft University of Technology, Delft, The Netherlands. ³IRBIM-CNR, Spianata S.Raineri 86, 98122 Messina, Italy. ⁴School of Natural Sciences, Bangor University, Gwynedd LL57 2UW, UK. ⁵Institute of Polar Sciences, ISP-CNR, Messina, Italy. ✉email: soroc@inmi.ru

Received: 3 October 2021 Revised: 3 January 2022 Accepted: 27 January 2022

Published online: 7 February 2022

presence of sulfate [18]. However, there is hardly any published evidence for anaerobic microorganisms that could directly oxidize CO to CO₂ in anaerobic respiratory mode as the only energy generation process. One of the possible candidates is the euryarchaeal genus *Thermococcus*, which is apparently capable of both proton-dependent hydrogenogenic carboxydrotrophy and lithoheterotrophic growth by direct oxidation of CO with elemental sulfur as an electron acceptor. The genome of *Thermococcus* sp. AM4 contains two different CODH loci, probably responsible for different modes of CO oxidation: one is associated with membrane-bound hydrogenase (MBH) and the other with a flavin disulfide oxidoreductase, a putative electron acceptor. Both clusters also contain CooF-like ferredoxins, which provide immediate electron transfer during CO oxidation. A close relative of AM4, *T. gammatolerans*, can only grow with CO in the presence of sulfur and has a single *cooS/cooF/cooC* locus in the genome not linked to any recognizable electron transfer complexes [19–21]. Further electron transfer from CODH to sulfur in *Thermococcus* species is still unresolved, although the genomes of both seem to encode the membrane-bound MBX complexes (not associated with CODH loci) with a proven potential for ferredoxin-dependent polysulfide reductase activity in thermophilic archaea [22]. Interestingly, two out of five different CooS-coding loci present in the CO-utilizing sulfate-reducing genus *Desulfofundulus* are structurally very similar to the hydrogenase-independent *Thermococcus* CODH locus and are likely responsible for the oxidation of CO with sulfate as an electron acceptor in these bacteria [17].

So far there is no evidence of anaerobic carboxydrotrophy (based on the Ni₂Fe-CODH) for extreme halophiles living in hypersaline pH neutral habitats. There are only two examples of anaerobic haloalkaliphilic members of *Firmicutes* utilizing CO during acetogenic growth: two species of the genus *Alkalibaculum* (the low salt and alkali-tolerant *A. bachii* and the moderately haloalkaliphilic *A. sporogenes*) - a member of the family *Eubacteriaceae* [23–25], and *Natranaerofaba carboxydovora*, - an extremely haloalkaliphilic member of the class *Natranaerobii*, which grows optimally in hypersaline soda brines at pH 10 and 50 °C [26]. It was enriched in a consortium with a triple extremophilic methyl-reducing methanogen of the genus *Methanonatronarchaeum*, previously found in hypersaline soda lakes [27]. In fact, methane was formed in this enrichment from formate, released as one of the products of CO conversion by acetogens.

The present work targeted a possibility of elemental sulfur-dependent anaerobic carboxydrotrophy in two types of hypersaline lakes, with neutral and alkaline pH, with a special attention to haloarchaea. Extremely halophilic archaea belonging to the class *Halobacteria* dominate prokaryotic communities in salt-saturated brines of various hypersaline habitats of marine and terrestrial origin. Most of the known species are aerobic organoheterotrophs. Recently, however, this point of view has changed. Using elemental sulfur as the terminal electron acceptor, we discovered a novel ecotype of haloarchaea living in anoxic sediments by anaerobic dissimilatory sulfur respiration. To date, four functional varieties of sulfur-reducing haloarchaea have been isolated and physiologically characterized from hypersaline habitats. The two groups isolated from salt lakes are obligate anaerobes and classified within closely related genera, which include acetate- and pyruvate oxidizing *Halanaeroarchaeum* [28–30] and lithoheterotrophic formate/H₂-oxidizing *Halodesulfurarchaeum* [31, 32]. The third group was found in hypersaline soda lakes. It consists of facultatively anaerobic haloalkaliphilic archaea, belonging to two new genera *Halalkaliarchaeum* and *Natrarchaeobaculum*, with a more versatile range of the electron donors for anaerobic sulfur respiration, including formate/H₂, pyruvate, C₄-C₉ fatty acids and peptone [33–36]. The last group includes facultatively aerobic haloarchaea, both neutrophilic and alkaliphilic, utilizing carbohydrates as an e-donor and carbon source for anaerobic sulfur respiration. Neutrophilic members form a new species in the

genus *Halapricum*. They can grow either by fermentation with the formation of H₂ or by sulfur respiration, while the single alkaliphilic strain belonging to a novel genus "*Natranaerarchaeum*" possesses a strictly respiratory metabolism growing by either O₂ or sulfur/thiosulfate-dependent respiration [37, 38].

In this paper, we describe the isolation, physiological and genomic properties of a fifth group of sulfur-respiring haloarchaea enriched from hypersaline lake sediments with CO as an electron donor. These lithoheterotrophic carboxydrotrophs directly oxidize CO to CO₂ with elemental sulfur as an electron acceptor, a type of metabolism that has not been known in haloarchaea and is a very rare catabolic trait in the whole archaeal domain.

EXPERIMENTAL PROCEDURES

Sample collection and initial treatments

Samples of the upper 15 cm sediment and near-bottom brine samples were taken from three pH-neutral hypersaline chloride-sulfate lakes (total salinity 220–340 g/l; pH 7.2–7.8) and three hypersaline soda lakes (total salinity from 150 to 320 g/l, total soluble carbonate alkalinity 1.5–3.8 M, pH 10.2–11.0) in the southern Kulunda Steppe (Altai region, Russia) in July 2015–2017. Samples were collected into a stratometric corer with an inner diameter of 25 mm, extruded into a sterile 100 ml Schott flask, sealed without air bubbles, transported to the laboratory in a insulated box with cooling elements within 3 days after sampling and stored at 10 °C until use. For initial enrichments, the sediment slurries from 3 different salt lakes and soda lakes were mixed in equal proportions to produce two "master mixes" that were used at 5% (v/v) as a combined inoculum.

Enrichment and cultivation conditions

Two mineral basic media (4 M total Na⁺) were used for enrichments and cultivation: a NaCl-base medium with a final pH 7.0 and a Na₂CO₃-base medium with a final pH 10.0. The former was used as such for the enrichment and cultivation of neutrophilic extreme halophiles, while for the soda lake cultivation was performed using a 1:1 mix of the NaCl- and Na₂CO₃-base media (final pH 9.7). The NaCl-base medium contained (g l⁻¹): NaCl, 240; KCl, 5; K₂HPO₄, 2.5; NH₄Cl, 0.5. The sodium carbonate-base medium contained (g l⁻¹): Na₂CO₃, 190; NaHCO₃, 30; NaCl, 16; KCl, 5.0; and K₂HPO₄, 1.0. After autoclave sterilization, both base media were supplemented with 1 mM MgCl₂, 1 ml l⁻¹ of acidic trace metal solution and 1 ml l⁻¹ vitamin mix [39], 1 ml l⁻¹ of alkaline Se/W solution [40] and 100 mg l⁻¹ of yeast extract. Elemental sulfur flour (J.T. Baker, Netherlands) wet paste was sterilized at 110 °C for 30 min and after cooling the clean water phase was removed. Sulfur was used at ~2 g l⁻¹. Cultivation was performed at 30 °C in 115 ml serum vials with butyl rubber stoppers filled with 30 ml medium in case of CO or H₂ as the electron donors or in 30 ml bottles with 25 ml media in case of soluble electron donors. Vials with sterile medium were subjected to 3 cycles of evacuation/flushing with sterile argon. Anaerobic conditions were achieved by final addition of 0.2 mM Na₂S from a 1.0 M filter sterilized stock solution. CO (20–50%) or H₂ (0.2 bar overpressure) were added through a sterile gas filter over an argon atmosphere. The final media reduction was done by adding about 20 µl of 10% dithionite in 1 M NaHCO₃ with a syringe. The cultures were incubated statically with occasional shaking of the flasks.

The strategy of isolation of pure cultures consisted of several rounds of decimal transfers to obtain sediment-free enrichments, followed by serial dilution-to-extinction. The maximum positive dilutions were determined by the formation of sulfides. Further purification was only possible using soft agar approach using maximum positive dilution cultures as starting material. The cultures were serially diluted in anoxic medium heated to 55 °C, mixed 4:1 with 4% washed agar, also kept at 55 °C, and the

resulted mix was poured into plates (15 ml total). Plates were incubated in 3.5 L anaerobic jars (Oxoid) under argon at 30 °C with an O₂-removing catalyzer (Oxoid). The colonies were picked into liquid medium with CO and sulfur. Finally, the purity of isolates was checked by 16S rRNA gene sequencing.

Potential catabolic activity experiments with washed (resting) cells were carried out in the mineral media (see above) at pH 7 or 9.7 as buffers lacking the N-source and containing 0.1 mM sulfide. 2 ml of the concentrated cell suspension was placed into 7 ml serum bottles containing 10 mg sulfur, made anoxic as described above, reduced by adding 20 µl of 1% dithionite in 1 M NaHCO₃ and incubated at 30 °C with various electron donors. Several 50 µl liquid samples were periodically taken during 5 days of incubation for sulfide determination and 0.5 ml gas samples for CO/CO₂ analyses as described below.

Analyses

Cumulative free sulfide + sulfane from polysulfide formed during elemental sulfur reduction were measured by the methylene blue method [41] after fixing the supernatant in 10% (w/v) zinc acetate. Gases were analyzed using a GC (Chromateck Crystall 5000 [Russia], Hayesep column 80–100 mesh, 2 m × 3 mm, 40 °C) equipped with a methanizer and thermal conductivity detector for H₂ and flame ionization detector (for CO and CO₂) (200 °C; carrier gas argon; flow rate 25 ml min⁻¹); the injection volume was 250 µl. Cellular protein was determined by the Lowry method in 1–2 ml culture samples after centrifugation 13,000 × *g* for 20 min. The cell pellets were washed with 4 M NaCl at pH 5.0 to remove iron (II) sulfide bound to the cells. Microbial growth was monitored by the increase in optical density at 600 nm (OD₆₀₀). Phase contrast microphotographs were obtained with a Zeiss Axioplan Imaging 2 microscope (Göttingen, Germany).

Genome sequencing and phylogenetic analysis

The complete genomes of two CO-utilizing sulfur-reducing haloarchaea were sequenced with the MiSeq System of Illumina Inc. (San Diego, CA, USA) using short insert paired-end library (2 × 250 bp, MiSeq Reagent Kit v2). Both genomes were assembled as a single circular chromosome and a small plasmid and analyzed as described previously [37]. The genome statistics are given in the Supplementary Table S1.

For the identification of the isolates phylogenetic reconstructions based on sequences of 122 archaeal single copy conservative marker genes [42] as well as of 16S rRNA gene sequences were used. For phylogenetic reconstruction of *CooS*, database consisting of 1942 non-redundant protein sequences described in Inoue et al., 2019 was used. In all cases phylogenetic trees were built with IQ-TREE 2 [43] and PhyML 3.0 [44] software with the approximate likelihood-ratio test for branches [45]. MAFFT version 7.487 was used for a multiple alignment [46]. Models for phylogenetic reconstructions were selected using ModelFinder software [47]. Five automated alignment trimming strategies were used in trimAl 1.2rev59 software [48]: (1) strictplus mode; (2) automated1 mode; (3) strictplus mode plus deletion of columns that contain more than 10%; (4) automated1 mode plus deletion of columns that contain more than 5%; (5) deletion of columns that contain more than 5%. After trimming all alignments were manually reviewed.

Whole cell lysate shotgun proteomic analysis

Cell lysis, protein extraction and proteolytic digestion was performed as described previously [26]. Briefly, ~10–20 mg of each cell pellet (biomass wet weight) were lysed using Complete Bacterial Protein Extraction Reagent (B-PER; Thermo-Fisher)/TEAB buffer and bead beating followed by centrifugation at 14,000 × *g* under cooling, using a bench top centrifuge. TCA was added to the protein supernatant (20% TCA, v/v), incubated at 4 °C and centrifuged to collect the protein precipitate. The pellet was

washed with ice cold acetone and dissolved in 200 mM ammonium bicarbonate containing 6 M urea. The disulfides were reduced with dithiothreitol (10 mM) and sulfhydryl groups were blocked with iodoacetamide (20 mM). The solution was diluted to below 1 M urea and digested using sequencing grade Trypsin (Promega). The proteolytic peptide digest was desalted using an Oasis HLB solid phase extraction well plate (Waters) according to the manufacturers protocol. An aliquot corresponding to ~500 ng of protein digest was analysed with a shotgun proteomics approach. The peptides were analysed using a nano-liquid-chromatography system consisting of an EASY nano LC 1200, equipped with an Acclaim PepMap RSLC RP C18 separation column (50 µm × 150 mm, 2 µm, 100 Å, Thermo Scientific) and a QE plus Orbitrap mass spectrometer (Thermo Fisher, Germany). The flow rate was maintained at 350 nL over a linear gradient from 5 to 30% solvent B over 38 min and finally to 60% B over 15 min. Solvent A was H₂O containing 0.1% formic acid, and solvent B consisted of 80% acetonitrile in H₂O and 0.1% formic acid. The Orbitrap was operated in data-dependent acquisition (DDA) mode acquiring peptide signals from 385 to 1250 m/z. The top 10 signals were isolated at a window of 2.0 m/z and fragmented at a NCE of 28. The collected mass spectrometric raw data were analysed against the constructed genome sequence database (described above), using PEAKS Studio 10 (Bioinformatics Solutions, Canada) allowing 20 ppm parent ion and 0.02 Da fragment ion mass error tolerance, 2 missed cleavages, carbamidomethylation as fixed and methionine oxidation and N/Q deamidation as variable modifications. Peptide spectrum matches were filtered for 1% false discovery rate (FDR) and protein identifications with at least 2 unique peptides were considered as significant.

Data deposition

16S rRNA gene sequences of strains HSR-CO and AArc-CO were deposited in the NCBI GenBank database with accession no. MZ466393-MZ466394 and their complete genomes were deposited in the GenBank database by the NCBI Genome submission portal with the accession numbers CP081498-CP081499 (chromosome-plasmid) for AArc-CO and CP081500-CP081501 (chromosome-plasmid) for HSR-CO. The proteomic data were deposited at the proteomics identification database (PRIDE Archive) and are available through the proteome exchange server (<http://www.proteomexchange.org/>, <http://www.ebi.ac.uk/pride>, (project code PXD028241); reviewer account details: reviewer_pxd028241@ebi.ac.uk).

RESULTS AND DISCUSSION

Enrichment and isolation of pure cultures

No CO consumption and H₂ formation was observed in primary anaerobic enrichments without electron acceptors and in presence of thiosulfate, nitrate/nitrite and ferrihydrite as the electron acceptors. CO consumption was only obtained when elemental sulfur served in this capacity. The salt-lake enrichment formed 2.5 mM sulfide and the soda-lake enrichment formed 6.3 mM sulfane (in polysulfides) after 2 months incubation (with endogenous controls without CO subtracted). After several stabilizing 1:100 transfers, sediment-free cultures were obtained in which the rates of CO-dependent sulfur reduction increased approximately two times for the salt-lake and four times for the soda lake cultivations. The two cultures were different in appearance. The salt-lake culture was only slightly turbid and colorless (Supplementary Fig. S1a). The soda-lake enrichment showed denser growth and was orange due to accumulation of polysulfide (S_n²⁻), spontaneously formed from biologically produced sulfide and the remaining sulfur, since S_n²⁻ is chemically stable at highly alkaline anoxic conditions (Supplementary Fig. S1b). Several rounds of dilution-to-extinction did not lead to the isolation of pure cultures: 16S rRNA gene sequencing

determined the microdiversity of archaea in the salt-lake enrichment, while the bacterial component was present along with the archaea in the soda-lake enrichment. Therefore, for the final purification, an attempt was made using soft agar plating under anaerobic conditions, although elemental sulfur could not be evenly distributed in the agar even after sonication of the suspended sulfur in liquid medium before mixing with the agar. Indeed, plating of the neutrophilic enrichment was difficult, since the culture did not form distinct colonies and grew as a diffuse cloud on the surface of sulfur conglomerates. However, the target organism was easy to differentiate from contamination by blackening of the sulfur particles with the *de novo* formed FeS. Microscopy showed that such changed conglomerates indeed contain a dense population of flattened cocci typical for haloarchaea, and their incubation in liquid medium with CO resulted in a pure carboxydrotrophic haloarchaeal culture designated as strain HSR-CO. Plating under microaerobic conditions (1% O₂ in the gas phase) on the medium with yeast extract and pyruvate did not yield any colonies for this culture after 2 months incubation. In contrast, plating of the soda lake highest positive dilution culture produced visible colored colonies both under anoxic condition in a CO-containing atmosphere and under microaerobic conditions with yeast extract and pyruvate. The surface microaerophilic colonies were bright red and had two shapes, but only one of them grew back anaerobically with CO. Anaerobic colonies formed under CO were of only a single type (pale pink lens-shaped). Notably, these colonies created a clearance zone around them due to the dissolution of fine sulfur particles and the conversion of the solubilized sulfur into a yellow polysulfide fringe outside the clearance zone (Supplementary Fig. S1c-d). Inoculation of the anaerobic colonies into liquid medium with CO and sulfur resulted in a positive culture designated AArc-CO. Both HSR-CO and AArc-CO cells were mostly angled flattened coccoids, typical for haloarchaea (Supplementary Fig. S2). The purity of the isolates was confirmed by 16S rRNA gene and whole genome sequencing. The strains were deposited in the Japanese Culture Collection of Microorganisms with the accession numbers JCM 34029 (HSR-CO) and JCM 34103 (AArc-CO).

Identification of the isolates

The HSR-CO genome contained a single *rrn* operon with its 16S rRNA gene 99.0% identical to halophilic *Halanaeroarchaeum sulfurireducens* HSR2^T, while the AArc-CO genome has two nearly identical *rrn* operons with 99.9% identity of 16S rRNA genes to each other and 98.4% identity to the haloalkaliphilic *Halalkaliarchaeum desulfuricum* AArc-SI^T 16S rRNA gene. Despite the high similarity of 16S rRNA genes of both these carboxydrotrophic strains to their nearest relatives (Supplementary Fig. S3), they represent new species in the respective genera of sulfur-respiring haloarchaea according to ANI, AAI and DDH analysis (Supplementary Table S2) and according to phylogenomic reconstruction based on 122 archaeal single copy conserved protein marker genes (Fig. 1).

Carboxydrotrophic growth and CO-dependent sulfidogenic activity

Already at the state of highly enriched cultures, it became clear that anaerobic utilization of CO is obligatory associated with the respiration on elemental sulfur, rather than more common pathways of anaerobic acetogenic or hydrogenogenic carboxydrotrophy. This was confirmed in growth experiments with pure haloarchaeal cultures, which also revealed another aspect—the obligate dependence of growth on the presence of yeast extract as a carbon source. Taken together, the carboxydrotrophic metabolism of isolated haloarchaea can be characterized as lithoheterotrophy assuming that CO is similar to H₂ in its capacity as an electron donor.




Fig. 1 Phylogenomic placement of carboxydrotrophic sulfur-reducing isolates AArc-CO and HSR-CO among their nearest relatives based on maximum likelihood inference and concatenated partial amino acid sequences of 122 archaeal single copy conserved marker proteins. Bootstrap consensus tree is shown with values above 90% placed at the nodes. Bar, 0.1 changes per position.

Both cultures grew and reduced sulfur only with yeast extract, but in the presence of CO a concentration-dependent increase in biomass (up to two times) and a significant increase in sulfidogenic activity (up to twelve times) were observed. There was a clear difference in the rate and intensity of carboxydrotrophic growth between the neutro- and alkali-philic isolates, including the following (Fig. 2): (i) the maximum cell density and sulfide formation during HSR-CO cultivation were reached after 35–40 days, while in case of AArc-CO it peaked in 20–23 days; (ii) the maximum CO-dependent increase in biomass (compared to the controls without CO) was similar in the two cultures, while sulfide/sulfane formation was three times higher in AArc-CO; (iii) the growth of neutrophilic culture was suppressed at CO concentrations above 50% in the gas phase, while the alkaliphilic culture was not inhibited at up to 80% CO, although growth and sulfidogenic activity did not increase further compared with the maximum reached at 40% CO; (iv) in abiotic control at neutral pH, there was no spontaneous sulfur reduction in the presence of CO in contrast to a weak abiotic reaction occurred at a pH above 9.5 at 30 °C, becoming significant (up to 10% of living cultures) upon increase of temperature to 37 °C and a CO concentration above 50% in the gas phase. Interestingly, no chemical reaction occurred in the activity tests (see below) using the same alkaline buffer applied in the AArc-CO cultivation medium (but without trace metals addition). This observation points to the possibility of an abiotic sulfur/polysulfide reduction by CO at high pH/temperature catalyzed by trace metals. The most likely mechanism here is the initial reaction of CO with polysulfide(s), leading to the formation of carbonyl sulfide (COS), followed by its hydrolysis to sulfide and CO₂. This reaction is favored by high pH, temperature and CO/polysulfide concentrations [49].

The optimal CO concentrations for growth and sulfidogenic activity in both strains was 30–40% in the gas phase. Stimulation of biomass yield with CO was clearly less effective than sulfidogenesis, indicating that CO was used only as an electron donor, and growth efficiency was limited by the assimilatory




Fig. 2 Anaerobic growth kinetics of carboxydrotrophic sulfur-reducing haloarchaea grown at 4.0 M total Na⁺/pH 7/37 °C (HSR-CO) or pH 9.5/30 °C (AArc-CO) with 0.5 g l⁻¹ yeast extract as the carbon source, CO as the electron donor and elemental sulfur as the electron acceptor. **a**, **b**, growth dynamics at 55 mM CO; (**c**), influence of CO concentration on growth and sulfur-reducing activity. The data are mean values from three parallel incubations.

metabolism. The same situation was observed for those sulfur-respiring haloarchaea that utilize H_2 or formate as the electron donors, including the genera *Halodesulfurarchaeum* and *Halalkaliarchaeum*. The stoichiometry of CO consumption in relation to the production of sulfide/sulfane was close to the theoretical 1:1. The only gaseous product of anaerobic CO oxidation in the neutrophilic culture was CO_2 , while in the alkaliphilic culture the accumulation of CO_2 in the gas phase was very low due to its absorption in the alkaline liquid phase (Fig. 2).

Both organisms were also able to grow lithoheterotrophically with two other low-potential electron donors, including H_2 and formate with sulfur as the acceptor. HSR-CO was significantly more active with H_2 (up to 11 mM HS^- and 3 mM on formate over 30 days of incubation), while AArc-CO showed the opposite trend (9 and 23 mM HS^- on H_2 and formate after 25 days, respectively).

Activity of CO-dependent sulfur reduction in resting cells

The cells of both strains, grown anaerobically with only yeast extract and sulfur, did not show any stimulation of sulfur-reducing activity in presence of CO during the first 5 days of incubation, but the activity began to manifest during further incubation. Since de novo protein synthesis in such incubations is unlikely, it is possible that the CODH is present but inactive in cells grown without CO and is activated after prolonged exposure to CO. In contrast, in cells grown with CO, the CO-stimulated sulfidogenesis began immediately, and the maximum CO-specific sulfur reduction in strain AArc-CO was twice that in HSR-CO, in line with what was observed in cultivation experiments (see above) (Fig. 3). Another difference between the cells of the two strains was that the CO-dependent sulfur reduction in neutrophilic HSR-CO was only active when dithionite was added as a second reductant to HS^- at the beginning of the experiment, while the cells of alkaliphilic AArc-CO were already active in presence of only 0.2 mM HS^- . This may indicate that under alkaline conditions the low redox potential required to activate (Ni-Fe) CODH can be more easily achieved even at low HS^- concentrations. Growth on CO also induced the high H_2 -dependent sulfur reducing activity observed in resting cells, while the effect of formate stimulation was much lower (Fig. 3). Overall, these experiments clearly demonstrated that CO and H_2 are strong electron donors for two novel isolates of sulfur-reducing haloarchaea.

Genomic analysis and comparative proteomics

Utilization of CO as the electron donor. A search for functional genes specific for anaerobic CODH was carried out in genomes of novel isolates. Both genomes contained a bicistron operon encoding the Ni-FeS catalytic subunit CooS and its Ni-chaperon CooC (CODH maturase). Using the CooS from AArc-CO and HSR-CO for a search of similar sequences within the class *Halobacteria* yielded a very limited number of matches including four closely related hits in the genomes of previously described sulfur-reducing haloarchaea. CooS from AArc-CO had the highest identity with the homolog from *Halalkaliarchaeum desulfuricum* AArc-S1^T [34], while CooS from HSR-CO had a high-level identity with CooS homologs encoded in the genomes of three of the four genome-sequenced strains (HSR-12-1, HSR12-2^T and HSR-Est) of carbohydrate-utilizing *Halapricum desulfuricans* [37, 38] (Supplementary Table S3). There were two additional hits including undescribed *Halorubrum* sp. CSM-61 (more similar to CooS of *Halalkaliarchaeum*) and *Salinarchaeum* sp. IM2453 (although it might be wrongly assigned to this genus, judging from the level of its 16S rRNA gene sequence). In addition, a *Halobacteria* metagenome-assembled genome (MAG) recovered from hypersaline industrial tailings in Canada, containing two copies of CooS with low homology to CooS from HSR-CO was identified. This indicates that anaerobic carboxydrotrophy is an extremely rare trait in haloarchaea with only a few examples bearing its functional determinates detected so far. Interestingly, in all seven cultivated




Fig. 3 Sulfidogenic activity of resting cells of carboxydrotrophic haloarchaea grown anaerobically with CO and sulfur. (a) HSR-CO; (b) AArc-CO. Cells of HSR-CO were incubated in 4 M NaCl containing 50 mM K-P buffer at pH 7. Cells of AArc-CO were incubated in an alkaline buffer containing 3.5 M NaCl and 0.5 M Na^+ as sodium carbonates, pH 9.5. The incubation temperature was 30 °C. The anoxic cell suspensions were reduced by 0.2 mM sulfide and 10 μ l of 10% dithionite in 1 M $NaHCO_3/2$ ml. The chemical controls were buffers without cells and were negative during the whole incubation period (up to 7 days). The data are mean values from duplicate experiments.

strains of haloarchaea, CooS was colocalized with CooC in one operon (with the exception of the second copy in the *Halobacteria* MAG), indicating the importance of this small maturase in the functionality of catalytic CooS, which is consistent with the results of biochemical studies [50]. The N-terminus of the CooC proteins of two *Halalkaliarchaeum* strains is particularly interesting in that it has a large locus of histidine-aspartate (HD) repeats, which are known to be present in Ni-chaperones, such as CooJ in *Rhodospirillum rubrum* [51], HypB (Ni,Fe hydrogenase assembly proteins) and in Ni-storage/transport proteins, Hpn and RcnA, present in some pathogens such as *Helicobacter pylori* [52]. Moreover, in AArc-CO, this HD-rich region was double that of the type strain of *Halalkaliarchaeum* (Supplementary Table S3). On the other hand, the CooC from the previously isolated neutrophilic HSR strains and the two undescribed haloarchaea lacks this characteristic pattern. It is likely, that the Ni-transport and handling in alkaliphiles is more difficult than under neutral hypersaline conditions.

Due to the presence of CooS/CooC operon in the genomes of three *Halapricum desulfuricans* strains, we tested these haloarchaea (stored and maintained in our personal culture collection) for their ability to grow anaerobically with CO and sulfur. Controls without CO in the presence of yeast extract exhibited no growth and sulfide formation, but in the presence of limited concentrations of glucose addition of 20% CO to the gas phase substantially increased sulfide formation in the type strain of *H. desulfuricans*




Fig. 4 Phylogeny of the CooS catalytic subunit from haloarchaeal Ni,Fe-CODH based on maximum likelihood inference. Bootstrap consensus tree is shown with values above 80% placed at the nodes. Bar, 0.1 changes per position. The clades are according to Inoue et al. [4].

HSR12-2, while growth stimulation was only marginal. The most pronounced effect of CO addition was observed at the lowest glucose concentration used (Supplementary Fig. S4). Interestingly, among the closely related *Halapricum desulfuricum* isolates, HSR12-2^T was the only one capable of lithoheterotrophic growth with H₂ as the electron donor and sulfur as the electron acceptor [37].

The carboxydrotrophic potential in the alkaliphilic genus *Halalkaliarchaeum*, was more prominent. In addition to the novel strain AArc-CO, enriched with CO, we have also recently described strain AArc-GB capable of using glycine betaine as a carbon source and electron donor and sulfur as an electron acceptor [53]. The complete genome of the strain was not yet sequenced, but because of its proximity to AArc-CO (99% of the 16S rRNA gene sequence identity), it was also included into the test for carboxydrotrophy. The results showed that both the type species *H. desulfuricum* AArc-SI¹ and strain AArc-GB, can grow anaerobically with CO as the electron donor, sulfur as electron acceptor and yeast extract as the carbon source in the following order of sulfidogenic intensity: AArc-CO > AArc-GB >> AArc-SI¹.

Phylogeny of CooS from sulfur-reducing carboxydrotrophic haloarchaea. In order to define the phylogenetic position of CooS from anaerobic haloarchaea, a multiple sequence alignment was applied included 1942 non-redundant protein sequences and the previously established Ni,Fe CODH clade structure [4, 7]. Sequences of CooS from haloarchaea are located in clade F (Fig. 4) and formed a separate deep cluster. Based on the resulting reconstruction, we can assume that these haloarchaea have

acquired CooS not during a recent horizontal gene transfer event, but evolved with it for a long time. All haloarchaeal CooS in this cluster belong to the structural group F-1 which mostly include the ECH-associated proteins. The F-1 Ni-CODHs were mainly found in bacteria, with a few exceptions [4]. Taking into account this fact and the fact, that all clades, except clade A, are mainly bacterial we can assume, that haloarchaea have initially acquired Ni-CODH from bacterial anaerobic carboxydrotrophs.

In general our phylogenetic reconstructions is consistent with previously published works including subdivision into clades A-F [4, 7]. However, during our work, we have also noticed an instability of clade F. Depending on the methods of alignment trimming (see Experimental procedures section), this biggest CooS clade in many cases became paraphyletic. Probably, a larger number of diverse sequences of CooS from various sources would improve our understanding of CooS phylogeny, specifically clade F.

Possible routes of electron transfer from CooS in sulfur-reducing haloarchaea. In addition to the functional CODH proteins, a most important question for understanding the anaerobic CO-oxidizing system in sulfur-reducing haloarchaea is the nature of electron transfer from the cytoplasmic CooS to the extracellular polysulfide reductase complex Psr. Since the electron transport system is well established only in hydrogenogenic carboxydrotrophs (via CooF ferredoxin to membrane-bound H₂-forming hydrogenases), it is difficult to predict what acts in the direct anaerobic oxidation of CO to CO₂ with sulfur as an actual electron acceptor. However, one thing is clear – the membrane-associated subunit of Psr (PsrC) can

accept as the direct electron donor only menaquinol, a membrane electron shuttle that is ubiquitous in all haloarchaea [54]. Each of the *Halalkaliarchaeum* genomes revealed the presence of at least three electron transfer flavoprotein (Etf) complexes, known to be coupled to several redox reactions in the cytoplasm including, ferredoxin-dependent bifurcation [55, 56]. Noteworthy, the four-subunit EtfABCX complex containing the ferredoxin EtfX and the quinone-reductase EtfC (AArcCO_1914-1917) and the three subunits EtfABC (AArcCO_1725-1727) may be involved in electron transfer from the cytoplasmic CODH to the membrane menaquinone pool (Fig. 5).

In the genome of neutrophilic HSR-CO, the CODH operon is immediately preceded by a membrane-bound Ni,Fe uptake hydrogenase HyaABC accompanied with its full set of maturation factors HypABCDE/HyaD. It would be tempting to suggest that both [Ni,Fe] catalytic subunits HyaB (located outside the CPM) and the cytoplasmic Ni,Fe CooS can use the common membrane menaquinol-reductase subunit HyaC to donate electrons to the terminal electron-accepting PsrABC complex (Fig. 5). Such a connection would mimic, at least superficially, the hydrogenogenic system with the CooS linked to membrane-bound H_2 -forming hydrogenases. The high H_2 -dependent rates of sulfur reduction, observed in cells of HSR-CO grown with CO supports the proposed electron transfer scheme. However, both schemes (for AArc-CO and HSR-CO) are purely hypothetical, and deep biochemical and genetic studies will be required to understand the actual electron transport routes from CO to sulfur Psr in carboxydrotrophic haloarchaea.

Another possibility of converting low-potential electrons from CO oxidation into a cytoplasmic redox active donor could be via ferredoxin-NAD⁺ oxidoreductase (Fnor) [57] with further menaquinone reduction through the NADH dehydrogenase (Nuo complex I) ($CO \rightarrow Fnor \rightarrow NADH \rightarrow Nuo \rightarrow$ menaquinol \rightarrow Psr). The genomes of both HSR-CO and AArc-CO do indeed contain homologs of both the Fnor and Nuo complexes (NuoAB-C/D-HIJ12KLMN). Notably, in both of them, the Nuo complexes lack three subunits of the NADH dehydrogenase N-module (NuoEFG), indicating that these carboxydrotrophic haloarchaea use other low-potential electron donors rather than NADH, such as F_{420} - H_2 or ferredoxin, potentially formed by oxidation of CO [58].





Fig. 5 Hypothetical scheme of electron transport from CooS to PsrABC in two anaerobic carboxydrotrophic haloarchaea (based on genomic content). HyaABC membrane-bound Ni, Fe uptake hydrogenase, PsrABC membrane-bound polysulfide reductase, EtfABCX cytoplasmic electron transfer flavoprotein complex, MK/MKH2 menaquinone-menaquinol, S polysulfide.

H₂ and formate as the electron donors. Apart from the CODH and consistent with the growth and cell activity tests, the genomes of HSR-CO and AArc-CO encode the type 1a membrane-bound Ni,Fe uptake hydrogenase Hya/HydABC and its maturation factors and two types of formate dehydrogenases: a membrane-bound FdoGHI and a cytoplasmic monosubunit FdhH. While both of those have previously been found in the genome of *Halkaliarchaeum desulfuricum* closely related to AArc-CO, neither are present in the type species of *Halanaeroarchaeum sulfurireducens* which utilize only acetate or pyruvate as the electron donors. Therefore, despite close phylogenetic relation, HSR-CO has a very different key catabolic properties from the type species, more reminiscent of the lithoheterotrophic *Halodesulfurarchaeum*, a sister genus to *Halanaeroarchaeum*.

Dissimilatory sulfur reduction. Apart of revealed capability for anaerobic CO oxidation, AArc-CO and HSR-CO exhibit anaerobic respiration patterns very similar to those of previously described sulfidogenic halo(natrono)archaea mentioned in the Introduction. In addition to the aforementioned formate dehydrogenases, both genomes encode an analogous set of DMSO reductase DmsABC, polysulfide reductase PsrABC and tetrathionate reductase TtrABC molybdopterin oxidoreductases from the CISM superfamily [59], acting as potential terminal electron acceptors (Fig. 6). Interestingly, while HSR-CO has two polysulfide- and four DMSO-reductases, the AArc-CO set of CISM enzymes contains only single Psr and DMSOr. Such a “minimal” respiratory suit, which determines CO- or formate-dependent respiration with elemental sulfur and DMSO, makes this organism an ideal model for unambiguously identifying the main mechanisms of the observed types of respiration. A single set of AArc-CO Psr polysulfide reductase genes is located in one polycistronic operon, which includes genes encoding MoPterin (*psrA*), FeS subunit (*psrB*), membrane quinol-interacting anchor protein (*psrC*) and system-specific chaperone (*psrD*). In addition to the *psrA-D* genes (AArcCO_1246-8), the AArc_1245 gene, which encodes rhodanese-like sulfurtransferase belonging to the SseA family, is located upstream. We recently hypothesized that this sulfurtransferase acts as a potential polysulfide-binding carrier that is responsible for the mobilization of sulfur during sulfur respiration in sulfur-respiring halo(natrono)archaea [28, 31, 33]. It is noteworthy that such a rhodanese-containing Psr operon was also found as a single set of polysulfide reductase genes in another sulfidogenic natronoarchaea, *Natranaerobaculum sulfurireducens* [33]. This finding strongly suggests that the products of this operon are sufficient to determine the type of respiration based on the use of elemental sulfur as a terminal electron acceptor. The catalytic subunits of CISM enzymes found in the carboxydrotrophic isolates AArc-CO and HSR-CO clustered together with the homologs from the previously described sulfur reducing halo (natrono)archaea (Fig. 6) forming deep branches within the respective CISM families and likely representing ancient forms of molybdopterin oxidoreductases acquired in the past from bacteria.

The genome of AArc-CO also contains three homologous gene clusters with a similar structure: (1) ferredoxin>NrfC1D1>membrane protein>NrfC2fD2; (2) ferredoxin>membrane protein>NrfC3D3; (3) ferredoxin>NrfC4D4>membrane protein; the membrane protein in all three loci is conserved but its function cannot be predicted, while the ferredoxin seems to belong to the family of ferredoxin-NAD(P) oxidoreductase family. The NrfC and D are homologs of PsrB and C, respectively, and recently this membrane associated quinol-dehydrogenase module was found to be widely present in various anaerobic quinol-interacting respiratory complexes, for example in the Qrc (as QrcCD) and Dsr (as DsrOP) complexes of sulfate-reducing bacteria [60]. The presence of four such modules in AArc-CO genome strongly suggest that they are an important part of the anaerobic respiration in this organism, but, once more, since they are not in a



direct association with the loci encoding either CO, H₂ or formate-oxidizing enzymes, their true function cannot be predicted for now.

Other electron transfer complexes. Both genomes contain four-gene operons encoding a full quinone-reducing electron transfer flavoprotein complex EtfABCX (archael homolog of the bacterial

FixBACX [61]), but since they are not located in proximity to the CooS/CooC operons, their possible link to CO oxidation cannot be predicted from only genomic content. In contrast to the classical FixC of bacteria, which is the integral membrane quinone-reductase, in the studied archaea this protein is predicted to be cytoplasmic having only a small hydrophobic patch at the

Fig. 6 Maximum likelihood phylogenetic tree of catalytic subunit A of molybdopterin oxidoreductases, members of the Complex iron-sulfur molybdoenzyme (CISM) superfamily according to Duval et al. [67]. **a** Respiratory chain components in sulfur-reducing haloarchaea, experimentally proven by us **(b)**. In total, 158 sequences were taken for CISM analysis, including 61 nonhaloarchaeal members (highlighted in gray). The filled circles at the nodes indicate the bootstrap values >70% (1000 bootstrap iterations) in the tree with the highest log likelihood. The loci of the CISM catalytic subunits A of new carboxydophilic sulfur-reducing halo(natrono)archaea of the genus *Halalkaliarchaeum* (red), and the genus *Halanaeroarchaeum* (blue) are highlighted in bold. The unknown CISM family is shown with an asterisk. Scale bar—0.06 amino acid substitutions per site. The purple rectangle in **(b)** highlights the polysulfide reductase PsrABCD, which is transcribed together with sulfurtransferase/rhodanese-like protein. Organisms containing only this PsrABCD polysulfide reductase, which thus represents the “minimum” respiratory outfit required for their detected respiration pattern, are shown in empty black circles. DMSO/Nar DMSO/nitrate reductase family, Fdh/Nas/Nap/ assimilatory nitrate reductase/formate dehydrogenase family, [NiFe]-H₂ase [NiFe] uptake hydrogenase, Psr/Phs polysulfide/thiosulfate reductase family, Ttr/Arr tetrathionate/arsenate reductase family.

N-terminus which would allow it to attach to the inner face of the CPM similar to the characterized EtfABCX from a hyperthermophilic crenarchaeon *Pyrobaculum aerophilum* [62].

Both AArc-CO and HSR-CO genomes encode a non-proton pumping quinol oxidase *bd*, frequently present in obligate anaerobes, including members of two obligate anaerobic sulfur-reducing haloarchaeal genera *Halanaeroarchaeum* and *Halodesulfurarchaeum*. Unexpectedly, in addition to this oxidase, the HSR-CO genome contains a cluster of 5 genes (HSRCO_1948-52), apparently encoding a full set of the heme-copper superfamily cytochrome *c*-oxidase *caa*₃ (COX I-III and COX cluster proteins 1 and 2), which theoretically should allow this organism to grow by aerobic respiration. Despite this, all attempts to grow HSR-CO under microaerophilic conditions with variety of electron donors, including CO, H₂, formate, organic acids and pepton/yeast extract, have been unsuccessful. In contrast, strain AArc-CO can be easily transferred from the sulfur-reducing cultures to full aerobiosis through intermediate microaerophily (2% of O₂ in gaseous phase) using a heterotrophic medium with pyruvate+yeast extract, similar to the other two strains of *Halalkaliarchaeum* (AArc-SI^T and AArc-GB) [34, 53]. In agreement with this result, AArc-CO genome contains gene clusters encoding for the proton-pumping quinol oxidases *bo*₃ (CyoAB) and *ba*₃ (CbaABDE) also present the genome of AArc-SI^T.

Another unexpected feature of the HSR-CO genome is the presence of an operon encoding an archaeal type of extracellularly located membrane-bound nitrate reductase, which includes NarGHI/BC [63]. Moreover, downstream of the *nar* cluster, there is a locus encoding an enigmatic complex reminiscent of the ammonifying nitrite reductase NrfABCD of the enteric gamma-proteobacteria [64]. This complex includes an octaheme cytochrome *c* (annotated as the hydroxylamine oxidoreductase), a tetraheme *c* of the NrfB/DmsE family, a membrane-bound quinol dehydrogenase module NrfCD (ferredoxine/membrane diheme cytochrome *b*) and eight cytochrome *c* biogenesis proteins. As far as we know, the presence of such a complex has never been observed in haloarchaeal genomes. Among other archaea, the octaheme *c* protein of HSR-CO has only a low identity (26%) to octaheme *c* hypothetical proteins from *Ca*. Methanoperedens nitroreducens. Also, there is an apparent resemblance of the complex to a part of the hypothetical scheme of extracellular electron transport from a quinol pool to MnO₂ in recently discovered Mn-reducing anaerobic methane oxidizing *Ca*. Methanoperedens manganicus and *Ca*. Methanoperedens manganireducens (Fig. 5 in [65]). This includes a membrane-bound quinol-dehydrogenase NrfCD-like module and several multiheme *c* cytochromes conducting extracellular electron transport to MnO₂ or Fe(III) oxides. Attempts to grow HSR-CO anaerobically at hetero- and lithoheterotrophic conditions with nitrate as the electron acceptor showed positive results when H₂ served as the electron donor and yeast extract as the C-source, but not with CO or formate/yeast extract or at heterotrophic conditions with pyruvate or lactate as substrates. But even with H₂, nitrate (5 mM) was only reduced to nitrite (Supplementary Fig. S5) and there was no growth with nitrite even at minimal tested concentration of 1 mM. Nevertheless, these are remarkable results, since, to our

knowledge, even incomplete nitrate respiration to nitrite with H₂ as the electron donor has never been demonstrated before in haloarchaea [66].

Anaerobic methylophony in the genus Halalkaliarchaeum. As previously mentioned, a new member of the genus *Halalkaliarchaeum*, strain AArc-GB, was enriched and isolated in pure culture from hypersaline soda lakes using glycine betaine (GB, trimethylglycine) as energy and carbon source with sulfur as an electron acceptor, being the first haloarchaeon with such anaerobic methylophytic metabolism [53]. Since this strain was closely related to AArc-CO, as well as to the type strain of *Halalkaliarchaeum*, *H. desulfuricum* AArc-SI^T, the latter two were also checked for their ability for anaerobic methylophony with GB in presence of sulfur as the electron acceptor. The results were positive with the following order of sulfidogenic activity: AArc-GB > AArc-CO > AArc-SI^T. Previous genome analysis of the type strain AArc-SI^T identified a substantial number of genomic loci which, taken together, suggest a hypothetical scheme of the sequential demethylation of GB to glycine with a concomitant methyl group oxidation via the methylophytic C₁ pathway. The first demethylation step driven by the GB-specific methyltransferases MtgABC is unique for anaerobes, while the second and the third steps catalyzed by the dimethylglycine dehydrogenase Dmg and monomeric sarcosine oxidase SoxB, respectively, are described for aerobic methylophytes utilizing GB as the carbon and energy source. Furthermore, a full complement of the tetrahydrofolate-dependent methyl group oxidative pathway is also present in the same genomic island [53 and the references therein]. The AArc-CO genome seems to encode similar proteins with the exception of a few variations in the number of the homologs (Supplementary Table S4) suggesting that both the anaerobic sulfur-dependent C₁ catabolism (carboxydo- and methylo- trophy) are inherent property in the members of genus *Halalkaliarchaeum* isolated in pure culture so far.

Whole cell lysate shotgun proteomic analysis. The whole cell lysates of the AArc-Co and HSR-CO cells grown at sulfur-reducing conditions either heterotrophically with yeast extract only or lithoheterotrophically with CO + yeast extract were analysed using a shotgun proteomics approach which confirmed the expression of the presumed key enzymes/pathways and moreover provided the following observations (summary of selected enzymes/pathways can be found on Supplementary Table S5):

- i. Catalytic subunits of the key catabolic enzymes—CooS of the CODH and PsrA of the polysulfide reductase were equally present among the most highly abundant proteins in cells grown with or without CO in both strains. While for the latter it is reasonable, since in both growth conditions sulfur was used as the electron acceptor, the high expression level of the CooS in cells grown without CO would signify the constitutive nature of this enzyme. This contradicts to what was observed in the resting cell experiments (see above). Presumably the observed activation following incubation with CO for several days might indicate the necessity for

maturation of constitutively expressed *CooS* by *CooC*. The apparently low expression level of the *CooC* protein in AArc-CO might be (artefactually) connected to interference of the large locus of multiple HD repeats (see above) with the standard sample preparation protocol.

- ii. In the working hypothesis on the nature of electron transfer complexes possibly involved in carboxydutrophy, the uptake Ni₂Fe hydrogenase is implicated as the CODH partner in HSR-CO (based on the proximity in the genome) and the EtfABC or EtfABCX in case of AArc-CO (on the basis of its quinone-reductase potential and the presence of a ferredoxin subunit). The proteomic data showed that the Ni₂Fe hydrogenase catalytic subunits HyaAB are among the highly expressed proteins in HSR-CO cells, although only HyaB had a higher expression level in the presence of CO. In this respect the proteomic results are well in agreement with the high hydrogenase activity observed in the CO-grown cells of HSR-CO cells (see Fig. 3). Among the four Etf complexes in AArc-CO (two EtfAB, EtfABC and EtfABCX), the EtfABC was definitely positively influenced by the presence of CO.
- iii. Regarding the other possibilities for channelling electrons from CO oxidation to P_{sr}, it is worth mentioning that cells of AArc-CO grown with CO had a higher level of expression of the ferredoxin-NAD⁺ reductase (Fnr) in comparison with the cells grown heterotrophically only with yeast extract (Supplementary Table S5). These cumulatively might be utilized for an alternative route of the menaquinone reduction with the low-potential electrons derived from CO oxidation. Furthermore, in AArc-CO, only the cells grown with CO expressed one of the four copies of the NrfCD/PsrBC-like membrane-bound complex which is supposed to be able to interact with the quinone pool in various respiratory complexes [60], indicating that it might potentially be involved in the electron transfer from CODH as well.
- iv. The overall proteomic results (Supplementary Table S5) showed that cells of alkaliphilic AArc-CO responded more actively to the presence of CO in comparison to the cells of neutrophilic HSR-CO.

CONCLUSIONS

In this work, we demonstrate for the first time that haloarchaea isolated from hypersaline lakes with neutral and alkaline pH are capable of anaerobic growth with CO as the electron donor and sulfur as the electron acceptor. The direct anaerobic sulfur-dependent carboxydutrophy of the new *Halanaeroarchaeum* and *Halalkaliarchaeum* isolates is a catabolic trait which has previously been shown only for a single archaeal genus *Thermococcus*. The exact mechanism of the electron transport from anaerobic CODH to a final electron acceptor (sulfur in this case) remains unclear yet either in bacteria or archaea. Genomic and proteomic analysis of carboxydutrophic haloarchaea suggest several candidates for linking anaerobic CO oxidation to polysulfide reductase via a menaquinone pool, such as the membrane-bound Ni₂Fe uptake hydrogenase HyaABCD and the electron transfer flavoprotein complex EtfABC(X), but both biochemical characterization and genetic manipulation analysis are still needed to confirm it. In addition to the anaerobic carboxydutrophy, members of the genus *Halalkaliarchaeum* are also capable of anaerobic sulfur-dependent methylotrophy with glycine betaine as a substrate, making this genus truly unique among haloarchaea.

REFERENCES

1. Ragsdale SW. Life with carbon monoxide. *Crit Rev Biochem Mol Biol.* 2004;39:165–95.
2. Svetlitchnyi V, Peschel C, Acker G, Meyer O. Two membrane-associated NiFeS-carbon monoxide dehydrogenases from the anaerobic carbon-monoxide-utilizing eubacterium *Carboxydotherrmus hydrogenoformans*. *J Bacteriol.* 2001;183:5134–44.
3. Techtmann SM, Lebedinsky AV, Colman AS, Sokolova TG, Woyke T, Goodwin L, et al. Evidence for horizontal gene transfer of anaerobic carbon monoxide dehydrogenases. *Front Microbiol.* 2012;3:132.
4. Inoue M, Nakamoto I, Omae K, Oguro T, Ogata H, Yoshida T, et al. Structural and phylogenetic diversity of anaerobic carbon-monoxide dehydrogenases. *Front Microbiol.* 2020;9:3353.
5. Jeoung J-H, Martins BM, Dobbek H. Carbon monoxide dehydrogenases. In: Hu Y (ed.), *Metalloproteins: Methods and Protocols.* Methods in Molecular Biology. 2019; 1876:37–54.
6. King GM, Weber CF. Distribution, diversity and ecology of aerobic CO-oxidizing bacteria. *Nat Rev Microbiol.* 2007;5:107–18.
7. Robb FT, Techtmann SM. Life on the fringe: microbial adaptation to growth on carbon monoxide. *F1000Research.* 2018;7. <https://doi.org/10.12688/f1000research.16059.1>.
8. King GM. Carbon monoxide as a metabolic energy source for extremely halophilic microbes: Implications for microbial activity in Mars regolith. *Proc Natl Acad Sci USA.* 2015;112:4465–70.
9. Sokolova TG, Henstra AM, Sipma J, Parshina SN, Stams AJM, Lebedinsky AV. Diversity and ecophysiological features of thermophilic carboxydutrophic anaerobes. *FEMS Microbiol Ecol.* 2009;68:131–41.
10. Diender M, Stams AJM, Sousa DZ. Pathways and bioenergetics of anaerobic carbon monoxide fermentation. *Front Microbiol.* 2015;6:1275.
11. Hoelt SE, Switzer Blum J, Stolz JF, Tabita FR, Witte B, King GM, et al. *Alkalilimnicola ehrlichii* sp. nov., a novel arsenite-oxidizing, haloalkaliphilic gamma-proteobacterium capable of chemoautotrophic or heterotrophic growth with nitrate or oxygen as the electron acceptor. *Int J Syst Evol Microbiol.* 2007;57:504–12.
12. Sorokin DY, Kovaleva OL, Tourova TP, Kuenen JG, Muyzer G. Aerobic carboxydutrophy at extremely haloalkaline conditions in *Alkalispirillum/Alkalilimnicola* strains isolated from soda lakes. *Microbiol (SGM).* 2010;156:819–27.
13. Myers MR, King GM. Perchlorate-coupled carbon monoxide (co) oxidation: evidence for a plausible microbe-mediated reaction in Martian brines. *Front Microbiol.* 2017;8:2571.
14. Jensen A, Finster K. Isolation and characterization of *Sulfurospirillum carboxydovorans* sp. nov., a new microaerophilic carbon monoxide oxidizing epsilon-proteobacterium. *Antonie van Leeuwenhoek.* 2005;87:339–53.
15. Parshina SN, Sipma J, Henstra AM, Stams AJM. Carbon monoxide as an electron donor for the biological reduction of sulphate. *Int J Microbiol.* 2010;2010:319527.
16. Fukuyama Y, Omae K, Yoneda Y, Yoshida T, Sako Y. Insight into energy conservation via alternative carbon monoxide metabolism in *Carboxydotherrmus pertinax* revealed by comparative genome analysis. *Appl Environ Microbiol.* 2018;84:e00458–18.
17. Alves JI, Visser M, Arantes AL, Nijssse B, Plugge CM, Alves MM, et al. Effect of sulfate on carbon monoxide conversion by a thermophilic syngas-fermenting culture dominated by a *Desulfofundulus* species. *Front Microbiol.* 2020;11:588468.
18. Henstra AM, Dijkema C, Stams AJ. *Archaeoglobus fulgidus* couples CO oxidation to sulfate reduction and acetogenesis with transient formate accumulation. *Environ Microbiol.* 2007;9:1836–41.
19. Kozhevnikova DA, Taranov EA, Lebedinsky AV, Bonch-Osmolovskaya EA, Sokolova TG. Hydrogenogenic and sulfidogenic growth of *Thermococcus* archaea on carbon monoxide and formate. *Microbiol (Mosc, Engl Translation).* 2016;85:400–10.
20. Oger P, Sokolova TG, Kozhevnikova DA, Chernyh NA, Bartlett DH, Bonch-Osmolovskaya EA, et al. Complete genome sequence of the hyperthermophilic archaeon *Thermococcus* sp. strain AM4, capable of organotrophic growth and growth at the expense of hydrogenogenic or sulfidogenic oxidation of carbon monoxide. *J Bacteriol.* 2011;193:7019–20.
21. Benvenuti M, Meneghello M, Guendon C, Jacq-Bailly A, Jeoung J-H, Dobbek H, et al. The two CO-dehydrogenases of *Thermococcus* sp. AM4. *Biochim Biophys Acta.* 2020;1861:148188.
22. Wu GJ, Schut FLP, Haja DK, Adams MWW. Characterization of membrane-bound sulfane reductase: A missing link in the evolution of modern day respiratory complexes. *J Biol Chem.* 2018;293:16687–96.
23. Allen TD, Caldwell ME, Lawson PA, Huhnke RL, Tanner RS. *Alkalibaculum bacchi* gen. nov., sp. nov., a CO-oxidizing, ethanol-producing acetogen isolated from livestock-impacted soil. *Int J Syst Evol Microbiol.* 2010;60:2483–2489.
24. Liu K, Atiyeh HK, Tanner RS, Wilkins MR, Hahnke RL. Fermentative production of ethanol from syngas using moderately alkaliphilic strains of *Alkalibaculum bacchi*. *Bioresour Technol.* 2012;104:336–41.
25. Khomyakova MA, Merkel AY, Petrova DA, Bonch-Osmolovskaya EA, Slobodkin AI. *Alkalibaculum sporogenes* sp. nov., isolated from a terrestrial mud volcano and emended description of the genus *Alkalibaculum*. *Int J Syst Evol Microbiol.* 2020;70:4914–4919.
26. Sorokin DY, Diender M, Merkel AY, Koenen M, Bale NJ, Pabst M, et al. *Natranaerofaba carboxydovora* gen. nov., sp. nov., an extremely haloalkaliphilic CO-utilizing acetogen

- from a hypersaline soda lake representing a novel deep phylogenetic lineage in the class "Natranaerobii". *Environ Microbiol.* 2021;23:3460–76.
27. Sorokin DY, Makarova KS, Abbas B, Ferrer M, Golyshin PN, Galinski EA, et al. Discovery of extremely halophilic, methyl-reducing euryarchaea provides insights into the evolutionary origin of methanogenesis. *Nat Microbiol.* 2017;2:17081.
 28. Sorokin DY, Kublanov IV, Gavrilo SN, Rojo D, Roman P, Golyshin PN, et al. Elemental sulfur and acetate can support life of a novel strictly anaerobic haloarchaeon. *ISME J.* 2016;10:240–252.
 29. Sorokin DY, Kublanov IV, Yakimov MM, Rijpstra WI, Sinninghe Damsté JS. *Halanaeroarchaeum sulfurireducens* gen. nov., sp. nov., the first obligately anaerobic sulfur-respiring haloarchaeon, isolated from a hypersaline lake. *Int J Syst Evol Microbiol.* 2016;66:2377–81.
 30. Sorokin DY, Yakimov MM, Kublanov IV, Oren A. *Halanaeroarchaeum*. In: Whitman WB (ed). *Bergey's Manual of Systematic of Bacteria and Archaea*. John Wiley & Sons, Ltd. 2017; <https://doi.org/10.1002/9781118960608.gbm01496>.
 31. Sorokin DY, Messina E, Smedile F, Roman P, Sinninghe Damsté JS, Ciordia S, et al. Discovery of anaerobic lithoheterotrophic haloarchaea, ubiquitous in hypersaline habitats. *ISME J.* 2017;11:1245–60.
 32. Sorokin DY, Yakimov MM. Genus *Halodesulfuriarchaeum*. In: Whitman WB (ed). *Bergey's Manual of Systematic of Bacteria and Archaea*. John Wiley & Sons, Ltd. 2018. <https://doi.org/10.1002/9781118960608.gbm01528>. 32
 33. Sorokin DY, Messina E, La Cono V, Ferrer M, Ciordia S, del Carmen Mena, et al. Sulfur respiration in a group of facultatively anaerobic natronoarchaea ubiquitous in hypersaline soda lakes. *Front Microbiol.* 2018;9:2359.
 34. Sorokin DY, Yakimov MM, Messina E, Merkel AY, Bale NJ, Sinninghe Damsté JS. *Natronolimnobius sulfurireducens* sp. nov., and *Halalkaliarchaeum desulfuricum* gen. nov., sp. nov., the first sulfur-respiring alkaliphilic haloarchaea from hypersaline alkaline lakes. *Int J Syst Evol Microbiol.* 2019;69:2662–73.
 35. Sorokin DY, Merkel AY, Messina E, Yakimov MM, Itoh T, Mesbah NM, et al. Reclassification of the genus *Natronolimnobius*: proposal of two new genera, *Natronolimnohabitans* gen. nov. to accommodate *Natronolimnobius innermongolicus* and *Natrarchaeobaculum* gen. nov. to accommodate *Natronolimnobius aegyptiacus* and *Natronolimnobius sulfurireducens*. *Int J Syst Evol Microbiol.* 2020;70:3399–405.
 36. Sorokin DY, Merkel AY, Yakimov MM, Oren A. *Halalkaliarchaeum*. In: Whitman WB (ed). *Bergey's Manual of Systematics of Archaea and Bacteria*. John Wiley & Sons, Ltd. 2020. <https://doi.org/10.1002/9781118960608.gbm01943>.
 37. Sorokin DY, Messina E, Smedile F, La Cono V, Hallsworth JE, Yakimov MM. Carbohydrate-dependent sulfur respiration in halo(alkali)philic euryarchaea from hypersaline lakes. *Environ Microbiol.* 2021;23:3779–808. 37
 38. Sorokin DY, Yakimov MM, Messina E, Merkel AY, Koenen M, Bale NJ, et al. *Halapiricum desulfuricans* sp. nov., carbohydrate-utilizing sulfur-reducing haloarchaea from hypersaline lakes. *Syst Appl Microbiol.* 2021;44:126249.
 39. Pfennig N, Lippert KD. Über das Vitamin B12-Bedürfnis phototropher Schwefelbakterien. *Arch Mikrobiol.* 1966;55:245–56.
 40. Plugge CM. Anoxic media design, preparation, and considerations. *Methods Enzymol.* 2005;397:3–16.
 41. Trüper HG, Schlegel HG. Sulfur metabolism in *Thiorhodaceae*. 1. Quantitative measurements on growing cells of *Chromatium okenii*. *Antonie van Leeuwenhoek.* 1964;30:225–38.
 42. Rinke C, Chuvochina M, Mussig AJ, Chaumeil P-A, Davin AA, Waite DW, et al. A standardized archaeal taxonomy for the Genome Taxonomy Database. *Nat Microbiol.* 2021;6:946–59.
 43. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Mol Biol Evol.* 2020;37:1530–1534.
 44. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol.* 2010;59:307–21.
 45. Anisimova M, Gascuel O. Approximate likelihood-ratio test for branches: A fast, accurate, and powerful alternative. *Syst Biol.* 2006;55:539–52.
 46. Nakamura T, Yamada KD, Tomii K, Katoh K. Parallelization of MAFFT for largescale multiple sequence alignments. *Bioinformatics* 2018;34:2490–2492.
 47. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haesele A, Jermini LS. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods.* 2019;14:587–589.
 48. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 2009;25:1972–1973.
 49. Kamyshny A, Goifman A, Rizkov D, Lev O. Formation of carbonyl sulfide by the reaction of carbon monoxide and inorganic polysulfides. *Environ Sci Technol.* 2003;37:1865–72.
 50. Zeer-Wanklyn CJ, Zamble DB. Microbial nickel: cellular uptake and delivery to enzyme centers. *Curr Opin Chem Biol.* 2017;37:80–88.
 51. Alfano M, Pérard J, Carpentier P, Basset C, Zambelli B, Timm J, et al. The carbon monoxide dehydrogenase accessory protein Cool is a histidine-rich multidomain dimer containing an unexpected Ni(II)-binding site. *J Biol Chem.* 2019;294:7601–14.
 52. Rowinska-Zyrek M, Zakrzewska-Czerwinska J, Zawilak-Pawlak A, Kozłowski H. Ni²⁺ chemistry in pathogens - a possible target for eradication. *Dalton Trans.* 2014;43:8976–89.
 53. Sorokin DY. Microbial utilization of glycine betaine in hypersaline soda lakes. *Microbiol (Mosc).* 2021;90:567–75.
 54. Elling FJ, Becker KW, Könneke M, Schröder JM, Kellermann MY, Thomm M, et al. Respiratory quinones in Archaea: phylogenetic distribution and application as biomarkers in the marine environment. *Environ Microbiol.* 2016;18:692–707.
 55. Buckel W, Thauer RK. Flavin-based electron bifurcation, ferredoxin, flavodoxin, and anaerobic respiration with protons (Ech) or NAD⁺ (Rnf) as electron acceptors: a historical review. *Front Microbiol.* 2018;9:401.
 56. Buckel W, Thauer RK. Flavin-based electron bifurcation, a new mechanism of biological energy coupling. *Chem Rev.* 2018;118:3862–86.
 57. Tian L, Lo J, Shao X, Zheng T, Olson DG, Lynda LR. Ferredoxin:NAD oxidoreductase of *Thermoanaerobacterium saccharolyticum* and its role in ethanol formation. *Appl Environ Microbiol.* 2016;82:7134–41.
 58. Friedrich T, Scheide D. The respiratory complex I of bacteria, archaea and eukarya and its module common with membrane bound multisubunit hydrogenases. *FEBS Lett.* 2000;479:1–5.
 59. Grimaldi S, Schoepp-Cothenet B, Ceccaldi P, Guigliarelli B, Magalon A. The prokaryotic Mo/W-bisPGD enzymes family: A catalytic workhorse in bioenergetic. *Biochim Biophys Acta.* 2016;1827:1048–85.
 60. Duarte AG, Barbosa ACC, Ferreira D, Manteigas G, Domingos RM, Pereira IAC. Redox loops in anaerobic respiration - The role of the widespread NrfD protein family and associated dimeric redox module. *Biochim Biophys Acta.* 2021;1862:148416.
 61. Ledbetter RN, Garcia Costas AM, Lubner CE, Mulder DW, Tokmina-Lukaszewska M, Artz JH, et al. The electron bifurcating FixABCX protein complex from *Azotobacter vinelandii*: generation of low-potential reducing equivalents for nitrogenase catalysis. *Biochemistry.* 2017;56:4177–4190. 60
 62. Schut GJ, Mohamed-Raseek N, Tokmina-Lukaszewska M, Mulder DW, Nguyen DMN, Lipscomb GL. The catalytic mechanism of electron-bifurcating electron transfer flavoproteins (ETFs) involves an intermediary complex with NAD⁺. *J Biol Chem.* 2019;294:3271–83.
 63. Torregrosa-Crespo J, Pire C, Richardson DJ, Martínez-Espinosa RM. Exploring the molecular machinery of denitrification in *Haloferax mediterranei* through proteomics. *Front Microbiol.* 2020;11:605859.
 64. Simon J, Klotz MG. Diversity and evolution of bioenergetic systems involved in microbial nitrogen compound transformations. *Biochim Biophys Acta.* 2013;1827:114–35.
 65. Leu AO, Cai C, McIlroy SJ, Southam G, Orphan VJ, Yuan Z, et al. Anaerobic methane oxidation coupled to manganese reduction by members of the *Methanoperedenaceae*. *ISME J.* 2020;14:1030–41.
 66. Miralles-Robledillo JM, Bernabeu E, Giani M, Martínez-Serna E, Martínez-Espinosa RM, Pire C. Distribution of denitrification among haloarchaea: a comprehensive study. *Microorganisms.* 2021;9:1669.
 67. Duval S, Ducluzeau A-L, Nitschke W, Schoepp-Cothenet B. Enzyme phylogenies as markers for the oxidation state of the environment: the case of respiratory arsenate reductase and related enzymes. *BMC Evol Biol.* 2008;8:206.

ACKNOWLEDGEMENTS

DS and AM were supported by the Russian Foundation for Basic Research (RFBR grant 19-04-00401) for field work and isolation of pure cultures and by the Russian Ministry of Science and Higher Education (genomic and phylogenomic analyses). DS and MP were also supported by the Gravitation-SIAM Program of the Dutch Ministry of Education, Culture and Science (grant 24002002). EM, MY and PG were supported by the European Commission's Horizon 2020 Program under FUTUREZYME Project (Contract 101000327). PG was supported by the Center for Environmental Biotechnology Project, partly funded by the European Regional Development Fund via the Welsh Assembly Government.

AUTHOR CONTRIBUTIONS

DS performed sampling, isolation and physiological characterisation of pure cultures. Bioinformatics analyses was carried out by DS, AM, EM and MY. MP and CT performed proteomic analysis. PG was responsible for the genome sequencing. The data were interpreted and manuscript was written by DS, AM and MY.

COMPETING INTERESTS

The authors declare no competing interests.

ADDITIONAL INFORMATION

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41396-022-01206-x>.

Correspondence and requests for materials should be addressed to Dimitry Y. Sorokin.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.