

Computation of topological phase diagram of disordered $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$ using the kernel polynomial method

Varjas, Dániel; Fruchart, Michel; Akhmerov, Anton R.; Perez-Piskunow, Pablo M.

DOI

[10.1103/PhysRevResearch.2.013229](https://doi.org/10.1103/PhysRevResearch.2.013229)

Publication date

2020

Document Version

Final published version

Published in

Physical Review Research

Citation (APA)

Varjas, D., Fruchart, M., Akhmerov, A. R., & Perez-Piskunow, P. M. (2020). Computation of topological phase diagram of disordered $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$ using the kernel polynomial method. *Physical Review Research*, 2(1), Article 013229. <https://doi.org/10.1103/PhysRevResearch.2.013229>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

Computation of topological phase diagram of disordered $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$ using the kernel polynomial method

Dániel Varjas,^{1,2,*} Michel Fruchart,^{3,4} Anton R. Akhmerov,² and Pablo M. Perez-Piskunow^{2,5,†}

¹*QuTech, Delft University of Technology, P.O. Box 4056, 2600 GA Delft, The Netherlands*

²*Kavli Institute of Nanoscience, Delft University of Technology, P.O. Box 4056, 2600 GA Delft, The Netherlands*

³*Instituut-Lorentz, Universiteit Leiden, P.O. Box 9506, 2300 RA Leiden, The Netherlands*

⁴*James Franck Institute and Department of Physics, University of Chicago, Chicago, Illinois 60637, USA*

⁵*Catalan Institute of Nanoscience and Nanotechnology (ICN2), CSIC and BIST, Campus UAB, Bellaterra, 08193 Barcelona, Spain*



(Received 5 September 2019; accepted 6 February 2020; published 28 February 2020)

We present an algorithm to determine topological invariants of inhomogeneous systems, such as alloys, disordered crystals, or amorphous systems. Based on the kernel polynomial method, our algorithm allows us to study samples with more than 10^7 degrees of freedom. Our method enables the study of large complex compounds, where disorder is inherent to the system. We use it to analyze $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$ and tighten the critical concentration for the phase transition. Moreover, we obtain the topological phase diagram for related alloys in the family of three-dimensional mirror Chern insulators.

DOI: [10.1103/PhysRevResearch.2.013229](https://doi.org/10.1103/PhysRevResearch.2.013229)

I. INTRODUCTION

Topological materials have attracted continuing interest from both the fundamental physics and the material science communities for the last decade [1,2]. The program to theoretically classify noninteracting crystalline insulators has been completed, tabulating possible topological phases in all space groups [3–5]. Recent efforts focus on automated high-throughput methods to discover and classify new topological materials [6–8], culminating in the production of comprehensive databases of topological insulators and semimetals [8–12].

However, not all topological insulators are compounds with perfect stoichiometry. The first three-dimensional (3D) topological insulator to be predicted [13] and experimentally realized [14] was an alloy— $\text{Bi}_x\text{Sb}_{1-x}$. These systems are usually studied using the virtual crystal approximation (VCA) or the coherent potential approximation, which approximate an alloy by a perfect crystal [15,16]. This approach ignores the intrinsic disorder in alloys, and it is insufficient to explain topological transitions that appear at strong disorder [17,18], or accurately find critical concentrations.

The topological invariant converges to its bulk value in samples larger than the localization length ξ . This is the main limitation in resolving topological phase transitions as ξ diverges. Therefore, the asymptotic scaling of the computational

cost with ξ is the main distinction between different numerical approaches.

Available methods to compute topological invariants either apply the momentum space Berry curvature formalism to periodic systems with a disordered supercell or use a real-space formulation on a large finite sample [19–27]. However, these methods involve solving at least one eigenvalue equation with size equal to the number of degrees of freedom, resulting in the complexity ξ^{3d} in d dimensions. This restricts the applicability of such methods to small system sizes, especially in three dimensions. To our knowledge, the most efficient method in three dimensions is the scattering matrix approach [28], with a complexity scaling as $\xi^{3(d-1)}$, allowing for maximum sizes of 5×10^5 degrees of freedom.

We present an algorithm to efficiently identify topological phases of strongly disordered systems using the kernel polynomial method (KPM) [29–31], an approximation based on a polynomial expansion of the quantities of interest. All topological properties of a noninteracting system of electrons are encoded in the projector on the occupied states (spectral projector), which is efficiently approximated using KPM. Our algorithm builds on the method of topological markers [22] to construct a topological invariant as the trace of an operator. Since topological invariants are integers, it is sufficient to reduce the statistical uncertainty below 1/2 to obtain the exact value. In particular, the stochastic evaluation of traces [29] from a small number of random vectors, combined with KPM, is suited for this task. With ξ^{d+1} scaling of the computational effort, it is the most efficient method in three dimensions.

As a concrete example, we apply our method to lead-tin-telluride $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$ alloys—three-dimensional topological crystalline insulators characterized by a mirror Chern number. Thanks to the efficiency of our algorithm, we are able to analyze the topological properties of large systems that are hardly accessible with previously existing techniques. We study 3D

*dvarjas@gmail.com

†pablo.perez.piskunow@gmail.com

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.




FIG. 1. Top panels: Surface spectra of a $20 \times 80 \times 80$ sample of $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$ in the trivial (a) and topological (b) phase. The presence of a gapless surface Dirac cone indicates the mirror Chern insulator phase. (c) Transition between trivial and mirror Chern phase when varying x for $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$ calculated using our method with various system sizes. Inset: Finite-size collapse of the curves with $x_c \simeq 0.28(3)$ and $\nu \simeq 0.9(6)$.

systems with linear sizes of over 100 lattice constants ($L > 100$), and more than 10^7 degrees of freedom (see Fig. 1). In contrast to previous theoretical estimations for the case of $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$ [32–34], we find a critical concentration that matches the one found experimentally [35–41].

II. REVIEW OF EXISTING ALGORITHMS

We focus on the vicinity of disorder-driven phase transitions, where the localization length ξ diverges. The finite but potentially large value of ξ defines the relevant length scale of our problem. Two regions that are closer than ξ feel each other's presence. Therefore, with open boundary conditions, the bulk has to be further than ξ to the edge; analogously, with periodic boundary conditions (PBC) when the linear size of the system is smaller than ξ it hosts states the extent of which is larger than the system size. These states span the whole system and overlap with themselves because of the finite size. In order to simulate the bulk, the system needs to have a linear size $L \gtrsim \xi$. Fluctuations of local quantities resulting from disorder scale with ξ , so averaging over a larger sample provides a good approximation of the thermodynamic limit.

The momentum space formalism of topological invariants applies to disordered systems by studying a periodic system with a large disordered supercell of volume ξ^d . This is equivalent to taking a finite torus and threading fluxes through its cycles [20,42,43]. The final formula of the invariant is identical to the momentum space Berry curvature treatment applied to the supercell. Other approaches include the Bott index [21], topological markers [22], pseudospectra [23], and noncommutative index theorems [24–27]. All of these methods involve diagonalization of a matrix of size proportional to the volume of the system. Diagonalization scales as N^3 with

the number of degrees of freedom N , so the computational cost of such methods is order ξ^{3d} , restricting them to small system sizes in three dimensions.

The scattering invariant formalism [44] avoids full diagonalization and only requires the knowledge of the scattering matrix at the Fermi level. The most efficient known algorithm for computing the scattering matrix is based on the nested dissection method [45] and scales as ξ^{3d-3} for $d > 1$.

An alternative way to obtain the Chern number in two-dimensional (2D) systems is to compute the Hall conductivity. An efficient method to compute the Kubo-Bastin conductivity is given by García *et al.* [30] for large disordered systems. This approach uses a two-dimensional KPM expansion to construct a matrix kernel that holds the information of the Hall conductivity for all energies, which is integrated to compute the total Hall conductivity for any Fermi energy. The time and memory requirements of constructing the matrix kernel are large, of order NM^2 for a system with N orbitals and a KPM expansion of order M . Using our method, however, we compute the Chern number for a single value of the Fermi energy with time scaling NM and memory requirement of order N .

III. GENERAL STRATEGY

Computing the exact spectral projector

$$\hat{P} = \theta(E_F - \hat{H}) = \sum_{n: E_n < E_F} |n\rangle \langle n| \quad (1)$$

by full diagonalization of the Hamiltonian is numerically expensive. Instead, we approximate the projector using the kernel polynomial method with the Jackson kernel [29], detailed in Appendix A. For a d -dimensional system of linear size L the computational cost scales linearly with the number of degrees of freedom L^d , and with the number of moments M —the order of the expansion. The order of the expansion sets a real-space cutoff in the approximate projector, which is an M th-order polynomial of the finite-range Hamiltonian. In an insulating system, the projector is a local operator with matrix elements $\langle \mathbf{x} | \hat{P} | \mathbf{x}' \rangle \propto \exp(-|\mathbf{x} - \mathbf{x}'|/\xi)$ that have a decay length ξ . Hence, the error of the approximation scales as $\exp(-M/\xi)$, and the number of moments necessary for fixed precision scales linearly with the localization length as $M \sim \xi$ [46,47]. See Appendix B.

We use the topological marker formalism introduced by Bianco and Resta [22]. All \mathbb{Z} -valued topological markers are a partial trace per unit volume of a local operator \hat{v} :

$$\nu = \text{Tr}_S(\hat{v}) = \frac{1}{|S|} \sum_{\lambda, \mathbf{x} \in S} \langle \mathbf{x}, \lambda | \hat{v} | \mathbf{x}, \lambda \rangle, \quad (2)$$

where the sum runs over the sites \mathbf{x} inside the subsystem S with volume $|S|$, and their internal degrees of freedom λ . The operator \hat{v} is a polynomial of the spectral projector, position, and symmetry operators, such that ν is dimensionless and independent of the detailed energetics or the overall length scale of the system. The marker coincides with the momentum space invariant in periodic systems, and converges to a quantized integer for large S in insulating homogeneous disordered systems [22].

An example of a topological marker is the real-space expression for the Chern number [22]:

$$C = \text{Tr}_A \hat{C} = 2\pi i \text{Tr}_A [\hat{P}\hat{x}\hat{P}, \hat{P}\hat{y}\hat{P}]. \quad (3)$$

Here, A is the area of the subsystem, \hat{x} and \hat{y} are the two components of the position operator, and $[\cdot, \cdot]$ is the commutator. Topological markers for all strong and weak \mathbb{Z} -valued topological invariants have similar algebraic expressions of the projected position operators [2,48,49], making a straightforward application of our method to these cases [50]. We are not aware of similar formulations of \mathbb{Z}_2 topological indices suitable for KPM.

To estimate the trace per volume, we use the stochastic trace approximation [29]:

$$\text{Tr}_S(\hat{O}) \approx \frac{1}{R|S|} \sum_{i=1}^R \langle r_i | \hat{O} | r_i \rangle, \quad (4)$$

where $|r_i\rangle$ are random-phase vectors localized in the region S . The standard error of this approximation scales as $\sqrt{\xi^d/(R|S|)}$, meaning that the number of random vectors R required for a given precision is constant if the system size is proportional to the localization length (see Appendix B).

We build a supercell of size $L > \xi$. To ensure that the invariant ν obtained with the trace of the local marker converges to the momentum space topological invariant we must repeat the supercell with the disorder realization over the whole space. Since the approximation of the local operator $\hat{\nu}$ is localized with $\xi < L$, it is sufficient to repeat two times the supercell in every spatial direction, and compute the topological invariant as an average of the topological marker over the central L^d volume. For a detailed description, see Appendix C.

The resulting complexity of the computation depends linearly on the number of random vectors R used (typically of order 1), on the number of moments M , and on the number of sites of the system L^d . We use a sparse representation of the short-ranged Hamiltonian. As a result the memory requirement scales linearly with the system size L^d and is independent of other parameters. Setting all quantities to their minimal values ($L \sim \xi$ and $M \sim \xi$) results in an algorithm with a computational cost scaling of ξ^{d+1} .

IV. APPLICATION TO MIRROR CHERN NUMBER

Our method provides better scaling than existing approaches in $d \geq 3$. We apply it to disordered 3D mirror Chern insulators. These are a widely studied class of topological crystalline materials with a \mathbb{Z} topological classification that relies on reflection symmetry [51]. Several experimental realizations are known, including alloys [34,37–41].

In a reflection-symmetric system of fermions, all wave functions are eigenstates of the mirror operator \hat{M}_z , with eigenvalues $\pm i$. The Chern numbers C_{\pm} for mirror-even and mirror-odd wave functions are

$$C_{\pm} = 2\pi i \text{Tr}_A [\tilde{x}_{\pm}, \tilde{y}_{\pm}], \quad (5)$$

where $\tilde{x}_{\pm} = \hat{M}_{\pm} \hat{P} \hat{x} \hat{P} \hat{M}_{\pm}$ and $\tilde{y}_{\pm} = \hat{M}_{\pm} \hat{P} \hat{y} \hat{P} \hat{M}_{\pm}$ are the projected position operators restricted to the mirror-even or mirror-odd subspaces. Here \hat{M}_{\pm} are the projectors on the

mirror-even and mirror-odd subspaces and A is the area in the xy plane. The total Chern number C is the sum of the Chern numbers for each subspace $C = C_+ + C_-$, while the mirror Chern number equals to their difference $C_M = (C_+ - C_-)/2$. In the presence of time-reversal invariance, the total Chern number vanishes ($C_+ = -C_-$), and the mirror Chern number $C_M = C_+$ counts the helical surface modes. Since the mirror operator $\hat{M}_z = i(\hat{M}_+ - \hat{M}_-)$ commutes with the projector \hat{P} and position operators \hat{x} and \hat{y} , we express the mirror Chern number as

$$C_M = \pi \text{Tr}_A (\hat{M}_z [\hat{P}\hat{x}\hat{P}, \hat{P}\hat{y}\hat{P}]). \quad (6)$$

In order to compute the mirror Chern number we consider a system with a disorder configuration that is mirror symmetric across the M_1 plane at $z = 0$. The PBC in the z direction results in another mirror plane M_2 across the boundary (see Appendix C). The bulk of this system is locally indistinguishable from a sample without reflection symmetry except for the two mirror planes M_1 and M_2 . Therefore as long as the two mirror planes do not undergo a two-dimensional topological transition, the mirror Chern number of the symmetric sample equals that of the bulk system [52].

V. TIGHT-BINDING MODEL OF $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$

Topological crystalline insulators protected by reflection symmetry [36] were theoretically predicted [51] and experimentally observed [34,37–41] in $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$ alloys. They host metallic surface states on the surfaces that are symmetric with respect to the mirror plane [2,51,53]. Lead-tin-telluride was studied using either the VCA [32,34] or *ab initio* methods [33], finding a gap closing and phase transition near $x = 0.35$ or 0.23, respectively. We use a tight-binding approach that captures long-range correlations, and find a different critical concentration.

We consider the substitutional disorder of the lead-tin-telluride alloy $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$ resulting from substituting some Pb ions for Sn ions. This disorder is nonmagnetic, and it preserves the reflection symmetry on average, which is sufficient to protect the gapless surface states [28,54]. We disregard other types of symmetry-breaking disorder appearing naturally in $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$ [54], such as ferroelectric structural distortion [51,54,55] and magnetic dopants [51,54,56].

In our investigation we use two atomistic tight-binding models. The first one includes 18 spinful s , p , and d orbitals on both sublattices, with 36 bands in total. This model accurately describes the energetics, using tight-binding parameters for both SnTe and PbTe derived from *ab initio* simulations [32].

To simulate the alloy, we substitute randomly Sn for Te with probability x . We incorporate substitutional disorder by using the hopping amplitudes of SnTe for Sn-Te bonds and PbTe amplitudes for Pb-Te bonds. The onsite parameters of Te atoms are slightly different in SnTe and PbTe; we therefore use a weighted average of these depending on the local environment. We also use the appropriate onsite terms, including $L \cdot S$ spin-orbit coupling (SOC), depending on the type of the Sn or Pb atom. Further details can be found in Appendix D.

When investigating the onsite energy dependent phase diagram of $X_{1-x}\text{Sn}_x\text{Te}$ alloys, we use a simplified model that




FIG. 2. Comparison of the critical concentration obtained with our method, against theoretical approximations (squares) and experimental measurements (circles). The shapes are empty for the trivial phase, half filled for the estimated transition point (transition range for Zhong *et al.* [41]), and fully filled for the topological phase.

only includes six spinful p orbitals, with 12 bands [51,57,58]. We include $L \cdot S$ SOC terms and first and second neighbor hoppings, with amplitudes that depend on the sublattices but not on the types of the atoms. We restrict the effect of disorder to different onsite energies on Sn and X sites. For more details, see Appendix E.

VI. RESULTS

We define the Hamiltonians and perform the KPM expansions using the `kwant` software package [59]. The code to reproduce the figures in this paper is available in Ref. [60]. First, we study the topological phase transition in the realistic 18-orbital model of $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$. We build a tight-binding model with PBC that preserves reflection symmetry and contains $W \times L_{1\bar{1}0} \times L_z$ unit cells, with 36 degrees of freedom each. For the largest system size used this means 13 824 000 degrees of freedom in total. This model accurately reproduces the energetics, resulting in full bandwidth of about 25 eV and band gap of less than 0.3 eV. In order to resolve the gap that is multiple orders of magnitude smaller than the bandwidth, we use $M = 5000$ moments in the calculation. We use $R = 5$ random vectors and 12 disorder realizations each. This increases the time cost but not the memory cost of the algorithm.

We perform finite-size collapse of the data [61] (see Appendix F) and find the transition point at $x_c = 0.28(3)$, with critical exponent $\nu = 0.9(6)$ accurately describing the transition (see Fig. 2). This result improves significantly from the VCA result [32,34,40] and *ab initio* [33], but is consistent with the most precise experimental data [39,41]. Experimental estimates are obtained with only two values of the concentration x for works [35,40,62] focused on characterizing the trivial and topological phases rather than determining the transition point.

To study a larger parameter space that includes other possible alloys of the $X_{1-x}\text{Sn}_x\text{Te}$ family that manifest the mirror Chern phase, we use the simplified six-orbital model. Besides the composition x , we also vary the onsite energy of the dopant cation X , approximating compounds with lighter or heavier ions and similar electronic structures. Figure 3 shows the phase diagram. We find that the phase boundary




FIG. 3. Mirror Chern number C_M (a) and localization length ξ (b) of the $X_{1-x}\text{Sn}_x\text{Te}$ model as a function of onsite energy m_X and composition x . In both panels the overlay shows the phase boundary in the VCA and the horizontal dashed line corresponds to $X = \text{Pb}$. ξ was calculated from the scaling of conductivity with sample sizes up to $L = 20$ with ten disorder realizations. C_M was calculated in a system of $40 \times 40 \times 60$ unit cells, 2 304 000 degrees of freedom, and $R = 5$ random vectors and averaged over four disorder realizations.

differs from the VCA method, where the topological index only depends on the average onsite energy of the cations.

VII. CONCLUSIONS

Our method allows the computation of topological invariants of realistic 3D alloys. Disorder in the crystalline structure is present in naturally found and artificially grown compounds, and it is inherent to substitutional alloys. However, a computationally efficient method to analyze topological properties of realistic disordered materials was missing.

We apply our method to study the critical concentration of $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$, and find an estimate that agrees with the most precise experimental data [63], whereas it is not possible to reconcile any of the other theoretical studies (known to us) with all the experimental data, especially Ref. [41].

The scaling of computational time with system size of our method is better than the scaling of other methods available in the literature. By using the kernel polynomial method, we achieve a computational time scaling of ξ^{d+1} with the localization length. Since we do not use eigenvalue solvers, only matrix-vector multiplication, the memory requirement only scales linearly with the sample volume as ξ^d .

Beyond Chern numbers, our formalism allows calculation of all \mathbb{Z} -valued strong and weak topological invariants in all dimensions [50]. This method makes the automated discovery of topological alloys feasible, and can guide synthesis of new alloys in the future. In this paper we use energetically accurate tight-binding models obtained from *ab initio* calculations performed on pure materials as input. Using these tight-binding amplitudes for the atoms and bonds that appear in the alloy, we generate large disordered samples with various concentrations. We are able to probe the topology, something that would not be accessible with other methods. Simulation of small clusters with various disorder configurations is feasible using *ab initio* methods [33]; tight-binding parameters obtained for all local environments would serve as a more accurate input for disordered models [47].

This paper opens several directions for future research. Our method is directly applicable to all types of disorder, as

well as quasicrystalline and amorphous systems in symmetry classes that admit the topological marker formalism [50]. This approach is not restricted to electrons in solids, and can be combined with finite element methods to analyze topology in disordered classical mechanical and photonic systems [64–67]. We expect our method to perform well in disordered time-reversal breaking Weyl semimetals with nonzero Hall conductivity; further refinements could extend it to the time-reversal invariant case. While we are not aware of a similar formulation of \mathbb{Z}_2 indices, KPM could be utilized to calculate quantized responses associated with these phases, such as the quantized magnetoelectric effect of 3D strong topological insulators [68]. A similar approach could also be applied to higher-order topological insulators to calculate multipole moments of the charge density [69]. We expect that KPM could be used to study a wide variety of related topics in condensed-matter physics, such as probing localization, topological Anderson insulators, or numerical renormalization-group studies of the topological markers. During the preparation of the paper, our method served as a base to compute higher-order topological invariants [70], and as an aide in the identification of topological phases of quasicrystals [71].

ACKNOWLEDGMENTS

We are grateful to M. D. Caio for discussions and insights on local topological markers, M. Hoskam for help with implementing the tight-binding models, and B. Nijholt for technical assistance with the numerical calculations. We thank S. Roche and A. Cummings for critical revision of the paper, and P. Brouwer for helpful discussions. This work was supported by European Research Council Starting Grant No. 638760, The Netherlands Organisation for Scientific Research (NWO/OCW) as part of the Frontiers of Nanoscience program, NWO VIDI Grant No. 680-47-53, the ONR, University of Chicago MRSEC through a Kadanoff-Rice postdoctoral fellowship, and the European Union’s Horizon 2020 Research and Innovation program under Grant No. 824140.

The project to study topological invariants using KPM was initiated by P.M.P.-P. and D.V., and the scope of the project was later refined using contributions from all authors. P.M.P.-P. and D.V. wrote the code used for the numerical calculations and performed calculations. D.V. performed the large-scale numerical calculations on the $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$ tight-binding models. All authors took part in the analysis of the method and writing the paper.

APPENDIX A: SPECTRAL PROJECTOR OPERATOR EXPANDED WITH THE KERNEL POLYNOMIAL METHOD

The band projector is expanded with the KPM [29] and it is a finite-range approximation, where the range depends linearly on the number of moments used in the expansion. Each moment of the expansion of the projector operator applied to a vector is obtained recursively, by applying the Hamiltonian to an expanded vector of the previous iteration. The recursive algorithm effectively *spreads* a local vector to the neighboring sites via the hopping terms of the Hamiltonian, as depicted in Fig. 4.




FIG. 4. Convergence of the projector operator, computed via a KPM expansion with M moments, applied to a vector $|x_0, y_0\rangle$ located at the center of a Chern insulator. The KPM expansion converges to the exact projector operator for $M = \infty$, and finite values of M yield finite range approximations of the projector. The error of the approximation scales with $\exp(-\xi/M)$ [29].

Concurrently, each order of the expansion increases the precision in energy of the approximation, therefore setting an energy resolution for the expansion. To apply KPM, operators are rescaled such that the spectrum is in the $[-1, 1]$ range. As a consequence, the order of the expansion required to resolve the mobility gap is proportional to W/Δ , the full bandwidth divided by the mobility gap. This can be a large ratio even if ξ is small, for example, in an atomic insulator with a large range of on-site energies and vanishing hoppings. Typically W/Δ and ξ increase together near a phase transition where the gap closes, but the bandwidth changes slowly, and does not affect the scaling of the computational cost with ξ .

The kernel polynomial method provides a stable and efficient method to expand the action of any function of an operator \hat{f} that depends on the Hamiltonian H and a set of parameters λ , on a vector $|v\rangle$ [29,30]. The expansion up to order M is

$$\begin{aligned} \hat{f}(\lambda, H) |v\rangle &= \sum_{m=0}^M g_m \mu_m(\lambda) T_m(H) |v\rangle \\ &= \sum_{m=0}^M g_m \mu_m(\lambda) |v_m\rangle, \end{aligned}$$

where g_m are the Jackson kernel coefficients [29]. The coefficients μ_m , called *moments* in the context of KPM expansions, are defined as

$$\mu_m(\lambda) = \frac{2}{\pi} \frac{1}{1 + \delta_{m,0}} \int_{-1}^1 \frac{\hat{f}(\lambda, E) T_m(E)}{\sqrt{1 - E^2}} dE,$$

and the vectors $|v_m\rangle$ satisfy the recursion relation

$$\begin{aligned} |v_0\rangle &= |v\rangle, \\ |v_1\rangle &= H |v_0\rangle, \\ |v_{m+1}\rangle &= 2H |v_m\rangle - |v_{m-1}\rangle. \end{aligned}$$

We approximate the projector operator defined as the step function:

$$\begin{aligned} \hat{P}(\varepsilon, H) &= \theta(\varepsilon - H), \\ \hat{P}(\varepsilon, H) &= \sum_{m=0}^M g_m \mu_m(\varepsilon) T_m(H), \end{aligned}$$

and in this case the coefficients take the form

$$\mu_m(\varepsilon) = \begin{cases} 1 - \frac{\arccos(\varepsilon)}{\pi} & m = 0 \\ \frac{-2 \sin[m \arccos(\varepsilon)]}{m\pi} & m \neq 0 \end{cases}.$$

Equipped with the KPM expanded projector, we proceed to evaluate matrix elements of topological markers. These are finite polynomials of \hat{P} and other sparse operators such as position and mirror. The matrix elements are evaluated by successive application of these operators to the states. The resulting memory cost scales linearly with the system size (number of degrees of freedom); by cumulatively summing up the expanded vectors for fixed E_F , only a small number of sparse matrices and dense vectors are stored at any given time. Most of the time cost comes from sparse matrix-vector multiplications, linear in the system size. The number of operations is proportional to the number of moments M .

APPENDIX B: SCALING OF STOCHASTIC TRACE

We optimize the calculation further by utilizing the stochastic trace approximation to evaluate the trace. We take R independent random-phase vectors $|r_i\rangle$ that are only nonzero inside the region S , $\langle x, l | r_i \rangle = \delta_{x \in S} \exp(i\phi_{x,l,i})$ with $\phi_{x,l,i} \in [0, 2\pi]$ independent random phases for all sites and orbitals. The trace of an operator \hat{O} equals the expectation value

$$\text{Tr} \hat{O} = \mathbb{E} \left(\frac{1}{R} \sum_{i=1}^R \langle r_i | \hat{O} | r_i \rangle \right) = \mathbb{E}(\text{Tr}_{\text{st}} \hat{O}), \quad (\text{B1})$$

where \mathbb{E} denotes the expectation value over random vector realizations and we introduced the notation $\text{Tr}_{\text{st}} \hat{O}$ for the random variable giving the stochastic trace of operator \hat{O} . The above equality is proved by using that the random phases are independent, hence $\mathbb{E}(e^{i(\phi_{x,l,i} - \phi_{x',l',i})}) = \delta_{x,x'} \delta_{l,l'}$ and only the diagonal entries contribute to the expectation value.

The standard deviation of the stochastic trace of an operator scales with the total square magnitude of the off-diagonal entries which enter in the expectation value with random phases [29]:

$$\sigma(\text{Tr}_{\text{st}} \hat{O}) = \sqrt{\frac{1}{R} \sum_{i \neq j} |\hat{O}_{ij}|^2}, \quad (\text{B2})$$

where $\sigma(X) = \sqrt{\mathbb{E}(|X|^2) - |\mathbb{E}(X)|^2}$ is the standard deviation and we used that $\sigma(e^{i\phi_{x,l,i}}) = 1$. We also use that the standard deviation of the sum of independent random variables obeys $\sigma(\sum_i X_i) = \sqrt{\sum_i \sigma(X_i)^2}$.

We are concerned with the stochastic trace of topological markers, such as the Chern and mirror Chern operators. In order to draw conclusions, we need to know the scaling of the off-diagonal matrix elements with respect to the relevant length scales in the problem. There are three length scales, the lattice constant a , the localization length ξ , and the system size L (this we take to be the linear size of the subsystem where we take the partial trace; the overall system size is a constant factor larger). We use units of a to measure the other two distances, and, as explained in the main text, we




FIG. 5. Off-diagonal matrix elements of the Chern operator in two dimensions as a function of real-space distance in the units of the localization length. Here we use a simple continuum model of a Chern insulator, discretized on a square lattice with various lattice constants a and fixed ξ . The collapse of the curves verifies the scaling form of the matrix elements that we use.

are interested in systems the size of which is proportional to the localization length, so we will set $L = c\xi$ in the end. We introduce a as the lattice constant here for clarity, but it is an arbitrary reference length scale we can define in fully disordered (e.g., amorphous) systems as well, for example, as the typical spacing of sites. As it cancels from the final result, this argument does not rely on the assumption of an underlying regular lattice.

We start with the Chern operator in two dimensions:

$$\hat{C} = \frac{2\pi i}{a^2} [\hat{P}\hat{x}\hat{P}, \hat{P}\hat{y}\hat{P}], \quad (\text{B3})$$

where the a^{-2} prefactor is included to measure all distances in units of a ; this way the sum of the diagonal entries of \hat{C} on a site coincides with the Chern number in a clean system. We numerically verify (see Fig. 5) that the off-diagonal matrix elements scale as

$$\langle x, y | \hat{C} | x', y' \rangle = f \left(\frac{x - x'}{\xi}, \frac{y - y'}{\xi} \right), \quad (\text{B4})$$

where f is a dimensionless function and we suppressed the dependence on the internal degrees of freedom. f is a quickly decaying function for $(x - x')/\xi \gg 1$ in insulating systems, as matrix elements of \hat{P} also decay at the length scale of ξ .

The Chern marker averaged over a square region S of size L around the origin is given by

$$C = \text{Tr}_{\text{st}} \left[\left(\frac{a}{L} \right)^2 \hat{C} \right], \quad (\text{B5})$$

where we still measure length in units of a . Substituting (B4) we find for the standard deviation of C

$$\begin{aligned} \sigma(C) &= \frac{1}{\sqrt{R}} \sqrt{\sum_{r \neq r' \in S} \left| \left(\frac{a}{L} \right)^2 f \left(\frac{\mathbf{r} - \mathbf{r}'}{\xi} \right) \right|^2} \\ &= \frac{1}{\sqrt{R}} \sqrt{\left(\frac{a}{L} \right)^4 \frac{1}{a^4} \int_S d^2 r d^2 r' \left| f \left(\frac{\mathbf{r} - \mathbf{r}'}{\xi} \right) \right|^2} \end{aligned}$$




FIG. 6. Contributions per layer to the mirror Chern number. The shade represents the standard deviation of the stochastic trace per layer, calculated with ten random vectors per layer.

$$\begin{aligned}
 &= \frac{1}{\sqrt{R}} \sqrt{\frac{1}{L^4} \xi^4 \int_{-c/2}^{c/2} d^2\tilde{\mathbf{r}} d^2\tilde{\mathbf{r}}' |f(\tilde{\mathbf{r}} - \tilde{\mathbf{r}}')|^2} \\
 &= \frac{1}{c^2 \sqrt{R}} \sqrt{\int_{-c/2}^{c/2} d^2\tilde{\mathbf{r}} d^2\tilde{\mathbf{r}}' |f(\tilde{\mathbf{r}} - \tilde{\mathbf{r}}')|^2}. \quad (\text{B6})
 \end{aligned}$$

In the second line we took the limit of $\xi \gg a$, so we can replace sums over sites with integrals as $\sum_{x \in [-L/2, L/2]} = 1/a \int_{-L/2}^{L/2} dx$. In the third line we changed the integration variables to $\tilde{\mathbf{r}} = \mathbf{r}/\xi$. The result is only dependent on the ratio of the localization length and the system size $c = L/\xi$ and the number of random vectors R , but not ξ . As the integral is proportional to the integration area, c^2 for $c \gg 1$, the overall scaling of the error with c is $1/\sqrt{Rc^2}$.

We find a similar scaling for other topological markers, such as the 3D winding number in chiral classes [48,49]. In general, the standard deviation of the stochastic trace evaluation of the invariant depends only on the ratio of the system size and the localization length:

$$\sigma(\hat{\nu}) \propto \sqrt{\frac{1}{R} \left(\frac{\xi}{L}\right)^d}. \quad (\text{B7})$$

For the mirror Chern operator

$$\hat{C}_M = \frac{\pi}{a^2} \hat{M}_z [\hat{P} \hat{x} \hat{P}, \hat{P} \hat{y} \hat{P}] \quad (\text{B8})$$

the scaling is

$$\langle x, y, z | \hat{C}_M | x', y', z' \rangle = \frac{a}{\xi_z} f\left(\frac{x-x'}{\xi}, \frac{y-y'}{\xi}, \frac{z+z'}{\xi_z}\right), \quad (\text{B9})$$

where ξ_z is the localization length in the z direction. This form is justified by the fact that the contributions to the mirror Chern number are centered on the invariant planes, but are spread out on layers in a thickness proportional to ξ_z (see Fig. 6). The total for all layers is, however, constant, hence the a/ξ_z prefactor. This is the key difference compared

to the Chern number; the mirror Chern number is effectively a 2D invariant that we evaluate on a thick slab. In a clean system every plane parallel to a mirror plane is also a mirror plane, hence the matrix element can only depend on $z+z'$. The mirror marker averaged over a square region of size L is given by

$$C_M = \text{Tr}_{\text{st}} \left[\left(\frac{a}{L}\right)^2 \hat{C}_M \right], \quad (\text{B10})$$

and we find using a similar derivation for the standard deviation (setting $\xi = \xi_z = L/c$)

$$\sigma(C_M) = \frac{1}{c^2 \sqrt{R}} \sqrt{\int_{-c/2}^{c/2} d^3\tilde{\mathbf{r}} d^3\tilde{\mathbf{r}}' |f(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}')|^2}, \quad (\text{B11})$$

which is only dependent on the ratio of the localization length and the system size c and the number of random vectors R .

In the numerical calculations we split the stochastic trace in two halves, using two sets of random vectors, each localized in one half of the system separated by mirror planes. This eliminates most of the large off-diagonal entries with $z = -z'$, resulting in a constant factor reduction in the error. Splitting the stochastic trace into more regions (e.g., separate for each layer parallel to the mirror plane) results in further reduction in the error, at the cost of increased computational effort. The overall scaling of the computational time with ξ for a fixed standard deviation is the same up to a constant factor for all of these schemes, ξ^{d+1} .

APPENDIX C: GEOMETRY USED IN THE NUMERICS

As described in the main text, we build a tight-binding model with PBC using translation vectors $W[1, 1, 0]$, $L_{1\bar{1}0}[1, -1, 0]$, and $L_z[0, 0, 1]$. This geometry preserves the reflection symmetry with $[1, 1, 0]$ normal and contains $W \times L_{1\bar{1}0} \times L_z$ unit cells with 36 degrees of freedom each. The averaging region of the stochastic trace extends the full width of the system in the $[1, 1, 0]$ direction and contains half of the linear size in the perpendicular directions, as depicted by the inner box in Fig. 7. Imposing PBC in all directions eliminates gapless surface states, and the (mobility) gap guarantees that the Fermi projector is short ranged.

However, imposing PBC in the direction normal to the mirror planes results in two mirror invariant planes. As argued in the main text, in a sample with open boundary conditions in the other directions, this results in a doubling of the interface modes, around the edges of the mirror invariant planes. On the other hand, C_M counts helical modes; while the 3D mirror Chern number is given by the number of chiral modes, this factor of 1/2 cancels the previous factor of 2. We conclude that Eq. (6) is applicable to this 3D M_z symmetric geometry with PBC in z .

APPENDIX D: 18-ORBITAL TIGHT-BINDING MODEL

We use the 18-orbital model of SnTe and PbTe derived in Ref. [32]. The cubic rock-salt structure has two sublattices A and C (referring to the anion and cation nature of the atoms occupying them), the first occupied by Te and the second by Sn or Pb atoms. Each site hosts spinful s , p , and d orbitals, 18




FIG. 7. Geometry used for calculating the mirror Chern number. The upper and lower halves of the slab have mirror image disorder configurations with mirror plane M_1 . Because of the PBC in the z direction, there is a second mirror plane M_2 . The central pane with stars is a schematic representation of the disorder, repeated in the x and y directions, and mirrored in z . The box in the center of the sample shows the averaging region where the partial trace is evaluated, containing one period of the disorder configuration in the x and y directions.

degrees of freedom in total, with annihilation operators c_l ($l = 0, 1, 2$ for s, p, d orbitals, respectively), which is a vector of length 2, 6, or 10 depending on the value of l .

The hopping terms are expressed as two-center integrals $H_{l'l'md}$ in the *linear combination of atomic orbitals* method [72], where l and l' are the total angular momenta of the orbitals connected on the two sites and m is the angular momentum of the bonding along the bonding axis d ($m = 0, 1, 2$ for σ, π, δ bonding, respectively). The matrices $H_{l'l'md}$ are $2(2l+1) \times 2(2l'+1)$ and are proportional to the identity in spin space. The onsite terms contain different onsite energies for the various orbitals E_l and $L \cdot S$ SOC terms with




FIG. 8. Density of states as a function of x near the bulk gap in the 18-band model of $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$. The red line shows the placement of the Fermi level.

TABLE I. Tight-binding parameters in electron volts for SnTe and PbTe from Lent *et al.* [32]. Note that we use opposite sign conventions for $V_{sp\sigma}$ and $V_{ps\sigma}$. All other parameters not listed here vanish.

	SnTe	PbTe
E_{sc}	-6.578	-7.612
E_{sa}	-12.067	-11.002
E_{pc}	1.659	3.195
E_{pa}	-0.167	-0.237
E_{dc}	8.38	7.73
E_{da}	7.73	7.73
λ_{pc}	0.592	1.500
λ_{pa}	0.564	0.428
$V_{ss\sigma}$	-0.510	-0.474
$V_{sp\sigma}$	-0.949	-0.705
$V_{ps\sigma}$	0.198	-0.633
$V_{pp\sigma}$	2.218	2.066
$V_{pp\pi}$	-0.446	-0.430
$V_{pd\sigma}$	-1.11	-1.29
$V_{pd\pi}$	0.624	0.835
$V_{dp\sigma}$	-1.67	-1.59
$V_{dp\pi}$	0.766	0.531
$V_{dd\sigma}$	-1.72	-1.35
$V_{dd\delta}$	0.618	0.668

strength λ_l . The tight-binding Hamiltonian reads

$$H = \sum_{l,r} E_{lr} c_{lr}^\dagger \cdot c_{lr} + \sum_{l,r} \lambda_{lr} c_{lr}^\dagger (\mathbf{L}_l \cdot \mathbf{S}) c_{lr} + \sum_{l,l',m,(r,r')} V_{l,l',m,r,r'} c_{l'r'}^\dagger H_{l'l'm(r-r')} c_{lr}. \quad (\text{D1})$$

The first term is the onsite energy, and it is the main source of disorder in our simulation. For sites on the C sublattice the type of the site (Sn or Pb) is chosen randomly with probability $1-x$ and x . The value of E_{lr} is assigned accordingly to be E_{lc}^{SnTe} and E_{lc}^{PbTe} , respectively. The superscripts SnTe and PbTe refer to the two sets of parameters for the two pure materials. If $\mathbf{r} \in A$, we use a weighted average $E_{lr} = [nE_{la}^{\text{SnTe}} + (6-n)E_{la}^{\text{PbTe}}]/6$ where n is the number of nearest-neighbor sites occupied by Sn atoms. The second term is the $L \cdot S$ spin-orbit coupling; \mathbf{L}_l is the vector of angular momentum l operators (zero for $l = 0$). The values of λ_{lr} are assigned in the same fashion, depending on the type of atoms. The third term describes nearest-neighbor hopping terms in the [001] and equivalent crystal directions; the sum runs over all nearest-neighbor pairs with $\mathbf{r} \in A$ and $\mathbf{r}' \in C$. Depending on the atoms at sites \mathbf{r} and \mathbf{r}' the value of $V_{l,l',m,r,r'}$ is set to $V_{l,l',m}^{\text{SnTe}}$ if one of the sites is Sn or $V_{l,l',m}^{\text{PbTe}}$ if one of the sites is Pb. All of the onsite energies and hopping terms are spin independent; SOC only enters through the onsite SOC terms. We summarize the parameter values used for numerical results in Table I.

Because of the identical outer-shell electronic structure of Sn and Pb, the alloy composition x does not affect the doping level, therefore we set the Fermi level E_F to ensure half filling for all compositions (see Fig. 8).

TABLE II. Tight-binding parameters in electron volts for SnTe used in the six-orbital model. We use the same Hamiltonian as Sessi *et al.* [74], but the numerical values of the parameters differ due to different normalization and sign conventions.

	SnTe
m_{Te}	-1.65
m_{Sn}	1.65
t_{aa}	-0.5
$t_{ac} = t_{ca}$	0.9
t_{cc}	0.5
λ_a	-0.3
λ_c	-0.3

APPENDIX E: SIX-ORBITAL TIGHT-BINDING MODEL

We adopt the six-orbital model of SnTe and PbTe originally described in Mitchell and Wallis [57] and used in Refs. [51,58,73,74]. Each site hosts spinful p orbitals, six degrees of freedom in total with a vector of annihilation operators c . This Hamiltonian is formally identical to (D1) but only includes p orbitals and $pp\sigma$ hopping, while the hopping range is extended to second neighbors. The tight-binding Hamiltonian reads

$$H = \sum_r m_r c_r^\dagger \cdot c_r + \sum_r \lambda_r c_r^\dagger (\mathbf{L} \cdot \mathbf{S}) c_r + \sum_{\langle(r,r')\rangle} t_{r,r'} c_r^\dagger [\mathbb{1} - (\hat{\mathbf{d}}_{r,r'} \cdot \mathbf{L})^2] c_r. \quad (\text{E1})$$

The first term is the onsite energy (also termed the “mass term”), and it is the main source of disorder in our simulation. m_r takes the value of m_{Te} on the A sublattice, while for the B sublattice a value is chosen between m_{Sn} and m_X with probability $1 - x$ and x . The second term is the $\mathbf{L} \cdot \mathbf{S}$ spin-orbit coupling; its value depends on the sublattice only (identical for Sn and X). The third term is a $pp\sigma$ type of hopping [72] that only connects p orbitals oriented along the direction of the bond $\hat{\mathbf{d}}_{r,r'}$.

We include first neighbor [001] and second neighbor [110] hoppings, with amplitudes that depend on the sublattices. We summarize the parameter values used for numerical results in Table II.

Because of the identical outer-shell electronic structure of Sn and Pb, the alloy composition x does not affect the doping level, therefore we set the Fermi level E_F to ensure half filling for all compositions.




FIG. 9. Logarithm of the cost function $\Phi(x_c, \nu)$ as a function of the parameters (x_c, ν) . The region D allowed by uncertainties around the minimum is delimited by a thick black dashed line.

APPENDIX F: EVALUATION AND UNCERTAINTY OF THE FINITE-SIZE SCALING PARAMETERS

We start from the data $C_M = f_i(x)$ of the mirror Chern number as a function of the concentration of $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$ where $i = 1, \dots, N$ correspond to different system sizes L_i . These data are obtained by averaging multiple disorder realizations. We wish to perform a finite-size scaling collapse of the data. The changes of variables $\tilde{x}_i = (x - x_c)L_i^{1/\nu}$ define new functions $\tilde{x} \mapsto \tilde{f}_i(\tilde{x})$, where both the critical concentration x_c and the correlation length exponent ν are parameters to estimate so that the curves for different L collapse onto a universal master curve. Hence, we seek to minimize the distance between $\tilde{f}_i(\tilde{x})$ over their common domain of definition. To do so, we interpolate the discrete data to perform the change of variable, evaluate $\tilde{f}_i(\tilde{x})$ over a fixed interval, and compute the variance of the data. This defines a cost function $\Phi(x_c, \nu)$ that is minimal at optimal parameters $(x_c^*, \nu^*) \simeq (0.28, 0.9)$. The cost function is plotted as a function of the parameters (x_c, ν) in Fig. 9. To evaluate the uncertainty on the optimal parameters, we evaluate the cost function $\Phi(x_c^*, \nu^*)$ at the optimal parameters (x_c^*, ν^*) for different disorder realizations [that give different values of the initial data $C_{M,L}(x)$]. The standard deviation gives an estimation of the uncertainty $u(\Phi)$ on the cost function. The domain $D = \{(x_c, \nu) \mid \Phi(x_c, \nu) \leq \Phi(x_c^*, \nu^*) + u(\Phi)\}$ where the cost function is closer to its minimum than the uncertainty $u(\Phi)$ is approximately an ellipse (see Fig. 9) from which the uncertainties $u(x_c) \simeq 0.03$ and $u(\nu) \simeq 0.6$ can be directly estimated.

- [1] X.-L. Qi and S.-C. Zhang, *Rev. Mod. Phys.* **83**, 1057 (2011).
 [2] C.-K. Chiu, J. C. Y. Teo, A. P. Schnyder, and S. Ryu, *Rev. Mod. Phys.* **88**, 035005 (2016).
 [3] J. Kruthoff, J. de Boer, J. van Wezel, C. L. Kane, and R.-J. Slager, *Phys. Rev. X* **7**, 041069 (2017).

- [4] H. C. Po, A. Vishwanath, and H. Watanabe, *Nat. Commun.* **8**, 50 (2017).
 [5] B. Bradlyn, L. Elcoro, J. Cano, M. G. Vergniory, Z. Wang, C. Felser, M. I. Aroyo, and B. A. Bernevig, *Nature (London)* **547**, 298 (2017).

- [6] K. Yang, W. Setyawan, S. Wang, M. B. Nardelli, and S. Curtarolo, *Nat. Mater.* **11**, 614 (2012).
- [7] S. Curtarolo, G. L. W. Hart, M. B. Nardelli, N. Mingo, S. Sanvito, and O. Levy, *Nat. Mater.* **12**, 191 (2013).
- [8] Z. Zhang, R.-W. Zhang, X. Li, K. Koepf, Y. Yao, and H. Zhang, *J. Phys. Chem. Lett.* **9**, 6224 (2018).
- [9] F. Tang, H. C. Po, A. Vishwanath, and X. Wan, *Nature (London)* **566**, 486 (2019).
- [10] F. Tang, H. C. Po, A. Vishwanath, and X. Wan, *Nat. Phys.* **15**, 470 (2019).
- [11] T. Zhang, Y. Jiang, Z. Song, H. Huang, Y. He, Z. Fang, H. Weng, and C. Fang, *Nature (London)* **566**, 475 (2019).
- [12] M. G. Vergniory, L. Elcoro, C. Felser, N. Regnault, B. A. Bernevig, and Z. Wang, *Nature (London)* **566**, 480 (2019).
- [13] L. Fu and C. L. Kane, *Phys. Rev. B* **76**, 045302 (2007).
- [14] D. Hsieh, D. Qian, L. Wray, Y. Xia, Y. S. Hor, R. J. Cava, and M. Z. Hasan, *Nature (London)* **452**, 970 (2008).
- [15] S. Chadov, J. Kiss, J. Kübler, and C. Felser, *Phys. Status Solidi: Rapid Research Letters* **7**, 82 (2013).
- [16] D. Sante, P. Barone, E. Plekhanov, S. Ciuchi, and S. Picozzi, *Sci. Rep.* **5**, 11285 (2015).
- [17] J. Li, R.-L. Chu, J. K. Jain, and S.-Q. Shen, *Phys. Rev. Lett.* **102**, 136806 (2009).
- [18] C. W. Groth, M. Wimmer, A. R. Akhmerov, J. Tworzydło, and C. W. J. Beenakker, *Phys. Rev. Lett.* **103**, 196805 (2009).
- [19] A. Kitaev, *Ann. Phys. (NY)* **321**, 2 (2006).
- [20] A. M. Essin and J. E. Moore, *Phys. Rev. B* **76**, 165307 (2007).
- [21] T. A. Loring and M. B. Hastings, *Europhys. Lett.* **92**, 67004 (2010).
- [22] R. Bianco and R. Resta, *Phys. Rev. B* **84**, 241106(R) (2011).
- [23] T. A. Loring, *Ann. Phys. (NY)* **356**, 383 (2015).
- [24] E. Prodan, *Appl. Math. Res. eXpress* **2013**, 176 (2013).
- [25] E. Prodan and H. Schulz-Baldes, *Bulk and Boundary Invariants for Complex Topological Insulators* (Springer, New York, 2016).
- [26] Y. Akagi, H. Katsura, and T. Koma, *J. Phys. Soc. Jpn.* **86**, 123710 (2017).
- [27] H. Katsura and T. Koma, *J. Math. Phys.* **59**, 031903 (2018).
- [28] I. C. Fulga, B. van Heck, J. M. Edge, and A. R. Akhmerov, *Phys. Rev. B* **89**, 155424 (2014).
- [29] A. Weiß, G. Wellein, A. Alvermann, and H. Fehske, *Rev. Mod. Phys.* **78**, 275 (2006).
- [30] J. H. García, L. Covaci, and T. G. Rappoport, *Phys. Rev. Lett.* **114**, 116602 (2015).
- [31] D. Carvalho, N. A. García-Martínez, J. L. Lado, and J. Fernández-Rossier, *Phys. Rev. B* **97**, 115453 (2018).
- [32] C. S. Lent, M. A. Bowen, J. D. Dow, R. S. Allgaier, O. F. Sankey, and E. S. Ho, *Superlattices Microstruct.* **2**, 491 (1986).
- [33] X. Gao and M. S. Daw, *Phys. Rev. B* **77**, 033103 (2008).
- [34] P. Dziawa, B. J. Kowalski, K. Dybko, R. Buczko, A. Szczerbakow, M. Szot, E. Łusakowska, T. Balasubramanian, B. M. Wojek, M. H. Berntsen, O. Tjernberg, and T. Story, *Nat. Mater.* **11**, 1023 (2012).
- [35] J. O. Dimmock, I. Melngailis, and A. J. Strauss, *Phys. Rev. Lett.* **16**, 1193 (1966).
- [36] J. C. Y. Teo, L. Fu, and C. L. Kane, *Phys. Rev. B* **78**, 045426 (2008).
- [37] S.-Y. Xu, C. Liu, N. Alidoust, M. Neupane, D. Qian, I. Belopolski, J. D. Denlinger, Y. J. Wang, H. Lin, L. A. Wray, G. Landolt, B. Slomski, J. H. Dil, A. Marcinkova, E. Morosan, Q. Gibson, R. Sankar, F. C. Chou, R. J. Cava, A. Bansil, and M. Z. Hasan, *Nat. Commun.* **3**, 1192 (2012).
- [38] Y. Tanaka, Z. Ren, T. Sato, K. Nakayama, S. Souma, T. Takahashi, K. Segawa, and Y. Ando, *Nat. Phys.* **8**, 800 (2012).
- [39] Y. Tanaka, T. Sato, K. Nakayama, S. Souma, T. Takahashi, Z. Ren, M. Novak, K. Segawa, and Y. Ando, *Phys. Rev. B* **87**, 155105 (2013).
- [40] C. Yan, J. Liu, Y. Zang, J. Wang, Z. Wang, P. Wang, Z.-D. Zhang, L. Wang, X. Ma, S. Ji, K. He, L. Fu, W. Duan, Q.-K. Xue, and X. Chen, *Phys. Rev. Lett.* **112**, 186801 (2014).
- [41] R. Zhong, X. He, J. A. Schneeloch, C. Zhang, T. Liu, I. Pletikosić, T. Yilmaz, B. Sinkovic, Q. Li, W. Ku, T. Valla, J. M. Tranquada, and G. Gu, *Phys. Rev. B* **91**, 195321 (2015).
- [42] J. E. Avron and R. Seiler, *Phys. Rev. Lett.* **54**, 259 (1985).
- [43] Q. Niu, D. J. Thouless, and Y.-S. Wu, *Phys. Rev. B* **31**, 3372 (1985).
- [44] I. C. Fulga, F. Hassler, and A. R. Akhmerov, *Phys. Rev. B* **85**, 165409 (2012).
- [45] A. George, *SIAM J. Numer. Anal.* **10**, 345 (1973).
- [46] M. Aizenman and G. M. Graf, *J. Phys. A* **31**, 6783 (1998).
- [47] E. Prodan and W. Kohn, *Proc. Natl. Acad. Sci. USA* **102**, 11635 (2005).
- [48] J. Song and E. Prodan, *Phys. Rev. B* **89**, 224203 (2014).
- [49] I. Mondragon-Shem, T. L. Hughes, J. Song, and E. Prodan, *Phys. Rev. Lett.* **113**, 046802 (2014).
- [50] P. Perez-Piskunow, D. Varjas, M. Fruchart, M. Caio, and A. Akhmerov (unpublished).
- [51] T. H. Hsieh, H. Lin, J. Liu, W. Duan, A. Bansil, and L. Fu, *Nat. Commun.* **3**, 982 (2012).
- [52] We expect that this will not happen with a regular short-range correlated disorder.
- [53] L. Fu and C. L. Kane, *Phys. Rev. Lett.* **109**, 246605 (2012).
- [54] Y. Ando and L. Fu, *Annu. Rev. Condens. Matter Phys.* **6**, 361 (2015).
- [55] Y. Okada, M. Serbyn, H. Lin, D. Walkup, W. Zhou, C. Dhital, M. Neupane, S. Xu, Y. J. Wang, R. Sankar, F. Chou, A. Bansil, M. Z. Hasan, S. D. Wilson, L. Fu, and V. Madhavan, *Science* **341**, 1496 (2013).
- [56] C. Fang, M. J. Gilbert, and B. A. Bernevig, *Phys. Rev. Lett.* **112**, 046801 (2014).
- [57] D. L. Mitchell and R. F. Wallis, *Phys. Rev.* **151**, 581 (1966).
- [58] I. C. Fulga, N. Avraham, H. Beidenkopf, and A. Stern, *Phys. Rev. B* **94**, 125405 (2016).
- [59] C. W. Groth, M. Wimmer, A. R. Akhmerov, and X. Waintal, *New J. Phys.* **16**, 063065 (2014).
- [60] D. Varjas, M. Fruchart, A. R. Akhmerov, and P. Perez-Piskunow (2019), doi:10.5281/zenodo.2667604.
- [61] M. D. Caio, G. Möller, N. R. Cooper, and M. J. Bhaseen, *Nat. Phys.* **15**, 257 (2019).
- [62] T. Phuphachong, B. A. Assaf, V. V. Volobuev, G. Bauer, G. Springholz, L. A. De Vaulchier, and Y. Guldner, *J. Phys.: Conf. Ser.* **864**, 012038 (2017).
- [63] In the case of Ref. [40], the ARPES data for $x = 0.3$ are not conclusive, but they are interpreted with the help of VCA theory as a trivial phase. We claim that for $x = 0.3$ the experimental data and the theoretical results presented in Ref. [40] are not sufficient to ensure that $x = 0.3$ lies on the trivial phase.

- [64] S. Stützer, Y. Plotnik, Y. Lumer, P. Titum, N. H. Lindner, M. Segev, M. C. Rechtsman, and A. Szameit, *Nature (London)* **560**, 461 (2018).
- [65] N. P. Mitchell, L. M. Nash, D. Hexner, A. M. Turner, and W. T. M. Irvine, *Nat. Phys.* **14**, 380 (2018).
- [66] J. Cha, K. Woo Kim, and C. Daraio, *Nature (London)* **564**, 229 (2018).
- [67] D. Zhou, L. Zhang, and X. Mao, *Phys. Rev. Lett.* **120**, 068003 (2018).
- [68] R. Bianco and R. Resta, *Phys. Rev. Lett.* **110**, 087202 (2013).
- [69] W. A. Benalcazar, B. A. Bernevig, and T. L. Hughes, *Science* **357**, 61 (2017).
- [70] O. Pozo, C. Repellin, and A. G. Grushin, *Phys. Rev. Lett.* **123**, 247401 (2019).
- [71] D. Varjas, A. Lau, K. Pöyhönen, A. R. Akhmerov, D. I. Pikulin, and I. C. Fulga, *Phys. Rev. Lett.* **123**, 196401 (2019).
- [72] J. C. Slater and G. F. Koster, *Phys. Rev.* **94**, 1498 (1954).
- [73] F. Schindler, A. M. Cook, M. G. Vergniory, Z. Wang, S. S. P. Parkin, B. A. Bernevig, and T. Neupert, *Sci. Adv.* **4**, eaat0346 (2018).
- [74] P. Sessi, D. Di Sante, A. Szczerbakow, F. Glott, S. Wilfert, H. Schmidt, T. Bathon, P. Dziawa, M. Greiter, T. Neupert, G. Sangiovanni, T. Story, R. Thomale, and M. Bode, *Science* **354**, 1269 (2016).