

Mapping groundwater dynamics in the Netherlands

Three dimensional insight in groundwater systems based on the Dutch national database of suburface information (DINOloket.nl)

Zaadnoordijk, Willem Jan; Lourens, Aris

Publication date

2019

Document Version

Final published version

Citation (APA)

Zaadnoordijk, W. J., & Lourens, A. (2019). Mapping groundwater dynamics in the Netherlands: Three dimensional insight in groundwater systems based on the Dutch national database of suburface information (DINOloket.nl). IAH 2019: 46th Congress of the Interational Association of Hydrogeologists, Malaga, Spain.

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.



Mapping groundwater dynamics in the Netherlands:

three dimensional insight in groundwater systems based on the Dutch national database of subsurface information (DINOloket.nl)

Willem J. Zaadnoordijk^{1,2}, Aris Lourens¹ willem_jan.zaadnoordijk@tno.nl

¹TNO Geological Survey of the Netherlands – team geomodelling, Utrecht

²Delft University of Technology - Water Resources Section, Faculty of Civil Engineering and Geosciences



ONDERGRONDGEGEVENS

60 km

TOELICHTING



W

1**T**|

 2D


Groundwater time series models




http://www.grondwatertools.nl/grondwatertools-viewer











Klimaatsrepresentatieve grondwaterdynamiek

	cm + NAP	cm - mv
GHG	315.0	680.0
GLG	257.6	737.4
GVG	296.1	698.9
RHG	310.1	684.9
RLG	260.9	734.1
Grondwatertrap	VIII	


Toon in grafiek:

✓ GxG

Maaiveld



Decomposition of influences in time series model









Aspects groundwater dynamics

OF GEOLOGICAL MAPPING

- 1. Response time of precipitation response
- 2. Yearly fluctuation regime curve
- 3. Long term trends







Characterization impulse response




- Temporal moments (raw moments M_k)
 - M₀ total response
 - $\mu = M_1/M_0$ average response time

Physical meaning – M_k fulfills differential equation

similar to groundwater head

Analytic solutions for simple situations like circular island


Useful characteristics of impulse response





Lateral variation response time

First regional aquifer










Lateral variation response time














Vertical variation


Difference between first and second regional aquifer



Vertical variation

Difference between first and second regional aquifer

with hydraulic resistance of aquitard



Concluding remarks

OF GEOLOGICAL MAPPING

- Groundwater head time series show system characteristics in head response to precipitation
- Mapping response time in three dimensions provides insight
- Response function may differ even if no head difference
- Response integrates over larger area than bore log →
 additional information for delineating aquitards
- System changes in time so use only last 8 years





Thank you for your attention

Check out:

https://www.dinoloket.nl/ondergrondgegevens

https://www.grondwatertools.nl/grondwatertools-viewer

Contact: willem_jan.zaadnoordijk@tno.nl

