

## Evaluation of the CO<sub>2</sub>-storage potential in depleted gas fields of the West Netherlands Basin

### Case study P18 gas field

van Toorenenburg, Koen; Donselaar, Rick; Flores Colmenares, Jonatan; Groenenberg, RM; Pluymaekers, MP; Vandeweijer, VP; van de Weerd, AA

#### Publication date

2015

#### Document Version

Final published version

#### Citation (APA)

van Toorenenburg, K., Donselaar, R., Flores Colmenares, J., Groenenberg, RM., Pluymaekers, MP., Vandeweijer, VP., & van de Weerd, AA. (2015). *Evaluation of the CO<sub>2</sub>-storage potential in depleted gas fields of the West Netherlands Basin: Case study P18 gas field*. Poster session presented at 8th Dutch CCS symposium, Rotterdam, Netherlands.

#### Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

#### Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

#### Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

# Evaluation of the CO<sub>2</sub>-storage potential in depleted gas fields of the West Netherlands Basin: Case study P18 gas field


K.A. van Toorenburg<sup>1</sup>, M.E. Donselaar<sup>1</sup>, J.S. Flores Colmenares<sup>2</sup>, R.M. Groenenberg<sup>3</sup>, M.P. Pluymaekers<sup>4</sup>, V.P. Vandeweyer<sup>4</sup>, and A.A. van de Weerd<sup>5</sup>

## Introduction

Carbon Capture and Sequestration (CCS) is expected to become a serious CO<sub>2</sub>-emission reduction technology in the Netherlands. This study aims to evaluate the CO<sub>2</sub>-storage potential of depleted gas reservoirs in the West Netherlands Basin (WNB) based on geological boundary conditions.


## Basin-scale evaluation

The identification of suitable reservoirs for CO<sub>2</sub> storage is based on predefined geological boundary conditions, such as trap type, reservoir architecture and capacity, seal quality, and depth. Depleted gas fields inherently meet most criteria and their depth is derived from a basin-scale structural model (Fig. 1).




## P18

The nearly-depleted P18 gas field is located 20 km NW of the E.ON MPP3 power station (Fig. 2) and comprises fluvial and aeolian deposits of the Triassic Main Buntsandstein Subgroup (MBS) in several fault-bounded compartments (Fig. 2A and 3) whose juxtaposition locally inhibits connectivity.



**Fig. 3** Seismic inline through the P18 field (NE-SW; inset top right) showing interpreted faults and the facies model of the reservoir interval. Note that juxtaposition may isolate fault-bounded compartments.


The reservoir meets the criteria for carbon sequestration with a production-derived capacity of ~40 Mt CO<sub>2</sub>. Monte-Carlo simulations of the reservoir model capacity average ~33 Mt CO<sub>2</sub> (Fig. 4) and identify bulk rock volume (BRV) and geological heterogeneities as the main uncertainties.




**Fig 4** A) Static reservoir model of the effective porosity distribution in the P18 field. B) Results of Monte-Carlo simulations of the modelled reservoir capacity.

## Conclusions

- Depleted gas reservoirs may have CO<sub>2</sub>-storage potential
- The P18 offshore field is geographically and geologically suitable for carbon sequestration
- Several compartments add to a capacity of 33 - 40 Mt CO<sub>2</sub>
- The main uncertainties are in BRV and geological heterogeneities



**Fig. 2** A) Overlay of oil and gas fields on Google Earth image (north is up). B) E.ON MPP3 power plant.



Oil and gas map of the Netherlands (modified from TNO Geological Survey of the Netherlands 2015, 2<sup>nd</sup> edn.). Producing gas fields in green, depleted gas fields in grey with green edge.