

Using Landsat land surface temperature as a proxy for air temperature in urban settings Experiments in the Netherlands

Beuster, Lukas; García-Sánchez, Clara; Ledoux, Hugo

Publication date

2023

Document Version

Final published version

Citation (APA)

Beuster, L., García-Sánchez, C., & Ledoux, H. (2023). *Using Landsat land surface temperature as a proxy for air temperature in urban settings: Experiments in the Netherlands*. Abstract from 11th International Conference on Urban Climate, Sydney, Australia.

<https://virtual.oxfordabstracts.com/#/event/3742/submission/225>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Source: Beuster, Lukas, García-Sánchez, Clara, & Ledoux, Hugo. (n.d.). *Using Landsat land surface temperature as a proxy for air temperature in urban settings: Experiments in the Netherlands*. OA Virtual Conferences. Retrieved April 22, 2024, from <https://virtual.oxfordabstracts.com/#/event/3742/submission/225>

225

Using Landsat land surface temperature as a proxy for air temperature in urban settings: Experiments in the Netherlands

Mr. Lukas Beuster^{1,2}, Prof. Clara García-Sánchez¹, Prof. Hugo Ledoux¹

¹ Delft University of Technology, Delft, Netherlands. ² Amsterdam Institute for Advanced Metropolitan Solutions, Amsterdam, Netherlands

Abstract

Understanding the UHI effect in any city requires high-resolution temperature data. This data is often difficult to obtain as cities usually have only a few ground sensors, leaving large data gaps. To fill these gaps, we compare Landsat-derived land surface temperature (LST) with air temperature (T_{air}) measurements from urban weather stations in the two largest cities in the Netherlands. Previous studies of this kind have often been limited due to a few main factors: low spatial resolution, limited weather station data and small sample sizes (Chung et al., 2020, Mutiibwa, 2015; Sheng 2017; Xiong, 2017; Yang, 2020). As a result, findings have been inconsistent, albeit mostly promising. Addressing these issues and adding to Burnett and Chen's (2021) extensive comparison on a regional scale in Ontario, Canada, we present a reproducible, code-based approach focusing on cities. Using 149 Landsat scenes and data from 33 urban weather stations in the Netherlands (24 in Amsterdam, 9 in Rotterdam) between 2013-2022, 1700 comparison points across all European seasons are established.

We find that there is a strong positive and significant linear relationship between LST and T_{air} across the dataset ($r = .89$). OLS regression results indicate 80% of the T_{air} variation can be explained by the LST, with T_{air} increasing by 0.62°C for every 1°C increase in LST. Analyses were repeated to account for seasonality, each station's local climate zone (Stewart and Oke, 2012) as well as mean absolute error and root mean square error to interrogate the discrepancy, all of which will be highlighted in the presentation. Overall, our evidence suggests that LST can indeed be a suitable proxy for T_{air} and could consequently form an additional decision-making layer to assist climate monitoring and urban planning in the Netherlands as well as similar climates.

