



Delft University of Technology

Optimally reconfiguring list and correspondence colourings

Cambie, Stijn; Cames van Batenburg, Wouter; Cranston, Daniel W.

DOI

[10.1016/j.ejc.2023.103798](https://doi.org/10.1016/j.ejc.2023.103798)

Publication date

2024

Document Version

Final published version

Published in

European Journal of Combinatorics

Citation (APA)

Cambie, S., Cames van Batenburg, W., & Cranston, D. W. (2024). Optimally reconfiguring list and correspondence colourings. *European Journal of Combinatorics*, 115, Article 103798. <https://doi.org/10.1016/j.ejc.2023.103798>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.



Contents lists available at ScienceDirect

European Journal of Combinatorics

journal homepage: www.elsevier.com/locate/ejc

Optimally reconfiguring list and correspondence colourings[☆]



Stijn Cambie^{a,b,c}, Wouter Cames van Batenburg^d,
Daniel W. Cranston^e

^a Department of Mathematics, Radboud University Nijmegen, The Netherlands

^b Mathematics Institute, University of Warwick, UK

^c Department of Computer Science, KU Leuven Campus Kulak-Kortrijk, 8500 Kortrijk, Belgium¹

^d Delft University of Technology, Delft Institute of Applied Mathematics, The Netherlands

^e Virginia Commonwealth University, Department of Computer Science, United States of America

ARTICLE INFO

Article history:

Received 6 March 2023

Accepted 13 August 2023

Available online xxxx

ABSTRACT

The reconfiguration graph $C_k(G)$ for the k -colourings of a graph G has a vertex for each proper k -colouring of G , and two vertices of $C_k(G)$ are adjacent precisely when those k -colourings differ on a single vertex of G . Much work has focused on bounding the maximum value of $\text{diam}C_k(G)$ over all n -vertex graphs G . We consider the analogous problems for list colourings and for correspondence colourings. We conjecture that if L is a list-assignment for a graph G with $|L(v)| \geq d(v) + 2$ for all $v \in V(G)$, then $\text{diam}C_L(G) \leq n(G) + \mu(G)$. We also conjecture that if (L, H) is a correspondence cover for a graph G with $|L(v)| \geq d(v) + 2$ for all $v \in V(G)$, then $\text{diam}C_{(L,H)}(G) \leq n(G) + \tau(G)$. (Here $\mu(G)$ and $\tau(G)$ denote the matching number and vertex cover number of G .) For every graph G , we give constructions showing that both conjectures are best possible, which also hints towards an exact form of Cereceda's Conjecture for regular graphs. Our first main result proves the upper bounds (for the list and correspondence versions, respectively) $\text{diam}C_L(G) \leq n(G) + 2\mu(G)$ and $\text{diam}C_{(L,H)}(G) \leq n(G) + 2\tau(G)$. Our second main result proves that both conjectured bounds hold, whenever all v satisfy $|L(v)| \geq 2d(v) + 1$. We conclude by proving one or both conjectures for various classes of graphs such as complete bipartite graphs,

[☆] Supported by a Vidi grant (639.032.614) of the Netherlands Organisation for Scientific Research (NWO) and the UK Research and Innovation Future Leaders Fellowship MR/S016325/1.

E-mail addresses: stijn.cambie@hotmail.com (S. Cambie), w.p.s.camesvanbatenburg@tudelft.nl

(W. Cames van Batenburg), dcranston@vcu.edu (D.W. Cranston).

¹ Current affiliation.

subcubic graphs, cactuses, and graphs with bounded maximum average degree. The full paper can also be found at arxiv.org/abs/2204.07928.

© 2023 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

In this paper we study questions of transforming one proper colouring of a graph G into another, by a sequence of *recolouring steps*. Each step recolours a single vertex, and we require that each resulting intermediate colouring is also proper. Our work fits into the broader context of *reconfiguration*, in which some object (in our case a proper colouring) is transformed into another object of the same type, via a sequence of small changes, and we require that after each change we again have an object of the prescribed type (for us a proper colouring). It is natural to pose reconfiguration questions for a wide range of objects: proper colourings, independent sets, dominating sets, maximum matchings, spanning trees, and solutions to 3-SAT, to name a few. For each type of object, we must define allowable changes between successive objects in the sequence (in our case, recolouring a single vertex). Typically, we ask four types of questions. (1) Given objects α and β , is there a reconfiguration sequence from α to β ? (2) If the first question is answered yes, what is the length of a shortest such sequence? (3) Is the first question answered yes for every pair of objects α and β ? (4) If yes, what is the maximum value of $\text{dist}(\alpha, \beta)$ over all α and β ? For an introduction to reconfiguration, we recommend surveys by van den Heuvel [1] and Nishimura [2]. Most of our definitions and notation are standard, but for completeness we include many of them at the start of Section 2.

For a graph G and a positive integer k , the k -colouring reconfiguration graph of G , denoted $C_k(G)$, has as its vertices all proper k -colourings of G and two vertices of $C_k(G)$ are adjacent if their corresponding colourings differ on exactly one vertex of G . Our goal here is to study the diameter of this reconfiguration graph, denoted $\text{diam } C_k(G)$. In general, $C_k(G)$ may be disconnected, in which case its diameter is infinite. For example $C_2(C_{2s})$ consists of two isolated vertices (here C_{2s} is a cycle of length $2s$). More strongly, for each integer $k \geq 2$ there exist k -regular graphs G such that $C_{k+1}(G)$ has isolated vertices. The simplest example is the clique K_{k+1} , but this is also true for every k -regular graph with a $(k+1)$ -colouring α such that all $k+1$ colours appear on each closed neighbourhood; such colourings are called *frozen*.

To avoid a colouring α being frozen, some vertex v must have some colour unused by α on its closed neighbourhood, $N[v]$. And to avoid $C_k(G)$ being disconnected, every induced subgraph H must contain such a vertex v with some colour unused by α on $N[v] \cap V(H)$. Thus, it is natural to consider the degeneracy of G , denoted $\text{degen}(G)$.

Our examples of frozen colourings above show that, if we aim to have $C_k(G)$ connected, then in general it is not enough to require $k \geq \text{degen}(G) + 1$. However, an easy inductive argument shows that a slightly stronger condition is sufficient: If $k \geq \text{degen}(G) + 2$, then $C_k(G)$ is connected. (Earlier, Jerrum [3] proved that $k \geq \Delta(G) + 2$ suffices, but the arguments are similar.) This inductive proof only yields that $\text{diam } C_k(G) \leq 2^{|V(G)|}$. But Cereceda [4, Conjecture 5.21] conjectured something much stronger.

Cereceda's Conjecture. *For an n -vertex graph G with $k \geq \text{degen}(G) + 2$, the diameter of $C_k(G)$ is $O(n^2)$.*

Bousquet and Heinrich [5] proved that $\text{diam } C_k(G) = O_d(n^{\text{degen}(G)+1})$, which is the current best known bound. When $k \geq \Delta(G) + 2$, Cereceda [4, Proposition 5.23] proved that $\text{diam } C_k(G) = O(n\Delta) = O(n^2)$. In particular, Cereceda's conjecture is true for regular graphs. But, as we show here, if $k \geq \Delta(G) + 2$, then in fact we have the stronger bound $\text{diam } C_k(G) \leq 2n$, and we conjecture $\text{diam } C_k(G) \leq \lfloor 3n/2 \rfloor$.

A folklorish result observes that $\text{diam } C_k(G) = O(n)$ when k is large relative to $\Delta(G)$ (for example, see [6, Section 3.1]). But until now, it seems that no one has investigated exact values of the diameter (say, when $k \geq \Delta(G)$). This is the main goal of our paper. Before stating our two main conjectures, and our results supporting them, we present an easy lower bound on $C_k(G)$ in terms of the matching number $\mu(G)$.

Proposition 1. *For a graph G , if $k \geq 2\Delta(G)$, then $\text{diam } C_k(G) \geq n(G) + \mu(G)$.*

Proof. Let M be a maximum matching in G . Form \widehat{G} from G as follows. If $vw, xy \in M$ and $wx \in E(G)$, then add vy to \widehat{G} . Note that $\Delta(\widehat{G}) \leq 2\Delta(G) - 1$. So $\chi(\widehat{G}) \leq 2\Delta(G)$. Let α be a $2\Delta(G)$ -colouring of \widehat{G} . Form β from α by swapping colours on endpoints of each edge in M and, for each $v \in V(G)$ not saturated by M , picking $\beta(v)$ outside of $\{\alpha(v)\} \cup \bigcup_{w \in N(v)} \beta(w)$. To recolour G from α to β , every vertex must be recoloured. Further, for each edge $e \in M$, the first endpoint v of e to be recoloured must initially receive a colour other than $\beta(v)$, so must be recoloured at least twice. Thus, $\text{diam } C_k(G) \geq n(G) + \mu(G)$, as desired. \square

At the end of Section 2, we mention various special cases in which the hypothesis of Proposition 1 can be weakened. We pose it as an open question whether the bound in Proposition 1 holds for all $k \geq \Delta(G) + 2$.

We study this reconfiguration problem in the more general contexts of list colouring and correspondence colouring, both of which we define in Section 2. These more general contexts offer the added advantage of enabling us to naturally prescribe fewer allowed colours for vertices of lower degree. Analogous to $C_k(G)$, for a graph G and a list-assignment L or correspondence cover (L, H) for G , we define the L -reconfiguration graph $C_L(G)$ or (L, H) -reconfiguration graph $C_{(L,H)}(G)$ of G . Generalising our constructions of frozen k -colourings above, it is easy to show that $C_L(G)$ and $C_{(L,H)}(G)$ can contain frozen L -colourings (and thus be disconnected) if we require only that all $v \in V(G)$ satisfy $|L(v)| = d(v) + 1$. Thus, we adopt a slightly stronger hypothesis: all $v \in V(G)$ satisfy $|L(v)| \geq d(v) + 2$. Now we can state our two main conjectures.

Conjecture 1 (List Colouring Reconfiguration Conjecture). *For a graph G , if L is a list-assignment such that $|L(v)| \geq d(v) + 2$ for every $v \in V(G)$, then $\text{diam } C_L(G) \leq n(G) + \mu(G)$.*

Conjecture 2 (Correspondence Colouring Reconfiguration Conjecture). *For a graph G , if (L, H) is a correspondence cover such that $|L(v)| \geq d(v) + 2$ for every $v \in V(G)$, then $\text{diam } C_{(L,H)}(G) \leq n(G) + \tau(G)$.*

Here $\tau(G)$ denotes the vertex cover number of G . For brevity, we often call these conjectures the List Conjecture and the Correspondence Conjecture. Our aim in this paper is to provide significant evidence for both conjectures. Due to Proposition 1, the List Conjecture is best possible, when the lists are large enough. We will soon give easy constructions showing that both conjectures are best possible whenever all v satisfy $|L(v)| \geq d(v) + 2$. But we defer these constructions until Section 2, where we formally define list and correspondence colourings.

Proposition 2. *For every graph G (i) there exists a list-assignment L such that $|L(v)| = d(v) + 2$ for all v for which $\text{diam } C_L(G) = n(G) + \mu(G)$ and (ii) there exists a correspondence cover (L, H) such that $|L(v)| = d(v) + 2$ for all v for which $\text{diam } C_{(L,H)}(G) = n(G) + \tau(G)$.*

For both the List Conjecture and the Correspondence Conjecture, it is trivial to construct examples that need at least $n(G)$ recolourings. We simply require that $\alpha(v) \neq \beta(v)$ for all $n(G)$ vertices v . So we view these conjectured upper bounds as consisting of a “trivial” portion, $n(G)$ recolourings, and a “non-trivial” portion, $\mu(G)$ or $\tau(G)$ recolourings. We give two partial results towards each conjecture. Our first result proves both conjectures up to a factor of 2 on the non-trivial portions of these upper bounds.

Theorem 1. *(i) For every graph G and list-assignment L with $|L(v)| \geq d(v) + 2$ for all $v \in V(G)$ we have $\text{diam } C_L(G) \leq n(G) + 2\mu(G)$. (ii) For every graph G and correspondence cover (L, H) with $|L(v)| \geq d(v) + 2$ for all $v \in V(G)$ we have $\text{diam } C_{(L,H)}(G) \leq n(G) + 2\tau(G)$.*

Bousquet, Feuilloley, Heinrich, and Rabie [7, following Question 1.3] asked about the diameter of $C_k(G)$ when $k = \Delta(G) + 2$. In this case, a result of Bonamy and Bousquet [8, Theorem 1] implies, for an n -vertex graph G and $k = \Delta(G) + 2$, that $\text{diam } C_k(G) = O(\Delta(G)n)$. The authors of [7] asked whether it is possible to remove this dependency on $\Delta(G)$. We answer their question affirmatively. Always $\mu(G) \leq n(G)/2$, so Theorem 1(i) implies that $\text{diam } C_k(G) \leq 2n(G)$ when $k = \Delta(G) + 2$.

To complement Theorem 1, we prove both conjectured upper bounds when $|L(v)|$ is sufficiently large.

Theorem 2. (i) For every graph G and list-assignment L with $|L(v)| \geq 2d(v) + 1$ for all $v \in V(G)$ we have $\text{diam } C_L(G) \leq n(G) + \mu(G)$. (ii) For every graph G and correspondence cover (L, H) with $|L(v)| \geq 2d(v) + 1$ for all $v \in V(G)$ we have $\text{diam } C_{(L,H)}(G) \leq n(G) + \tau(G)$.

In Section 2 we present definitions and notation, as well as some easy lower bounds on diameters of reconfiguration graphs. In Section 3 we prove the list colouring portions of Theorems 1 and 2; and in Section 4 we prove the correspondence colouring portions. In Section 5 we prove more precise results when G is a tree (in this case the correspondence problem is identical to the list problem).

In Sections 6 and 7, we conclude by proving one or both conjectures exactly for various graph classes, such as complete bipartite graphs, subcubic graphs, cactuses, and graphs with low maximum average degree. These graphs are all sparse or have low chromatic number. On the other end of the sparsity spectrum, the List Conjecture is also true for all complete graphs, due to an argument of Bonamy and Bousquet [8, Lemma 5].

2. Definitions, notation, and easy lower bounds

For a graph $G = (V, E)$, we denote its order by $n(G)$. A *matching* is a set of vertex disjoint edges in G . A matching M of G is *perfect* if it saturates every vertex of G , and *near-perfect* if it saturates every vertex of G but one. A *vertex cover* S is a vertex subset such that every edge of G has at least one endpoint in S . The *matching number* $\mu(G)$ of G is the size of a largest matching. The *vertex cover number* $\tau(G)$ of G is the size of a smallest vertex cover. (Recall that if G is bipartite, then $\mu(G) = \tau(G)$.)

The *distance* $\text{dist}(v, w)$ between two vertices $v, w \in V(G)$ is the length of a shortest path in G between v and w . The *eccentricity* of a vertex v , denoted $\text{ecc}(v)$, is $\max_{w \in V} \text{dist}(v, w)$. The *diameter* $\text{diam}(G)$ of G is $\max_{v, w \in V} \text{dist}(v, w)$, which is equal to $\max_{v \in V} \text{ecc}(v)$, while the *radius* $\text{rad}(G)$ of G is $\min_{v \in V} \text{ecc}(v)$. A diameter can also refer to a shortest path between v and w for which $\text{dist}(v, w) = \text{diam}(G)$. A graph G is *k-degenerate* if every non-empty subgraph H contains a vertex v such that $d_H(v) \leq k$. The *degeneracy* of G is the minimum k such that G is k -degenerate.

A directed graph or *digraph* $D = (V, A)$ is analogous to a graph, except that each edge is directed. The directed edges in A are ordered pairs of vertices, and are called *arcs*.

Definition 3. For a digraph D , the vertex cover number $\tau(D)$ and matching number $\mu(D)$ are defined to be $\tau(H_D)$ and $\mu(H_D)$, where H_D is the undirected graph with vertex set $V(D)$ whose edges correspond to the bidirected edges in D .

Colourings are mappings from V to \mathbb{N} , and we denote them by greek letters such as α, β , and γ . A colouring α of G is *proper* if for every edge $vw \in E(G)$ we have $\alpha(v) \neq \alpha(w)$. Let $[k] := \{1, 2, \dots, k\}$. A *k-colouring* is a colouring using at most k colours, which are generally from the set $[k]$. The *chromatic number* $\chi(G)$ of G is the smallest k such that G has a proper k -colouring.

A *list-assignment* L , for a graph G , assigns to each $v \in V(G)$ a set $L(v)$ of natural numbers (“list” of allowable colours). A *proper L-colouring* is a proper colouring $\alpha: V(G) \rightarrow \mathbb{N}$ such that $\alpha(v) \in L(v)$ for every $v \in V(G)$. The *list chromatic number* (or *choice number*) $\chi_\ell(G)$ is the least k such that G admits a proper L -colouring whenever every $v \in V(G)$ satisfies $|L(v)| \geq k$.

A *correspondence cover* (L, H) for a graph G assigns to each vertex $v \in V(G)$ a set $L(v)$ of colours $\{(v, 1), \dots, (v, f(v))\}$ and to each edge $vw \in E(G)$ a matching between $L(v)$ and $L(w)$. (In this paper, we typically consider either $f(v) = d(v) + 2$ or $f(v) = 2d(v) + 1$ for all $v \in V(G)$.) Given a correspondence cover (L, H) , an (L, H) -colouring of G is a function α such that $\alpha \in L(v)$ for all

v and whenever $vw \in E(G)$ the edge $\alpha(v)\alpha(w)$ is not an edge of the matching assigned to vw . Here H denotes the union of the matchings assigned to all edges of G . (It is easy to check that correspondence colouring generalises list colouring.)

The *reconfiguration graph* $C_k(G)$ has as its vertices the proper k -colourings of G , and two k -colourings α and β are adjacent in $C_k(G)$ if they differ on exactly one vertex of G . In particular, if $k < \chi(G)$, then $C_k(G)$ has no vertices. (The reconfiguration graph was first defined in [9], where it was called the k -colour graph.) The distance between k -colourings α and β is at most j if we can form β , starting from α , by recolouring at most j vertices, one at a time, so that after each recolouring the current k -colouring of G is proper. Similarly, for a graph G and list-assignment L , the *reconfiguration graph* $C_L(G)$, or reconfiguration graph of the L -colourings of G , is the graph whose vertices are the proper L -colourings of G . Again, two L -colourings α and β are adjacent in $C_L(G)$ if they differ on exactly one vertex. This extension was first defined in [10]. For a correspondence cover (L, H) , the *reconfiguration graph* $C_{(L,H)}(G)$ is defined analogously; its vertices are the correspondence colourings of G and two vertices of $C_{(L,H)}(G)$ are adjacent precisely when their colourings differ on exactly one vertex of G . We will also use the associated colour-shift digraph for a graph G and two proper colourings, which is defined as follows.

Definition 4. Let $G = (V, E)$ be a graph and α and β be two proper colourings of G . We define an associated colour-shift digraph $D_{\alpha,\beta} = (V, A)$ where $\vec{vw} \in A$ if and only if $vw \in E$ and $\beta(v) = \alpha(w)$.

2.1. Improved lower bounds

In this subsection we prove lower bounds which show that our upper bounds in the rest of the paper are sharp or nearly sharp. These will not be explicitly needed elsewhere, so the impatient reader should feel free to skip to Section 3, where we prove [Theorems 1\(i\)](#) and [2\(i\)](#).

Observation 5. If G is a graph with list colourings α and β , then $\text{dist}(\alpha, \beta) \geq \mu(D_{\alpha,\beta}) + \sum_{v \in V} \mathbf{1}_{\alpha(v) \neq \beta(v)}$.

Proof. Whenever $\alpha(v) \neq \beta(v)$, vertex v must be recoloured. Further, if v and w are neighbours for which $\alpha(v) = \beta(w)$ and $\alpha(w) = \beta(v)$, then we cannot recolour both v and w only once, since after every recolouring step the resulting colouring must be proper. So at least $\sum_{v \in V} \mathbf{1}_{\alpha(v) \neq \beta(v)}$ vertices must be recoloured, and at least $\mu(D_{\alpha,\beta})$ vertices must be recoloured at least twice. \square

Does the lower bound in [Observation 5](#) always hold with equality? Our next example shows that it does not; see [Fig. 1](#). Specifically, this example exhibits, for the 4-cycle, colourings α and β such that $\mu(D_{\alpha,\beta}) < \mu(G)$ but still $\text{dist}(\alpha, \beta) = n(G) + \mu(G)$.

Example 6. Let $G = C_4$ and denote $V(G)$ by [4]. If $\alpha(i) = i$ and $\beta(i) \equiv i + 1 \pmod{4}$ and the colour set is precisely [4], then transforming α to β uses at least 6 recolourings, i.e. $\text{dist}(\alpha, \beta) = 6$. Part of this reconfiguration graph is shown in [Fig. 1](#).

Proof. Starting from α , each vertex i has a single colour with which it can be recoloured. But each possible recolouring creates a colouring α' for which [Observation 5](#) gives $\text{dist}(\alpha', \beta) \geq 5$. \square

Next we present the (previously promised) proof of [Example 6](#). For easy reference, we restate it.

Proposition 2. For every graph G (i) there exists a list-assignment L such that $|L(v)| = d(v) + 2$ for all v for which $\text{diam } C_L(G) = n(G) + \mu(G)$ and (ii) there exists a correspondence cover (L, H) such that $|L(v)| = d(v) + 2$ for all v for which $\text{diam } C_{(L,H)}(G) = n(G) + \tau(G)$.

Proof. For a graph G , fix a maximum matching M . Assign to each vertex v a list of $d(v) + 2$ colours such that $|L(v) \cap L(w)| = 2$ if $vw \in M$ and otherwise $|L(v) \cap L(w)| = 0$. Pick α and β such that for each $vw \in M$, we have $\alpha(v) = \beta(w)$ and $\alpha(w) = \beta(v)$, and for each $v \in V(G)$ we have $\alpha(v) \neq \beta(v)$.




Fig. 1. Part of the reconfiguration graph $C_4(C_4)$.

The lower bound holds by [Observation 5](#). The upper bound is trivial, since $L(x) \cap L(y) = \emptyset$ for all $xy \in E(G) \setminus M$. This proves (i).

Let (L, H) be a correspondence cover of G such that $L(v) = \{(v, 1) \dots (v, d(v) + 2)\}$ for every $v \in V(G)$. For every $vw \in E(G)$, let the matching (from H) consist of the two edges $(v, 1)(w, 2)$ and $(v, 2)(w, 1)$; otherwise, (v, i) and (w, j) are unmatched. Let $\alpha(v) := (v, 1)$ and $\beta(v) := (v, 2)$ for all $v \in V(G)$. Starting from α , we can recolour a vertex v with $\beta(v)$ only if all neighbours of v have already been recoloured. Thus, the set of vertices recoloured only once must be an independent set; equivalently, the set of vertices recoloured at least twice must be a vertex cover. So we must use at least $n(G) + \tau(G)$ recolourings, which proves the lower bound. For the upper bound, fix a minimum vertex cover S . First recolour each $v \in S$ with $(v, 3)$; afterward, recolour each vertex with $(v, 2)$, first the vertices in $V(G) \setminus S$ and then those in S . So $n(G) + \tau(G)$ recolourings suffice. This proves (ii). \square

It is helpful to note that the proof of [Example 6](#) actually works (for both list colouring and correspondence colouring) whenever all v satisfy $|L(v)| \geq 3$.

Clearly, the graph \widehat{G} constructed in [Proposition 1](#) depends on our choice of a perfect matching M . It is interesting to note that some choices of M work quite well, while others work rather poorly.

Example 7. [Fig. 2](#) shows a graph G built from two cliques K_p and a complete bipartite graph $K_{p,p}$, with parts U and W , by adding a perfect matching M between one copy of K_p and the vertices of U and a perfect matching between W and the other copy of K_p . Now $\chi(\widehat{G}) = 2p$ if we choose our perfect matching to be M . If p is even, then we can instead choose a perfect matching M' that is the disjoint union of perfect matchings in the two copies of K_p and in the $K_{p,p}$ so that the resulting \widehat{G} instead satisfies $\chi(\widehat{G}) = p + 2$.

Although [Example 7](#) implies that our choice of M can effect $\chi(\widehat{G})$ dramatically, for many graphs, any choice of M is fine. To conclude this section, we present various cases in which (for all choices of M) we can weaken the hypothesis of [Proposition 1](#).

Proposition 3. For a graph G and positive integer k we have

$$\text{diam } C_k(G) \geq n(G) + \mu(G)$$

whenever (a) $\chi(\widehat{G}) \leq k - 1$ or (b) $\chi(\widehat{G}) \leq k$ and $k \geq \Delta(G) + 2$, where \widehat{G} is as in [Proposition 1](#). In particular, this is true in the following cases.

- (1) $1 + \chi(G)(\chi(G) - 1) \leq k$ (in particular, if G is bipartite and $k \geq 3$).
- (2) $\Delta(G) = 3$ and $k \geq \Delta(G) + 2 = 5$.




Fig. 2. A graph G (shown on both left and right) consisting of a complete bipartite graph $K_{p,p}$ in the centre, two copies of K_p on the sides, and a matching from each copy of K_p to one part of $K_{p,p}$. The left and right figures specify different perfect matchings M (in bold). On the left, $\chi(\widehat{G}) \geq \omega(\widehat{G}) = |V(K_{p,p})| = 2p = 2\Delta(G) - 2$. On the right, $\omega(\widehat{G}) = \omega(G) = p$, and in fact $\chi(\widehat{G}) \leq p + 2 = \Delta(G) + 1$. (We use colours $1, \dots, p$ on each clique and colours $p + 1$ and $p + 2$ in the centre.)

- (3) $\chi(G) \leq 3$ and $k \geq \Delta(G) + 2$ (in particular, if G is outerplanar and $k \geq \Delta(G) + 2$).
- (4) $k \geq \Delta(G) + 3$ and G is triangle-free, if Reed's Conjecture² holds for every graph \widehat{G} with $\omega(\widehat{G}) \leq 5$.
- (5) $k \geq \Delta(G) + 2$ and $\Delta(G) > f(\omega(G))$ for some function f .

Proof. Recall the definition of \widehat{G} from Proposition 1. Let M be a maximum matching in G . Form \widehat{G} from G as follows. If $vw, xy \in M$ and $wx \in E(G)$, then add vy to \widehat{G} . To prove each case of the proposition, we construct the graph \widehat{G} , let α be a $\chi(\widehat{G})$ -colouring of \widehat{G} (which is also a $\chi(\widehat{G})$ -colouring of G), and form a k -colouring β from α by swapping the colours on the endpoints of each edge e in the specified maximum matching M ; for each v not saturated by M , we choose $\beta(v)$ to avoid $\alpha(v) \cup \bigcup_{x \in N(v)} \beta(x)$. The latter is possible because we assumed that k is bounded from below by $\chi(\widehat{G}) + 1$ or by $\Delta(G) + 2$. By Observation 5, we have $\text{dist}(\alpha, \beta) \geq \mu(D_{\alpha, \beta}) + \sum_{v \in V} \mathbf{1}_{\alpha(v) \neq \beta(v)} = \mu(G) + n(G)$. Note that $\Delta(\widehat{G}) \leq 2\Delta(G) - 1$. So $\chi(\widehat{G}) \leq 1 + \Delta(\widehat{G}) \leq 2\Delta(G)$. Now we show that in each case listed above we have $\chi(\widehat{G}) \leq k$ or (in the first case) $\chi(\widehat{G}) \leq k - 1$.

- (1) Fix a $\chi(G)$ -colouring α of G . We assume that M is a perfect matching; if not, then we add a pendent edge at each vertex that is unsaturated by M and add all these new edges to M . To colour \widehat{G} , we give each vertex v the colour (i, j) , where $\alpha(v) = i$, $\alpha(w) = j$, and $vw \in M$. It is easy to check that the resulting colouring of \widehat{G} is proper.
- (2) Suppose that $\Delta(G) = 3$. As noted above, we have $\Delta(\widehat{G}) \leq 2\Delta(G) - 1 = 5$. If $\Delta(\widehat{G}) \leq 4$, then $\chi(\widehat{G}) \leq 5$ as desired. So assume instead that $\Delta(\widehat{G}) = 5$. By Brooks' Theorem it suffices to show that no component of \widehat{G} is K_6 . Suppose, to the contrary, that there exists $S \subseteq V(G)$ such that $\widehat{G}[S] = K_6$. It is easy to check that $G[S]$ must be 3-regular, and every vertex of S must be saturated by M . Observe that if two edges of M , say v_1v_2, v_3v_4 lie on a 4-cycle in G , then $d_{\widehat{G}}(v_i) \leq 4$, for each $i \in \{1, 2, 3, 4\}$. Now we need only to check that such a configuration occurs for every 3-regular graph G on 6 vertices and every perfect matching M . This is straightforward to verify: $G - M$ is either a 6-cycle or two 3-cycles; in the first case, M can be added in two (non-isomorphic) ways and in the second, M can be added in only one way.
- (3) If $\Delta(G) \geq 4$, then we are done by condition (1). If $\Delta(G) = 3$, then we are done by condition (2). If $\Delta(G) = 2$, then we are done by Proposition 1. And if $\Delta(G) \leq 1$, then we are done trivially.
- (4) We show that if G is triangle-free, then $\omega(\widehat{G}) \leq 5$. Suppose, to the contrary, that $\omega(\widehat{G}) \geq 6$. Consider $S \subseteq V(G)$ such that $\widehat{G}[S] = K_6$, and colour each edge of $\widehat{G}[S]$ with 1 if it appears in

² Recall that Reed conjectured that $\chi(G) \leq \lceil (\Delta(G) + 1 + \omega(G))/2 \rceil$ for every graph G .

G and with 2 otherwise. Since G is triangle-free, $\widehat{G}[S]$ has no triangle coloured 1. Let $R(t, t)$ denote the diagonal Ramsey number for cliques of order t . Since $R(3, 3) = 6$, we must have a triangle in $\widehat{G}[S]$ that is coloured 2; denote its vertices by v, w, x . Let vv', ww', xx' denote the edges of M incident with v, w, x . Since $vw, vx, wx \in E(\widehat{G}) \setminus E(G)$, we conclude that $v'w', v'x', w'x' \in E(G)$. This contradicts that G is triangle-free. Hence, $\omega(\widehat{G}) \leq 5$, as claimed. Finally, by Reed's Conjecture, $\chi(\widehat{G}) \leq \lceil (\Delta(\widehat{G}) + \omega(\widehat{G}) + 1)/2 \rceil \leq \lceil (2\Delta(G) - 1 + 5 + 1)/2 \rceil \leq \Delta(G) + 3 \leq k$.

- (5) Let t be a positive integer and consider any graph G with $\omega(G) = t$. Note that \widehat{G} can be decomposed into two (edge-disjoint) copies of G ; here \widehat{G} is formed from the two copies of G by identifying, for each edge $vw \in M$, the instance of v in one copy of G with the instance of w in the other copy. This implies that $\omega(\widehat{G}) < R(t + 1, t + 1)$. We also know that $\Delta(\widehat{G}) \leq 2\Delta(G)$. Now by e.g. [11, Theorem 2] we have

$$\chi(\widehat{G}) \leq 200R(t + 1, t + 1) \frac{2\Delta(G) \ln \ln(2\Delta(G))}{\ln(2\Delta(G))}.$$

This is smaller than $\Delta(G) + 2$ whenever $\Delta(G)$ is sufficiently large. \square

We also consider the graph \widetilde{G} formed from G by contracting each edge of a maximum matching M .

Proposition 4. *For a graph G and positive integer k we have*

$$\text{diam } C_k(G) \geq n(G) + \mu(G)$$

whenever $k \geq 2\chi(\widetilde{G})$ for some maximum matching of G . In particular, this is

- (1) true a.a.s. for $G_{n,p}$, for fixed $p \in (0, 1)$, and $k \geq \Delta(G_{n,p}) + 2$.
- (2) true if G is K_{t+1} -minor free and $k \geq 2t$, provided Hadwiger's Conjecture³ holds for K_{t+1} -minor free graphs. The latter is true for $t \leq 5$.

Proof. Let $\tilde{\alpha}$ be a $\chi(\widetilde{G})$ -colouring of \widetilde{G} . We construct a $2\chi(\widetilde{G})$ -colouring α of G such that if $v \in V(\widetilde{G})$ and v arises from contracting edge $wx \in M$ and $\tilde{\alpha}(v) = i$, then $\{\alpha(w), \alpha(x)\} = \{2i - 1, 2i\}$; if v does not arise from contracting an edge, then we choose $\alpha(v) = 2\tilde{\alpha}(v) = 2i$. Form β from α by swapping the colours on the endpoints of each edge of M and letting $\beta(v) := \alpha(v) - 1 = 2i - 1$ for each $v \in V(G) \setminus M$.

Hence $\text{diam } C_k(G) \geq \text{dist}(\alpha, \beta) \geq n(G) + \mu(G)$ by [Observation 5](#). We now prove that in the two mentioned cases, we (a.a.s.) have $k \geq 2\chi(\widetilde{G})$, regardless of the maximum matching of G we choose to form \widetilde{G} .

- (1) Let $G := G_{n,p}$. By [12], the Hadwiger number h (or contraction clique number) of G for fixed $p \in (0, 1)$ is $\Theta\left(\frac{n}{\sqrt{\ln n}}\right)$ a.a.s., so G is K_{h+1} -minor free. Note that \widetilde{G} is also K_{h+1} -minor free. By recent progress towards Hadwiger's conjecture, culminating in [13], this implies that (still a.a.s.) $\chi(\widetilde{G}) = O(h \ln \ln h) = O\left(\frac{n \ln \ln n}{\sqrt{\ln n}}\right)$. So $\chi(\widetilde{G})$ is a.a.s. smaller than $\Delta(G_{n,p}) + 2$, which is $np(1 + o(1)) = \Theta(n)$.
- (2) Note that \widetilde{G} is also K_{t+1} -minor free. So, by assumption, $\chi(\widetilde{G}) \leq t$. \square

Question 1. *Is it true, for every graph G and every $k \geq \Delta(G) + 2$, that $\text{diam } C_k(G) \geq n(G) + \mu(G)$?*

If the answer to this question is yes, then the [List Conjecture](#) (if true) would imply that $\text{diam } C_k(G) = n(G) + \mu(G)$ for every graph G and every $k \geq \Delta(G) + 2$. In particular, this would constitute a precise version for Cereceda's Conjecture in the case of regular graphs.

³ Recall that Hadwiger conjectured that $\chi(G) \leq t$ for every graph G with no K_{t+1} -minor.

3. Proving the main results for lists

In this section, we prove the list colouring portions of our main results: [Theorems 1\(i\)](#) and [2\(i\)](#). Recall that a graph G is *factor-critical* if $|V(G)|$ is odd and $G - v$ has a perfect matching for each $v \in V(G)$. Gallai showed that if every vertex is unsaturated by some maximum matching, i.e., $\mu(G - v) = \mu(G)$ for every vertex v , then G is factor-critical. So, in each proof we begin with a lemma to handle factor-critical graphs, and thereafter proceed to the general case. The following lemma serves as the base case in an inductive proof of [Theorem 1\(i\)](#). (We will only need it when G is factor-critical, but we prove it for all G .)

Lemma 8. *Let G be graph and let L be a list-assignment for G such that $|L(v)| \geq d(v) + 2$ for all $v \in V(G)$. If α and β are L -colourings of G , then we can recolour G from α to β in at most $2n(G) - 1$ steps.*

Proof. We use induction on $|\beta(V(G))|$. The case $|\beta(V(G))| = 1$, is trivial, since then G is an independent set; starting from α , we can recolour each vertex v to $\beta(v)$. We use at most $n(G)$ recolouring steps, which suffices since $n(G) \leq 2n(G) - 1$.

Now assume $|\beta(V(G))| \geq 2$. For each colour c , let $\alpha^{-1}(c) := \{v \in V(G) \text{ s.t. } \alpha(v) = c\}$ and $\beta^{-1}(c) := \{v \in V(G) \text{ s.t. } \beta(v) = c\}$. Since $\sum_{c \in \alpha(V(G))} |\alpha^{-1}(c)| = n(G) = \sum_{c \in \beta(V(G))} |\beta^{-1}(c)|$, there exists c such that $|\alpha^{-1}(c)| \leq |\beta^{-1}(c)|$. Starting from α , for each $v \in \alpha^{-1}(c)$, recolour v to avoid colour c . Now for each $v \in \beta^{-1}(c)$, recolour v with c . After at most $2|\beta^{-1}(c)|$ steps, we have $|\beta^{-1}(c)|$ more vertices that are coloured to agree with β . Now we delete these $|\beta^{-1}(c)|$ vertices, and delete the colour c from $L(v)$ whenever v had some neighbours in $\beta^{-1}(c)$, and finish by induction. \square

Corollary 9. *For every graph G and every $k \geq \Delta(G) + 2$, the diameter of $\mathcal{C}_k(G)$ is at most $2n(G) - 1$.*

Now we prove [Theorem 1\(i\)](#). For easy reference, we restate it below.

Theorem 10. *Let G be an n -vertex graph and L be a list-assignment with $|L(v)| \geq d(v) + 2$ for all $v \in V(G)$. If α and β are L -colourings of G , then we can recolour G from α to β in at most $n(G) + 2\mu(G)$ steps.*

Proof. We use induction on $n(G)$. We assume G is connected; otherwise, we handle each component separately. (This suffices because, for a graph G with components G_1, \dots, G_s , we have $n(G) = \sum_{i=1}^s n(G_i)$ and $\mu(G) = \sum_{i=1}^s \mu(G_i)$.) If $\mu(G - v) = \mu(G)$ for all $v \in V(G)$, then G is factor-critical. This is an easy result of Gallai, but also follows from [Theorem 12\(2\)](#). Now $n(G) + 2\mu(G) = 2n(G) - 1$, so we are done by [Lemma 8](#).

Assume instead that some vertex v is saturated by every maximum matching. Note that $2d(v) < 2|L(v)|$. By Pigeonhole there exists $c \in L(v)$ such that $|\beta^{-1}(c) \cap N(v)| + |\alpha^{-1}(c) \cap N(v)| \leq 1$. (We handle the case when equality holds, since the other case is easier.) Assume there exists $w \in N(v)$ such that $\alpha(w) = c$ and $c \notin \cup_{x \in N(v) \setminus \{w\}} \{\alpha(x), \beta(x)\}$. (If instead $\beta(w) = c$, then we swap the roles of α and β .) Form $\tilde{\alpha}$ from α by recolouring w from $L(w) \setminus (\alpha(N(w)) \cup \{c\})$ and recolouring v with c . Let $G' := G - v$. Let $L'(x) := L(x) - c$ for all $x \in N(v)$ and $L'(x) := L(x)$ for all other $x \in V(G')$. Denote the restrictions to G' of $\tilde{\alpha}$ and β by $\tilde{\alpha}'$ and β' . By induction, we can recolour G' from $\tilde{\alpha}'$ to β' in at most $n(G') + 2\mu(G')$ steps. After this, we can finish by recolouring v with $\beta(v)$. Thus, $\text{dist}_L(\alpha, \beta) \leq 2 + \text{dist}_{L'}(\tilde{\alpha}', \beta') + 1 \leq 2 + n(G') + 2\mu(G') + 1 = 2 + n(G) - 1 + 2(\mu(G) - 1) + 1 = n(G) + 2\mu(G)$. \square

To prove [Theorem 2\(i\)](#), we will use the following lemma to handle the case when G is factor-critical.

Lemma 11. *Let G be a graph and let L be a list-assignment for G such that $|L(v)| \geq 2d(v) + 1$ for all $v \in V(G)$. If α and β are L -colourings of G , then we can recolour G from α to β in at most $\lfloor \frac{3n(G)}{2} \rfloor$ steps.*

Proof. Let α and β be arbitrary L -colourings of G . (Recall that all colours are positive integers.) Let V_1 and V_2 be the subsets of $V(G)$, respectively, for which $\alpha(v) > \beta(v)$ and $\alpha(v) < \beta(v)$. Let $n_1 := |V_1|$

and $n_2 := |V_2|$. Since $n_1 + n_2 \leq n(G)$, we assume by symmetry that $n_1 \leq \frac{n(G)}{2}$; otherwise, we interchange α and β . We show how to recolour G from α to β using at most $n(G) + n_1 \leq n(G) + \lfloor \frac{n(G)}{2} \rfloor$ steps.

First, iteratively for every $v \in V_1$ we recolour v from $L(v) \setminus \cup_{w \in N(v)} \{\beta(w), \gamma(w)\}$, where γ is the current colouring of G . By definition this recolouring is proper (we actually do not need to recolour v if $\gamma(v) \neq \beta(w)$ for all $w \in N(v)$). Next, iteratively for j from $\max\{\cup_{v \in V} L(v)\}$ to $\min\{\cup_{v \in V} L(v)\}$, we recolour with j all vertices $v \in V(G)$ for which $\beta(v) = j$. In these steps, we only have proper colourings, as we will now show. Consider a neighbour w of v . If w has already been recoloured with some $\gamma(w)$, then $\gamma(w) \neq \beta(v)$, by construction. If w has not been recoloured, then $\alpha(w) \leq \beta(w) < \beta(v)$. \square

The following classical result is helpful in our proof of [Theorem 2\(i\)](#).

Theorem 12 (Edmonds–Gallai Decomposition Theorem). *For a graph G , let*

$$\begin{aligned} V_1(G) &:= \{v \in V(G) \text{ such that some maximum matching avoids } v\}, \\ V_2(G) &:= \{v \in V(G) \text{ such that } v \text{ has a neighbour in } V_1(G), \text{ but } v \notin V_1(G)\}, \\ V_3(G) &:= V(G) \setminus (V_1(G) \cup V_2(G)). \end{aligned}$$

Now the following statements hold.⁴

- (1) *The subgraph induced by $V_3(G)$ has a perfect matching.*
- (2) *The components of the subgraph induced by $V_1(G)$ are factor-critical.*
- (3) *If M is any maximum matching, then M contains a perfect matching of each component of $V_3(G)$, and M contains a near-perfect matching of each component of $V_1(G)$, and M matches all vertices of $V_2(G)$ with vertices in distinct components of $V_1(G)$.*
- (4) $\mu(G) = \frac{1}{2} (|V(G)| - c(V_1(G)) + |V_2(G)|)$, where $c(V_1(G))$ denotes the number of components of the graph spanned by $V_1(G)$.

Now we prove [Theorem 2\(i\)](#). For easy reference, we restate it below.

Theorem 13. *Let G be a graph and L be a list-assignment for G with $|L(v)| \geq 2d(v) + 1$ for all $v \in V(G)$. If α and β are L -colourings of G , then we can recolour G from α to β in at most $n(G) + \mu(G)$ steps.*

Proof. We refer to $V_1(G)$, $V_2(G)$, and $V_3(G)$, as defined in [Theorem 12](#). Starting from α , recolour each vertex $v \in V_2(G)$ to avoid each colour used currently on $N(v)$ and also to avoid each colour used on $N(v)$ in β ; call the resulting colouring $\tilde{\alpha}(G)$. Let $G' := G - V_2(G)$. For each $v \in V(G')$, let $L'(v) := L(v) \setminus (\cup_{w \in N(v) \cap V_2(G)} \tilde{\alpha}(w))$. Note that $|L'(v)| \geq 2d_{G'}(v) + 1$. Thus, for each component H of G' , we can recolour H from α to β in at most $\lfloor 3n(H)/2 \rfloor$ steps, by [Lemma 11](#). Finally, we recolour each vertex of $V_2(G)$ to its colouring β . Let $\text{comp}(G')$ denote the set of components of G' . The number of recolouring steps used is at most $2|V_2(G)| + \sum_{H \in \text{comp}(G')} \lfloor 3n(H)/2 \rfloor = 2|V_2(G)| + \frac{3}{2}(|V(G)| - |V_2(G)|) - \frac{1}{2}c(V_1(G)) = |V(G)| + \frac{1}{2}(|V(G)| - c(V_1(G)) + |V_2(G)|) = |V(G)| + \mu(G)$. The final equality uses [Theorem 12\(4\)](#). The first equality uses that the vertices of a component $H \in \text{comp}(G')$ are contained either in $V_1(G)$ or in $V_3(G)$. If $V(H) \subseteq V_3(G)$ then $n(H)$ is even by [Theorem 12\(3\)](#). On the other hand, if $V(H) \subseteq V_1(G)$ then $n(H)$ is odd by [Theorem 12\(2\)](#). \square

By the comment after [Example 6](#), [Theorem 13](#) is sharp. For (non-list) colouring, we get the following.

Corollary 14. *For every graph G and every $k \geq 2\Delta(G) + 1$, we have $\text{diam } C_k(G) = n(G) + \mu(G)$.*

Proof. The upper and lower bounds follow, respectively, from [Theorem 13](#) and [Proposition 1](#). \square

⁴ We prefer the names $V_1(G)$, $V_2(G)$, $V_3(G)$ over the standard terminology $D(G)$, $A(G)$, and $C(G)$, since the latter terms conflict with our use of $D = (V, A)$ for a digraph D with vertex set V and arc set A .

4. Proving the main results for correspondences

In this section, we prove the correspondence colouring portions of our main results: [Theorems 1\(ii\)](#) and [2\(ii\)](#).

Theorem 15. *Let G be a graph and (L, H) be a correspondence cover with $|L(v)| \geq 2d(v) + 1$ for all $v \in V(G)$. If α and β are (L, H) -colourings of G , then G can be recoloured from α to β in at most $n(G) + \tau(G)$ steps.*

Proof. Let S be a minimum vertex cover. For every vertex $v \in S$, we recolour v with a colour that, for every $w \in N(v)$, conflicts (under cover (L, H)) with neither the current colour $\gamma(w)$ nor the final colour $\beta(w)$. This is possible because $|L(v)| \geq 2d(v) + 1$. Now we recolour every vertex $v \in V(G) \setminus S$ with $\beta(v)$ and then do this also for every vertex in S . Note that we use at most $n(G) + \tau(G)$ recolourings. \square

Theorem 16. *Let G be a graph and (L, H) be a correspondence cover for G such that $|L(v)| \geq d(v) + 2$ for all $v \in V(G)$. If α and β are (L, H) -colourings of G , then we can recolour G from α to β in at most $n(G) + 2\tau(G)$ steps.*

Proof. We use induction on $\tau(G)$. If $\tau(G) = 0$, i.e., G is an independent set, then we simply recolour every vertex v to $\beta(v)$. So assume the theorem is true for all graphs G' with $0 \leq \tau(G') < \tau(G)$.

Let S be a minimum vertex cover of G and pick $v \in S$. Consider the union over all $w \in N(v)$ of the edges in H from $\alpha(w)$ and $\beta(w)$ to $L(v)$. By Pigeonhole, since $2d(v) < 2|L(v)|$, some colour $c \in L(v)$ has at most one incident edge in this union. (We handle the case when c has exactly one incident edge, since the other case is easier.) Assume there exists $w_0 \in N(v)$ such that $\alpha(w_0)$ is matched to c and no other $\alpha(w)$ or $\beta(w)$ is matched to c . (If instead $\beta(w_0)$ is matched to c , then we swap the roles of α and β .) Form $\tilde{\alpha}$ from α by first recolouring w_0 with a colour different from $\alpha(w_0)$, that still gives a proper (L, H) -colouring, and afterward recolouring v with c . Let $G' := G - v$. For every $w \in N(v)$, remove from $L(w)$ the colour c' matched with c ; call the resulting correspondence cover (L', H') . Denote the restrictions to G' of $\tilde{\alpha}$ and β by $\tilde{\alpha}'$ and β' . By induction, since $\tau(G') < \tau(G)$, we can recolour G' from $\tilde{\alpha}'$ to β' in at most $n(G') + 2\tau(G')$ steps. After this, we can finish by recolouring v with $\beta(v)$. Thus, $\text{dist}_{(L,H)}(\alpha, \beta) \leq 2 + \text{dist}_{(L',H')}(\tilde{\alpha}', \beta') + 1 \leq 2 + n(G') + 2\tau(G') + 1 = 2 + n(G) - 1 + 2(\tau(G) - 1) + 1 = n(G) + 2\tau(G)$. This proves the theorem. \square

5. Distances in reconfiguration graphs of trees

For each tree T , it is straightforward to check that reconfiguration of correspondence colouring is no harder than reconfiguration of list colouring. Given a tree T and a correspondence cover (L, H) , it is easy to construct a list-assignment L' for T such that L' -colourings of T are in bijection with (L, H) -colourings of T . We can do this by induction on $n(G)$, by deleting a leaf v and extending the list-assignment L' , given by hypothesis, for $T - v$. So, for simplicity, we phrase all results in this section only in terms of list colourings.

Theorem 17. *Fix a tree T and a list-assignment L with $|L(v)| \geq d(v) + 2$ for every $v \in V(T)$. If α and β are proper L -colourings of T , then the distance between α and β in the reconfiguration graph $C_L(T)$ is equal to $\mu(D_{\alpha,\beta}) + \sum_{v \in V} \mathbf{1}_{\alpha(v) \neq \beta(v)}$.*

Proof. The lower bound holds by [Observation 5](#). Now we prove that this lower bound is also an upper bound.

We use induction on $n(T)$. The base case, $n(T) = 1$, is trivial. Assume the theorem is true whenever T has order at most $s - 1$. We will prove it for an arbitrary tree T on s vertices. If $\alpha(v) = \beta(v)$ for some $v \in V(T)$, then we use induction on the components of $T - v$, with $\alpha(v)$ removed from the lists of the neighbours of v .

Suppose instead there are neighbours v and w for which $\vec{vw} \notin A(D_{\alpha,\beta})$. Now $T - vw$ has two components; so let C_v and C_w be the components, respectively, containing v and w . By deleting $\alpha(w)$ from $L(v)$, we can first recolour C_v . Afterward, we delete $\beta(v)$ from $L(w)$ and recolour C_w . By using the induction hypothesis (twice), we get the desired upper bound, since $\mu(D_{\alpha,\beta}) = \mu(D_{\alpha,\beta}[C_v]) + \mu(D_{\alpha,\beta}[C_w])$.

In the remaining case, we have two colour classes, say with colours 1 and 2, which need to be swapped. We pick a smallest vertex cover S , which has size $\tau(D_{\alpha,\beta}) = \mu(D_{\alpha,\beta})$. Iteratively, we recolour every vertex v in S with a colour different from 1, 2, and all colours used to recolour neighbours of v in S . Note that v has at most $d(v) - 1$ neighbours in S , since otherwise S would not be a minimal vertex cover. Since $|L(v)| \geq d(v) + 2$, we can recolour as desired. Next, for each $w \in V(T) \setminus S$, we recolour w with $\beta(w)$. Finally, we recolour each $v \in S$ with $\beta(v)$. This proves the induction step, which finishes the proof. \square

Corollary 18. *If T is a tree and L is a list-assignment with $|L(v)| \geq d(v) + 2$ for every $v \in V(T)$, then*

$$\text{diam } C_L(T) \leq n(T) + \mu(T) \leq \lfloor 3n(T)/2 \rfloor.$$

Proof. Theorem 17 implies that $\text{diam } C_L(T) \leq n(T) + \mu(T)$, and the bound $\mu(T) \leq \lfloor \frac{n(T)}{2} \rfloor$ holds trivially, since each edge of a matching saturates two vertices. \square

Recall that $[k]$ denotes $\{1, \dots, k\}$. Thus, we write $[k]$ -colouring to mean a k -colouring from the colours $[k]$.

Proposition 5. *For every tree T and $k \geq \Delta(T) + 2$, we have*

$$\text{diam } C_k(T) = n(T) + \mu(T) \text{ and } \text{rad } C_k(T) \geq n(T) + \left\lceil \frac{\mu(T)}{2} \right\rceil.$$

Proof. The inequality $\text{diam } C_k(T) \leq n(T) + \mu(T)$ holds by Corollary 18; and this inequality actually holds with equality, since the two $[2]$ -colourings of T are at distance exactly $n(T) + \mu(T)$. That is, $\text{diam } C_k(T) = n(T) + \mu(T)$. Now we consider $\text{rad } C_k(T)$. If we contract all edges of a maximum matching M in T , the result is also a tree, T' . When we 2-colour T' , one colour is used on at least half of the contracted edges. Denote the set of these contracted edges by E' . Note that E' is an induced matching in T . Fix an arbitrary proper k -colouring α of T . Now we construct a proper k -colouring β of T by swapping the colours on the endpoints of each edge $vw \in E'$, i.e., let $\beta(v) := \alpha(w)$ and $\beta(w) := \alpha(v)$ for every edge $vw \in E'$. For every vertex v not belonging to an edge in E' , choose a colour in $[k]$ different from the colours already assigned (in β) to neighbours of v and different from $\alpha(v)$. Now $\text{dist}(\alpha, \beta) \geq n(T) + |E'| \geq n(T) + \lceil \frac{\mu(T)}{2} \rceil$ by Theorem 17. \square

We construct trees T for which certain colourings are more “central” in the reconfiguration graph than others. That is, we construct trees T for which $\text{rad } C_k(T) \neq \text{diam } C_k(T)$. We also study the maximum possible diameter and minimum possible radius of reconfiguration graphs of trees of given order, and the maximum possible difference of these quantities.


Proposition 6. *For every $k \geq 4$, the path P_n satisfies*

$$\text{diam } C_k(P_n) = \left\lfloor \frac{3n}{2} \right\rfloor \text{ and } \text{rad } C_k(P_n) = \left\lceil \frac{4n - 1}{3} \right\rceil.$$

Furthermore, there exist n -vertex trees T , with maximum degree 3, such that for every $k \geq 5$ we have

$$\text{diam } C_k(T) = \left\lfloor \frac{3n}{2} \right\rfloor \text{ and } \text{rad } C_k(T) = \left\lceil \frac{5n - 1}{4} \right\rceil.$$

All such T maximise, over all n -vertex trees, the difference $\text{diam } C_k(T) - \text{rad } C_k(T)$.



$$\alpha(v_i) = \begin{cases} 1, & \text{for } i \equiv 3 \pmod{4} \\ 2, & \text{for } i \equiv 1 \pmod{4} \\ 3, & \text{for } i \equiv 0, 2 \pmod{4} \end{cases} \quad \alpha(w_i) = \begin{cases} 1, & \text{for } i \equiv 1, 2 \pmod{4} \\ 2, & \text{for } i \equiv 3, 4 \pmod{4}. \end{cases}$$

Fig. 3. The comb graph T_{16} with colouring α . The shaded region denotes four edges no two of which, for any colouring β , yield disjoint bidirected edges of $D_{\alpha,\beta}$.

Proof. Consider P_n with vertex set $\{v_1, v_2, \dots, v_n\}$. Fix $k \geq 4$, and let α and β be the colourings with ranges $[3]$ and $[2]$, respectively, such that

$$\alpha(v_i) \equiv i \pmod{3} \quad \text{and} \quad \beta(v_i) \equiv i \pmod{2} \quad \forall i \in [n].$$

Note that $\text{ecc}(\beta) = \lfloor \frac{3n}{2} \rfloor$ by Theorem 17, since switching colours 1 and 2 in β requires $\lfloor \frac{3n}{2} \rfloor$ recolouring steps. Furthermore, Corollary 18 implies that $\text{diam } C_k(P_n) = \text{ecc}(\beta) = \lfloor \frac{3n}{2} \rfloor$. On the other hand, for every proper colouring γ , we know that $D_{\alpha,\gamma}$ cannot have a pair of bidirected edges of the form $v_i v_{i+1}$ and $v_{i+2} v_{i+3}$, since $\alpha(v_i) = \alpha(v_{i+3})$. So $\mu(D_{\alpha,\gamma}) \leq \lceil \frac{n-1}{3} \rceil$; hence, $\text{ecc}(\alpha) \leq n + \lceil \frac{n-1}{3} \rceil$, which implies that $\text{rad } C_k(P_n) \leq \lceil \frac{4n-1}{3} \rceil$.

Next we prove the lower bound on $\text{rad } C_k(P_n)$. For every colouring α of P_n , we can choose at least $\lceil \frac{n-1}{3} \rceil$ disjoint edges such that swapping the colours in α on the endpoints of each edge (and possibly recolouring vertices not in any of these edges) yields another proper colouring. To see this, first select edge $v_1 v_2$. Whenever $v_i v_{i+1}$ has been selected, there exists $j \in \{i+3, i+4\}$ for which $\alpha(v_i) \neq \alpha(v_j)$. Now add edge $v_{j-1} v_j$ to the set of selected edges. When our selection ends, the set E' contains at least $\lceil \frac{n-1}{3} \rceil$ edges. We can now construct a $[k]$ -colouring β where $\beta(w) = \alpha(v)$ and $\beta(v) = \alpha(w)$ for every edge $vw \in E'$, and $\beta(v) \neq \alpha(v)$ for every $v \in V(P_n)$. Theorem 17 gives $\text{dist}(\alpha, \beta) \geq n + \lceil \frac{n-1}{3} \rceil$.

Now let T_n be the n -vertex comb graph; see Fig. 3. Here T_n is formed from a path $v_1 v_2 \dots v_t$, where $t = \lceil \frac{n}{2} \rceil$, by adding, for each v_i (except v_t when n is odd), an additional neighbour w_i . Since $\mu(T_n) = \lfloor \frac{n}{2} \rfloor$, by Proposition 5 the diameter of $C_k(T_n)$ is $\lfloor \frac{3n}{2} \rfloor$. Let α be the colouring shown and described in Fig. 3.

For any proper colouring β of T_n , note that $\mu(D_{\alpha,\beta}) \leq \lceil \frac{n-1}{4} \rceil$. This holds because, for each odd i , the subgraph $D_{\alpha,\beta}[\{v_i, v_{i+1}, v_{i+2}, w_i, w_{i+1}\}]$ contains no two disjoint bidirected edges. Thus, $\text{ecc}(\alpha) \leq \lceil \frac{5n-1}{4} \rceil$, by Theorem 17. Since $\mu(T_n) = \lfloor \frac{n}{2} \rfloor$, Proposition 5 implies that $\text{rad } C_k(T_n) = n + \lceil \lfloor n/2 \rfloor / 2 \rceil = \lceil \frac{5n-1}{4} \rceil$. Thus, $\text{diam } C_k(T_n) - \text{rad } C_k(T_n) = \lfloor \frac{3n}{2} \rfloor - \lceil \frac{5n-1}{4} \rceil = \lfloor \frac{n}{4} \rfloor$. For every n -vertex tree T , Proposition 5 implies that $\text{diam } C_k(T) - \text{rad } C_k(T) \leq (n + \mu(T)) - (n + \lceil \frac{\mu(T)}{2} \rceil) = \lfloor \frac{\mu(T)}{2} \rfloor \leq \lfloor \frac{n}{4} \rfloor$. Thus, T_n maximises this difference. \square

For list colourings, the difference $\text{diam } C_l(T) - \text{rad } C_l(T)$ can be even larger.

Proposition 7. For every tree T , there exists a list-assignment L , with $|L(v)| \geq d(v)+2$ for all $v \in V(T)$, such that $\text{diam } C_l(G) = n(T) + \mu(T)$ and $\text{rad } C_l(G) = n(T)$. These values are, respectively, the maximum and minimum possible over all such list-assignments L .

Proof. Choose a maximum matching M of T and fix L such that

$$|L(v) \cap L(w)| = \begin{cases} 0, & \text{when } vw \notin M \\ 2, & \text{when } vw \in M. \end{cases}$$

Let α and β be L -colourings such that, for every edge $vw \in M$, we have $\alpha(v) = \beta(w)$ and $\alpha(w) = \beta(v)$. For every vertex v not saturated by M , we pick $\alpha(v)$ and $\beta(v)$ to be arbitrary distinct colours in $L(v)$. By [Theorem 17](#), we have $\text{dist}(\alpha, \beta) = n(T) + \mu(T)$. [Corollary 18](#) implies that $\text{diam } C_L(T) = n(T) + \mu(T)$. Further, this diameter is the maximum possible over all such list-assignments L .

To show that $\text{rad } C_L(T) \leq n(T)$, we construct an L -colouring γ such that $\text{dist}(\gamma, \delta) \leq n(T)$ for every L -colouring δ . Since $|L(v) \setminus (\cup_{w \in N(v)} L(w))| \geq 3 - 2 = 1$ for every vertex v , we can form an L -colouring γ such that every $v \in V(T)$ satisfies $\gamma(v) \notin \cup_{w \in N(v)} L(w)$. For every proper colouring δ , by our construction of γ , we can recolour δ into γ greedily; hence, $\text{dist}(\delta, \gamma) \leq n(T)$.

Now we show that $\text{rad } C_L(T) \geq n(T)$. For every L -colouring γ , there exists an L -colouring δ such that $\delta(v) \neq \gamma(v)$ for all v . To see this, let $L'(v) := L(v) \setminus \gamma(v)$. Since $|L'(v)| \geq d(v) + 1$ for all v , we form δ by colouring G greedily (in any order) from L' . Clearly, $\text{dist}(\gamma, \delta) \geq n(G)$. Thus, $\text{rad } C_L(T) = n(T)$. In fact, this lower bound does not depend on the specific choice of L , but only uses that $|L(v)| \geq d(v) + 2$ for all v . Thus, this radius is the minimum over all such list-assignments L . \square

6. Proving the list conjecture for complete bipartite graphs and cactuses

A *cactus* is a connected graph in which each edge lies on at most one cycle. In this section, we prove the [List Conjecture](#) for all complete bipartite graphs and for all cactuses. Both proofs are by induction, and rely on the following helpful lemma.

Lemma 19. Fix a positive integer b . Let $G = (V, E)$ be a graph for which there exists a list-assignment L and two proper L -colourings α, β such that $|L(v)| \geq d(v) + b$ for every $v \in V(G)$, and $\text{dist}_L(\alpha, \beta) > n(G) + \mu(G)$, but, for every proper induced subgraph of G , no such list-assignment exists. Now for every partition $V(G) = V_1 \cup V_2$ into two non-empty subsets of vertices, $D_{\alpha, \beta}$ has at least one arc from V_1 to V_2 and at least one arc from V_2 to V_1 , i.e., $D_{\alpha, \beta}$ is strongly connected.

Proof. Assume the lemma is false; by symmetry, assume $D_{\alpha, \beta}$ has no arcs from V_1 to V_2 . Let $G_1 := G[V_1]$ and $G_2 := G[V_2]$. Let γ be the L -colouring where $\gamma(v) := \beta(v)$ when $v \in V_1$ and $\gamma(v) := \alpha(v)$ when $v \in V_2$.

For every $v \in V_1$, let $L'(v) := L(v) \setminus \{\alpha(w) \mid w \in N(v) \cap V_2\}$. Note that $|L'(v)| \geq d_{G_1}(v) + b$. Also note that still $\gamma(v) \in L'(v)$ for every $v \in V_1$; this is where we use that there is no arc from V_1 to V_2 . Since G_1 is a proper induced subgraph of G , by hypothesis $\text{diam } C_{L'}(G_1) \leq n(G_1) + \mu(G_1)$; thus $\text{dist}_L(\alpha, \gamma) \leq n(G_1) + \mu(G_1)$. Now for every $v \in V_2$, let $L'(v) := L(v) \setminus \{\beta(w) \mid w \in N(v) \cap V_1\}$. Similarly to before, $|L'(v)| \geq d_{G_2}(v) + b$, so $\text{diam } C_{L'}(G_2) \leq n(G_2) + \mu(G_2)$; thus $\text{dist}_L(\gamma, \beta) \leq n(G_2) + \mu(G_2)$. By the bounds above and the triangle inequality, we have

$$\begin{aligned} \text{dist}_L(\alpha, \beta) &\leq \text{dist}_L(\alpha, \gamma) + \text{dist}_L(\gamma, \beta) \\ &\leq n(G_1) + \mu(G_1) + n(G_2) + \mu(G_2) \\ &\leq n(G) + \mu(G). \end{aligned}$$

The last step uses that the union of a matching in G_1 and a matching in G_2 is a matching in G . So G is not a counterexample, and the lemma is true. \square

It is helpful to note that an analogous statement holds for correspondence colouring. In fact, its proof is nearly identical to that given above. We use this observation in our proof of [Theorem 27](#).

Using [Lemma 19](#), we prove that the [List Conjecture](#) holds for all complete bipartite graphs.

Theorem 20. The [List Conjecture](#) is true for all complete bipartite graphs.

Proof. Suppose the theorem is false and choose a counterexample $K_{p,q}$ minimising $p + q$. By symmetry, we assume that $p \leq q$. Denote the parts of G by U and W , with $|U| = p$ and $|W| = q$.

By Lemma 19, we have $\alpha(U) \subseteq \beta(W)$ and $\alpha(W) \subseteq \beta(U)$. By swapping the roles of α and β , we also have $\beta(U) \subseteq \alpha(W)$ and $\beta(W) \subseteq \alpha(U)$; thus, $\alpha(U) = \beta(W)$ and $\alpha(W) = \beta(U)$. Note that $\alpha(U) \cap \beta(U) = \alpha(U) \cap \alpha(W) = \emptyset$, because the graph is complete bipartite.

Case 1: $|W| \geq |\alpha(U)| + |\beta(U)| - 1$. For each $u \in U$, we have $|L(u)| \geq |W| + 2 \geq |\alpha(U)| + |\beta(U)| + 1 = |\beta(W)| + |\alpha(W)| + 1$. First recolour each $u \in U$ from $L(u) \setminus (\alpha(W) \cup \beta(W))$. Now recolour each $w \in W$ with $\beta(w)$. Finally, recolour each $u \in U$ with $\beta(u)$. The number of steps that we use is at most $2|U| + |W| = n(G) + \mu(G)$.

Case 2: $|W| \leq |\alpha(U)| + |\beta(U)| - 2$. If $|W| \geq 2|\alpha(U)|$ and $|W| \geq 2|\beta(U)|$, then $2|W| \geq 2|\alpha(U)| + 2|\beta(U)|$, which contradicts the case. So assume, by symmetry, that $|W| \leq 2|\beta(U)| - 1$; if not, then simply interchange the roles of α and β . Now, by Pigeonhole, there exists $c \in \beta(U)$ such that $|\alpha^{-1}(c)| = |\alpha^{-1}(c) \cap W| \leq 1$. So, recolour $w \in \alpha^{-1}(c)$, if such w exists, to avoid $\alpha(U) \cup \{c\}$, and then recolour every $u \in \beta^{-1}(c)$ with c . Now we delete every u such that $\beta(u) = c$, we delete c from $L(w)$ for every $w \in W$, and we finish on the resulting smaller graph G_2 (with an assignment of smaller lists) by the minimality of G . In total, the number of recolouring steps we use is at most $|\alpha^{-1}(c)| + |\beta^{-1}(c)| + |V(G_2)| + \mu(G_2) \leq |\alpha^{-1}(c)| + |\beta^{-1}(c)| + (n(G) - |\beta^{-1}(c)|) + (\mu(G) - |\beta^{-1}(c)|) \leq n(G) + \mu(G)$. \square

Using Lemma 19, we prove that the List Conjecture holds for all cycles.

Lemma 21. Let G be a cycle $v_1v_2 \dots v_n$ and L be a list-assignment such that, for all v_i , we have $|L(v_i)| \geq 4$. If α and β are proper L -colourings of G , then we can recolour G from α to β in at most $\lfloor 3n/2 \rfloor$ steps.

Proof. By Lemma 19, with $b = 2$, if $\text{dist}(\alpha, \beta) > \lfloor 3n/2 \rfloor$, then $D_{\alpha,\beta}$ is strongly connected. This implies that $D_{\alpha,\beta}$ contains a directed cycle C_n as a subdigraph. (If $D_{\alpha,\beta}$ contains a bidirected path P , then $|\alpha(V(P)) \cup \beta(V(P))| = 2$. So, if P is spanning, then its order must be even, to avoid a conflict between the colours of its endpoints. But then $D_{\alpha,\beta}$ contains an additional arc, so $D_{\alpha,\beta}$ contains a directed cycle, as claimed.)

If $D_{\alpha,\beta}$ is a bidirected cycle, then n must be even, as in the previous paragraph, and $\frac{3n}{2}$ recolouring steps suffice. Suppose instead that $D_{\alpha,\beta}$ is precisely a directed cycle. When $n = 3$, we recolour one vertex v in a colour absent from $\alpha(V(G)) \cup \beta(V(G))$, recolour the other two vertices in order (to match β), and finally recolour v with $\beta(v)$. When $n \geq 4$, we recolour one vertex v in a colour absent from $\alpha(v) \cup \alpha(N(v))$. Now we can recolour correctly (in order) all vertices of G except for v and one of its neighbours. Correctly colouring these final two vertices takes at most 3 recolouring steps. Now we are done, since $1 + (n - 2) + 3 \leq \lfloor 3n/2 \rfloor$.

In the remaining case, we have $n \geq 4$ and some directed edge is adjacent to a bidirected edge. Without loss of generality, we assume $\overrightarrow{v_1v_2}, \overrightarrow{v_2v_3}, \overrightarrow{v_2v_1} \in A(D_{\alpha,\beta})$ but $\overrightarrow{v_3v_2} \notin A(D_{\alpha,\beta})$. Recolour v_1 with a colour c different from $\alpha(v_1), \alpha(v_2)$, and $\alpha(v_n)$. Now delete c from $L(v_2)$ and $L(v_n)$, and let $G' := G - v_1$. By Theorem 17, we can recolour G' from α to β (both restricted to G') using at most $(n - 1) + \lfloor \frac{n-2}{2} \rfloor$ recolouring steps. Here we use that $\overrightarrow{v_3v_2}$ is not an arc, so $\mu(D_{\alpha,\beta}[V(G')]) \leq \lfloor \frac{n-2}{2} \rfloor$. Finally, we recolour v_1 with $\beta(v_1)$. \square

Using Lemmas 19 and 21, we can prove that the List Conjecture is true for every cactus.

Theorem 22. The List Conjecture is true for every cactus.

Proof. Instead assume the theorem is false. Let G be a counterexample minimising $n(G)$ and let α and β be L -colourings with $\text{dist}(\alpha, \beta) > n(G) + \mu(G)$. Every proper induced subgraph H of G is a disjoint union of cactuses, say H_1, \dots, H_r ; since $n(H) = \sum_{i=1}^r n(H_i)$ and $\mu(H) = \sum_{i=1}^r \mu(H_i)$, the List Conjecture must hold for H , by the minimality of G . Lemma 19 implies that $D_{\alpha,\beta}$ is strongly connected. And Lemma 21 implies that G is not a cycle. The following claim restricts the structure of G, α , and β .

Claim 23. *If $v \in V(G)$ and $\mu(G - v) = \mu(G) - 1$, then $L(v) \subseteq \cup_{w \in N(v)} \{\alpha(w), \beta(w)\}$. In particular, each block of G containing v gives rise to exactly two arcs in $D_{\alpha, \beta}$ incident to v , and v has no adjacent leaf in G .*

Proof. Instead assume there exists $c \in L(v) \setminus \cup_{w \in N(v)} \{\alpha(w), \beta(w)\}$. Recolour v with c , let $G' := G - v$, let $L'(w) := L(w) \setminus \{c\}$ for all $w \in N(v)$, and otherwise let $L'(w) := L(w)$. Denote by α' and β' the restrictions to G' of α and β . Since G is minimal, we can recolour G' from α' to β' , using L' , in at most $n(G') + \mu(G') = n(G) - 1 + \mu(G) - 1$ recolouring steps. We finish by recolouring v with $\beta(v)$. This proves the first statement.

Since $D_{\alpha, \beta}$ is strongly connected, each block B of G containing v gives rise to at least two arcs in $D_{\alpha, \beta}$ incident to v , one in each direction. Since G is a cactus, each such B contains at most 2 neighbours of v . So at least half of the neighbours of v are coloured $\beta(v)$ by α , and also at least half of the neighbours of v are coloured $\alpha(v)$ by β . Therefore $|\cup_{w \in N(v)} \{\alpha(w), \beta(w)\}| \leq 2 \cdot (d(v)/2 + 1) = d(v) + 2$. If any such B is either K_2 or gives rise to at least three arcs in $D_{\alpha, \beta}$ incident with v , then $|\cup_{w \in N(v)} \{\alpha(w), \beta(w)\}| \leq d(v) + 1 < |L(v)|$, which contradicts the first statement. This proves the second statement. \diamond

It is easy to check that every neighbour v of a leaf w in G satisfies $\mu(G - v) = \mu(G) - 1$. Thus, Claim 23 implies that G has no leaf vertex. This implies that all endblocks of G are cycles. Let C_s be an endblock. Denote its vertices by w_1, \dots, w_s , such that w_s is the unique cut-vertex in the block.

Claim 24. *The endblock C_s is not an even cycle.*

Proof. Assume instead that C_s is an even cycle. It is easy to check that $\mu(G - w_i) = \mu(G) - 1$ whenever i is even. That is, every maximum matching saturates w_i whenever i is even. By Claim 23, every w_i is incident in $D_{\alpha, \beta}$ to exactly two arcs arising from C_s . Since $D_{\alpha, \beta}$ is strongly connected, this implies that $D_{\alpha, \beta}[V(C_s)]$ is a directed cycle; by symmetry, we assume that it is oriented as $\overrightarrow{w_s w_1}, \overrightarrow{w_1 w_2}, \dots, \overrightarrow{w_{s-1} w_s}$.

We first recolour w_s with a colour absent from $\alpha(N[w_s]) = (\cup_{x \in N(w_s)} \alpha(x)) \cup \beta(w_{s-1})$. Now we recolour each w_i to $\beta(w_i)$, with i decreasing from $s - 1$ to 2. Let $G' := G - \{w_1, \dots, w_{s-1}\}$, let $L'(w_s) := L(w_s) \setminus \{\alpha(w_1), \beta(w_{s-1})\}$, and otherwise let $L'(v) := L(v)$. Since G is minimal, we can recolour G' from its current colouring to β (restricted to G') using at most $n(G') + \mu(G') \leq (n(G) - (s - 1)) + (\mu(G) - \frac{s}{2} + 1)$ steps. Finally, recolour w_1 to $\beta(w_1)$. The total number of recolouring steps is at most $1 + (s - 2) + (n(G) - (s - 1) + \mu(G) - \frac{s}{2} + 1) + 1 = n(G) + \mu(G) + 2 - \frac{s}{2}$. Since $s \geq 4$, this is at most $n(G) + \mu(G)$. \diamond

Claim 25. *The endblock C_s is not an odd cycle.*

Proof. Assume instead that C_s is an odd cycle. If $D_{\alpha, \beta}[C_s - w_s]$ contains fewer than $\lfloor \frac{s}{2} \rfloor$ disjoint digons (for example, this is true when $s = 3$), then we can easily finish, as follows. We recolour w_s in a colour different from $\alpha(N(w_s)) \cup \beta(\{w_1, w_{s-1}\})$. This is possible because $\beta(w_s)$ is used by α on at least two neighbours of w_s , since $D_{\alpha, \beta}$ is strongly connected. By Theorem 17, we recolour $C_s - w_s$ to β in at most $(s - 1) + \lfloor \frac{s}{2} \rfloor - 1$ steps, and we then recolour $G \setminus (C_s - w_s)$ to β in at most $n(G) - (s - 1) + \mu(G) - \lfloor \frac{s}{2} \rfloor$ steps. Thus, we recolour G from α to β in at most $n(G) + \mu(G)$ steps.

Now we consider the other case, when $w_i w_{i+1}$ is a digon of $D_{\alpha, \beta}$ for every odd i with $i < s$. Since $D_{\alpha, \beta}[C_s]$ is strongly connected, we assume $D_{\alpha, \beta}$ contains arc $w_i w_{i+1}$ for every $0 \leq i \leq s - 1$. Similar to the proof of Lemma 21, here $D_{\alpha, \beta}[C_s]$ cannot contain a spanning bidirected path, since s is odd. This implies, for some even j , that $D_{\alpha, \beta}$ does not contain $w_{j+1} w_j$. Now we will recolour some set of w_i 's to reach a new colouring $\tilde{\alpha}$, such that $D_{\tilde{\alpha}, \beta}[C_s]$ is either acyclic or contains a single directed cycle, $w_{s-2} w_{s-1}$. From $\tilde{\alpha}$, we can recolour $G \setminus (C_s - w_s)$ by the minimality of G , and afterward finish on $C_s - w_s$. The details follow.

First suppose that $\beta(w_s) = \alpha(w_{s-1})$. For every i such that either (a) $i < j$ and i is odd or (b) $i > j$ and i is even, recolour w_i with a colour different from those in $\{\alpha(w_{i-1}), \alpha(w_i), \alpha(w_{i+1})\}$. Here we take the indices modulo s , i.e., $w_0 = w_s$. If $\beta(w_s) \neq \alpha(w_{s-1})$, then we recolour the same set of w_i 's,

except for w_{s-1} . Let $G' := G - \{w_1, \dots, w_{s-1}\}$. By the minimality of G , we can recolour G' in at most $n(G') + \mu(G') = (n(G) - (s - 1)) + (\mu(G) - \lfloor \frac{s}{2} \rfloor)$ steps. In the case that $\beta(w_s) \neq \alpha(w_{s-1})$, we now recolour w_{s-1} with a colour different from those in $\{\alpha(w_{s-2}), \alpha(w_{s-1}), \beta(w_s)\}$. For every odd i such that $j + 1 \leq i \leq s - 2$, we recolour w_i with $\beta(w_i)$. Next, for every even i such that $2 \leq i \leq j$, we recolour w_i with $\beta(w_i)$. Finally, the remaining vertices in C_s can also be recoloured with their colour in β . This process uses at most $n(G) + \mu(G)$ steps. \diamond

Recall that every endblock of G is a cycle, as observed following Claim 23. Thus, Claims 24 and 25 yield a contradiction, which proves the theorem. \square

7. Proving the correspondence conjecture for cactuses, subcubic graphs, and graphs with low maximum average degree

The *maximum average degree* of a graph G , denoted $\text{mad}(G)$, is the maximum, taken over all subgraphs H , of the average degree of H . That is, $\text{mad}(G) := \max_{H \subseteq G} 2|E(H)|/|V(H)|$. Let $d^1(v)$ denote the number of neighbours w of v such that $d(w) = 1$. In this section we prove the Correspondence Conjecture for all subcubic graphs, cactuses, and graphs G with $\text{mad}(G) < 2.4$. To do so, we often implicitly use the following observation. We omit its proof, which is easy.

Observation 26. *If G is a graph with a minimum vertex cover S and $v \in S$, then $\tau(G - v) = \tau(G) - 1$.*

Theorem 27. *Let G be a graph and (L, H) a correspondence cover for G such that, for all $v \in V(G)$ we have $|L(v)| \geq d(v) + 2$. Now $\text{diam}_{C_{(L,H)}}(G) \leq n(G) + \tau(G)$ if at least one of the following holds.*

- (a) $\Delta(G) \leq 3$.
- (b) G is a cactus.
- (c) $\text{mad}(G) < 2.4$.

Proof. Fix G satisfying (a), (b), or (c) and (L, H) as in the theorem. Fix arbitrary (L, H) -colourings α and β . Our proof is by a double induction: primarily on $\tau(G)$ and secondarily on $|V(G)|$. The base case, $\tau(G) = 0$, is trivial, since G is an independent set and we can greedily recolour G from α to β . For the induction step, assume $\tau(G) \geq 1$.

We first show that G contains either (i) a vertex v such that $d(v) \leq 3$ and v lies in some minimum vertex cover or (ii) a vertex v such that $d(v) \geq 4$ and $d(v) - d^1(v) \leq 2$. Next we show how to proceed by induction in each of cases (i) and (ii).

If G satisfies (a), then clearly G contains an instance of (i). Suppose G satisfies (b), and consider an endblock B of G . If B is a cycle, then B contains adjacent non-cut vertices, v and w ; note that $d(v) = d(w) = 2$. Further, every vertex cover of G contains v or w , so G contains (i). Assume instead that every endblock of G is an edge.

Form a graph J with a vertex for every block in G , where two vertices of J are adjacent if their corresponding blocks share a vertex in G . Now the endpoints of a diameter in J correspond with pendent edges in G . For the leaf v corresponding to such a pendent edge, call its neighbour w . Clearly w lies in some minimum vertex cover of G . If $d(w) \leq 3$, then G contains (i). Otherwise, $d(w) - d^1(w) \leq 2$, since by the choice of w at most one block incident to w is not an endblock; so G contains (ii). This concludes the case that G satisfies (b).

Assume instead that G satisfies (c). Suppose, to reach a contradiction, that G contains neither (i) nor (ii). Form G' from G by deleting all vertices with degree at most 1. We will show that $2|E(G')|/|V(G')| \geq 2.4$, a contradiction. Note that G' has minimum degree (at least) 2. If an arbitrary vertex v satisfies $d_{G'}(v) = 2$, then $d_G(v) = 2$, since otherwise v is an instance of (i) or (ii) in G . Further, if $d_{G'}(v) = 2$, then $d(w) \geq 3$ for all $w \in E(G')$, since otherwise v or w is an instance of (i) in G . Now we will reach a contradiction with discharging. Give each vertex v in G' charge $\text{ch}(v) := d_{G'}(v)$. We use a single discharging rule: Each 2-vertex in G' takes 0.2 from each neighbour. Now each 2-vertex v in G' finishes with charge $\text{ch}^*(v) = 2 + 2(0.2) = 2.4$. And each vertex v with $d_{G'}(v) \geq 3$ finishes with charge $\text{ch}^*(v) \geq d_{G'}(v) - 0.2d_{G'}(v) = 0.8d_{G'}(v) \geq 2.4$. This yields the desired contradiction, which proves that G contains either (i) or (ii) if $\text{mad}(G) < 2.4$.

Now we prove the induction step in the cases that G contains (i) or (ii).

Suppose G contains (i). Suppose at most one neighbour, say w , of v has $\alpha(w)$ matched with $\beta(v)$. Now recolour w with a colour not matched to $\alpha(x)$, for every $x \in N(w)$, and not matched to $\beta(v)$. Next, recolour v with $\beta(v)$, and proceed on $G - v$ by induction (with the colour matched to $\beta(v)$ deleted from the lists of all vertices in $N(v)$). So instead assume there exist at least two such neighbours, say w_1 and w_2 . By interchanging the roles of α and β (and repeating this argument), we see that also there exist two neighbours, say x_1 and x_2 , such that $\beta(x_1)$ and $\beta(x_2)$ are matched to $\alpha(v)$. The total number of colours in $L(v)$ matched to $\alpha(w)$ or $\beta(w)$ for some $w \in N(v)$ is at most $2d(v) - 2 < d(v) + 2$, since $d(v) \leq 3$. Hence, there exists $c \in L(v)$ that is not matched to $\alpha(w)$ or $\beta(w)$ for all $w \in N(v)$. Recolour v with c . Proceed on $G - v$ by induction, with that colour that c is matched to deleted from the list $L(w)$ for each $w \in N(v)$. After finishing on $G - v$, recolour v with $\beta(v)$. The number of steps we use is at most $1 + n(G - v) + \tau(G - v) + 1 = n(G) + \tau(G)$.

Suppose instead G contains (ii). The proof is almost the same as for (i). If there exists $w \in N(v)$ such that $d(w) = 1$ and $\alpha(v)$ is not matched to $\beta(w)$, then simply recolour w with $\beta(w)$, and proceed by induction; here we use the secondary induction hypothesis, since possibly $\tau(G - w) = \tau(G)$. So assume that no such w exists. By interchanging the roles of α and β , we also assume that $\beta(v)$ is matched to $\alpha(w)$ for each $w \in N(v)$ with $d(w) = 1$. Further, if v has a non-leaf neighbour w , then at least one such neighbour has $\alpha(v)$ matched to $\beta(w)$ and at least one (possibly the same one) has $\alpha(w)$ matched to $\beta(v)$. (This follows from the correspondence colouring analog of Lemma 19.) But now there exists $(v, i) \in L(v)$ such that for all $w \in N(v)$, colour (v, i) is not matched to either $\alpha(w)$ or $\beta(w)$. Recolour v to (v, i) , and let $G' := G - v$. Form (L', H) from (L, H) by deleting from $L(w)$, for every $w \in N(v)$, the colour matched with (v, i) . By induction, we can recolour G' from α to β (both restricted to G') using at most $n(G') + \tau(G') = n(G) + \tau(G) - 2$ steps. Finally, recolour v to $\beta(v)$. This uses at most $n(G) + \tau(G)$ steps, which finishes the proof. \square

Corollary 28. *The List Conjecture holds for all bipartite graphs G with $\Delta(G) \leq 3$.*

Proof. When G is bipartite, recall that $\tau(G) = \mu(G)$. \square

8. Concluding remarks

In this paper, we give evidence for both the List Conjecture and the Correspondence Conjecture. We also give evidence for an affirmative answer to Question 1. The List Conjecture and Question 1 would together determine the precise diameter for $\mathcal{C}_{\Delta(G)+2}(G)$ and, as such, a precise bound when Cereceda’s Conjecture is restricted to regular graphs. So we explicitly conjecture the following.

Conjecture 3 (Regular Cereceda’s Conjecture). *For a d -regular graph G , if $k = d + 2$, then $\text{diam } \mathcal{C}_k(G) = n(G) + \mu(G)$.*

In Theorem 10 we prove, when $|L(v)| \geq d(v) + 2$ for all $v \in V(G)$, that $\text{diam } \mathcal{C}_L(G) \leq n(G) + 2\mu(G) \leq 2n(G)$. In a similar vein, it would be interesting to show that $\text{diam } \mathcal{C}_L(G)$ is linear when $|L(v)| \geq \lceil \text{mad}(G) + 2 \rceil$. The following conjecture can be viewed as a “balanced” version of the List Conjecture.

Conjecture 4 (Mad Colouring Reconfiguration Conjecture). *For a graph G with $\text{mad}(G) = d$, if L is a list-assignment such that $|L(v)| \geq \lceil d + 2 \rceil$ for every $v \in V(G)$, then $\text{diam } \mathcal{C}_L(G) = O_d(n)$.*

Feghali [14] proved that if $\epsilon > 0$ and $k \geq d + 1 + \epsilon$, then $\text{diam } \mathcal{C}_k(G) = O_d(n(2 \ln n)^d)$. Conjecture 4 aims to prove a similar result for list colouring, with one more colour available for each vertex, and with a somewhat stronger bound on diameter. All planar graphs G have $\text{mad}(G) < 6$, and all triangle-free planar graphs G have $\text{mad}(G) < 4$, so we note that Conjecture 4 would imply stronger forms of [15, Conjecture 22]. If G is regular, then $\text{mad}(G) = \text{degen}(G)$, so Conjecture 4 is true by Theorem 10. Conjecture 4 is also related to⁵ a conjecture of Bartier et al. [16, Conjecture 1.6] that graphs G with $\text{degen}(G) = d$ satisfy $\text{diam } \mathcal{C}_{d+3}(G) = O_d(n)$.

⁵ If $\text{mad}(G)$ is not an integer, then $\text{degen}(G) + 3 \leq \lceil \text{mad}(G) + 2 \rceil$. But if G has a regular subgraph of degree $\text{mad}(G)$, then this inequality fails.

We do not know if [Conjecture 4](#) might be true with a linear bound of the form $O(n)$ instead of $O_d(n)$. But we do note, for this version of the conjecture and every constant c , that no bound of the form $n(G) + c\mu(G)$ can hold. This is shown by the star $K_{1,3c+3} = (\{u\} \cup \{w_1, \dots, w_{3c+3}\}, E)$ and the colourings α, β with $\alpha(u) = 1, \beta(u) = 4, \beta(w_i) = 1$ and $\alpha(w_i) = 1 + \lceil i/(c+1) \rceil$. Here $\lceil \text{mad}(G) + 2 \rceil = 4$ and $\text{dist}(\alpha, \beta) > n + c$.

For a correspondence cover (L, H) such that $|L(v)| = d(v) + 2$ for all $v \in V(G)$, in [Theorem 1\(ii\)](#) we proved that $\text{diam}_{C_{(L,H)}(G)} \leq n(G) + 2\tau(G)$. We view this as modest evidence that Cereceda's Conjecture might remain true in the more general context of correspondence colourings.

8.1. Open problems

The three focuses of this paper are the [List Conjecture](#), the [Correspondence Conjecture](#), and (to a lesser extent) [Question 1](#). All of these remain open. However, each of them seems rather hard. So, to motivate further research, below we identify some specific classes of graphs for which we believe that each conjecture may be approachable. We begin with some graph classes for which it would be particularly interesting to make further progress on the [List Conjecture](#).

- (1) Complete r -partite graphs for each $r \geq 3$. (We know the [List Conjecture](#) is true for both complete graphs and complete bipartite graphs, so this is a natural common generalisation.)
- (2) Bipartite graphs, not necessarily complete.
- (3) Outerplanar graphs and, more generally, planar graphs.
- (4) Subcubic graphs. (We already proved the [Correspondence Conjecture](#) for this class; nonetheless, the [List Conjecture](#) remains open.)

Conversely, the [Correspondence Conjecture](#) remains open for the following basic graph classes.

- (1) Complete graphs. (The argument of Bonamy and Bousquet for complete graphs directly yields the [List Conjecture](#), but does not yield the [Correspondence Conjecture](#).)
- (2) Complete bipartite graphs.
- (3) Bipartite graphs, not necessarily complete. (This would imply the [List Conjecture](#) for the same class.)
- (4) Outerplanar graphs and, more generally, planar graphs.

Finally, we stress that it will be interesting to improve on [Theorems 1](#) and [2](#). For example, find the smallest ϵ such that for every graph G and list-assignment L with $|L(v)| \geq d(v) + 2$ for all $v \in V(G)$, we have $\text{diam}_{C_L(G)} \leq n(G) + (1 + \epsilon)\mu(G)$. We proved $\epsilon = 1$ suffices, while $\epsilon = 0$ would resolve the [List Conjecture](#).

Acknowledgements

We thank the organisers of the online workshop *Graph Reconfiguration* of the Sparse Graphs Coalition,⁶ where this project started. We also thank the two referees for their careful reading and valuable feedback.

References

- [1] J. van den Heuvel, The complexity of change, in: *Surveys in Combinatorics 2013*, in: London Math. Soc. Lecture Note Ser., Vol. 409, Cambridge Univ. Press, Cambridge, 2013, pp. 127–160, [arXiv:1312.2816](https://arxiv.org/abs/1312.2816).
- [2] N. Nishimura, *Introduction to reconfiguration*, *Algorithms (Basel)* 11 (4) (2018) 25, Paper No. 52.
- [3] M. Jerrum, A very simple algorithm for estimating the number of k -colorings of a low-degree graph, *Random Struct. Algorithms* 7 (2) (1995) 157–165, [http://dx.doi.org/10.1002/rsa.3240070205](https://doi.org/10.1002/rsa.3240070205).
- [4] L. Cereceda, *Mixing Graph Colourings* (Ph.D. thesis), The London School of Economics and Political Science (LSE), 2007.
- [5] N. Bousquet, M. Heinrich, A polynomial version of Cereceda's conjecture, *J. Combin. Theory Ser. B* 155 (2022) 1–16, [http://dx.doi.org/10.1016/j.jctb.2022.01.006](https://doi.org/10.1016/j.jctb.2022.01.006).

⁶ For more information, visit <https://sparse-graphs.mimuw.edu.pl/doku.php>.

- [6] C.M. Mynhardt, S. Nasserar, Reconfiguration of colourings and dominating sets in graphs, 2020, pp. 171–191, <http://dx.doi.org/10.1201/9780429280092-10>.
- [7] N. Bousquet, L. Feuilloley, M. Heinrich, M. Rabie, Short and local transformations between $(\Delta+1)$ -colourings, 2022, [arXiv:2203.08885](https://arxiv.org/abs/2203.08885).
- [8] M. Bonamy, N. Bousquet, Recoloring graphs via tree decompositions, *European J. Combin.* 69 (2018) 200–213, [arXiv:1403.6386](https://arxiv.org/abs/1403.6386).
- [9] L. Cereceda, J. van den Heuvel, M. Johnson, Connectedness of the graph of vertex-colourings, *Discrete Math.* 308 (5–6) (2008) 913–919, <http://dx.doi.org/10.1016/j.disc.2007.07.028>.
- [10] P. Bonsma, L. Cereceda, Finding paths between graph colourings: PSPACE-completeness and superpolynomial distances, *Theoret. Comput. Sci.* 410 (50) (2009) 5215–5226, <http://dx.doi.org/10.1016/j.tcs.2009.08.023>.
- [11] M. Molloy, The list chromatic number of graphs with small clique number, *J. Combin. Theory Ser. B* 134 (2019) 264–284, <http://dx.doi.org/10.1016/j.jctb.2018.06.007>.
- [12] B. Bollobás, P.A. Catlin, P. Erdős, Hadwiger’s conjecture is true for almost every graph, *European J. Combin.* 1 (3) (1980) 195–199, [http://dx.doi.org/10.1016/S0195-6698\(80\)80001-1](http://dx.doi.org/10.1016/S0195-6698(80)80001-1).
- [13] M. Delcourt, L. Postle, Reducing linear hadwiger’s conjecture to coloring small graphs, 2021, [arXiv:2108.01633](https://arxiv.org/abs/2108.01633).
- [14] C. Feghali, Reconfiguring colorings of graphs with bounded maximum average degree, *J. Combin. Theory Ser. B* 147 (2021) 133–138, <http://dx.doi.org/10.1016/j.jctb.2020.11.001>.
- [15] Z. Dvořák, C. Feghali, A Thomassen-type method for planar graph recoloring, *European J. Combin.* 95 (2021) 12, <http://dx.doi.org/10.1016/j.ejc.2021.103319>, Paper No. 103319.
- [16] V. Bartier, N. Bousquet, C. Feghali, M. Heinrich, B. Moore, T. Pierron, Recoloring planar graphs of girth at least five, *SIAM J. Discrete Math.* (ISSN: 0895-4801) 37 (1) (2023) 332–350, <http://dx.doi.org/10.1137/21M1463598>, <https://doi-org.proxy.library.vcu.edu/10.1137/21M1463598>, 4550677.