

Key indicators of Arctic climate change: 1971–2017

Box, Jason Eric; Colgan, William T; Røjle Christensen, Torben ; Schmidt, Niels Martin ; Lund, Magnus; Parmentier, Frans-Jan W; Brown, Ross; Bhatt, Uma S; Wouters, Bert; More Authors

DOI

[10.1088/1748-9326/aafc1b](https://doi.org/10.1088/1748-9326/aafc1b)

Publication date

2019

Document Version

Final published version

Published in

Environmental Research Letters

Citation (APA)

Box, J. E., Colgan, W. T., Røjle Christensen, T., Schmidt, N. M., Lund, M., Parmentier, F.-J. W., Brown, R., Bhatt, U. S., Wouters, B., & More Authors (2019). Key indicators of Arctic climate change: 1971–2017. *Environmental Research Letters*, 14(4), Article 045010. <https://doi.org/10.1088/1748-9326/aafc1b>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Environmental Research Letters



LETTER

Key indicators of Arctic climate change: 1971–2017

OPEN ACCESS

RECEIVED
12 October 2018

REVISED
21 December 2018

ACCEPTED FOR PUBLICATION
4 January 2019

PUBLISHED
8 April 2019

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Jason E Box¹, William T Colgan¹, Torben Røjle Christensen^{2,3}, Niels Martin Schmidt², Magnus Lund^{2,4}, Frans-Jan W Parmentier^{3,5}, Ross Brown⁶, Uma S Bhatt⁷, Eugénie S Euskirchen⁸, Vladimir E Romanovsky⁹, John E Walsh¹⁰, James E Overland¹¹, Muyin Wang^{11,12}, Robert W Corell^{13,14,15}, Walter N Meier¹⁶, Bert Wouters^{17,18}, Sebastian Mernild^{19,20,21}, Johanna Mård²², Janet Pawlak²³ and Morten Skovgård Olsen²⁴

¹ Geological Survey of Denmark and Greenland (GEUS), Copenhagen, Denmark

² Department of Bioscience, Arctic Research Centre, Aarhus University, Denmark

³ Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden

⁴ Department of Soil Quality and Climate Change, Norwegian Institute of Bioeconomy Research (Nibio), Ås, Norway

⁵ Department of Geosciences, University of Oslo, Oslo, Norway

⁶ Climate Research Division, Environment and Climate Change Canada

⁷ Department of Atmospheric Sciences, Geophysical Institute, University of Alaska Fairbanks

⁸ Institute of Arctic Biology, University of Alaska Fairbanks

⁹ Geophysical Institute, University of Alaska Fairbanks, United States of America

¹⁰ International Arctic Research Center, University of Alaska

¹¹ NOAA/Pacific Marine Environmental Laboratory, Seattle WA, United States of America

¹² University of Washington/Joint Institute for the Study of the Atmosphere and Ocean, Seattle WA, United States of America

¹³ University of Miami, Miami, United States of America

¹⁴ University of the Arctic, Tromsø, Norway

¹⁵ Global Environment and Technology Foundation, Arlington, United States of America

¹⁶ National Snow and Ice Data Center, University of Colorado, Boulder, CO, United States of America

¹⁷ Institute for Marine and Atmospheric Research, Utrecht University, The Netherlands

¹⁸ Delft University of Technology, The Netherlands

¹⁹ Nansen Environmental and Remote Sensing Center, Bergen, Norway

²⁰ Department of Environmental Sciences, Western Norway University of Applied Sciences, Sogndal, Norway

²¹ Direction of Antarctic and Sub-Antarctic Programs, Universidad de Magallanes, Punta Arenas, Chile

²² Department of Earth Sciences, Uppsala University, Sweden

²³ Arctic Monitoring and Assessment Program (AMAP) secretariat

²⁴ Danish Ministry of Energy, Efficiency and Climate, Copenhagen, Denmark

E-mail: jeb@geus.dk

Keywords: Arctic climate change, observational records, AMAP

Abstract

Key observational indicators of climate change in the Arctic, most spanning a 47 year period (1971–2017) demonstrate fundamental changes among nine key elements of the Arctic system. We find that, coherent with increasing air temperature, there is an intensification of the hydrological cycle, evident from increases in humidity, precipitation, river discharge, glacier equilibrium line altitude and land ice wastage.

Downward trends continue in sea ice thickness (and extent) and spring snow cover extent and duration, while near-surface permafrost continues to warm. Several of the climate indicators exhibit a significant statistical correlation with air temperature or precipitation, reinforcing the notion that increasing air temperatures and precipitation are drivers of major changes in various components of the Arctic system. To progress beyond a presentation of the Arctic physical climate changes, we find a correspondence between air temperature and biophysical indicators such as tundra biomass and identify numerous biophysical disruptions with cascading effects throughout the trophic levels. These include: increased delivery of organic matter and nutrients to Arctic near-coastal zones; condensed flowering and pollination plant species periods; timing mismatch between plant flowering and pollinators; increased plant vulnerability to insect disturbance; increased shrub biomass; increased ignition of wildfires; increased growing season CO₂ uptake, with counterbalancing increases in shoulder season and winter CO₂ emissions; increased carbon cycling, regulated by local hydrology and permafrost thaw; conversion between terrestrial and aquatic ecosystems; and shifting animal distribution and demographics. The Arctic

biophysical system is now clearly trending away from its 20th Century state and into an unprecedented state, with implications not only within but beyond the Arctic. The indicator time series of this study are freely downloadable at AMAP.no.

1. Introduction

Rising concentrations of greenhouse gases are driving widespread changes in global physical climate and its ecosystems (IPCC 2014a, 2014b). This article assembles nine diverse observational records that serve as key indicators of Arctic climate and ecosystem status. This review of physical changes is accompanied by a discussion of links with the Arctic biological systems. We present and discuss each indicator in turn and where possible, we discuss ecosystem impacts. A statistical evaluation of correlations between the indicators and various time series of pan-Arctic, Arctic regional or hemispheric surface air temperatures (or precipitation) is made in effort to identify, quantify and further illuminate potential interactions. Key findings are listed in the conclusion section, including a commentary on observational gaps with recommendations for future work.

2. Key indicators

While ‘indicator’ has been defined in various ways in the literature, this study will follow the definition of Kenney *et al* (2016) by regarding indicators as ‘reference tools that can be used to regularly update status, rates of change, or trends of a phenomenon using measured data, modeled data or an index’. We apply the notion of indicators to capture the state of the Arctic environment through observational data series that span various components of the Arctic system. Figure 1 illustrates nine key Arctic indicators, updated and expanded from the AMAP 2017 assessment. Each indicator is discussed in the following subsections and where considered appropriate, their inter-relations are further examined.

3. Methodology

3.1. Period of analysis

While homogeneous datasets for some variables pre-date 1971, such datasets for other indicator variables (e.g. sea ice, permafrost temperature, wildfire area) are not available until the 1970s. The 1971–2017 period used in this synthesis spans the decades prior to and during the Arctic’s systemwide changes starting in the mid-1980s (Overland *et al* 2004) and unprecedented extremes that have occurred since the mid-1990s (e.g. Overland *et al* 2018).

3.2. Temperature and precipitation data

Here, near surface air temperature data time series are taken from the NCEP/NCAR Re-analysis (updated from Kalnay *et al* 1996). Justification for the use of these data are prompt updates and consistent performance versus other reanalysis products (Overland and Wang 2016). The data are not separated between land and ocean because our aim is to include changes both over the land and above the ocean for an integrated ‘indicator’, which is associated with other indicators we are studying in this study (e.g. sea ice (ocean), permafrost (land), snow cover (land)). Our coverage is pan-Arctic, regional and Northern Hemispheric. Nevertheless, the relative contribution of the land versus ocean stations to e.g. air temperature, is not the same, and this sampling bias is a possible source of uncertainty.

3.3. Seasonal and regional variable definitions

We define temperature and precipitation variables for both seasonal or annual and pan-Arctic or regional areal averages. By ‘warm season’ we refer to the June through September period that often includes above freezing air and surface temperatures. By ‘cold season’ we refer to the October through May period that is characterized by below-freezing temperatures over much of the Arctic. ‘Arctic’ here refers to the area north of 60 degrees latitude. The Northern Hemisphere is abbreviated as ‘N. Hem.’ For example, when referring to Arctic air temperature we use the following variable: $T_{\text{Arctic warm season}}$. Regional polygons cover land ice areas and hence we refer to e.g. $T_{\text{N. Hem. Warm Season}}$. For a snow cover statistical evaluation, we choose the May to June (MJ) period since parts of the Arctic can be snow free by June.

3.4. Regression methodology, definition of trend units

Potential relationships to air temperatures are quantified by regressions between Northern Hemisphere (or Arctic) seasonal or annual temperatures and other climate parameters against the assembled collection of Arctic climate indicators for the 1971–2017 period. The regressions that reveal signals of interconnection with high confidence ($1 - p > 0.9$) in correlations (R) are emphasized using bold text in table 1.

When assessing confidence in regressions between two time series, e.g. air temperature and precipitation, one year lagged temporal correlations (r_{lag1}) are evaluated to control for serial autocorrelation, in which we compute effective degrees of freedom not as $n_{\text{time-series_years-2}}$ but as ($n_{\text{effective}}$):




Figure 1. (Left) Arctic climate observational indicator records. Multi-year running averages are illustrated using Gaussian smoothing lines, (right) data sources are described.

$$n_{\text{effective}} = -n_{\text{timeseries years}} \log(r_{\text{lag 1}}).$$

The lowest $n_{\text{effective}}$ is then used to determine the p -value of the correlation. The effect of controlling for serial autocorrelation is to give more realistic (lower) ‘confidence’ ($1 - p$) values.

While a correlation is indicative of a possible relationship, it does not distinguish the contributions of the trends and the interannual variations superimposed on the trend. Physically meaningful relationships should be manifested in interannual variations as well as in corresponding trends. Since trends are apparent in figure 1, we focus our correlation analysis on the interannual timescale. To do so, we temporally

detrend the data prior to computing cross-correlations. Further, here, we make no lagged correlation analysis.

Our use of the term ‘change’ refers to the magnitude of linear trends assessed by standard least squares regression (Chatterjee and Hadi 2006), that is, the regression slope multiplied by the duration of data in years.

3.5. Smoothing of series

To illustrate longer-term variations than that of single years and recognizing that some temporal autocorrelation can occur, figure 1 presents normal distribution

Table 1. Linear trends and temporally detrended correlation of figure 1 Arctic climate indicators versus air temperature from NCEP/NCAR re-analysis.

| Indicator | Units | Trend per decade | Change during period | 1 - p | Versus variable | Corr. coef. | 1 - p | Earliest year | Latest year |
|--|---------------------------------|------------------|----------------------|------------------|---------------------------|---------------|------------------|---------------|-------------|
| $T_{Arctic Annual}$ | °C | 0.6 | 2.7 | >0.999 | $T_{N. Hem. Annual}$ | 0.660 | >0.999 | 1971* | 2017 |
| $T_{Arctic Warm Season}$ | °C | 0.4 | 1.8 | >0.999 | $T_{N. Hem. Warm Season}$ | 0.295 | 0.950 | 1971* | 2017 |
| $T_{Arctic Cold Season}$ | °C | 0.7 | 3.1 | >0.999 | $T_{N. Hem. Cold Season}$ | 0.660 | >0.999 | 1971* | 2017 |
| $P_{Arctic Annual}$ | % | 1.3 | 6.2 | >0.999 | $T_{Arctic Annual}$ | 0.503 | 0.998 | 1971* | 2017 |
| $P_{Arctic Cold Season}$ | % | 1.4 | 6.8 | 0.994 | $T_{N. Hem. Annual}$ | 0.303 | 0.948 | 1971* | 2017 |
| | | | | | $T_{Arctic Cold Season}$ | 0.447 | 0.996 | 1971* | 2017 |
| | | | | | $T_{N. Hem. Cold Season}$ | 0.447 | 0.996 | 1971* | 2017 |
| $P_{Arctic Warm Season}$ | % | 1.0 | 4.7 | 0.935 | $T_{Arctic Warm Season}$ | 0.061 | 0.314 | 1971* | 2017 |
| | | | | | $T_{N. Hem. Warm Season}$ | 0.061 | 0.314 | 1971* | 2017 |
| | | | | | $T_{Arctic Annual}$ | -0.024 | 0.122 | 1971* | 2015 |
| Ob river annual | % | -0.6 | -2.6 | 0.255 | $T_{N. Hem. Annual}$ | 0.162 | 0.695 | 1971* | 2015 |
| Pechora River annual | % | 1.7 | 5.8 | 0.564 | $T_{Arctic Annual}$ | 0.191 | 0.704 | 1981 | 2014 |
| Severnaya Dvina River | % | 5.9 | 25.9 | 0.956 | $T_{N. Hem. Annual}$ | 0.271 | 0.864 | 1981 | 2014 |
| | | | | | $T_{Arctic Annual}$ | 0.035 | 0.175 | 1971* | 2014 |
| Yenisei River | % | 1.6 | 7.0 | 0.844 | $T_{N. Hem. Annual}$ | -0.036 | 0.180 | 1971* | 2014 |
| | | | | | $T_{Arctic Annual}$ | 0.065 | 0.326 | 1971* | 2015 |
| Lena River | % | 3.3 | 13.6 | 0.945 | $T_{N. Hem. Annual}$ | 0.035 | 0.179 | 1971* | 2015 |
| | | | | | $T_{Arctic Annual}$ | 0.114 | 0.514 | 1971* | 2011 |
| Kolyma River | % | 7.4 | 33.3 | 0.987 | $T_{N. Hem. Annual}$ | 0.074 | 0.348 | 1971* | 2011 |
| | | | | | $T_{Arctic Annual}$ | -0.001 | 0.004 | 1971* | 2015 |
| Yukon River | % | 3.1 | 13.0 | 0.989 | $T_{N. Hem. Annual}$ | -0.133 | 0.604 | 1971* | 2015 |
| | | | | | $T_{Arctic Annual}$ | -0.133 | 0.604 | 1971* | 2015 |
| Mackenzie River | % | 1.1 | 4.8 | 0.575 | $T_{Arctic Annual}$ | 0.090 | 0.394 | 1975 | 2016 |
| | | | | | $T_{N. Hem. Annual}$ | 0.048 | 0.213 | 1975 | 2016 |
| Eurasian Arctic rivers composite | km ³ y ⁻¹ | 18.7 | 56.1 | 0.996 | $T_{Arctic Annual}$ | -0.019 | 0.093 | 1973 | 2015 |
| | | | | | $T_{N. Hem. Annual}$ | -0.135 | 0.601 | 1973 | 2015 |
| North American Arctic rivers composite | km ³ y ⁻¹ | 5.9 | 23.6 | 0.972 | $T_{Arctic Annual}$ | 0.152 | 0.555 | 1981 | 2011 |
| | | | | | $T_{N. Hem. Annual}$ | 0.075 | 0.294 | 1981 | 2011 |
| Tundra Max NDVI | st. dev. | 0.1 | 0.05 | >0.999 | $T_{Arctic Annual}$ | -0.266 | 0.862 | 1975 | 2015 |
| | | | | | $T_{N. Hem. Annual}$ | -0.048 | 0.213 | 1975 | 2015 |
| Tundra time-integrated NDVI | st. dev. | 0.1 | 0.23 | 0.999 | $T_{Arctic Warm Season}$ | 0.406 | 0.982 | 1982 | 2017 |
| Alaska burned area | 1e6 Ha | 0.1 | 0.5 | 0.863 | $T_{Arctic Warm Season}$ | 0.555 | 0.999 | 1982 | 2017 |
| Canada burned area | 1e6 Ha | -0.0 | -0.0 | 0.077 | $T_{Arctic Warm Season}$ | -0.054 | 0.251 | 1980 | 2017 |
| September Arctic Sea Ice extent | 1e6 sq. km | -0.8 | -3.3 | >0.999 | $T_{Arctic Warm Season}$ | 0.010 | 0.044 | 1980 | 2017 |
| Spring Snow Covered Area | days | -3.4 | -15.5 | >0.999 | $T_{Arctic Warm Season}$ | -0.623 | >0.999 | 1979 | 2017 |
| | | | | | $T_{Arctic MJ}$ | -0.464 | 0.998 | 1972 | 2017 |

Table 1. (Continued.)

| Indicator | Units | Trend per decade | Change during period | $1 - p$ | Versus variable | Corr. coef. | $1 - p$ | Earliest year | Latest year |
|--------------------------|----------|------------------|----------------------|------------------|---------------------------------|---------------|--------------|---------------|-------------|
| Greenland Mass Balance | st. dev. | -0.7 | -3.2 | >0.999 | $T_{\text{Arctic Warm Season}}$ | -0.472 | 0.999 | 1971* | 2017 |
| Canada Mass Balance | st. dev. | -0.3 | -1.5 | >0.999 | $T_{\text{Arctic Warm Season}}$ | -0.332 | 0.974 | 1971* | 2017 |
| Alaska Mass Balance | st. dev. | -0.3 | -1.6 | 0.999 | $T_{\text{Arctic Warm Season}}$ | 0.115 | 0.551 | 1971* | 2017 |
| Scandinavia Mass Balance | st. dev. | -0.2 | -0.8 | 0.937 | $T_{\text{Arctic Warm Season}}$ | -0.184 | 0.776 | 1971* | 2017 |
| Svalbard Mass Balance | st. dev. | -0.2 | -0.7 | 0.879 | $T_{\text{Arctic Warm Season}}$ | -0.339 | 0.977 | 1971* | 2017 |

Note. Bold values highlight high confidence ($1 - p > 0.9$) correlations. An asterisk beside the year indicates data that begin before then but are not analyzed here.

weighted running average values, i.e. smoothing. The chosen envelope is ± 5 years and the Gaussian width has 1.5 standard deviations per 11-year sample. Within 4 years of the time series beginning or end, the tail on the Gaussian sample is truncated by one in each year toward the end of the series until the sample size is 6 years. While we present smoothed data, in all cases, all presented statistics are computed only from the unsmoothed raw data. In order to detrend the data, we subtract the linear trend resulting from temporal regression.

4. Arctic climate indicators

4.1. Air temperature

Arctic air temperature change (ΔT) from 1971 to 2017 measured by the regression slope (multiplied by 47 years) indicate warming by: 2.7 °C at the annual scale ($\Delta T_{\text{Arctic Annual}}$); 3.1 °C in the cold season (October–May) ($\Delta T_{\text{Arctic Cold Season}}$) and 1.8 °C in the warm season (June–September) ($\Delta T_{\text{Arctic Warm Season}}$) (table 1, figure 1(a)). A number of processes contribute to amplified Arctic temperature variations as compared to global temperatures (Pithan and Mauritsen 2014). As a metric of Arctic Amplification (AA), comparing the change in Arctic temperatures with those from the Northern Hemisphere, we find $AA_{\text{Annual}} = \Delta T_{\text{Arctic Annual}} / \Delta T_{\text{N. Hem. Annual}} = 2.4$, $AA_{\text{Cold Season}} = \Delta T_{\text{Arctic Cold Season}} / \Delta T_{\text{N. Hem. Cold Season}} = 2.8$, and $AA_{\text{Warm Season}} = \Delta T_{\text{Arctic Warm Season}} / \Delta T_{\text{N. Hem. Warm Season}} = 1.7$. Thus, similar to the observed increase in temperature changes from 1971 to 2017, AA is greatest in the cold season and smallest in the warm season (June through September).

Later freeze up of sea ice (e.g. Markus *et al* 2009) and advection of moisture into the Arctic (Zhang *et al* 2013, Neff *et al* 2014) are key contributors to the rise in cold season air temperatures, producing maximum Arctic warming in autumn and winter. For the 1959–2008 period, Bekryaev *et al* (2010) conclude annual AA to be 1.52 for 1959–2008. The values of AA depend on the region considered, e.g. Arctic Ocean else land, distance from the coast (Bekryaev *et al* (2010) and altitude (Hernandez-Henriquez *et al* 2015). See Serreze and Barry 2011) for further review. Evaluating AA using paleo data, Miller *et al* (2010) concluded a higher AA, between 3 and 4. However, during the last glacial maximum, AA was negative due to a stronger northern latitude insolation increase as compared to the present Anthropogenic warming driven by excessive greenhouse gas concentrations.

4.2. Permafrost and carbon cycling

New record-high annual average temperatures in the upper 10–20 m of the ground have been observed at many permafrost observatories with the greatest temperature increases (>2 °C) occurring in the colder permafrost of the northern Arctic (Romanovsky *et al*

2017). Here, at 20 m depth for three North Slope of Alaska sites (West Dock, Deadhorse and Franklin Bluffs) we find a 2.5 °C permafrost temperature increase in the past 47 years (figure 1(b)). In northern Alaska, the active layer freeze-up date in the 2010s (mid-December) was almost two months later than in the mid-1980s (mid-October). In Zackenberg, north-east Greenland, maximum thaw depths increased by c. 1.6 cm yr⁻¹ between 1997 and 2010 (Lund *et al* 2014). Reduced permafrost area contributes to amplified warming because of a reduced ground latent heat sink (Lund *et al* 2014, Parazoo *et al* 2018).

The impact of thawing permafrost on ecosystem processes is dependent on permafrost type and local hydrology. In areas with discontinuous permafrost, thawing can lead to permafrost collapse with major implications for hydrology, vegetation composition and biogeochemical cycling (Johansson *et al* 2006). Bring *et al* (2016) suggest that permafrost thaw may increase hydrological connectivity between ground-water and surface water systems and change water storage in lakes and soils, which will influence exchange of moisture with the atmosphere. Jorgenson *et al* (2001) document permafrost degradation causing ecosystem shifts from birch forests to fens and bogs. In upland tundra areas with continuous permafrost, increasing active layer depths may on the other hand lead to soil drying (Liljedahl *et al* 2016), limiting vegetation growth.

As a response to increased air and ground temperature, there are now clear signs of permafrost thaw (Nicolosky *et al* 2017, Romanovsky *et al* 2017). In combination with warming-induced impacts on Arctic tundra vegetation, these landscape-scale structural changes will affect tundra-atmosphere interactions including both biogeophysical and biogeochemical feedback effects on the climate system (Lund 2018).

Jeong *et al* (2018) find accelerating rates of carbon cycling revealed by 42 years of atmospheric CO₂ measurements from Barrow, Alaska (71.29 N, 156.79 W). They conclude that: ‘Temperature dependencies of respiration and carbon uptake suggest that increases in cold season Arctic labile carbon release will likely continue to exceed increases in net growing season carbon uptake under continued warming trends’. See also section 4.4. Tundra greening and terrestrial ecosystems, below.

For the Canadian boreal forest, Price *et al* (2013) document how ‘approximately 40% of the forested area is underlain by permafrost, some of which is already degrading irreversibly, triggering a process of forest decline’. Through modeling, Schuur *et al* (2015) suggest that Arctic climate warming will cause an increasingly large net upward flux of terrestrial carbon to the atmosphere via microbial release of carbon from decomposition of accumulated surface biomass. Observational data from Zackenberg, NE Greenland, combined with ecosystem modeling for the period 2000–2014 also shows trends towards increased

overall carbon cycling but of a variable nature differing between time periods 2000–2008 and 2008–2014 (Zhang *et al* 2018). Long-term observational records are needed to verify any possible consistent trends in possible Arctic tundra carbon emissions as the potential releases are hypothetically an extremely important feedback given that it would likely amplify future climate warming. Hugelius *et al* (2014) estimate that Arctic soils contain ~50% of the world's global soil carbon and hence the potential release is enormous.

While Arctic seafloor methane (CH₄) release is observed (Shakova *et al* 2013, Andreassen *et al* 2017), there is no conclusive proof that hydrate-derived CH₄ is reaching the atmosphere today (Ruppel and Kessler 2017). Most of the CH₄ is oxidized or dissolved into the sediments or water column before reaching the atmosphere, especially in deeper waters (>50 m) (Parmentier *et al* 2017). Nonetheless, the idea of warming-triggered carbon release is hypothetically an extremely important feedback given that it would likely amplify future climate warming. This effect was recently quantified for CH₄ only to potentially cause a more than 20% increase in the CH₄ radiative forcing on top of anthropogenic 'business as usual' scenario. However, it is also shown that with serious mitigation of anthropogenic emissions or a 'maximum feasible reduction' scenario the effect of even extreme natural arctic CH₄ emission increase will be neutralized and even still maintain a lower radiative forcing by 2100 than a business as usual scenario will lead to (Christensen *et al* 2019).

Recent changes in biogeophysical energy exchange and transport within the Arctic, and between this region and the rest of the globe, now exceed even extreme projections. There is now clear evidence for both the marine and terrestrial Arctic environments that winter is not, as has previously been assumed, a dormant time for ecosystem processes (Mastepanov *et al* 2008, Christensen, 2014, Pirk *et al* 2016, Commane *et al* 2017). The winter includes carbon exchange through sea ice (Parmentier *et al* 2013). Terrestrial carbon exchange is complicated by the interaction of thawing permafrost, intensified hydrological cycle, vegetation change, and coupling between the land and ocean.

There is now mounting evidence for increasing gross primary production and ecosystem respiration with warming, however, the net effect on land-atmosphere CO₂ exchange remains unclear (Lund *et al* 2010, Lopez-Blanco *et al* 2017). The sea ice decline associated with late-summer-focused warming impacts terrestrial processes and ecosystems and greenhouse gas exchange (Parmentier *et al* 2013, Post *et al* 2013). The greening of the Arctic is expected to result in stronger growing season carbon uptake as well as lower albedo and higher turbulent heat fluxes (Chapin *et al* 2005, Lund 2018). Conversely, thawing permafrost mobilizes carbon through both vertical (Schuur *et al* 2015) and lateral pathways (Spencer *et al* 2015).

The CO₂:CH₄ emission ratio from thawing permafrost soils is dependent on soil moisture conditions (Schadel *et al* 2016). While higher temperatures promote CH₄ production within Arctic soils, the net flow into the atmosphere is constrained by the water table depth. Whether the Arctic surface will become wetter or drier may thus determine the net atmospheric CH₄ exchange (Watts *et al* 2014). Under climate change, trends in the net carbon flux may thus be damped (Parmentier *et al* 2011, Lund *et al* 2012) and possibly offset by increases in early winter respiration when plants have senesced (Commane *et al* 2017).

4.3. Changes to arctic hydroclimatology

4.3.1. Arctic humidification

Available observations from land and coastal stations indicate a humidity increase at the Arctic surface (Hartmann *et al* 2013, Vihma *et al* 2016) and in the mid troposphere (Serreze *et al* 2012). The humidification is in part related to increased advection of moist air from mid-latitudes (Zhang *et al* 2013) and longer sea ice-free seasons (Markus *et al* 2009, Serreze *et al* 2012). Walsh *et al* (2011) find increases in cloudiness over the Arctic, especially in low clouds during the warm season. The higher humidity increases downward longwave radiation (Zhang *et al* 2001), contributing to amplification of warming (Pithan and Mauritsen 2014).

4.3.2. Precipitation increase

While there is considerable uncertainty in precipitation trends over the Arctic, the available observations and reanalysis datasets (Rawlins *et al* 2010, Rapaic *et al* 2015) suggest increases of 1.5%–2.0% per decade in annual precipitation which is consistent with the estimated temperature sensitivity of Arctic precipitation of 4.5% per K (Bintanja and Selten 2014).

Here, according to NCEP/NCAR Reanalysis, the increase in annual total precipitation for the area north of 50 deg. N latitude 1971–2017 (47 years) is strongest during the cold season (October through May), increasing from 1971 to 2017 by 6.8% about an average rate of 225 mm during the eight-month period with high confidence ($1 - p = 0.994$). The increase during the June through September warm season is less; 4.7% about an average rate of 168 mm during the four-month period ($1 - p = 0.935$). The 1971–2017 period of Arctic precipitation exhibits inter-decadal fluctuations with a prominent increase from the mid 1980s to the late 2000s (figure 1(c)). Annually, the increase is 6.2% ($1 - p > 0.999$) about an average of rate of 393 mm per year.

Consistent with precipitation enhancement from water vapour feedback theory (e.g. Trenberth 2011), Box *et al* (2013) find a +6.8% °C⁻¹ increase in Greenland snow accumulation. Here, regression of annual NCEP/NCAR reanalysis precipitation for the Arctic region (north of 50° latitude) for the 1971–2017 (47 year) period with Northern Hemispheric air

temperatures suggest a $+7.5\% \text{ } ^\circ\text{C}^{-1}$ sensitivity ($R = 0.276$, $1 - p > 0.937$)²⁵. Seasonally, the precipitation sensitivity is $7.1\% \text{ } ^\circ\text{C}^{-1}$ ($R = 0.191$, $1 - p = 0.795$) for the Arctic warm season and $6.5\% \text{ } ^\circ\text{C}^{-1}$ ($R = 0.203$, $1 - p = 0.823$) for the cold season. When using Arctic temperatures (instead of the Hemispheric temperatures), the precipitation sensitivity values range from 3.3% to 3.7%, roughly a factor of two lower, presumably because the amplitude of Arctic temperature variability is roughly $2\times$ higher than hemispheric air temperature. The associated correlations: 0.270 ($1 - p = 0.930$) in the warm season, $R = 0.447$ ($1 - p = 0.998$) in the cold season and $R = 0.510$ ($1 - p > 0.999$) annually, suggest that interannual variations in air temperature is not the only process controlling precipitation.

Increased precipitation does not necessarily mean that the Arctic surface will become wetter, since increased temperature tends to increase evapotranspiration (Zhang *et al* 2009). For example, drying conditions result in areas where changes in evapotranspiration exceed precipitation inputs. Increased drainage following permafrost thaw may also lead to drier conditions (Liljedahl *et al* 2016), and reductions in water availability will limit vegetation growth and CO_2 uptake.

4.3.3. Rainfall increase

Decreasing snowfall at the expense of increasing rainfall is observed around the Greenland ice sheet margin (Doyle *et al* 2015) and in regions with warmer winter climates such as Scandinavia and the Baltic Sea basin (Rasmus *et al* 2015, Irannezhad *et al* 2016). Increasing snowfall is documented in colder regions such as northern Canada and Siberia (Kononova, 2012, Vincent *et al* 2015) and the lower elevations of the Greenland ice sheet (Box *et al* 2013, Hawley *et al* 2014, Wong *et al* 2015).

4.3.4. Soil moisture

Spatial variability in soil moisture may be an important driver of local-scale plant composition (Nabe-Nielsen *et al* 2017). On a larger scale, the spatial variability in soil moisture may explain the heterogeneous pattern of vegetation growth as deduced from remotely-sensed vegetation greenness indices (Bhatt *et al* 2017). Changes in precipitation patterns (e.g. shifts from snow to rain) will impact animal distribution and demographics both directly (e.g. Schmidt *et al* 2015, Kankaanpaa *et al* 2018) and indirectly through changes in plant composition and productivity. Increased winter snow fall will accelerate permafrost warming from increased insulation (Zhang 2005). Increasing cloudiness decreases tundra ecosystem photosynthesis and, contrary to the effect over snow- and ice-covered surfaces, it reduces surface energy availability (Lund

et al 2017). Any summer drying may be outweighed by enhanced winter precipitation (Serreze *et al* 2002). Further, changes in evaporation only exceed those in precipitation in a limited area of the Arctic oceanic domain and not over land areas (Jakobson and Vihma 2010).

4.3.5. Arctic river discharge increase

An increase in the discharge of major rivers terminating in the Arctic is well documented (e.g. Peterson *et al* 2002, Serreze *et al* 2006, Rawlins *et al* 2010, Haine *et al* 2015, Holmes *et al* 2015, Vihma *et al* 2016), with Eurasian rivers showing the greatest increase. Here, we assess Arctic river discharge using Global Runoff Data Centre (GRDC) data, providing 91% complete temporal coverage of six Eurasian rivers (Ob, Pechora, Severnaya Dvina, Yenisei, Lena, and Kolyma) during 1981–2011 and 86% complete coverage from the two major North American Arctic rivers (Mackenzie and Yukon) during 1975–2015 (figure 1(d)). By volume, the six-Eurasian river discharge is 1.8 times the average of the assessed two-North American river discharge. The combined river basin area cover 70% of the pan-Arctic drainage area (Holmes *et al* 2015).

For the limited set of cases when all rivers are reporting data, we find the average discharge increasing in Eurasian rivers by $56.1 \text{ km}^3 \text{ yr}^{-1}$ or . The North American river discharge increased by $23.6 \text{ km}^3 \text{ yr}^{-1}$ over the $1.32\times$ longer 1975–2015 period (table 1). While for a different period, the Eurasian discharge increase about a six-river 1981–2011 average of $467 \text{ km}^3 \text{ yr}^{-1}$ is $+12\%$, the North American Arctic river discharge about a 1975–2015 average of $253 \text{ km}^3 \text{ yr}^{-1}$ is lower; $+9\%$. We find no high confidence correlations of individual nor composite river discharge with Arctic nor hemispheric temperatures (table 1).

4.3.6. Arctic sedimentation increase

Increased delivery of organic matter and nutrients is evident in Arctic near-coastal zones (Bring *et al* 2016). Increases in Greenland ice sheet meltwater runoff during the 20th Century are linked to increased sedimentation rates (Bendixen *et al* 2017). Hawkings *et al* (2016) estimated that the Greenland ice sheet contributes about 15% of total bioavailable phosphorus input to the Arctic oceans ($\sim 11 \text{ Gt yr}^{-1}$) and dominates the total phosphorus input (408 Gt yr^{-1}), which is more than three times that estimated from Arctic rivers (126 Gt yr^{-1}).

4.3.7. Arctic ocean freshening

Arctic Ocean freshening is being driven by increases in Arctic precipitation and river discharge (Vihma *et al* 2016), with enhanced oceanic heat inflows from both the North Atlantic and the North Pacific playing a role in the retreat of sea ice in the Arctic Ocean. Increased ocean heat storage in newly sea-ice-free ocean areas

²⁵ Both temperature and precipitation time series are temporally detrended to avoid spurious correlation.

has been confirmed from recent shipboard observations (Walsh *et al* 2011).

4.4. Tundra greening and terrestrial ecosystems

Arctic greening (overall increases in vegetation biomass as deduced from satellite observations of land surface reflectance via NDVI, the normalized difference vegetation index) has been observed across tundra ecosystems over the past 30 years (e.g. Bhatt *et al* 2017) (figure 1(e)). Since Arctic tundra vegetation is temperature-limited, summers with above average summer warmth correspond to higher NDVI values and vice versa. Here, the increase of Arctic tundra average and maximum NDVI both correlate with high confidence with $T_{\text{Arctic Warm Season}}$ (table 1). The NDVI covariability with air $T_{\text{Arctic Warm Season}}$ is most likely related to greater amounts of photosynthetically active radiation during warmer-than-normal summers. Further, Martin *et al* (2017) link shrub biomass with air temperature, soil moisture, herbivory, and snow dynamics. Declines in the NDVI, i.e. ‘browning’, may be related to water or nutrient limitation, permafrost degradation, and extreme winter events (Phoenix and Bjerke 2016, Bhatt *et al* 2017).

Tundra-atmosphere CO_2 exchange, as presented by observation-based modelling (Zhang *et al* 2018), indicates a trend towards increased tundra CO_2 sink functioning (more negative net ecosystem exchange) during 2000–2008, caused by a stronger increase in gross primary production compared with ecosystem respiration (Lund *et al* 2012). However, this trend reversed from 2008 to 2014. As discussed above, high-latitude CH_4 emissions from Arctic tundra ecosystems represent a potentially important biogeochemical climate feedback, and are related to changes in temperature, moisture, and shifts in vegetation composition (e.g. Olefeldt *et al* 2013). Long-term observations of CH_4 emissions at Arctic sites are still relatively rare, and in particular few studies include non-growing season CH_4 emissions, which may represent up to 50% of annual CH_4 emissions (Treat *et al* 2018). The rather stable interannual variation in ecosystem respiration, as indicated by CH_4 emissions is observed at Zackenberg, NE Greenland. However, when comparing with other sites where similar monitoring is taking place in West Greenland and on Svalbard, a clear relationship is found with an increasing annual CH_4 emission with growing degree days (figure 3; Pirk *et al* 2017). Differing local tundra CH_4 dynamics points towards the importance of comparable observations being made at multiple sites for an improvement of our understanding of the potential CH_4 tundra emission changes (Christensen 2014).

In terms of floral population dynamics, there is now strong evidence that the summer warming trend is causing an earlier and more condensed flowering period of key plant species in the interaction web, including pollination. A condensed flowering period

leaves a progressively shorter time-window for the pollinators with possible subsequent cascading effects through the ecosystem (Hoye *et al* 2013, Schmidt *et al* 2016).

4.5. Fire

Fire clearly causes dramatic short-term changes in vegetation and ecosystem function (Bret-Harte *et al* 2013). Drier conditions and an increase in maximum air temperatures contribute to increased fire risk (Jolly *et al* 2015). Price *et al* (2013), conclude that increases in the average North American area burned will be gradual, despite periodic extremes. The fire data analyzed here (figure 1(f)) do exhibit non-normal distributions, containing a relatively small number of severe years. Burned area does not exhibit any collinearity with $T_{\text{Arctic Warm Season}}$ in this analysis (table 1). Rather, the fire-climate relationship is related to sub-seasonal dry/warm episodes and to increasing lightning ignition that is shown to correlate with air temperature and precipitation (Veraverbeke *et al* 2017). That study finds an increase in lightning ignitions since 1975, and that the large 2014 and 2015 events (figure 1(f)) ‘coincided with a record number of lightning ignitions and exceptionally high levels of burning near the northern treeline. Indeed, lightning ignition explains the majority of the interannual variability in burned area’. Supportive of a climate driven fire relationship, for Alaska, Young *et al* (2017) find ‘summer temperature and annual moisture availability as the most influential controls of historical fire regimes’ and ‘a nonlinear increase in the probability of fire above an average July temperature’.

4.6. Disturbance

Physical disturbance events such as wildfire and abrupt permafrost thaw and insects are becoming more frequent and could accelerate biome shifts, including increasing tree density in taiga, expansion of tall shrubs and trees into tundra, and conversion between terrestrial and aquatic ecosystems. For example, shrubs and trees have been observed to increase in upland tundra ecosystems when permafrost thaw increases soil drainage. Price *et al* (2013) make the following synthesis ‘Maladaptation commonly occurs when climate becomes significantly different from that to which the local genotypes have adapted. The climatic effects may be direct (e.g. effects of increased temperature on respiration rates) or indirect (e.g. increased drought stress owing to decreased soil water availability resulting from increased evapotranspiration and (or) reduced precipitation). These climatic effects often render trees more susceptible to additional stressors and their interactions, including insect pests (Frey *et al* 2004, Hogg *et al* 2008, Morin *et al* 2009), disease (Kliejunas *et al* 2009), and fire (e.g. Bergeron and Leduc 1998, Volney and Hirsch 2005).’

4.7. Terrestrial snow cover decrease

Seasonal snow covers part of the Arctic for up to ten months each year. Through its unique physical properties of high reflectivity and low thermal conductivity, as well as its water storage effects, snow cover plays critical roles in energy and water exchanges, ice growth, hydrology, ground thermal regime, carbon cycling, and ecosystem services (Brown *et al* 2017). The start and end dates of snow cover, and hence its duration, are closely linked to air temperature with spring snow cover duration anomalies significantly correlated with May–June (MJ) Arctic air temperatures ($R = -0.464$, $1 - p = 0.998$) (table 1).

There is widespread evidence of a reduced snow cover duration in the Arctic; by two to four days per decade over the past 30–40 years (figure 1(g)). The largest downward trends are occurring at high latitudes and elevations, a pattern that is consistent with Arctic amplification of warming and enhanced albedo feedbacks (Hernandez-Henriquez *et al* 2015, Pepin *et al* 2015). Most of the decrease in snow cover duration results from earlier snow melt, but delayed snow onset is more important to the snow duration decline in e.g. eastern Canadian Arctic (Brown *et al* 2018). Climate change attribution studies have detected the influence of greenhouse gas induced climate warming in the observed decreases of spring snow cover (Najafi *et al* 2016) and snow water equivalent (Jeong *et al* 2017).

Arctic spring (May through June) snow cover extent on land has now decreased by more than 30% since 1971 (figure 1(g)). Trends in annual maximum snow accumulation are more uncertain but suggest a decreasing trend of pan-Arctic land areas in the amount of water stored in seasonal snow cover over the past ~20 years. There is evidence of increased ice layer development in snowpacks in some regions of the Arctic in response to more frequent winter thaw and rain-on-snow events (Langlois *et al* 2016).

Snow is a major driver for Arctic ecosystem functioning, affecting the surface energy balance, permafrost thaw, hydrology, plant phenology and greenhouse gas exchange. Longer snow-free periods will strongly affect tundra energy budgets, with increasing surface energy availability and higher turbulent heat fluxes to the atmosphere (Chapin *et al* 2005, Stiegler *et al* 2016). The timing of snow melt is key for both growing season CO₂ (Parmentier *et al* 2011, Lund *et al* 2012) and CH₄ emissions (Mastepanov *et al* 2013, Pirk *et al* 2016). Longer snow-free seasons will further extend the period of plant growth, enhancing CO₂ uptake, but at the same time respiration increases too. Changes in the net carbon balance may, therefore, not be as strong (Parmentier *et al* 2011, Lund *et al* 2012) and possibly offset by increases in early winter respiration when plants have senesced (Commane *et al* 2017).

Changes in snow cover can also have large impacts on ecosystems outside of the growing season. Snow cover is a good insulator and protects plants from extreme winter temperatures. Winter warm spells,

however, may remove this protective cover and cause plant damage (Phoenix and Bjerke 2016). Rain-on-snow events can lead to thick ground ice while a complete melt of snow cover exposes vegetation to a return to cold conditions. The damage caused by these extreme winter events can affect vegetation growth and carbon cycling in the following growing season (Parmentier *et al* 2018) and is linked with mass caribou mortality (Tyler 2010).

Snow cover sensitivities are complex and may include timing dependencies that create transient phenological and trophic mismatches from rapidly changing snow cover, e.g. Doiron *et al* (2015). Rapid advance in snowmelt timing can cause a timing mismatch between Arctic plant flowering and pollinating species, with cascading effects throughout the trophic levels (Hoye *et al* 2013, Schmidt *et al* 2016).

The relation between declining Arctic spring snow cover and lower latitude climate is unclear, as most of the available evidence suggest that potential linkages are more likely during the snow cover onset period in the fall (Cohen *et al* 2014). Observations of increasing Arctic snow cover in the fall period from the NOAA-CDR dataset (e.g. Cohen *et al* 2012) have been shown to be inconsistent with multiple lines of observational evidence and climate model simulations (Brown and Derksen, 2013, Mudryk *et al* 2017).

The loss of the perennial snow banks that buffer low flow periods in dry Arctic environments is evident (Woo and Young 2014). Traditional activities of northern residents such as hunting are sensitive to snow conditions (Bokhorst *et al* 2016). The Arctic-wide trend towards a shorter snow season is adversely impacting access to food sources with implications for health and disposable income (Furgal *et al* 2012).

4.7.1. Sea ice

The recent decade continues the unprecedented change in Arctic sea ice, in both the rates and magnitude of change in extent, area, thickness, spatial distribution, and most aspects of temporal and spatial variability (e.g. Overland and Wang 2013, Meier *et al* 2014, Comiso *et al* 2017). The Arctic has transformed from an environment dominated by thick multi-year sea ice to one dominated by thinner first-year sea ice (Tschudi *et al* 2016), with an earlier melt onset (Bliss *et al* 2017), later freeze-up (Markus *et al* 2009, Stroeve *et al* 2014), and longer open water period (Parkinson 2014, Stroeve *et al* 2016, Peng *et al* 2018, Wang *et al* 2018). Sea-ice extent (figure 1(h)), thickness and volume (Kwok and Cunningham 2015) are continuing their downward trends. The past six years have seen high variability, with record-low extent in summer 2012, low extents in 2015 through 2017, but relatively higher extent and thickness in 2013 and 2014 (though still much lower than values in the 1980s and 1990s). Here, the highest correlation among the compared variables with $T_{\text{Arctic Warm Season}}$ is for September sea

ice extent (table 1), strongly suggesting that further sea ice loss is to be expected from a warming Arctic.

The Pacific sector of the Arctic Ocean, and Hudson Bay and Baffin Bay, are showing increased open water from August through December. This autumn extension of the open-water period (Stroeve *et al* 2016; Peng *et al* 2018) is dominated by the ice albedo feedback (Perovich and Polashenski 2012; Stroeve *et al* 2014) and heat capture in the upper ocean (Serreze and Barry 2011; Lien *et al* 2017). The Atlantic sector shows increased open water in winter. The open-water period is dominated by horizontal ocean heat fluxes. Understanding the evolution of snow on sea ice remains a significant challenge and basin-wide estimates of snow are rare (Webster *et al* 2014). The increasing presence of very young ice types results in high salinity ice covers (e.g. frost flowers) that are reactive in chemical exchanges with the atmosphere and ocean.

Along with Arctic sea ice decline, there is emerging evidence for a loss of biodiversity in sea-ice habitats (Meier *et al* 2014), including that of the polar bear (Amstrup *et al* 2010). Open-water species, here whales (cetaceans), may see new habitats opening. According to Meier *et al* (2014), ‘Killer whales (*Orcinus orca*) sightings have increased markedly in the eastern Canadian Arctic over a period of decades; associated with changing ice patterns (Higdon *et al* 2012), blue whales (*Balaenoptera musculus*) have been acoustically recorded in Fram Strait over an extended seasonal period, covering June through until October (Moore *et al* 2011), and North Atlantic right whales (*Eubalaena glacialis*) appear to have spread north as southeast Greenland (Mellinger *et al* 2011). Similarly, in the Pacific regions, fin whales (*Balaenoptera physalus*) are present in the Bering Sea almost year-round now (Stafford *et al* 2010) and gray whales (*Eschrichtius robustus*) are spending increasingly long periods in Arctic waters, delaying the southward migrations [Moore 2008]. White whales (*Delphinapterus leucas*) in West Greenland have shifted their summer distribution westward as sea ice has declined [Heidi-Jørgensen *et al* 2010]. Sea surface temperature changes (intimately linked to sea ice formation) have also been implicated in changing phonologies of movements in this species in the Canadian Arctic [Bailleaul *et al* 2012]. Bowhead whale (*B. mysticetus*) distribution has also shifted recently, with significant population level implications; Alaskan and Greenlandic populations, which have been separated by ice in the past, are now overlapping spatially in the Northwest Passage [Heidi-Jørgensen *et al* 2012].

4.7.2. Land ice

Observational records of Arctic land ice mass balance indicate stability or growth from 1971 until the mid 1980s, followed by a strong increase in ice loss. In the 47 year period (1971–2017), the Arctic was the largest global source of land ice to sea-level rise, accounting for 48% of the contribution during 2003–2010 (AMAP 2017) and 30% of the total sea-level rise since

1992 (Box *et al* 2018). After Greenland, the largest northern contributions are from Alaska, Arctic Canada and the Russian High Arctic. Glacier mass balance deficit increased in the Alaskan sector in the late 1980s followed by Arctic Canada then Greenland (figure 1(i)).

Persistent extremes in warm season atmospheric circulation are very influential for the observed mass balance changes. In figure 1(i), note for example periods of anti-correlation between Alaska and Arctic Canada mass balance, after 21st century, linked to persistent regional atmospheric circulation extremes (Box *et al* 2018). A shift to more negative Arctic Canada glacier mass balance occurred after 1986 (Gardner and Sharp 2007), linked to increased July air temperatures related to variations in the position and strength of the July circumpolar vortex. The years since 2013 have been a mix of extremes; Ahlstrøm *et al* (2017) suggest evidence for a regime shift in atmospheric circulation after 2006.

Increases in the post-2005 equilibrium line altitude by >250 m relative to the pre-2005 levels (Thomson and Copland 2017, Burgess 2017) coincide with enhanced warming of ice cap surfaces above 1400 m a.s.l. (Mortimer *et al* 2016). Densification of ice cap firn areas due to warming has reduced or eliminated the refreezing storage capacity of the many ice caps in this region, thus increasing their sensitivity to future warming (Noël *et al* 2018). Of the global glacier mass loss between 1991 and 2010, 70% has been attributed to anthropogenic climate change by Marzeion *et al* (2014).

Here, NCEP/NCAR reanalysis region-specific warm season (June through September) ($T_{\text{regional Warm Season}}$) and cold season (October through May) precipitation ($P_{\text{regional Cold Season}}$) is compared with Greenland, Alaskan, Canadian, Scandinavian and Svalbard land ice mass balance (table 2). First, glacier mass balance is closely varying with $T_{\text{regional Warm Season}}$. The connection is through the long demonstrated simple melting degree days relationship (e.g. Braithwaite 1995) but also connected with downward longwave irradiance (Ohmura 2001) and surface albedo reduction associated with increased melting (Hock 2003). Of the glacier mass balance regions compared, the Greenland ice sheet exhibits the strongest correlation with $T_{\text{regional Warm Season}}$, followed by Arctic Canada and Svalbard land ice mass balance. Scandinavian mass balance records, though more numerous and thus being expected to yield robust statistical sampling, do not correlate with $T_{\text{regional Warm Season}}$. Scandinavian mass balance variability has been more influenced by precipitation variability (Dowdeswell *et al* 1997). Norwegian glacier positive mass balance anomalies in the 1990s are attributed to North Atlantic Oscillation (NAO) extremes (Nesje *et al* 2000). The NAO is regarded as internal variability that is not well correlated with $T_{\text{regional Warm Season}}$. The lacking Alaskan sensitivity to $T_{\text{regional Warm Season}}$ is similar to the low correlation also found comparing with the Northern Hemisphere air temperature series.

Table 2. Regional land ice mass balance comparison with regional warm season temperature and regional cold season precipitation.

| Region | Correlation coefficient, mass balance versus $T_{\text{regional Warm Season}}$ | Correlation coefficient, mass balance versus $P_{\text{regional Cold Season}}$ | Multiple correlation coefficient, mass balance versus $T_{\text{regional Warm Season}}$ and $P_{\text{regional Cold Season}}$ |
|---------------|--|--|---|
| Greenland | −0.612 | −0.038 | 0.620 |
| Alaska | −0.715 | 0.281 | 0.744 |
| Arctic Canada | −0.760 | 0.087 | 0.760 |
| Scandinavia | −0.674 | 0.627 | 0.823 |
| Svalbard | −0.633 | 0.032 | 0.656 |

Including precipitation totals in the regression analysis suggests a dominance of warm season climate on mass balance, consistent with Dowdeswell *et al* (1997) and Østby *et al* (2017). Note how there is a consistent pattern of larger magnitude negative mass balance correlation with $T_{\text{regional Warm Season}}$ and a smaller magnitude positive mass balance correlation with $P_{\text{regional Cold Season}}$ (table 2).

Taking the reasonable assumption that $T_{\text{regional Warm Season}}$ is independent of $P_{\text{regional Cold Season}}$ yields multiple regression correlation coefficients that are larger than single regressions with $T_{\text{regional Warm Season}}$ or $P_{\text{regional Cold Season}}$ (table 2). Explained variance is highest for Scandinavia ($\text{Correlation}^2 = 0.674$) where precipitation correlates with mass balance nearly as highly as with $T_{\text{regional Warm Season}}$. Dowdeswell *et al* (1997) similarly found that Scandinavia had a stronger mass balance response from the relatively more variable precipitation rate for other Arctic glaciated regions.

Arctic Canada has no apparent precipitation sensitivity yet exhibits the strongest $T_{\text{regional Warm Season}}$ sensitivity ($\text{Correlation} = -0.760$), consistent with low precipitation rates (under 300 mm yr^{-1} , Cogley *et al* 1996, Dyurgerov 2002) based on the reanalysis product. Arctic Canada snow accumulation rates are similar to other High Arctic glacier regions. With few exceptions, using annual or warm season precipitation degrades the correlations, reinforcing the expectation that mass balance may be best represented by integrating cold season precipitation, i.e. the accumulation season part of the so-called ‘winter balance’.

4.8. Ecosystems

Long-term observational data to identify ecosystem trends in the Arctic are few, due to the remoteness of the region. However, in the past decade, newly available contributions through sustained long-term research have begun to enhance our ability to document ecological change in the Arctic. Some of these contributions are through research programs associated with Arctic observatories, including Zackenberg in Greenland (Schmidt *et al* 2017), northern Sweden at Abisko (Callaghan *et al* 2013), and the Alaskan Arctic near Toolik Lake (Hobbie and Klings 2014, Hobbie *et al* 2017). Other long-term ecological data are available through coordinated networks spanning multiple sites, such as the International Tundra

Experiment, that aims to evaluate the long-term effects of increases in temperature on plant growth, phenology, and community composition (Oberbauer *et al* 2013). Moreover, long-term Arctic vertebrate data have been compiled and routinely updated based on contributions from individuals and organizations to identify trends across 35% of the known Arctic vertebrates since 1970 (Barry and Helgason 2016). While these newly available contributions are essential for reaching a better understanding of long-term ecological Arctic change, new initiatives are also needed, particularly for data collected during the critical spring and fall shoulder seasons, as well as the winter period, to gain a better understanding of change over the full annual cycle (e.g. Bokhorst *et al* 2012; Blume-Werry *et al* 2016).

5. Summary and conclusions

5.1. Key messages

Arctic air temperature: Arctic annual average air temperatures 1971–2017 increased $2.7 \text{ }^\circ\text{C}$, at 2.4 times the rate of the Northern Hemisphere average. The $3.1 \text{ }^\circ\text{C}$ increase in the cold season (October–May) is the largest by season, 2.8 times the rate of the Northern Hemisphere cold season average. Arctic warm season (June through September) temperatures increased $1.8 \text{ }^\circ\text{C}$, 1.7 times the rate of Northern Hemisphere summer.

Alaskan permafrost: New record-high annual average temperatures in the upper 10–20 m of the ground have been observed at many permafrost observatories. At 20 m depth for three North Slope of Alaska sites (West Dock, Deadhorse and Frankliln Bluffs) we find a $2.5 \text{ }^\circ\text{C}$ permafrost temperature increase in the past 47 years.

Arctic hydroclimatology: Observations from land and coastal stations indicate widespread increases in humidity, low-level clouds, precipitation, rainfall (at the expense of snowfall), river discharge, sedimentation and delivery of organic matter to the Arctic ocean, freshening of the Arctic Ocean, and reductions in snow cover, all of which are controlling factors in Arctic terrestrial and probably marine ecosystems.

Snow cover: Arctic snow cover is responding to multiple environmental drivers and feedbacks (such as warming, increased moisture availability, changing

atmospheric circulation, changing vegetation, increased frequency of winter thaws, rain-on-snow events). There is widespread multi-dataset evidence of declining snow cover over the Arctic with the annual duration of snow on the ground shortening by 2 to 4 days per decade with the largest negative trends occurring at high latitudes and elevations consistent with AA of warming and enhanced albedo feedbacks.

Arctic Ocean sea ice: Sea ice extent and volume are continuing their downward trends. The past decade had record-low extent in summer 2012, and it is the lowest decade ever in satellite era beginning in the 1970s. These are unprecedented change in Arctic sea ice, in both the rates and magnitude of change in extent, area, thickness, and spatial distribution. Along with Arctic sea ice decline, there is emerging evidence for a loss of biodiversity in sea-ice habitats.

Arctic land ice: In the 47 year period (1971–2017), the Arctic was the largest global source of sea-level rise contribution, 48% of the global land ice contribution 2003–2010 and 30% of the total sea-level rise since 1992. Temperature effects are dominant in land ice mass balance; precipitation represents a source of either damping or amplifying feedbacks respectively via snow and rain.

Arctic region wildfires: Drier conditions and an increase in maximum air temperatures contribute to increased fire risk. Fire clearly causes dramatic short-term changes in vegetation and ecosystem function. The fire-climate relationship is related to increasing lightning ignition that is shown to correlate with air temperature and precipitation, thus linking Arctic warming with the likelihood for increased fire.

Tundra and terrestrial ecosystems: Arctic greening has been observed across tundra ecosystems over the past 30 years. The increase of Arctic tundra average and maximum NDVI both correlate with Arctic warm season air temperature with high confidence.

Carbon cycling: The changes in the global climate system are already affecting biogeophysical energy exchange and transport within the Arctic. The response of the carbon cycle in northern high latitude regions is influenced by terrestrial carbon exchange and by coupling between the land and ocean, which has worldwide consequences. Importantly, there are substantial organic matter stocks of carbon in the Arctic contained in permafrost and within the methane hydrates that exist beneath both subterranean and subsea permafrost of the Arctic, all of which can affect carbon cycling. Observational data indicate increased tundra ecosystem CO₂ uptake during the growing season. Further temperature increase will affect tundra CO₂ and CH₄ emissions, their ratio being dependent on local hydrology and permafrost thaw.

5.2. Closing remarks

Increasingly clear linkages are evident within and between multiple Arctic climate indicators, having

cascading effects, from condensed flowering and pollination plant species periods; timing mismatch between plant flowering and pollinators; increased plant vulnerability to insect disturbance; increased shrub biomass; increased ignition of wildfires; increased growing season CO₂ uptake, with counterbalancing increases in shoulder season and winter CO₂ emissions; increased carbon cycling, regulated by local hydrology and permafrost thaw; conversion between terrestrial and aquatic ecosystems; and shifting animal distribution and demographics.

The Arctic biophysical system is now clearly trending away from its previous state and into a period of unprecedented change, with implications not only within but also beyond the Arctic. These indicator-based observations also provide a foundation for the research that is needed to address the gaps in knowledge and to support a more integrated understanding of the Arctic region and its role in the global dynamics of the Earth's biogeophysical systems.

5.3. Recommendations for future work

Future work should be concerned with further unifying our understanding of physical and biological elements of the Arctic system.

In situ observations must be maintained, especially where verifying higher spatial coverage satellite observation, in data assimilation and for model verification studies. Further, *in situ* observations should be extended to include the critical winter period.

There is a need to quantify ecosystem impacts of changes and their relationships to physical drivers in the Arctic system.

Indicators that capture changes in extreme events (winds, extreme temperatures, intense precipitation events, droughts, fires) are needed to complement indicators based on mean values, especially in the context of impacts on humans and ecosystems.

Socioeconomic indicators are largely absent from this study, primarily because their development has lagged the compilation of physical and biological indicators.

Major gaps include: poor knowledge of Arctic precipitation; Arctic snow water equivalent; Arctic freshwater budget, lacking high resolution homogeneous reanalysis datasets; hydrological and biophysical processes in mountain regions; etc.

The period since SWIPA 2011 has seen important advances in snow science and greater understanding of the role and interactions of snow in Arctic soil-climate-vegetation systems. However, there are still fundamental knowledge gaps and scaling issues that need to be addressed to narrow uncertainties in observing, understanding, and predicting Arctic snow cover and snow-cover processes.










Critical areas for further work include: documenting and narrowing the uncertainties in snow observing systems over the Arctic (snow water equivalent in

particular); more realistic treatment of sub grid-scale processes and snow-vegetation interactions in land surface models; and the development of fully-coupled snow chemistry and physics models.

Acknowledgments

This work is in support of the Arctic Monitoring and Assessment Program (AMAP). Financing for this study is primarily by DANCEA (Danish Cooperation for Environment in the Arctic) under the Danish Ministry of Energy, Buildings and Climate. Wang is partially funded by the Joint Institute for the Study of the Atmosphere and Ocean (JISAO) under NOAA Cooperative Agreement NA15OAR4320063, Contribution No. 2018–0169, NOAA grant (NA17OAR4310160). Parmentier was supported by the Norwegian Research Council under grant agreement 274711, and the Swedish Research Council under registration nr. 2017-05268.

ORCID iDs

Jason E Box  <https://orcid.org/0000-0003-0052-8705>
 William T Colgan  <https://orcid.org/0000-0001-6334-1660>
 Torben Røjle Christensen  <https://orcid.org/0000-0002-4917-148X>
 Magnus Lund  <https://orcid.org/0000-0003-1622-2305>
 Frans-Jan W Parmentier  <https://orcid.org/0000-0003-2952-7706>
 Uma S Bhatt  <https://orcid.org/0000-0003-1056-3686>
 James E Overland  <https://orcid.org/0000-0002-2012-8832>
 Walter N Meier  <https://orcid.org/0000-0003-2857-0550>
 Bert Wouters  <https://orcid.org/0000-0002-1086-2435>

References

- Ahlström A P, Petersen D, Langen P L, Citterio M and Box J E 2017 Abrupt shift in the observed runoff from the southwestern Greenland ice sheet *Sci. Adv.* **3** e1701169
- AMAP 2017 *Snow, Water, Ice and Permafrost in the Arctic (SWIPA) 2017* (Oslo, NO: Arctic Monitoring and Assessment Programme (AMAP)) p 269 xiv +
- Amstrup S C, DeWeaver E T, Douglas D C, Marcot B G, Durner G M, Bitz C M and Bailey D A 2010 Greenhouse gas mitigation can reduce sea-ice loss and increase polar bear persistence *Nature* **468** 955–8
- Andreassen K *et al* 2017 Massive blow-out craters formed by hydrate-controlled methane expulsion from the Arctic seafloor *Science* **356** 948–52
- Barry T and Helgason H 2016 Arctic Species Trend Index (ASTI): Terrestrial. Version 1.2. Conservation of Arctic Flora and Fauna. Occurrence dataset <https://doi.org/10.15468/gd0xmy>
- Bekryaev R V, Polyakov I V and Alexeev V A 2010 Role of polar amplification in long-term surface air temperature variations and modern arctic warming *J. Clim.* **23** 3888–906
- Bendixen M *et al* 2017 Delta progradation in greenland driven by increasing glacial mass loss *Nature* **550** 101
- Bergeron Y and Leduc A 1998 Relationships between change in fire frequency and mortality due to spruce budworm outbreak in the southeastern Canadian boreal forest *Journal of Vegetation Science* **9** 493–500
- Bhatt U S *et al* 2017 Changing seasonality of panarctic tundra vegetation in relationship to climatic variables *Environ. Res. Lett.* **12** 055003
- Bintanja R and Selten F M 2014 Future increases in Arctic precipitation linked to local evaporation and sea-ice retreat *Nature* **509** 479
- Bliss A C, Miller J A and Meier W N 2017 Comparison of passive microwave-derived early melt onset records on Arctic sea ice *Remote Sens.* **9** 199
- Blume-Werry G, Wilson S D, Kreyling J and Milbau A 2016 The hidden season: growing season is 50% longer below than above ground along an arctic elevation gradient *New Phytol.* **209** 978–86
- Bokhorst S *et al* 2016 Changing Arctic snow cover: a review of recent developments and assessment of future needs for observations, modelling, and impacts *Ambio* **45** 516–37
- Bokhorst S, Bjerke J W, Tommervik H, Preece C and Phoenix G K 2012 Ecosystem response to climatic change: the importance of the cold season *Ambio* **41** 246–55
- Box J E *et al* 2013 Greenland ice sheet mass balance reconstruction: I. Net snow accumulation (1600–2009) *J. Clim.* **26** 3919–34
- Box J E, Colgan W T, Wouters B, Burgess D O, O'Neil S, Thomson L I and Mernild S H 2018 Global sea-level contribution from Arctic land ice: 1971–2017 *Environ. Res. Lett.* **13** 105795
- Braithwaite R J 1995 Positive degree-day factors for ablation on the greenland ice-sheet studied by energy-balance modeling *J. Glaciol.* **41** 153–60
- Bret-Harte M S, Mack M C, Shaver G R, Huebner D C, Johnston M, Mojica C A, Pizano C and Reiskind J A 2013 The response of Arctic vegetation and soils following an unusually severe tundra fire *Phil. Trans. R. Soc. B* **368**
- Bring A, Fedorova I, Dibike Y, Hinzman L, Mard J, Mernild S H, Prowse T, Semenova O, Stuefer S L and Woo M K 2016 Arctic terrestrial hydrology: a synthesis of processes, regional effects, and research challenges *J. Geophys. Res.* **121** 621–49
- Brown R D, Barrette C, Brown L, Chaumont D, Grenier P, Howell S and Sharp M 2018 Climate variability, trends and projected change for the Eastern Canadian Arctic *From Science to Policy in the Eastern Canadian Arctic: An Integrated Regional Impact Study (IRIS) of Climate Change and Modernization* ed T Bell and T M Brown (Quebec City: ArcticNet) ch 2 p 560
- Brown R D and Derksen C 2013 Is Eurasian october snow cover extent increasing? *Environ. Res. Lett.* **8**
- Brown R D, Vikhamar-Schuler D, Bulygina O, Derksen C, Luoju K, Mudryk L, Wang L and Yang D 2017 Arctic terrestrial snow cover *AMAP 2017 Snow, Water, Ice and Permafrost in the Arctic (SWIPA) 2017* (Oslo, NO: Arctic Monitoring and Assessment Programme (AMAP)) ch 3 p 269 xiv +
- Burgess D O 2017 Mass balance of ice caps in the Queen Elizabeth Islands, Arctic Canada: 2014–2015 Open File 8223 Geological Survey of Canada p38
- Callaghan T V *et al* 2013 Ecosystem change and stability over multiple decades in the Swedish sub-Arctic: complex processes and multiple drivers *Phil. Trans. R. Soc. B* **368** 20120488
- Chapin F S *et al* 2005 Role of land-surface changes in Arctic summer warming *Science* **310** 657–60
- Chatterjee S and Hadi A S 2006 *Regression Analysis by Example* Fourth Edition (John Wiley & Sons) (<https://doi.org/10.1002/0470055464>)
- Christensen T R 2014 Climate science: understand Arctic methane variability *Nature* **509** 279–81

- Christensen T R, Arora V K, Gauss M, Höglund-Isaksson L and Parmentier F-J W 2019 Tracing the climate signal: mitigation of anthropogenic methane emissions can outweigh a large Arctic natural emission increase *Sci. Rep.* in press **9** 1146
- Cogley J G, Adams W P, Ecclestone M A, JungRothenhausler F and Ommanney C S L 1996 Mass balance of white glacier, Axel Heiberg island, NWT, Canada, 1960–91 *Journal of Glaciology* **42** 548–63
- Cohen J *et al* 2014 Recent Arctic amplification and extreme mid-latitude weather *Nat. Geosci.* **7** 627–37
- Cohen J L, Furtado J C, Barlow M A, Alexeev V A and Cherry J E 2012 Arctic warming, increasing snow cover and widespread boreal winter cooling *Environmental Research Letters* **7**
- Comiso J C, Meier W N and Gersten R 2017 Variability and trends in the Arctic sea ice cover: results from different techniques *J. Geophys. Res.-Oceans* **122** 6883–900
- Commune R *et al* 2017 Carbon dioxide sources from Alaska driven by increasing early winter respiration from Arctic tundra *Proc. Natl Acad. Sci. USA* **114** 5361–6
- Doiron M, Gauthier G and Levesque E 2015 Trophic mismatch and its effects on the growth of young in an Arctic herbivore *Glob. Change Biol.* **21** 4364–76
- Dowdeswell J A *et al* 1997 The mass balance of circum-Arctic glaciers and recent climate change *Quaternary Research* **48** 1–14
- Doyle S H *et al* 2015 Amplified melt and flow of the Greenland ice sheet driven by late-summer cyclonic rainfall *Nat. Geosci.* **8** 647
- Dyrugerov M 2002 Glacier mass balance and regime: data of measurements and analysis. Boulder, CO, University of Colorado. Institute of Arctic and Alpine Research. (INSTAAR Occasional Paper 55)
- Frey B R, Lieffers V J, Hogg E H and Landhauser S M 2004 Predicting landscape patterns of aspen dieback: mechanisms and knowledge gaps *Canadian Journal of Forest Research-Revue Canadienne De Recherche Forestiere* **34** 1379–90
- Furgal C, Chan L, Tremblay M, Rajdev V, Barrett M and Sheldon T 2012 Impacts of climate change on food security in Nunavik and Nunatsiavut *Nunavik and Nunatsiavut: From Science to Policy: An Integrated Regional Impact Study (IRIS) of Climate Change and Modernization* ed M Allard and M Lemay (ArcticNet) pp 156–69
- Gardner A S and Sharp M 2007 Influence of the arctic circumpolar vortex on the mass balance of Canadian high arctic glaciers *J. Clim.* **20** 4586–98
- Haine T W N *et al* 2015 Arctic freshwater export: Status, mechanisms, and prospects *Glob. Planet. Change* **125** 13–35
- Hartmann D L *et al* 2013 Observations: atmosphere and surface *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* IPCC, 2014 (Cambridge: Cambridge University Press)
- Hawkings J, Wadham J, Tranter M, Telling J, Bagshaw E, Beaton A, Simmons S L, Chandler D, Tedstone A and Nienow P 2016 The Greenland ice sheet as a hot spot of phosphorus weathering and export in the Arctic *Glob. Biogeochem. Cycles* **30** 191–210
- Hawley R L, Courville Z R, Kehrl L M, Lutz E R, Osterberg E C, Overly T B and Wong G J 2014 Recent accumulation variability in northwest Greenland from ground-penetrating radar and shallow cores along the Greenland inland traverse *J. Glaciol.* **60** 375–82
- Heide-Jorgensen M P, Laidre K L, Borchers D, Marques T A, Stern H and Simon M 2010 The effect of sea-ice loss on beluga whales (*Delphinapterus leucas*) in West Greenland *Polar Research* **29** 198–208
- Heide-Jorgensen M P, Laidre K L, Quakenbush L T and Citta J J 2012 The Northwest Passage opens for bowhead whales *Biology Letters* **8** 270–3
- Hernandez-Henriquez M A, Dery S J and Derksen C 2015 Polar amplification and elevation-dependence in trends of Northern Hemisphere snow cover extent, 1971–2014 *Environ. Res. Lett.* **10**
- Higdon J W, Hauser D D W and Ferguson S H 2012 Killer whales (*Orcinus orca*) in the Canadian Arctic: Distribution, prey items, group sizes, and seasonality *Marine Mammal Science* **28** E93–E109
- Hobbie J E and Klings G W (ed) 2014 *Alaska's Changing Arctic: Ecological Consequences for Tundra, Streams, and Lakes* (New York, NY: Oxford University Press)
- Hobbie J E, Shaver G R, Rastetter E B, Cherry J E, Goetz S J, Guay K C, Gould W A and Kling G W 2017 Ecosystem responses to climate change at a low arctic and a high arctic long-term research site *Ambio* **46** S160–73
- Hock R 2003 Temperature index melt modelling in mountain areas *J. Hydrol.* **282** 104–15
- Hogg E H, Brandt J P and Michaellian M 2008 Impacts of a regional drought on the productivity, dieback, and biomass of western Canadian aspen forests *Canadian Journal of Forest Research* **38** 1373–84
- Holmes R M, Shiklomanov A I, Tank S E, McClelland J W and Tretiakov M 2015 River Discharge *Arctic Report Card: Update for* (<https://arctic.noaa.gov/Report-Card/Report-Card-2015/ArtMID/5037/ArticleID/227/River-Discharge>)
- Hoye T T, Post E, Schmidt N M, Trojelsgaard K and Forchhammer M C 2013 Shorter flowering seasons and declining abundance of flower visitors in a warmer Arctic *Nat. Clim. Change* **3** 759–63
- Hugelius G *et al* 2014 Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps *Biogeosciences* **11** 6573–93
- IPCC 2013, 2014 *Climate Change 2013: The Physical Science Basis*
- IPCC 2014a Summary for policymakers *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed C B Field (Cambridge and New York, NY: Cambridge University Press) pp 1–32
- IPCC 2014b *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge: Cambridge University Press)
- Irannezhad M, Ronkanen A K and Klove B 2016 Wintertime climate factors controlling snow resource decline in Finland *International Journal of Climatology* **36** 110–31
- Jakobson E and Vihma T 2010 Atmospheric moisture budget in the Arctic based on the ERA-40 reanalysis *Int. J. Climatol.* **30** 2175–94
- Jeong D, Sushama L and Khaliq M N 2017 Attribution of spring snow water equivalent (SWE) changes over the northern hemisphere to anthropogenic effects *Clim. Dyn.* **48** 3645–58
- Jeong S-J *et al* 2018 Accelerating rates of Arctic carbon cycling revealed by long-term atmospheric CO₂ measurements *Sci. Adv.* **4** eaao1167
- Johansson M, Christensen T R, Akerman H J and Callaghan T V 2006 What determines the current presence or absence of permafrost in the Tornetrask region, a sub-arctic landscape in Northern Sweden? *Ambio* **35** 190–7
- Jolly W M, Cochrane M A, Freeborn P H, Holden Z A, Brown T J, Williamson G J and Bowman D 2015 Climate-induced variations in global wildfire danger from 1979 to 2013 *Nat. Commun.* **6**
- Jorgenson M T, Racine C H, Walters J C and Osterkamp T E 2001 Permafrost degradation and ecological changes associated with a warming climate in central Alaska *Clim. Change* **48** 551–79
- Kalnay E *et al* 1996 The NCEP/NCAR 40-year reanalysis project *Bull. Am. Meteorol. Soc.* **77** 437–71
- Kliejunas J T, Geils B W, Micales Glaeser J, Michaels Goheen E, Hennon P, Kim M-S, Kope H, Stone J, Sturrock R and Frankel S J 2009 Review of Literature on Climate Change and Forest Diseases of Western North America, United States Department of Agriculture Forest Service Pacific Southwest Research Station General Technical Report PSW-GTR-225 October 2009

- Kankaanpaa T, Skov K, Abrego N, Lund M, Schmidt N M and Roslin T 2018 Spatiotemporal snowmelt patterns within a high Arctic landscape, with implications for flora and fauna *Arctic Antarct. Alpine Res.* **50**
- Kenney M A, Janetos A C and Lough G C 2016 Building an integrated US national climate indicators system *Clim. Change* **135** 85–96
- Kononova N K 2012 The influence of atmospheric circulation on the formation of snow cover on the north eastern Siberia *Ice and Snow* **1** 38–53 (In Russian, English summary)
- Kwok R and Cunningham G F 2015 Variability of Arctic sea ice thickness and volume from CryoSat-2 *Phil Trans. R. Soc. A* **373**
- Lien V S, Schlichtholz P, Skagseth O and Vikebo F B 2017 Wind-driven Atlantic Water flow as a direct mode for reduced barents sea ice cover *J. Clim.* **30** 803–12
- Liljedahl A K *et al* 2016 Pan-Arctic ice-wedge degradation in warming permafrost and its influence on tundra hydrology *Nature Geoscience* **9** 312 –+
- Lopez-Blanco E, Lund M, Williams M, Tamstorf M P, Westergaard-Nielsen A, Exbrayat J F, Hansen B U and Christensen T R 2017 Exchange of CO₂ in Arctic tundra: impacts of meteorological variations and biological disturbance *Biogeosciences* **14** 4467–83
- Lund M 2018 Uncovering the unknown–climate interactions in a changing arctic tundra *Environ. Res. Lett.* **13**
- Lund M *et al* 2010 Variability in exchange of CO₂ across 12 northern peatland and tundra sites *Glob. Change Biol.* **16** 2436–48
- Lund M, Falk J M, Friborg T, Mbufong H N, Sigsgaard C, Soegaard H and Tamstorf M P 2012 Trends in CO₂ exchange in a high Arctic tundra heath, 2000–2010 *J. Geophys. Res.* **117**
- Lund M, Hansen B U, Pedersen S H, Stiegler C and Tamstorf M P 2014 Characteristics of summer–time energy exchange in a high Arctic tundra heath 2000–2010 *Tellus B* **66**
- Lund M, Stiegler C, Abermann J, Citterio M, Hansen B U and van As D 2017 Spatiotemporal variability in surface energy balance across tundra, snow and ice in greenland *Ambio* **46** S81–93
- Markus T, Stroeve J C and Miller J 2009 Recent changes in Arctic sea ice melt onset, freezeup, and melt season length *Journal of Geophysical Research-Oceans* **114**
- Martin A C, Jeffers E S, Petrokofsky G, Myers-Smith I and Macias-Fauria M 2017 Shrub growth and expansion in the Arctic tundra: an assessment of controlling factors using an evidence-based approach *Environ. Res. Lett.* **12**
- Marzeion B, Cogley J G, Richter K and Parkes D 2014 Attribution of global glacier mass loss to anthropogenic and natural causes *Science* **345** 919–21
- Mastepanov M, Sigsgaard C, Dlugokencky E J, Houweling S, Strom L, Tamstorf M P and Christensen T R 2008 Large tundra methane burst during onset of freezing *Nature* **456** 628–U58
- Mastepanov M, Sigsgaard C, Strom L, Tamstorf M P, Lund M and Christensen T R 2013 Revisiting factors controlling methane emissions from high-Arctic tundra *Biogeosciences* **10** 5139–58
- Meier W N *et al* 2014 Arctic sea ice in transformation: a review of recent observed changes and impacts on biology and human activity *Rev. Geophys.* **52** 185–217
- Mellinger D K, Nieuwkirk S L, Klinck K, Klinck H, Dziak R P, Clapham P J and Brandsdottir B 2011 Confirmation of right whales near a nineteenth-century whaling ground east of southern Greenland *Biology Letters* **7** 411–3
- Miller G H, Alley R B, Brigham-Grette J, Fitzpatrick J J, Polyak L, Serreze M C and White J W C 2010 Arctic amplification: can the past constrain the future? *Quat. Sci. Rev.* **29** 1779–90
- Moore S E 2008 Marine mammals as ecosystem sentinels *J. Mamm.* **89** 534–40
- Moore S E, Van Parijs S M, Southall B L and Stafford K M 2011 Listening to marine mammals at basin to local scales *J. Acoustical Soc. America* **129** 2371
- Morin X and Thuiller W 2009 Comparing niche- and process-based models to reduce prediction uncertainty in species range shifts under climate change *Ecology* **90** 1301–13
- Mortimer C A, Sharp M and Wouters B 2016 Glacier surface temperatures in the Canadian high Arctic, 2000–15 *Journal of Glaciology* **62** 963–75
- Mudryk L R, Kushner P J, Derksen C and Thackeray C 2017 Snow cover response to temperature in observational and climate model ensembles *Geophys. Res. Lett.* **44** 919–26
- Nabe-Nielsen J, Normand S, Hui F K C, Stewart L, Bay C, Nabe-Nielsen L I and Schmidt N M 2017 Plant community composition and species richness in the high Arctic tundra: from the present to the future *Ecol. Evol.* **7** 10233–42
- Najafi M R, Zwiers F W and Gillett N P 2016 Attribution of the spring snow cover extent decline in the northern hemisphere, Eurasia and north America to anthropogenic influence *Clim. Change* **136** 571–86
- Neff W, Compo G P, Ralph F M and Shupe M D 2014 Continental heat anomalies and the extreme melting of the Greenland ice surface in 2012 and 1889 *J. Geophys. Res.-Atmos.* **119** 6520–36
- Nesje A, Lie O and Dahl S O 2000 Is the north Atlantic oscillation reflected in scandinavian glacier mass balance records? *J. Quat. Sci.* **15** 587–601
- Nicolsoy D J, Romanovsky V E, Panda S K, Marchenko S S and Muskett R R 2017 Applicability of the ecosystem type approach to model permafrost dynamics across the Alaska north slope *J. Geophys. Res.* **122** 50–75
- Noel B, van de Berg W J, Lhermitte S, Wouters B, Schaffer N and van den Broeke M R 2018 Six decades of glacial mass loss in the Canadian Arctic Archipelago *Journal of Geophysical Research-Earth Surface* **123** 1430–49
- Oberbauer S F *et al* 2013 Phenological response of tundra plants to background climate variation tested using the international tundra experiment *Phil. Trans. R. Soc. B* **368**
- Ohmura A 2001 Physical basis for the temperature-based melt-index method *Journal of Applied Meteorology* **40** 753–61
- Olefeldt D, Turetsky M R, Crill P M and McGuire A D 2013 Environmental and physical controls on northern terrestrial methane emissions across permafrost zones *Glob. Change Biol.* **19** 589–603
- Ostby T I, Schuler T V, Hagen J O, Hock R, Kohler J and Reijmer C H 2017 Diagnosing the decline in climatic mass balance of glaciers in Svalbard over 1957–2014 *Cryosphere* **11** 191–215
- Overland J E, Dunlea E, Box J E, Corell R, Forsius M, Kattsov V, Olsen M, Pawlak J, Reiersen L-O and Wang M 2018 The urgency of arctic change *Polar Sci.* In press (<https://doi.org/10.1016/j.polar.2018.11.008>)
- Overland J E, Spillane M C, Percival D B, Wang M Y and Mofjeld H O 2004 Seasonal and regional variation of pan-Arctic surface air temperature over the instrumental record *J. Clim.* **17** 3263–82
- Overland J E and Wang M Y 2016 Recent extreme arctic temperatures are due to a split polar vortex *J. Clim.* **29** 5609–16
- Parazoo N C, Koven C D, Lawrence D M, Romanovsky V and Miller C E 2018 Detecting the permafrost carbon feedback: talik formation and increased cold-season respiration as precursors to sink-to-source transitions *Cryosphere* **12** 123–44
- Parkinson C L 2014 Spatially mapped reductions in the length of the Arctic sea ice season *Geophys. Res. Lett.* **41** 4316–22
- Parmentier F-J W, Christensen T R, Rysgaard S, Bendtsen J, Glud R N, Else B, van Huissteden J, Sachs T, Vonk J E and Sejr M K 2017 A synthesis of the arctic terrestrial and marine carbon cycles under pressure from a dwindling cryosphere *Ambio* **46** 53–69
- Parmentier F J W, Christensen T R, Sørensen L L, Rysgaard S, McGuire A D, Miller P A and Walker D A 2013 The impact of lower sea-ice extent on Arctic greenhouse-gas exchange *Nat. Clim. Change* **3** 195–202
- Parmentier F J W, Rasse D P, Lund M, Bjerke J W, Drake B G, Weldon S, Tommervik H and Hansen G H 2018 Vulnerability and resilience of the carbon exchange of a subarctic peatland to an extreme winter event *Environ. Res. Lett.* **13**

- Parmentier F J W, van der Molen M K, van Huissteden J, Karsanaev S A, Kononov A V, Suzdalov D A, Maximov T C and Dolman A J 2011 Longer growing seasons do not increase net carbon uptake in the northeastern Siberian tundra *J. Geophys. Res.* **116**
- Peng G, Steele M, Bliss A C, Meier W N and Dickinson S 2018 Temporal means and variability of arctic sea ice melt and freeze season climate indicators using a satellite *Clim. Data Record Remote Sens.* **10**
- Pepin N *et al* Mt Res Initiative E D W W G 2015 Elevation-dependent warming in mountain regions of the world *Nat. Clim. Change* **5** 424–30
- Perovich D K and Polashenski C 2012 Albedo evolution of seasonal Arctic sea ice *Geophys. Res. Lett.* **39**
- Peterson B J, Holmes R M, McClelland J W, Vorosmarty C J, Lammers R B, Shiklomanov A I, Shiklomanov I A and Rahmstorf S 2002 Increasing river discharge to the arctic ocean *Science* **298** 2171–3
- Phoenix G K and Bjerke J W 2016 Arctic browning: extreme events and trends reversing arctic greening *Glob. Change Biol.* **22** 2960–2
- Pirk N, Mastepanov M, Loez-Blanco E, Christensen L H, Christiansen H H, Hansen B U, Lund M, Parmentier F J W, Skov K and Christensen T R 2017 Toward a statistical description of methane emissions from arctic wetlands *Ambio* **46** 570–80
- Pirk N, Mastepanov M, Parmentier F J W, Lund M, Crill P and Christensen T R 2016 Calculations of automatic chamber flux measurements of methane and carbon dioxide using short time series of concentrations *Biogeosciences* **13** 903–12
- Pithan F and Mauritsen T 2014 Arctic amplification dominated by temperature feedbacks in contemporary climate models *Nat. Geosci.* **7** 181–4
- Post E, Bhatt U S, Bitz C M, Brodie J F, Fulton T L, Hebblewhite M, Kerby J, Kutz S J, Stirling I and Walker D A 2013 Ecological consequences of sea-ice decline *Science* **341** 519–24
- Price D T *et al* 2013 Anticipating the consequences of climate change for Canada's boreal forest ecosystems *Environ. Rev.* **21** 322–65
- Rapaic M, Brown R, Markovic M and Chaumont D 2015 An evaluation of temperature and precipitation surface-based and reanalysis datasets for the canadian arctic, 1950–2010 *Atmos.-Ocean* **53** 283–303
- Rasmus S *et al* 2015 *Recent Change-Terrestrial Cryosphere*
- Rawlins M A *et al* 2010 Analysis of the arctic system for freshwater cycle intensification: observations and expectations *J. Clim.* **23** 5715–37
- Romanovsky V E, Smith S L, Shiklomanov N I, Streletskiy D A, Isaksen K, Kholodov A L, Christiansen H H, Drozdov D S, Malkova G V and Marchenko S S 2017 [The Arctic] terrestrial permafrost [in “State of the Climate in 2016”] State of the Climate in 2016 *Bull. Am. Meteorol. Soc.* **98** S1
- Ruppel C D and Kessler J D 2017 The interaction of climate change and methane hydrates *Rev. Geophys.* **55** 126–68
- Schadel C *et al* 2016 Potential carbon emissions dominated by carbon dioxide from thawed permafrost soils *Nat. Clim. Change* **6** 950
- Schmidt N M, Christensen T R and Roslin T 2017 A high arctic experience of uniting research and monitoring *Earths Future* **5** 650–4
- Schmidt N M, Mosbacher J B, Nielsen P S, Rasmussen C, Hoyer T T and Roslin T 2016 An ecological function in crisis? The temporal overlap between plant flowering and pollinator function shrinks as the Arctic warms *Ecography* **39** 1250–2
- Schmidt N M, Pedersen S H, Mosbacher J B and Hansen L H 2015 Long-term patterns of muskox (*Ovibos moschatus*) demographics in high arctic greenland *Polar Biol.* **38** 1667–75
- Schuur E A G *et al* 2015 Climate change and the permafrost carbon feedback *Nature* **520** 171–9
- Serreze M C, Barrett A P and Stroeve J 2012 Recent changes in tropospheric water vapor over the Arctic as assessed from radiosondes and atmospheric reanalyses *J. Geophys. Res.* **117**
- Serreze M C and Barry R G 2011 Processes and impacts of Arctic amplification: a research synthesis *Glob. Planet. Change* **77** 85–96
- Serreze M C, Barrett A P, Slater A G, Woodgate R A, Aagaard K, Lammers R B, Steele M, Moritz R, Meredith M and Lee C M 2006 The large-scale freshwater cycle of the Arctic *Journal of Geophysical Research-Oceans* **111**
- Serreze M C, Bromwich D H, Clark M P, Etringer A J, Zhang T J and Lammers R 2002 Large-scale hydro-climatology of the terrestrial Arctic drainage system *J. Geophys. Res.* **108**
- Spencer R G M, Mann P J, Dittmar T, Eglinton T I, McIntyre C, Holmes R M, Zimov N and Stubbins A 2015 Detecting the signature of permafrost thaw in Arctic rivers *Geophys. Res. Lett.* **42** 2830–5
- Stafford K M, Moore S E, Stabeno P J, Holliday D V, Napp J M and Mellinger D K 2010 Biophysical ocean observation in the southeastern Bering Sea *Geophysical Research Letters* **37**
- Stiegler C, Lund M, Christensen T R, Mastepanov M and Lindroth A 2016 Two years with extreme and little snowfall: effects on energy partitioning and surface energy exchange in a high-Arctic tundra ecosystem *Cryosphere* **10** 1395–413
- Stroeve J C, Crawford A D and Stammerjohn S 2016 Using timing of ice retreat to predict timing of fall freeze-up in the Arctic *Geophys. Res. Lett.* **43** 6332–40
- Stroeve J C, Markus T, Boisvert L, Miller J and Barrett A 2014 Changes in Arctic melt season and implications for sea ice loss *Geophys. Res. Lett.* **41** 1216–25
- Thomson L I and Copland L 2017 Changing contribution of peak velocity events to annual velocities following a multi-decadal slowdown at White Glacier *Annals of Glaciology* **58** 145–54
- Treat C C, Bloom A A and Marushchak M E 2018 Nongrowing season methane emissions—a significant component of annual emissions across northern ecosystems *Glob. Change Biol.* **24** 3331–43
- Trenberth K E 2011 Changes in precipitation with climate change *Clim. Res.* **47** 123–38
- Tschudi M A, Stroeve J C and Stewart J S 2016 Relating the age of arctic sea ice to its thickness, as measured during NASA's ICESat and IceBridge *Campaigns Remote Sens.* **8**
- Tyler N J C 2010 Climate, snow, ice, crashes, and declines in populations of reindeer and caribou (*Rangifer tarandus* L.) *Ecol. Monogr.* **80** 197–219
- Veraverbeke S, Rogers B M, Goulden M L, Jandt R R, Miller C E, Wiggins E B and Randerson J T 2017 Lightning as a major driver of recent large fire years in North American boreal forests *Nat. Clim. Change* **7** 529
- Vihma T, Screen J, Tjernstrom M, Newton B, Zhang X D, Popova V, Deser C, Holland M and Prowse T 2016 The atmospheric role in the Arctic water cycle: a review on processes, past and future changes, and their impacts *J. Geophys. Res.* **121** 586–620
- Vincent L A, Zhang X, Brown R D, Feng Y, Mekis E, Milewska E J, Wan H and Wang X L 2015 Observed trends in Canada's climate and influence of low-frequency variability modes *J. Clim.* **28** 4545–60
- Volney W J A and Hirsch K G 2005 Disturbing forest disturbances *Forestry Chronicle* **81** 662–8
- Walsh J E, Overland J E, Groisman P Y and Rudolf B 2011 Arctic climate: recent variations. Snow, water, ice and permafrost in the Arctic *Arctic Monitoring and Assessment Programme (Oslo, Norway)* xii+ p 538
- Wang M Y, Yang Q, Overland J E and Stabeno P 2018 Sea-ice cover timing in the pacific Arctic: the present and projections to mid-century by selected CMIP5 models *Deep-Sea Res. II* **152** 22–34
- Watts J D, Kimball J S, Bartsch A and McDonald K C 2014 Surface water inundation in the boreal-Arctic: potential impacts on regional methane emissions *Environ. Res. Lett.* **9**
- Webster M A, Rigor I G, Nghiem S V, Kurtz N T, Farrell S L, Perovich D K and Sturm M 2014 Interdecadal changes in snow depth on Arctic sea ice *J. Geophys. Res.-Oceans* **119** 5395–406
- Wong G J, Osterberg E C, Hawley R L, Courville Z R, Ferris D G and Howley J A 2015 Coast-to-interior gradient in recent northwest Greenland precipitation trends (1952–2012) *Environ. Res. Lett.* **10**

- Woo M K and Young K L 2014 Disappearing semi-permanent snow in the High Arctic and its consequences *J. Glaciol.* **60** 192–200
- Young A M, Higuera P E, Duffy P A and Hu F S 2017 Climatic thresholds shape northern high-latitude fire regimes and imply vulnerability to future climate change *Ecography* **40** 606–17
- Zhang K, Kimball J S, Mu Q Z, Jones L A, Goetz S J and Running S W 2009 Satellite based analysis of northern ET trends and associated changes in the regional water balance from 1983 to 2005 *J. Hydrol.* **379** 92–110
- Zhang T, Stamnes K and Bowling S A 2001 Impact of the atmospheric thickness on the atmospheric downwelling longwave radiation and snowmelt under clear-sky conditions in the Arctic and subarctic *J. Clim.* **14** 920–39
- Zhang T J 2005 Influence of the seasonal snow cover on the ground thermal regime: an overview *Rev. Geophys.* **43**
- Zhang X D, He J X, Zhang J, Polyakov I, Gerdes R, Inoue J and Wu P L 2013 Enhanced poleward moisture transport and amplified northern high-latitude wetting trend *Nat. Clim. Change* **3** 47–51
- Zhang Y *et al* 2018 On the relationship between sub-daily instantaneous and daily total gross primary production: implications for interpreting satellite-based SIF retrievals *Remote Sens. Environ.* **205** 276–89