

Martingale solutions to the stochastic thin-film equation in two dimensions

Sauerbrey, Max

DOI

[10.1214/22-AIHP1328](https://doi.org/10.1214/22-AIHP1328)

Publication date

2024

Document Version

Final published version

Published in

Annales de l'institut Henri Poincaré (B) Probability and Statistics

Citation (APA)

Sauerbrey, M. (2024). Martingale solutions to the stochastic thin-film equation in two dimensions. *Annales de l'institut Henri Poincaré (B) Probability and Statistics*, 60(1), 373-412. <https://doi.org/10.1214/22-AIHP1328>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

Martingale solutions to the stochastic thin-film equation in two dimensions

Max Sauerbrey^a

Delft Institute of Applied Mathematics, TU Delft, Mekelweg 4, 2628 CD Delft, The Netherlands, ^aM.Sauerbrey@tudelft.nl

Received 12 August 2021; revised 13 September 2022; accepted 26 September 2022

Abstract. We construct solutions to the stochastic thin-film equation with quadratic mobility and Stratonovich gradient noise in the physically relevant dimension $d = 2$ and allow in particular for solutions with non-full support. The construction relies on a Trotter–Kato time-splitting scheme, which was recently employed in $d = 1$. The additional analytical challenges due to the higher spatial dimension are overcome using α -entropy estimates and corresponding tightness arguments.

Résumé. Nous construisons des solutions de l'équation aux dérivées partielles stochastique des couches minces avec une mobilité quadratique et un forçage stochastique gradient de type Stratonovich en dimension $d = 2$, physiquement pertinente. Les solutions à support non plein sont autorisées. La construction repose sur une méthode de Trotter–Kato en fractionnant l'intervalle de temps, récemment utilisée dans le cas $d = 1$. Les difficultés supplémentaires, dues à la dimension spatiale supérieure, sont surmontées à l'aide d'estimations de l' α -entropie et d'arguments de tension correspondants.

MSC2020 subject classifications: 35R60; 76A20

Keywords: Thin-film equation; Noise; α -Entropy estimates; Stochastic compactness method

1. Introduction

The general stochastic thin-film equation is of the form

$$(1.1) \quad \partial_t u = -\operatorname{div}(m(u)\nabla(\Delta u - F'(u))) + \operatorname{div}(\sqrt{m(u)}\partial_t W)$$

and models the height of a thin liquid film $u(t, x)$ on a surface under the influence of thermal fluctuations. The function m is the mobility function, F is an interface potential and $\partial_t W$ is spatio-temporal noise. The stochastic thin-film equation was independently introduced in [12] and [22] as a lubrication approximation of the stochastic Navier–Stokes equation and explains, as shown in [22], discrepancies between simulations of deterministic thin-films and experiments for films with small heights.

The mobility function has typically the form $m(u) = u^n$ for some real exponent n depending on the boundary condition imposed at the boundary between the film and the substrate. In particular, imposing the no-slip condition for the Navier–Stokes equations, one obtains the mobility $m(u) = u^3$ and using the Navier-slip condition with slip length l one obtains $m(u) = lu^2 + u^3$, which is up to scaling reasonably well approximated by $m(u) = u^2$ for small film heights $u \ll l$. The interface potential F models forces between the molecules of the fluid film and the surface and takes for example the form $F(u) = u^{-8} - u^{-2} + 1$ in case of the 6–12 Lennard–Jones potential. If one neglects these forces and allows in particular for the qualitatively interesting situation of solutions without full support, one chooses $F(u) = 0$ instead.

The first existence result for solutions to the stochastic thin-film equation was proved in [16] for $m(u) = u^2$ with non-zero interface potential and the Itô interpretation of the noise term in dimension $d = 1$ using a Galerkin approximation. The interface potential $F(u)$ is there assumed to become singular at $u = 0$, which ensures that the constructed solutions stay strictly positive for all times. In [18], it was pointed out that the Stratonovich interpretation of (1.1) is more natural, since the use of Itô noise would yield additional correction terms in the lubrication approximation. Moreover, the use of Stratonovich noise allowed the authors to employ a splitting of the deterministic and stochastic dynamics and construct

solutions to (1.1) with $m(u) = u^2$ and $F(u) = 0$, again in $d = 1$. In particular, the result allows for initial values without full support, which is from a mathematical viewpoint more challenging, due to the loss of parabolicity of (1.1) in this case. Based on a Galerkin approximation, solutions to (1.1) in $d = 1$ with $F(u) = 0$ and $m(u) = u^n$ with $n \in [\frac{8}{3}, 4)$ were recently constructed in [11]. Although no interface potential is assumed, the mathematical analysis is based on an entropy estimate which ensures that the constructed solution stays positive almost everywhere. We also mention that in [10] a-priori estimates for (1.1) were deduced at the cost of adding additional compensation terms on the right-hand side and it was shown that a local existence result would extend to a global one based on these estimates. The first existence result in the physically relevant dimension $d = 2$ was recently proved in [31] for $m(u) = u^2$ and a non-zero interface potential as an adaption of [16] to the higher-dimensional setting. Finally, in [21] a result similar to the one in [18] was shown using a different regularization procedure and additionally so-called α -entropy estimates were derived. Both, the higher dimensional setting as well as the α -entropy estimates are central elements of the current article and we point out that it was developed independently of [31] and [21].

In the current article, we generalize the approach from [18] to the case $d = 2$, i.e. we show existence of martingale weak solutions to (1.1) with $m(u) = u^2$, $F(u) = 0$ and Stratonovich noise on the two-dimensional torus. The higher spatial dimension leads to additional mathematical challenges due to the reduced gain of integrability after employing the Sobolev embedding theorem. Indeed, in [18] the control of the surface energy

$$\int_{\mathbb{T}} |u'|^2 dx$$

suffices to show convergence of the nonlinear terms from the sequence of approximate solutions. As apparent from the deterministic setting [33], the additional control of the dissipation terms of the α -entropy

$$- \int_{\mathbb{T}^2} u^{\alpha+1} dx, \quad \alpha \in (-1, 0)$$

is necessary to deduce convergence of the nonlinear terms in the two-dimensional case. Hence, to adapt the time splitting approach from [18], we have to additionally control the α -entropy along the splitting scheme and use the more delicate limiting procedure from [33] compared to the one-dimensional case [4]. Combining this with the stochastic compactness method is the key challenge of this article. Moreover, compared to the independently proven result in [31], where an existence result in the two-dimensional case based on the dissipation of the classical entropy

$$- \int_{\mathbb{T}^2} \log(u) dx$$

is given, we do allow for solutions with a contact line between the fluid film and the solid, which is a qualitatively interesting object to study, see for example [13] and the references therein.

1.1. Main Result

We state the existence result which we will prove in the course of this article. Choosing $m(u) = u^2$, $F(u) = 0$ and interpreting the noise in the Stratonovich sense, we obtain the SPDE

$$(1.2) \quad du_t = - \operatorname{div}(u_t^2 \nabla \Delta u_t) dt + \operatorname{div}(u_t \circ dW_t) \quad \text{on } \mathbb{T}^2,$$

where W_t is specified as follows. We let $(\psi_l)_{l \in \mathbb{N}}$ be the orthonormal basis in $H^2(\mathbb{T}^2, \mathbb{R}^2)$ consisting of the eigenfunctions to the periodic Laplacian in the first and second component respectively, i.e. every ψ_l is of the form $(\xi_k, 0)$ or $(0, \xi_k)$ for some $k \in \mathbb{Z}^2$, where

$$(1.3) \quad \xi_k(x, y) = \frac{\tilde{\xi}_{k_1}(x) \tilde{\xi}_{k_2}(y)}{\sqrt{1 + (2\pi |k|)^2 + (2\pi |k|)^4}}$$

and

$$(1.4) \quad \tilde{\xi}_j(x) = \begin{cases} \sqrt{2} \cos(2\pi jx), & j < 0, \\ 1, & j = 0, \\ \sqrt{2} \sin(2\pi jx), & j > 0. \end{cases}$$

Moreover, we let $\Lambda = (\lambda_l)_{l \in \mathbb{N}} \in l^2(\mathbb{N})$ satisfy the symmetry relation

$$(1.5) \quad \lambda_l = \lambda_{\bar{l}} \text{ whenever } \psi_l = (\xi_k, 0) \wedge \psi_{\bar{l}} = (0, \xi_k) \text{ for some } k \in \mathbb{Z}^2.$$

Then

$$(1.6) \quad W_\Lambda(t) = \sum_{l=1}^{\infty} \lambda_l \beta_t^{(l)} \psi_l$$

for independent Brownian motions $(\beta^{(l)})_{l \in \mathbb{N}}$ defines a centered Gaussian process on $H^2(\mathbb{T}^2, \mathbb{R}^2)$ with the covariance operator $Qf = \sum_{l=1}^{\infty} \lambda_l^2 (f, \psi_l)_{H^2(\mathbb{T}^2, \mathbb{R})} \psi_l$. Inserting W_Λ as the driving process in (1.2), writing the Stratonovich integral in Itô-form, and writing $J = u^2 \nabla \Delta u$ in the weak form from [33, Eq. (3.2)] yields the following notion of weak martingale solutions to (1.2).

Definition 1.1. Let $T \in (0, \infty)$ and $q \in (2, \infty)$. A weak martingale solution to (1.2) with q' -regular non linearity on $[0, T]$ consists out of a filtered probability space satisfying the usual conditions, a family of independent Brownian motions $(\beta^{(l)})_{l \in \mathbb{N}}$, a continuous process $(u(t))_{t \in [0, T]}$ in $H_w^1(\mathbb{T}^2)$ together with a random variable J with values in $L^2(0, T; L^{q'}(\mathbb{T}^2, \mathbb{R}^2))$ such that

- (i) $u(t), J|_{[0, t]}$ are \mathfrak{F}_t -measurable as random variables in $H^1(\mathbb{T}^2)$ and $L^2(0, t; L^{q'}(\mathbb{T}^2, \mathbb{R}^2))$, respectively, for every $t \in [0, T]$,
- (ii) $|\nabla u| \in L^3(\{u > 0\})$ and for all $\eta \in L^\infty(0, T; W^{2, \infty}(\mathbb{T}^2))$ it holds almost surely

$$(1.7) \quad \begin{aligned} \int_0^T \int_{\mathbb{T}^2} J \cdot \eta \, dx \, dt &= \int_0^T \int_{\{u(t) > 0\}} |\nabla u|^2 \nabla u \cdot \eta \, dx \, dt + \int_0^T \int_{\{u(t) > 0\}} u |\nabla u|^2 \operatorname{div} \eta \, dx \, dt \\ &+ 2 \int_0^T \int_{\{u(t) > 0\}} u \nabla^T u D \eta \nabla u \, dx \, dt + \int_0^T \int_{\mathbb{T}^2} u^2 \nabla u \cdot \nabla \operatorname{div} \eta \, dx \, dt \end{aligned}$$

- (iii) and for all $\varphi \in W^{1, q}(\mathbb{T}^2)$ we have

$$(1.8) \quad \begin{aligned} \langle u(t), \varphi \rangle - \langle u_0, \varphi \rangle &= - \int_0^t \langle \operatorname{div}(J), \varphi \rangle \, ds + \frac{1}{2} \sum_{l=1}^{\infty} \int_0^t \lambda_l^2 \langle \operatorname{div}(\operatorname{div}(u(s) \psi_l) \psi_l), \varphi \rangle \, ds \\ &+ \sum_{l=1}^{\infty} \lambda_l \int_0^t \langle \operatorname{div}(u(s) \psi_l), \varphi \rangle \, d\beta_s^{(l)} \end{aligned}$$

almost surely for all $t \in [0, T]$.

Remark 1.2.

- (i) By the weak continuity in $H^1(\mathbb{T}^2)$ any solution u in the sense of Definition 1.1 satisfies

$$(1.9) \quad \sup_{0 \leq t \leq T} \|u(t)\|_{H^1(\mathbb{T}^2)} < \infty$$

almost surely.

- (ii) The measurability assumption on J in item (i) ensures that all the terms on the right-hand side of (1.8) are adapted. Interpreting J as an element of the distribution space $\mathcal{D}'(\mathbb{R} \times \mathbb{T}^2)$, one can equivalently demand that J is adapted to \mathfrak{F} in the sense of distributions [8, Definition 2.2.13]. This follows by density of $C_c^\infty((0, t) \times \mathbb{T}^2)$ in $L^2(0, t; L^q(\mathbb{T}^2, \mathbb{R}^2))$, separability of $L^2(0, t; L^{q'}(\mathbb{T}^2, \mathbb{R}^2))$, and the equivalence of weak and Borel measurability in separable Banach spaces [25, Proposition 1.1.1].

In course of this article, we will derive the following existence result.

Theorem 1.3. *Let μ be a probability distribution on $H^1(\mathbb{T}^2)$ supported on the non-negative functions, $T \in (0, \infty)$, $q \in (2, \infty)$ and $\alpha \in (-1, 0)$. Then there exists a weak martingale solution to (1.2) on $[0, T]$ with q' -regular non linearity satisfying $u(0) \sim \mu$. Moreover,*

- (i) $u(t) \geq 0$ almost surely for all $t \in [0, T]$,
- (ii) we have for $p \in (0, \infty)$ the estimates

$$(1.10) \quad E \left[\sup_{0 \leq t \leq T} \|u\|_{H^1(\mathbb{T}^2)}^p \right] \lesssim_{\Lambda, p, T} \int \|\cdot\|_{H^1(\mathbb{T}^2)}^p d\mu,$$

$$(1.11) \quad E \left[\|J\|_{L^2(0, T; L^{q'}(\mathbb{T}^2, \mathbb{R}^2))}^{\frac{p}{2}} \right] \lesssim_{\Lambda, p, q, T} \int \|\cdot\|_{H^1(\mathbb{T}^2)}^p d\mu$$

- (iii) and it holds the additional spatio-temporal regularity

$$(1.12) \quad u^{\frac{\alpha+3}{2}} \in L^2(0, T; H^2(\mathbb{T}^2)) \quad \text{and} \quad u^{\frac{\alpha+3}{4}} \in L^4(0, T; W^{1,4}(\mathbb{T}^2))$$

almost surely.

Remark 1.4.

- (i) We point out that we allow for the right-hand sides of (1.10) and (1.11) to be infinite, in which case the corresponding estimate trivializes.
- (ii) We note that the differential identity

$$\nabla u = \frac{4}{\alpha + 3} u^{\frac{1-\alpha}{4}} \nabla u^{\frac{\alpha+3}{4}},$$

the Sobolev embedding theorem, (1.12) and (1.9) imply that $|\nabla u| \in L^{4-}([0, T] \times \mathbb{T}^2)$ almost surely and in particular the integrability condition from Definition 1.1(ii).

1.2. Discussion of the result

Theorem 1.3 generalizes [18, Theorem 1.2] to the setting in two dimensions and is therefore together with the independently developed result [31, Theorem 3.5] the first existence result for the stochastic thin-film equation in higher dimension. As in the one-dimensional case, the time splitting approach is not only suitable to construct solutions to the stochastic thin-film equation, but suggests a numerical approach for their simulation as well. The assumption $\Lambda \in \mathcal{I}^2(\mathbb{N})$ on the noise (1.6) is the same as in [11, 16, 18], where we refer the reader for an interpretation of the expansion (1.6) in terms of a spatial correlation function of the noise to the exposition in [5]. The additionally imposed symmetry condition (1.5) expresses that the coordinate-wise noise processes are distributed according to the same Gaussian law in $H^2(\mathbb{T}^2)$. This is a physically reasonable assumption since the noise is induced by thermal fluctuations and its distribution depends consequently on its position but not on its direction. The same symmetry condition appears in [31, Eq. (2.19)], where the use of Stratonovich noise is discussed, which indicates that it is an important assumption to treat the stochastic thin-film equation in higher dimensions. We point out that in [31], the expansion (1.6) in terms of eigenfunctions of the periodic Laplacian is relaxed to a slightly more general assumption.

In contrast to the existence results from the mentioned articles, there is no integrability assumption on the initial distribution required in Theorem 1.3. This is achieved by using a decomposition of the initial value in countably many parts which are each almost surely bounded in $H^1(\mathbb{T}^2)$. Then one can construct approximate solutions and apply tightness arguments for each of these parts separately and add them together afterwards. The only important feature of (1.2) for this to work is that $u(t) = 0$ is a solution to it. We remark that these kind of reductions to bounded or integrable initial values are well-known and can be achieved in the setting of probabilistically strong solutions via localization or changing the probability measure, see [2, Proposition 4.13] or [29, Theorem 6.9.2] for examples. We use the decomposition of the initial value instead, since it is more compatible with the stochastic compactness method as well as the estimates (1.10) and (1.11). The moment estimates (1.10) and (1.11) for $p < 2$ are also new for the stochastic thin-film equation and are obtained from the estimates for higher moments.

1.3. Outline and discussion of the proof

In Section 2 we review the existence result for weak solutions to the deterministic thin-film equation in two dimensions from [33] and state properties of the obtained solutions which are immediate from their construction. Additionally, we show that there is a measurable solution operator using the measurable selection theorem, which is important to combine these results with the stochastic setting. This approach is to the author’s knowledge new and might be of interest also for other situations, where a measurable solution operator is required.

In Section 3 we consider the regularized Stratonovich SPDE

$$(1.13) \quad du_t = \epsilon \Delta u_t dt + \operatorname{div}(u_t \circ dW_t) \quad \text{on } \mathbb{T}^2$$

and establish well-posedness in $H^1(\mathbb{T}^2)$ using the monotone operator approach to SPDEs. The coercivity estimates (3.7), (3.8) are obtained analogously to the one-dimensional case [18, Eq. (A.9)] and require only some adaptations to multi-variable calculus, where the symmetry condition (1.5) is used. Their uniformity in ϵ is key to letting later on $\epsilon \searrow 0$ and eliminating the regularization term $\epsilon \Delta u_t$ from (1.13). We note that this procedure is well-known and refer the reader to the article [17] and the references therein for more information on degenerate parabolic SPDEs. However, the general result [17, Theorem 2.1] does not directly apply to

$$(1.14) \quad du_t = \operatorname{div}(u_t \circ dW_t) \quad \text{on } \mathbb{T}^2$$

and the coercivity estimates are unique to our particular situation.

In Section 4, we start constructing approximate solutions to (1.2) by splitting the stochastic and deterministic dynamics along a time stepping scheme with step length δ . Using the properties of the solutions to the deterministic thin-film equation and the solutions to (1.13), we derive estimates on the approximate solutions which are uniform in ϵ and δ . The procedure is analogous to the one-dimensional case, but we note that we take the slightly different approach to let $\epsilon \searrow 0$ afterwards to be able to apply Itô’s formula to the whole time splitting scheme. After these estimates are obtained, it is straightforward to deduce tightness statements on the approximating sequence in ϵ and employ the Skohorod–Jakubowski theorem to obtain an almost surely convergent, equally distributed subsequence. Usually, the parabolic regularization procedure does not require to pass to another probability space, see again [17], but it is in our case convenient to ensure convergence of the solutions to the deterministic equation as well.

Finally, in Section 5, we derive additional estimates on the approximating sequence by controlling the entropy production along the stochastic dynamics by means of Itô’s formula. Using the obtained estimates, we show additional tightness properties of powers of the solution by an adaption of the compactness argument in [33, Lemma 2.5], which is compatible with our splitting scheme. These arguments are unique to the higher-dimensional setting and distinguish our approach from the one-dimensional case. We employ the Skohorod–Jakubowski theorem once more to let $\delta \searrow 0$ and identify the limit as a solution to (1.2) combining the methods from [33, Theorem 3.2] and [18, Section 5.2]. As a result of the construction the additional estimates (1.10), (1.11), and the regularity properties (1.12) follow.

The reason to use the time-splitting approach instead of a linear parabolic regularization is that it directly yields non negative solutions, because the deterministic result [33] provides non negative solutions and the regularized stochastic part of the equation admits a maximum principle. Since we are dealing with a fourth order equation, a linear parabolic regularization of the whole equation would yield possibly negative solutions, which lack a reasonable physical interpretation. However, a more delicate, nonlinear regularization is possible as demonstrated in the one-dimensional case [11] or [21], but would require a longer proof.

1.4. Notation

We use the notation \lesssim to indicate that an inequality holds up to a universal constant and write $\lesssim_{p_1, \dots}$ if the constant depends on nothing but the parameters p_1, \dots . Similarly, we write C for a universal constant and $C_{p_1, \dots}$, if the constant depends on p_1, \dots . We write

$$G_\alpha(t) = \int_1^t \int_1^s \tau^{\alpha-1} d\tau ds, \quad \alpha \in \mathbb{R}$$

for the (mathematical) α -entropy, and point out for later reference that

$$(1.15) \quad G_\alpha(t) = \frac{t^{\alpha+1}}{\alpha(\alpha+1)} + r_\alpha(t), \quad t \geq 0,$$

if $\alpha \in (-1, 0)$, where r_α is a first order polynomial. We use classical notation for differential operators, i.e. write ∇f , $\operatorname{div}(f)$, Δf for the gradient, divergence and Laplacian of a function or a vector field f , respectively. Moreover, we write Hf for the Hessian matrix and use the notational convention that a differential operator is only applied to the first function appearing afterwards such that e.g.

$$\nabla fg = g(\nabla f), \quad \text{but } \nabla(fg) = f(\nabla g) + g(\nabla f).$$

We denote our domain, the 2-torus, by \mathbb{T}^2 . We write $L^p(\mathbb{T}^2)$, $W^{k,p}(\mathbb{T}^2)$ and $H^k(\mathbb{T}^2)$ for the Lebesgue, Sobolev and Bessel potential spaces on \mathbb{T}^2 with integrability and smoothness exponents p, k , where more information on periodic spaces can be found in [34, Section 3]. We note that if k is an integer, we equip $H^k(\mathbb{T}^2)$ with the equivalent $W^{k,2}(\mathbb{T}^2)$ -inner product. We write $L^p(\mathbb{T}^2, \mathbb{R}^2)$, $W^{k,p}(\mathbb{T}^2, \mathbb{R}^2)$ and $H^k(\mathbb{T}^2, \mathbb{R}^2)$ for the corresponding spaces of vector fields and equip them with the direct sum norm and set for the special case $p = 2$

$$\|(f_1, f_2)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 = \|f_1\|_{L^2(\mathbb{T}^2)}^2 + \|f_2\|_{L^2(\mathbb{T}^2)}^2, \|(f_1, f_2)\|_{H^k(\mathbb{T}^2, \mathbb{R}^2)}^2 = \|f_1\|_{H^k(\mathbb{T}^2)}^2 + \|f_2\|_{H^k(\mathbb{T}^2)}^2$$

to preserve the Hilbert space structure. We write $\langle f, g \rangle$ for the dual pairing in $L^2(\mathbb{T}^2)$ and in $L^2(\mathbb{T}^2, \mathbb{R}^2)$ depending on f, g being functions or vector fields. If (S, ν) is a measure space and X is a Banach space, we write $L^p(S, \nu, X)$ for the Bochner space of strongly measurable, p -integrable, X -valued functions on (S, ν) . For details we refer to [25, Section 1]. If it is clear which measure is considered, we use also the notation $L^p(S, X)$ and if $S = [s, t]$ and ν the Lebesgue measure $L^p(s, t; X)$. Moreover, we write $C(0, T; X)$, $H^1(0, T; X)$, $C^\gamma(0, T; X)$ and $W^{\gamma,p}(0, T; X)$, for the space of continuous functions, first-order Sobolev space, Hölder and Sobolev–Slobodetskii space on $[0, T]$ with values in X , where we will only consider fractional exponents $\gamma \in (0, 1)$. The corresponding Hölder semi-norm is denoted by $[\cdot]_{\gamma, X}$ and for precise definitions of these spaces we refer to [3, Section 2]. If a Banach space X is considered with its weak or weak-* topology, we express this by writing X_w or X_{w*} respectively. Lastly we mention that we write $L_2(H_1, H_2)$ for the space of Hilbert–Schmidt operators between two Hilbert spaces H_1 and H_2 .

2. The deterministic thin-film equation

In this section we summarize the existence result for weak solutions to the deterministic thin-film equation in the special case of quadratic mobility

$$(2.1) \quad \partial_t v = -\operatorname{div}(v^2 \nabla v)$$

from [33]. Moreover, we show that the solutions can be chosen in a measurable way, which will be important later. We remark that in [33] solutions to (2.1) are constructed on a domain with Neumann boundary conditions, but the arguments translate verbatim to the periodic setting. First, we recall the definition of weak solutions to (2.1) from [33, Definition 3.1].

Definition 2.1. Let $q \in (2, \infty]$ and $T > 0$. A weak solution to the (deterministic) thin-film equation on $[0, T]$ with q' -regular non linearity is a tuple

$$(v, J) \in L^\infty(0, T; H^1(\mathbb{T}^2)) \cap H^1(0, T; W^{-1,q'}(\mathbb{T}^2)) \times L^2(0, T; L^{q'}(\mathbb{T}^2, \mathbb{R}^2)),$$

such that $\partial_t v = -\operatorname{div} J$ in $L^2(0, T; W^{-1,q'}(\mathbb{T}^2))$, $\nabla v \in L^3(\{v > 0\}, \mathbb{R}^2)$ and

$$(2.2) \quad \begin{aligned} \int_0^T \int_{\mathbb{T}^2} J \cdot \eta \, dx \, dt &= \int_0^T \int_{\{v(t)>0\}} |\nabla v|^2 \nabla v \cdot \eta \, dx \, dt + \int_0^T \int_{\{v(t)>0\}} v |\nabla v|^2 \operatorname{div} \eta \, dx \, dt \\ &+ 2 \int_0^T \int_{\{v(t)>0\}} v \nabla^T v D\eta \nabla v \, dx \, dt + \int_0^T \int_{\mathbb{T}^2} v^2 \nabla v \cdot \nabla \operatorname{div} \eta \, dx \, dt \end{aligned}$$

for all $\eta \in L^\infty(0, T; W^{2,\infty}(\mathbb{T}^2, \mathbb{R}^2))$.

Remark 2.2. By Rellich’s theorem, see [1, Theorem 6.3, p.168], and the Aubin–Lions lemma [35, Corollary 5] there is a compact embedding

$$L^\infty(0, T; H^1(\mathbb{T}^2)) \cap H^1(0, T; W^{-1,q'}(\mathbb{T}^2)) \hookrightarrow C(0, T; L^r(\mathbb{T}^2))$$

for any $r \in [1, \infty)$. In the following, we will always identify a weak solution to the thin-film equation with its $L^r(\mathbb{T}^2)$ -continuous version. By [7, Lemma II.5.9] this version is weakly continuous as a mapping with values in $H^1(\mathbb{T}^2)$.

The identity (2.2) is a weak formulation of $J = u^2 \nabla \Delta u$. The following existence statement is given in [33, Theorem 3.2], where we add some quantitative estimates which follow from the construction in [33] and are proved in detail in Appendix A.

Theorem 2.3. *Let $v_0 \in H^1(\mathbb{T}^2)$ be non negative, $q \in (2, \infty]$, $T > 0$ and $\alpha \in (-1, 0)$. Then there exists a weak solution (v, J) to the thin-film equation on $[0, T]$ with q' -regular non linearity and $v(0) = v_0$, which satisfies the following properties for universal constants $0 < C_\alpha, C_q < \infty$.*

(i) *We have for all $t \in [0, T]$ that*

$$\int_{\mathbb{T}^2} v(t, \cdot) dx = \int_{\mathbb{T}^2} v_0 dx \quad \text{and} \quad v(t, \cdot) \geq 0.$$

(ii) *It holds the energy estimate*

$$\sup_{0 \leq t \leq T} \|\nabla v(t)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)} \leq \|\nabla v_0\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}.$$

(iii) *It holds that*

$$\begin{aligned} & \|J\|_{L^2(0, T; L^{q'}(\mathbb{T}^2, \mathbb{R}^2))}^2 + C_q \|\nabla v(T)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 \left(\|\nabla v(T)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 + \left| \int_{\mathbb{T}^2} v_0 dx \right|^2 \right) \\ & \leq C_q \|\nabla v_0\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 \left(\|\nabla v_0\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 + \left| \int_{\mathbb{T}^2} v_0 dx \right|^2 \right). \end{aligned}$$

(iv) *We have the α -entropy estimate*

$$\int_{\mathbb{T}^2} G_\alpha(v(T, \cdot)) dx + \frac{1}{C_\alpha} \int_0^T \int_{\mathbb{T}^2} |Hv^{\frac{\alpha+3}{2}}|^2 + |\nabla v^{\frac{\alpha+3}{4}}|^4 dx dt \leq \int_{\mathbb{T}^2} G_\alpha(v_0) dx.$$

The following result can be proved along the lines of [33, Lemma 2.5, Proposition 2.6, Corollary 2.7, Theorem 3.2].

Proposition 2.4. *Let $q \in (2, \infty]$, $T > 0$ and $(v_n, J_n)_{n \in \mathbb{N}}$ be a sequence of non negative weak solutions to the deterministic thin-film equation on $[0, T]$ with q' -regular non linearity. Assume that there is an $\alpha \in (-1, 0)$ such that $v_n, J_n, v_n^{\frac{\alpha+3}{2}}$ and $v_n^{\frac{\alpha+3}{4}}$ are uniformly bounded in*

$$(2.3) \quad L^\infty(0, T; H^1(\mathbb{T}^2)), L^2(0, T; L^{q'}(\mathbb{T}^2, \mathbb{R}^2)), L^2(0, T; H^2(\mathbb{T}^2)), L^4(0, T; W^{1,4}(\mathbb{T}^2))$$

respectively. Then for a subsequence we have

- (i) $v_n \rightharpoonup^* v$ in $L^\infty(0, T; H^1(\mathbb{T}^2))$,
- (ii) $J_n \rightharpoonup J$ in $L^2(0, T; L^{q'}(\mathbb{T}^2, \mathbb{R}^2))$,
- (iii) $v_n^{\frac{\alpha+3}{2}} \rightharpoonup v^{\frac{\alpha+3}{2}}$ in $L^2(0, T; H^2(\mathbb{T}^2))$,
- (iv) $v_n^{\frac{\alpha+3}{4}} \rightharpoonup v^{\frac{\alpha+3}{4}}$ in $L^4(0, T; W^{1,4}(\mathbb{T}^2))$

and the limit (v, J) is a non-negative weak solution to the thin-film equation with q' -regular non-linearity.

Finally, we give proof to the existence of a measurable solution operator. To this end, we define the set $\mathcal{X}_{q,T}$ as the topological product of the spaces (2.3) equipped with the respective weak and weak-* topologies. Moreover, we write $B_X(r)$ for the ball in X centered at the origin with radius r , if X is a normed space.

Corollary 2.5. *Let $q \in (2, \infty]$, $T > 0$ and $\alpha \in (-1, 0)$. There is a Borel-measurable mapping*

$$(2.4) \quad \mathcal{S}_{\alpha,q,T} : \{v_0 \in H^1(\mathbb{T}^2) | v_0 \geq 0\} \rightarrow \mathcal{X}_{q,T}, \quad v_0 \mapsto (v, J, v^{\frac{\alpha+3}{2}}, v^{\frac{\alpha+3}{4}}),$$

which assigns to every initial value a weak solution to the thin-film equation on $[0, T]$, which satisfies the properties (i)-(iv) of Theorem 2.3.

Proof. We define for v_0 in the domain of (2.4) the set of all weak solutions to the stochastic thin-film equation with initial value v_0 and q' -regular non linearity satisfying (i)-(iv) from Theorem 2.3 together with its corresponding powers

by $\text{Sol}(v_0) \subset \mathcal{X}_{q,T}$. We write X_i for the i -th space in (2.3) and observe that if $\|v_0\|_{H^1(\mathbb{T}^2)} \leq n$ for some $n \in \mathbb{N}$ the a-priori bounds of Theorem 2.3 yield that

$$\text{Sol}(v_0) \subset \mathcal{X}_{q,T}^{(n)} := \prod_{i=1}^4 B_{X_i}(r_{i,n})$$

for suitably chosen $r_{i,n}$. We equip each $B_{X_i}(r_{i,n})$ again with the weak (weak-*) topology of the respective space X_i and $\mathcal{X}_{q,T}^{(n)}$ with the resulting product topology. We note that each $B_{X_i}(r_{i,n})$ is metrizable by the separability of the (pre-) dual of X_i , see [25, Proposition 1.2.29, Corollary 1.3.22] and consequently also the topological product $\mathcal{X}_{q,T}^{(n)}$. Moreover, $\mathcal{X}_{q,T}^{(n)}$ is compact as a consequence of Tychonoff's and the Banach–Alaoglu theorem and therefore in particular a Polish space. Let $(v_{0,j})_{j \in \mathbb{N}}$ be a sequence in

$$\{v_0 \in H^1(\mathbb{T}^2) \mid v_0 \geq 0, \|v_0\|_{H^1(\mathbb{T}^2)} \leq n\}$$

converging to $v_{0,*}$ in $H^1(\mathbb{T}^2)$ and

$$(2.5) \quad (v_j, J_j, v_j^{\frac{\alpha+3}{2}}, v_j^{\frac{\alpha+3}{4}}) \in \text{Sol}(v_{0,j}).$$

Then the measurable selection theorem as in [15, Corollary 103, p.506] yields a Borel-measurable solution map

$$(2.6) \quad \mathcal{S}_{\alpha,q,T}^{(n)} : \{v_0 \in H^1(\mathbb{T}^2) \mid v_0 \geq 0, \|v_0\|_{H^1(\mathbb{T}^2)} \leq n\} \rightarrow \mathcal{X}_{q,T}^{(n)}, \quad v_0 \mapsto (v, J, v^{\frac{\alpha+3}{2}}, v^{\frac{\alpha+3}{4}}) \in \text{Sol}(v_0),$$

if we can verify that a subsequence of

$$(v_j, J_j, v_j^{\frac{\alpha+3}{2}}, v_j^{\frac{\alpha+3}{4}})_{j \in \mathbb{N}}$$

converges to an element of $\text{Sol}(v_{0,*})$. Since (2.5) lies in $\mathcal{X}_{q,T}^{(n)}$, its components are uniformly bounded in (2.3). Therefore, we can apply Proposition 2.4 and obtain that

$$(v_j, J_j, v_j^{\frac{\alpha+3}{2}}, v_j^{\frac{\alpha+3}{4}}) \rightarrow (v, J, v^{\frac{\alpha+3}{2}}, v^{\frac{\alpha+3}{4}})$$

for a subsequence in $\mathcal{X}_{q,T}^{(n)}$, where (v, J) is a non negative weak solution to the thin-film equation with q' -regular non linearity. By [35, Corollary 5] we have $v_j \rightarrow v$ in $C(0, T; L^2(\mathbb{T}^2))$ and in particular $v_j(0) \rightarrow v(0)$ in $L^2(\mathbb{T}^2)$. Consequently we must have $v(0) = v_{0,*}$. By lower semi-continuity of the norm with respect to weak and weak-* convergence we deduce that (v, J) satisfies all the properties (i)-(iv) of Theorem 2.3 and therefore

$$(v, J, v^{\frac{\alpha+3}{2}}, v^{\frac{\alpha+3}{4}}) \in \text{Sol}(v_{0,*}).$$

Hence, the measurable selection theorem indeed yields a Borel measurable map (2.6). Finally, we set $\mathcal{S}_{\alpha,q,T} v_0 = \mathcal{S}_{\alpha,q,T}^{(n)} v_0$, if $n - 1 \leq \|v_0\| < n$. Since balls in $H^1(\mathbb{T}^2)$ are Borel sets, $\mathcal{S}_{\alpha,q,T}$ has the desired properties. \square

3. A regularized linear Stratonovich SPDE on $H^1(\mathbb{T}^2)$

In this section we show that the regularized version of the stochastic part in (1.2)

$$(3.1) \quad dw_t = \epsilon \Delta w_t dt + \text{div}(w_t \circ dW_t)$$

is well-posed using the variational approach to SPDEs [30, Chapter 4]. A key ingredient to checking the sufficient conditions for well-posedness is the spatial isotropy condition on the noise (1.5). Throughout this section, we fix a filtered probability space $(\Omega, \mathfrak{A}, P)$ satisfying the usual conditions with a sequence of independent real-valued Brownian motions $(\beta^{(l)})_{l \in \mathbb{N}}$ and an $\epsilon \in (0, 1)$. The main statement of this section reads as follows.

Theorem 3.1. *Let $p \in [2, \infty)$, $T \in [0, \infty)$ and $w_0 \in L^p(\Omega, H^1(\mathbb{T}^2))$ be \mathfrak{F}_0 -measurable. Then there exists a unique continuous, adapted $H^1(\mathbb{T}^2)$ -valued process w such that $w \in L^2([0, T] \times \Omega, H^2(\mathbb{T}^2))$ and*

$$(3.2) \quad w(t) = w_0 + \int_0^t \epsilon \Delta w(s) + \frac{1}{2} \sum_{l=1}^{\infty} \lambda_l^2 \operatorname{div}(\operatorname{div}(w(s)\psi_l)\psi_l) ds + \sum_{l=1}^{\infty} \lambda_l \int_0^t \operatorname{div}(w(s)\psi_l) d\beta_s^{(l)}$$

for every $t \in [0, T]$. Moreover, w satisfies

$$(3.3) \quad E \left[\sup_{0 \leq t \leq T} \|w(t)\|_{H^1(\mathbb{T}^2)}^p \right] \lesssim_{p,T} E \left[\|w_0\|_{H^1(\mathbb{T}^2)}^p \right],$$

almost surely we have

$$(3.4) \quad \int_{\mathbb{T}^2} w(t) dx = \int_{\mathbb{T}^2} w_0 dx$$

and if $w_0 \geq 0$ also $w(t) \geq 0$ for all $t \in [0, T]$.

Remark 3.2. We convince ourselves that all the terms from (3.2) are well-defined. By (1.3) it holds

$$(3.5) \quad \sup_{|\alpha| \leq 2} \sup_{k \in \mathbb{Z}^2} \|\partial_\alpha \xi_k\|_{L^\infty(\mathbb{T}^2)} < \infty.$$

and therefore we have

$$(3.6) \quad \|\operatorname{div}(\operatorname{div}(w\psi_l)\psi_l)\|_{L^2(\mathbb{T}^2)} \lesssim \|w\|_{H^2(\mathbb{T}^2)} \quad \text{and} \quad \|\operatorname{div}(w\psi_l)\|_{H^1(\mathbb{T}^2)} \lesssim \|w\|_{H^2(\mathbb{T}^2)}$$

for every $w \in H^2(\mathbb{T}^2)$. Using the first estimate we derive that

$$E \left(\int_0^T \left\| \epsilon \Delta w(t) + \frac{1}{2} \sum_{l=1}^{\infty} \lambda_l^2 \operatorname{div}(\operatorname{div}(w\psi_l)\psi_l) \right\|_{L^2(\mathbb{T}^2)}^2 dt \right) \lesssim_\Delta \|w\|_{L^2([0,T] \times \Omega, H^2(\mathbb{T}^2))}^2,$$

and consequently the deterministic integral in (3.2) exists almost surely as a Bochner integral in $L^2(\mathbb{T}^2)$. Using the second estimate from (3.6), one derives by the martingale moment inequality and Itô's isometry that

$$\begin{aligned} E \left(\sup_{0 \leq t \leq T} \left\| \sum_{l=n}^m \lambda_l \int_0^t \operatorname{div}(w(s)\psi_l) d\beta_s^{(l)} \right\|_{H^1(\mathbb{T}^2)}^2 \right) &\lesssim E \left(\left\| \sum_{l=n}^m \lambda_l \int_0^T \operatorname{div}(w(t)\psi_l) d\beta_t^{(l)} \right\|_{H^1(\mathbb{T}^2)}^2 \right) \\ &= E \left[\sum_{l=n}^m \lambda_l^2 \int_0^T \|\operatorname{div}(w(t)\psi_l)\|_{H^1(\mathbb{T}^2)}^2 dt \right] \\ &\lesssim \left(\sum_{l=n}^m \lambda_l^2 \right) \|w\|_{L^2([0,T] \times \Omega, H^2(\mathbb{T}^2))}^2 \end{aligned}$$

and the latter part converges to 0 as $n, m \rightarrow \infty$. Therefore, the series of stochastic integrals in (3.2) converges to a continuous square-integrable martingale in $H^1(\mathbb{T}^2)$.

In order to treat the equation (3.2) within the variational setting [30, Chapter 4], we introduce the operators

$$\begin{aligned} A^\epsilon : H^2(\mathbb{T}^2) &\rightarrow L^2(\mathbb{T}^2), \quad w \mapsto \epsilon \Delta w + \frac{1}{2} \sum_{l=1}^{\infty} \lambda_l^2 \operatorname{div}(\operatorname{div}(w\psi_l)\psi_l), \\ B : H^2(\mathbb{T}^2) &\rightarrow L_2(H^2(\mathbb{T}^2, \mathbb{R}^2), H^1(\mathbb{T}^2)), \quad w \mapsto \left[v \mapsto \sum_{l=1}^{\infty} \lambda_l (v, \psi_l)_{H^2(\mathbb{T}^2, \mathbb{R}^2)} \operatorname{div}(w\psi_l) \right]. \end{aligned}$$

As in Remark 3.2 we conclude that the operators A^ϵ and B are well-defined, linear and bounded. In the following lemma we verify coercivity of (A^ϵ, B) . Its proof is similar to [18, Lemma A.3], but nevertheless contained to stress the necessity of assumption (1.5).

Lemma 3.3. *There exists a constant $C_\Lambda < \infty$ such that*

$$(3.7) \quad 2\langle A^\epsilon w, w \rangle + \sum_{l=1}^{\infty} \|B(w)[\psi_l]\|_{L^2(\mathbb{T}^2)}^2 \leq C_\Lambda \|w\|_{L^2(\mathbb{T}^2)}^2 - 2\epsilon \|w\|_{H^1(\mathbb{T}^2)}^2,$$

$$(3.8) \quad 2\langle \nabla A^\epsilon w, \nabla w \rangle + \sum_{l=1}^{\infty} \|\nabla B(w)[\psi_l]\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 \leq C_\Lambda \|w\|_{H^1(\mathbb{T}^2)}^2 - 2\epsilon \|\nabla w\|_{H^1(\mathbb{T}^2, \mathbb{R}^2)}^2$$

for all $w \in H^2(\mathbb{T}^2)$.

Proof. By continuity of the involved operators, it suffices to verify (3.7) and (3.8) for $w \in C^\infty(\mathbb{T}^2)$. We first observe that

$$\begin{aligned} \langle A^\epsilon w, w \rangle &= -\epsilon \|\nabla w\|_{L^2(\mathbb{T}^2)}^2 - \frac{1}{2} \sum_{l=1}^{\infty} \lambda_l^2 \langle \operatorname{div}(w\psi_l)\psi_l, \nabla w \rangle \\ &= -\epsilon \|\nabla w\|_{L^2(\mathbb{T}^2)}^2 - \frac{1}{2} \sum_{l=1}^{\infty} \lambda_l^2 \|\psi_l \cdot \nabla w\|_{L^2(\mathbb{T}^2)}^2 + \frac{1}{4} \sum_{l=1}^{\infty} \lambda_l^2 \langle w^2, \operatorname{div}(\operatorname{div}(\psi_l)\psi_l) \rangle, \end{aligned}$$

where we have used the identity $\frac{1}{2} \nabla w^2 = w \nabla w$ in the second line. Utilizing the same identity again, we obtain

$$\begin{aligned} \|B(w)[\psi_l]\|_{L^2(\mathbb{T}^2)}^2 &= \lambda_l^2 \|\operatorname{div}(w\psi_l)\|_{L^2(\mathbb{T}^2)}^2 \\ &= \lambda_l^2 (\|\psi_l \cdot \nabla w\|_{L^2(\mathbb{T}^2)}^2 + 2\langle w \nabla w, \operatorname{div}(\psi_l)\psi_l \rangle + \langle w^2, \operatorname{div}(\psi_l)^2 \rangle) \\ &= \lambda_l^2 (\|\psi_l \cdot \nabla w\|_{L^2(\mathbb{T}^2)}^2 - \langle w^2, \psi_l \cdot \nabla \operatorname{div}(\psi_l) \rangle). \end{aligned}$$

Considering the bound (3.5) we can calculate

$$\begin{aligned} 2\langle A^\epsilon w, w \rangle + \sum_{l=1}^{\infty} \|B(w)[\psi_l]\|_{L^2(\mathbb{T}^2)}^2 &= -2\epsilon \|\nabla w\|_{L^2(\mathbb{T}^2)}^2 + \frac{1}{2} \sum_{l=1}^{\infty} \lambda_l^2 \langle w^2, \operatorname{div}(\operatorname{div}(\psi_l)\psi_l) - \psi_l \cdot \nabla \operatorname{div}(\psi_l) \rangle \\ &\leq C_\Lambda \|w\|_{L^2(\mathbb{T}^2)}^2 - 2\epsilon \|\nabla w\|_{L^2(\mathbb{T}^2)}^2 \end{aligned}$$

for a suitable constant $C_\Lambda < \infty$. Enlarging C_Λ by 2 yields (3.7). For (3.8) we observe that

$$\langle \nabla A^\epsilon w, \nabla w \rangle = -\frac{1}{2} \sum_{l=1}^{\infty} \lambda_l^2 \langle \operatorname{div}(\operatorname{div}(w\psi_l)\psi_l), \Delta w \rangle - \epsilon \|\Delta w\|_{L^2(\mathbb{T}^2)}^2.$$

To further analyze the involved series, we set $\mu_k = \lambda_l$ in the situation of (1.5) and rewrite

$$-\frac{1}{2} \sum_{l=1}^{\infty} \lambda_l^2 \langle \operatorname{div} \cdot (\operatorname{div}(w\psi_l)\psi_l), \Delta w \rangle = -\frac{1}{2} \sum_{k \in \mathbb{Z}^2} \mu_k^2 \langle \operatorname{div}(\xi_k \nabla(w\xi_k)), \Delta w \rangle.$$

Before moving on, we notice that

$$(3.9) \quad \tilde{\xi}_j^2(x) \stackrel{(1.4)}{=} \begin{cases} 1 + \cos(4\pi jx), & j < 0, \\ 1, & j = 0, \\ 1 - \cos(4\pi jx), & j > 0 \end{cases}$$

and therefore

$$\xi_k^2(x, y) = \frac{\tilde{\xi}_{k_1}^2(x) \tilde{\xi}_{k_2}^2(y)}{1 + (2\pi|k|)^2 + (2\pi|k|)^4}$$

yields the bound

$$(3.10) \quad \sup_{|\alpha| \leq 4} \sup_{k \in \mathbb{Z}^2} \|\partial_\alpha \xi_k^2\|_{L^\infty(\mathbb{T}^2)} < \infty.$$

Using this estimate, the product rule for Δ , integration by parts, as well as the differential identities

$$(3.11) \quad \nabla(\nabla f \cdot \nabla g) = Hf \nabla g + Hg \nabla f \quad \text{and} \quad Hf \nabla f = \frac{1}{2} \nabla |\nabla f|^2$$

we calculate

$$\begin{aligned} \langle \operatorname{div}(\xi_k \nabla(w \xi_k)), \Delta w \rangle &= \langle \nabla \xi_k \cdot \nabla(w \xi_k) + \xi_k \Delta(w \xi_k), \Delta w \rangle \\ &= \left\langle \xi_k^2 \Delta w + \frac{3}{2} \nabla w \cdot \nabla \xi_k^2 + \frac{1}{2} w \Delta \xi_k^2, \Delta w \right\rangle \\ &= \|\xi_k \Delta w\|_{L^2(\mathbb{T}^2)}^2 - \frac{3}{2} \langle Hw \nabla \xi_k^2 + H \xi_k^2 \nabla w, \nabla w \rangle + \frac{1}{2} \left\langle \Delta \xi_k^2, \frac{1}{2} \Delta w^2 - \nabla w \cdot \nabla w \right\rangle \\ &\geq \|\xi_k \Delta w\|_{L^2(\mathbb{T}^2)}^2 + \frac{3}{4} \langle \Delta \xi_k^2, |\nabla w|^2 \rangle + \frac{1}{2} \langle \Delta^2 \xi_k^2, w^2 \rangle - C \|w\|_{H^1(\mathbb{T}^2)}^2 \\ &\geq \|\xi_k \Delta w\|_{L^2(\mathbb{T}^2)}^2 - C \|w\|_{H^1(\mathbb{T}^2)}^2. \end{aligned}$$

Here, we have enlarged the constant $C < \infty$ from the second last to the last line. Concerning the other summand in (3.8), we observe that by integration by parts

$$\|\nabla B(w)[(\xi_k, 0)]\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 = \mu_k^2 \|\nabla \partial_1(w \xi_k)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 = \mu_k^2 \langle \Delta(w \xi_k), \partial_{11}(w \xi_k) \rangle.$$

Rewriting the expression $\|\nabla B(w)[(0, \xi_k)]\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2$ analogously yields that

$$\|\nabla B(w)[(\xi_k, 0)]\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 + \|\nabla B(w)[(0, \xi_k)]\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 = \mu_k^2 \|\Delta(w \xi_k)\|_{L^2(\mathbb{T}^2)}^2.$$

Using again the product rule for Δ , the bound (3.5), integration by parts and the formulas from (3.11), we can estimate the latter term by

$$\begin{aligned} \|\Delta(w \xi_k)\|_{L^2(\mathbb{T}^2)}^2 &= \|\xi_k \Delta w + 2 \nabla w \cdot \nabla \xi_k + w \Delta \xi_k\|_{L^2(\mathbb{T}^2)}^2 \\ &= \|\xi_k \Delta w\|_{L^2(\mathbb{T}^2)}^2 + \|2 \nabla w \cdot \nabla \xi_k + w \Delta \xi_k\|_{L^2(\mathbb{T}^2)}^2 + 2 \langle \xi_k \Delta w, 2 \nabla w \cdot \nabla \xi_k + w \Delta \xi_k \rangle \\ &\leq \|\xi_k \Delta w\|_{L^2(\mathbb{T}^2)}^2 + C \|w\|_{H^1(\mathbb{T}^2)}^2 - 2 \langle \nabla w, \nabla[\xi_k w \Delta \xi_k + 2 \xi_k \nabla w \cdot \nabla \xi_k] \rangle \\ &= \|\xi_k \Delta w\|_{L^2(\mathbb{T}^2)}^2 + C \|w\|_{H^1(\mathbb{T}^2)}^2 - 2 \langle \nabla w, \xi_k \Delta \xi_k \nabla w + w \nabla[\xi_k \Delta \xi_k] \rangle \\ &\quad - 4 \langle \nabla w, (\nabla \xi_k \otimes \nabla \xi_k) \nabla w + \xi_k Hw \nabla \xi_k + \xi_k H \xi_k \nabla w \rangle \\ &\leq \|\xi_k \Delta w\|_{L^2(\mathbb{T}^2)}^2 + C \|w\|_{H^1(\mathbb{T}^2)}^2 + \langle w^2, \Delta(\xi_k \Delta \xi_k) \rangle + 2 \langle |\nabla w|^2, \operatorname{div}(\xi_k \nabla \xi_k) \rangle \\ &\leq \|\xi_k \Delta w\|_{L^2(\mathbb{T}^2)}^2 + C \|w\|_{H^1(\mathbb{T}^2)}^2. \end{aligned}$$

We enlarged again the constant $C < \infty$ from line to line. Moreover, in the last line we have employed that

$$\|\Delta(\xi_k \Delta \xi_k)\|_{L^\infty(\mathbb{T}^2)} = (2\pi |k|)^2 \|\Delta \xi_k^2\|_{L^\infty(\mathbb{T}^2)} \leq \frac{2(2\pi |k|)^2 (4\pi |k|)^2}{1 + (2\pi |k|)^2 + (2\pi |k|)^4} \leq 8$$

by $\Delta \xi_k = -(2\pi |k|^2) \xi_k$ and (3.9). Combining all the previous estimates we finally obtain that

$$2 \langle \nabla A^\varepsilon w, \nabla w \rangle + \sum_{l=1}^{\infty} \|\nabla B(w)[\psi_l]\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2$$

$$\begin{aligned} &\leq -2\epsilon \|\Delta w\|_{L^2(\mathbb{T}^2)}^2 - \sum_{k \in \mathbb{Z}^2} \mu_k^2 [\|\xi_k \Delta w\|_{L^2(\mathbb{T}^2)}^2 - C \|w\|_{H^1(\mathbb{T}^2)}^2] + \sum_{k \in \mathbb{Z}^2} \mu_k^2 [\|\xi_k \Delta w\|_{L^2(\mathbb{T}^2)}^2 + C \|w\|_{H^1(\mathbb{T}^2)}^2] \\ &\leq C_\Lambda \|w\|_{H^1(\mathbb{T}^2)}^2 - 2\epsilon \|\Delta w\|_{L^2(\mathbb{T}^2)}^2. \end{aligned}$$

We arrive at (3.8) by enlarging C_Λ by 2. □

Proof of Theorem 3.1. The existence and uniqueness assertion follows, if we verify the assumptions of [30, Theorem 4.2.4] on the couple (A^ϵ, B) considered on the Gelfand triple

$$H^2(\mathbb{T}^2) \subset H^1(\mathbb{T}^2) \subset L^2(\mathbb{T}^2).$$

Here we equip $H^2(\mathbb{T}^2)$ with the equivalent Bessel potential norm to ensure that the usual norm in $L^2(\mathbb{T}^2)$ coincides with the norm of the dual of $H^2(\mathbb{T}^2)$ under the pairing in $H^1(\mathbb{T}^2)$, for details see Appendix B. Hemicontinuity and boundedness of A^ϵ follow from $A^\epsilon \in L(H^2(\mathbb{T}^2), L^2(\mathbb{T}^2))$. Coercivity is obtained by adding (3.7) and (3.8) together. By linearity, coercivity implies weak monotonicity. The proof of (3.3) translates verbatim from the one-dimensional case [18, Proposition A.2] and (3.4) follows from testing (3.2) with $\mathbb{1}_{\mathbb{T}^2}$. The claim regarding non negativity of w is a consequence of the maximum principle for second-order parabolic SPDEs [28, Theorem 4.3], which holds by analogous reasoning also on \mathbb{T}^2 . □

4. Time discretization scheme with degenerate limit

In this section we fix $N \in \mathbb{N}$. The goal of this section is to construct for a given end time T and an initial value u_0 a weak martingale solution to the split-up problem

$$(4.1) \quad \begin{cases} u(t) = v(2(t - j\delta) + j\delta) & j\delta \leq t < (j + \frac{1}{2})\delta, \\ u(t) = w(2(t - (j + \frac{1}{2})\delta) + j\delta) & (j + \frac{1}{2})\delta \leq t < (j + 1)\delta, \\ \partial_t v = -\operatorname{div}(v^2 \nabla \Delta v) & \text{on } [j\delta, (j + 1)\delta), \\ dw_t = \operatorname{div}(w_t \circ dW_t) & \text{on } [j\delta, (j + 1)\delta), \end{cases}$$

where $\delta = \frac{T}{N+1}$ and $j \in \{0, \dots, N\}$. Starting at the initial value u_0 the process $u(t)$ satisfies alternately the deterministic thin-film equation and the purely stochastic equation (1.14) on time intervals of length $\frac{\delta}{2}$ and yields thus a time splitting scheme for the stochastic thin-film equation (1.2). During the construction we derive bounds which are uniform in N , and will be important in the final section, where we take the time step limit $N \rightarrow \infty$ to construct a solution to the original problem. We refer the interested reader for more information on the time-splitting procedure to [23]. The main statement of this section is the following.

Theorem 4.1. *Let $T \in (0, \infty)$, $q \in (2, \infty)$ and $\alpha \in (-1, 0)$. We assume that u_0 is a non negative random variable in $H^1(\mathbb{T}^2)$ and set $R^{(k)} = \{k - 1 \leq \|u_0\|_{H^1(\mathbb{T}^2)} < k\}$ and $u_0^{(k)} = \mathbb{1}_{R^{(k)}} u_0$ for every $k \in \mathbb{N}$. Then there exists a probability space $(\tilde{\Omega}, \tilde{\mathfrak{A}}, \tilde{P})$ with a filtration $\tilde{\mathfrak{F}}$ satisfying the usual conditions, a family of independent Brownian motions $(\tilde{\beta}^{(l)})_{l \in \mathbb{N}}$, random variables $\mathbb{1}_{\tilde{R}^{(k)}}$, $H_w^1(\mathbb{T}^2)$ -continuous processes $\tilde{u}^{(k)}$ and $L^2(0, T; L^{q'}(\mathbb{T}^2))$ -valued random variables $\tilde{J}^{(k)}$ for $k \in \mathbb{N}$, such that $\tilde{u}^{(k)}$, $\tilde{J}^{(k)}$ and the processes $\tilde{v}^{(k)}$ and $\tilde{w}^{(k)}$ defined by*

$$\begin{cases} \tilde{u}^{(k)}(t) = \tilde{v}^{(k)}(2(t - j\delta) + j\delta), & j\delta \leq t < (j + \frac{1}{2})\delta, \\ \tilde{u}^{(k)}(t) = \tilde{w}^{(k)}(2(t - (j + \frac{1}{2})\delta) + j\delta), & (j + \frac{1}{2})\delta \leq t < (j + 1)\delta, \end{cases}$$

satisfy the following.

- (i) *The sequence $(\mathbb{1}_{\tilde{R}^{(k)}}, \tilde{u}^{(k)}(0))_{k \in \mathbb{N}}$ has the same distribution as $(\mathbb{1}_{R^{(k)}}, u_0^{(k)})_{k \in \mathbb{N}}$, in particular we have that $\sum_{k=1}^\infty \tilde{u}^{(k)}(0) \sim u_0$. Moreover, $\tilde{u}^{(k)}$ and $\tilde{J}^{(k)}$ are \tilde{P} -almost surely zero outside of the set $\tilde{R}^{(k)}$.*
- (ii) *$\tilde{u}^{(k)}(t)$ and $\tilde{J}^{(k)}|_{[0,t]}$ are $\tilde{\mathfrak{F}}_t$ -measurable as random variables in $H^1(\mathbb{T}^2)$ and $L^2(0, t; L^{q'}(\mathbb{T}^2))$ for every $t \in [0, T]$ and $k \in \mathbb{N}$.*
- (iii) *The tuples $(\tilde{v}^{(k)}, \tilde{J}^{(k)})$ are \tilde{P} -almost surely solutions to the deterministic thin-film equation on $[j\delta, (j + 1)\delta)$ satisfying property (iv) from Theorem 2.3 with initial value $\tilde{u}^{(k)}(j\delta)$ for every $j = 0, \dots, N$.*

(iv) For $k \in \mathbb{N}$, $\varphi \in H^1(\mathbb{T}^2)$ and $t \in [j\delta, (j+1)\delta)$ we have that

$$\langle \tilde{w}^{(k)}(t), \varphi \rangle - \langle \tilde{w}^{(k)}(j\delta), \varphi \rangle = \frac{1}{2} \sum_{l=1}^{\infty} \lambda_l^2 \int_{j\delta}^t \langle \operatorname{div}(\operatorname{div}(\tilde{w}^{(k)}(s)\psi_l)\psi_l), \varphi \rangle ds + \sum_{l=1}^{\infty} \lambda_l \int_{j\delta}^t \langle \operatorname{div}(\tilde{w}^{(k)}(s)\psi_l), \varphi \rangle d\beta_s^{(l)}.$$

(v) For every $k \in \mathbb{N}$, $p \in (0, \infty)$ we have

$$\begin{aligned} \tilde{E} \left[\sup_{0 \leq t \leq T} \|\tilde{u}^{(k)}\|_{H^1(\mathbb{T}^2)}^p \right] &\lesssim_{\Lambda, p, T} E \left[\|u_0^{(k)}\|_{H^1(\mathbb{T}^2)}^p \right], \\ \tilde{E} \left[\|\tilde{J}^{(k)}\|_{L^2(0, T; L^{q'}(\mathbb{T}^2))}^{\frac{p}{2}} \right] &\lesssim_{\Lambda, p, q, T} E \left[\|u_0^{(k)}\|_{H^1(\mathbb{T}^2)}^p \right]. \end{aligned}$$

(vi) Moreover, for any $\gamma \in (0, \frac{1}{2})$ and $K \in (1, \infty)$ it holds

$$\tilde{P}(\{ \|\tilde{u}^{(k)}\|_{\gamma, W^{-1, q'}(\mathbb{T}^2)} > K \}) \lesssim_{\Lambda, q, \gamma, T} \frac{1 + E[\|u_0^{(k)}\|_{H^1(\mathbb{T}^2)}^2]}{K}.$$

4.1. Construction and analysis of a regularized scheme

Let $u_0 \in L^\infty(\Omega, H^1(\mathbb{T}^2))$ be non negative. Up to extension and completion of the probability space we can assume that there exists a filtration \mathfrak{F} satisfying the usual conditions with a family of independent Brownian motions $(\beta^{(l)})_{l \in \mathbb{N}}$ such that u_0 is \mathfrak{F}_0 -measurable.

Remark 4.2. The construction with initial value u_0 within this subsection will in Section 4.2 be applied to each of the cut-off parts $u_0^{(k)}$ from Theorem 4.1. This justifies the strong assumption $u_0 \in L^\infty(\Omega, H^1(\mathbb{T}^2))$ here.

We fix for the rest of this subsection also $T \in (0, \infty)$, $q \in (2, \infty)$, $\alpha \in (-1, 0)$, $\epsilon \in (0, 1)$ and apply the operator $S_{\alpha, q, \delta}$ from Corollary 2.5 to the initial value u_0 . We define $v_\epsilon|_{[0, \delta)}$, $J_\epsilon|_{[0, \delta)}$ as the version of the solution which is in $C(0, \delta; L^2(\mathbb{T}^2))$ and in particular continuous in $H_w^1(\mathbb{T}^2)$, see Remark 2.2. Moreover, we define $w_\epsilon|_{[0, \delta)}$ as the solution to (3.2) with initial value $\lim_{t \nearrow \delta} v_\epsilon(t)$. Notice that since $v_\epsilon|_{[0, \delta)}$ fulfills the properties (i) and (ii) of Theorem 2.3, we have

$$E \left[\left\| \lim_{t \nearrow \delta} v_\epsilon \right\|_{H^1(\mathbb{T}^2)}^p \right] \lesssim E \left[\|u_0\|_{H^1(\mathbb{T}^2)}^p \right]$$

for any $p \in [2, \infty)$, and therefore Theorem 3.1 is indeed applicable and yields a non-negative solution $w_\epsilon|_{[0, \delta)}$. In particular, the terminal value $\lim_{t \nearrow \delta} w_\epsilon(t)$ lies again in $L^p(\Omega, H^1(\mathbb{T}^2))$. We repeat this and obtain inductively weak solutions $v|_{[j\delta, (j+1)\delta)}$ to (2.1) and variational solutions $w_\epsilon|_{[j\delta, (j+1)\delta)}$ to (3.1) for $j \in \{1, \dots, N\}$. Finally, we define the $H_w^1(\mathbb{T}^2)$ -continuous, adapted process

$$u_\epsilon(t) = \begin{cases} v_\epsilon(2(t - j\delta) + j\delta), & j\delta \leq t < (j + \frac{1}{2})\delta, \\ w_\epsilon(2(t - (j + \frac{1}{2})\delta) + j\delta), & (j + \frac{1}{2})\delta \leq t < (j + 1)\delta, \end{cases}$$

for $t \in [0, T)$. We note that we set for the final time $u_\epsilon(T) = \lim_{t \nearrow \delta} w_\epsilon(t)$. The divergence form of (2.1), (3.1), and an application of Itô's formula yield the following estimates along the whole time-splitting scheme.

Lemma 4.3. *It holds almost surely that*

$$(4.2) \quad \int_{\mathbb{T}^2} u_\epsilon(t) dx = \int_{\mathbb{T}^2} u_0 dx.$$

for all $t \in [0, T]$. Moreover, we have additionally

$$(4.3) \quad E \left[\sup_{0 \leq t \leq T} \|u_\epsilon\|_{H^1(\mathbb{T}^2)}^p \right] \lesssim_{\Lambda, p, T} E \left[\|u_0\|_{H^1(\mathbb{T}^2)}^p \right]$$

for $p \in (0, \infty)$.

Proof. The equality (4.2) follows from its respective counterparts from Theorem 2.3(i) and (3.4). Next, we apply Itô's formula to the composition of the functional $\|\nabla \cdot\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2$ with the process w_ϵ , which yields that

$$(4.4) \quad \begin{aligned} \|\nabla w_\epsilon(t)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 &= \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 + 2 \int_{j\delta}^t \langle \nabla w_\epsilon(s), \nabla A^\epsilon(w_\epsilon(s)) \rangle ds \\ &+ \sum_{l=1}^{\infty} \lambda_l \int_{j\delta}^t 2 \langle \nabla \operatorname{div}(w_\epsilon(s) \psi_l), \nabla w_\epsilon(s) \rangle d\beta_s^l + \sum_{l=1}^{\infty} \lambda_l^2 \int_{j\delta}^t \|\nabla \operatorname{div}(w_\epsilon(s) \psi_l)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 ds. \end{aligned}$$

for $t \in [j\delta, (j+1)\delta)$. A justification of the applicability of Itô's formula is given in Appendix C. As pointed out in (C.7), the martingale given by the series of stochastic integrals, which we denote by $M_{2,j}$, has quadratic variation

$$4 \sum_{l=1}^{\infty} \lambda_l^2 \int_{j\delta}^t \langle \nabla \operatorname{div}(w_\epsilon(s) \psi_l), \nabla w_\epsilon(s) \rangle^2 ds.$$

Combining (4.4) with (3.8) we conclude that

$$\|\nabla w_\epsilon(t)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 - \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 - M_{2,j}(t) \lesssim_\Lambda \int_{j\delta}^t \|w_\epsilon(s)\|_{H^1(\mathbb{T}^2)}^2 ds$$

and for the endpoint $t = (j+1)\delta$

$$(4.5) \quad \|\nabla w_\epsilon((j+1)\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 - \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 - M_{2,j}((j+1)\delta) \lesssim_\Lambda \int_{j\delta}^{(j+1)\delta} \|w_\epsilon(s)\|_{H^1(\mathbb{T}^2)}^2 ds.$$

By Theorem 2.3(ii) we have

$$\|\nabla w_{N,\epsilon}(j\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R})}^2 \leq \|\nabla v_{N,\epsilon}(j\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R})}^2,$$

such that a telescoping sum argument yields

$$(4.6) \quad \|\nabla w_\epsilon(t)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 - \|\nabla u_0\|_{L^2(\mathbb{T}^2, \mathbb{R})}^2 - M_2(t) \lesssim_\Lambda \int_0^t \|w_\epsilon(s)\|_{H^1(\mathbb{T}^2)}^2 ds$$

for $t \in [0, T]$. The appearing process M_2 is defined by the sum of orthogonal martingales

$$M_2(t) = \sum_{k=0}^{j-1} M_{2,k}((k+1)\delta) + M_{2,j}(t), \quad t \in [j\delta, (j+1)\delta)$$

and has therefore quadratic variation

$$4 \sum_{l=1}^{\infty} \lambda_l^2 \int_0^t \langle \nabla \operatorname{div}(w_\epsilon(s) \psi_l), \nabla w_\epsilon(s) \rangle^2 ds.$$

For $p \geq 2$ we deduce from (4.6) with help of the inequality $(a + b + c)^{\frac{p}{2}} \lesssim_p a^{\frac{p}{2}} + b^{\frac{p}{2}} + c^{\frac{p}{2}}$ and the Burkholder–Davis–Gundy inequality that

$$(4.7) \quad \begin{aligned} &E \left[\sup_{0 \leq s \leq t} \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^p \right] - C_p E \left[\|\nabla u_0\|_{L^2(\mathbb{T}^2, \mathbb{R})}^p \right] \\ &\lesssim_{\Lambda, p} E \left[\left(\int_0^t \|w_\epsilon(s)\|_{H^1(\mathbb{T}^2)}^2 ds \right)^{\frac{p}{2}} + \left(\int_0^t \langle \nabla \operatorname{div}(w_\epsilon(s) \psi_l), \nabla w_\epsilon(s) \rangle^2 ds \right)^{\frac{p}{4}} \right]. \end{aligned}$$

To estimate the latter expression we observe that

$$\nabla \operatorname{div}(w \psi_l) = H w \psi_l + D \psi_l \nabla w + w \nabla \operatorname{div} \psi_l + \operatorname{div}(\psi_l) \nabla w$$

and due to (3.5) and (3.11) consequently

$$(4.8) \quad \left| \langle \nabla \operatorname{div}(w\psi_l), \nabla w \rangle \right| \lesssim \|\nabla w\|_{L^2(\mathbb{T}^2, \mathbb{R})} \|w\|_{H^1(\mathbb{T}^2)}$$

for $w \in H^2(\mathbb{T}^2)$. We conclude with help of Young's inequality that

$$\begin{aligned} & E \left[\left(\int_0^t \langle \nabla \operatorname{div}(w_\epsilon(s)\psi_l), \nabla w_\epsilon(s) \rangle^2 ds \right)^{\frac{p}{4}} \right] \\ & \lesssim E \left[\sup_{0 \leq s \leq t} \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^{\frac{p}{2}} \left(\int_0^t \|w_\epsilon(s)\|_{H^1(\mathbb{T}^2)}^2 ds \right)^{\frac{p}{4}} \right] \\ & \leq \frac{\kappa}{2} E \left[\sup_{0 \leq s \leq t} \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^p \right] + \frac{1}{2\kappa} E \left[\left(\int_0^t \|w_\epsilon(s)\|_{H^1(\mathbb{T}^2)}^2 ds \right)^{\frac{p}{2}} \right] \end{aligned}$$

for any $\kappa > 0$. An appropriate choice of κ and (4.7) yield that

$$\begin{aligned} & \frac{1}{2} E \left[\sup_{0 \leq s \leq t} \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^p \right] - C_p E \left[\|\nabla u_0\|_{L^2(\mathbb{T}^2, \mathbb{R})}^p \right] \\ & \lesssim_{\Lambda, p} E \left[\left(\int_0^t \|w_\epsilon(s)\|_{H^1(\mathbb{T}^2)}^2 ds \right)^{\frac{p}{2}} \right] \lesssim_{p, T} E \left[\int_0^t \|w_\epsilon(s)\|_{H^1(\mathbb{T}^2)}^p ds \right] \\ & \lesssim_p E \left[\left(\int_0^t \left(\int_{\mathbb{T}^2} u_0 dx \right)^p + \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^p ds \right) \right] \\ & \lesssim_T E \left[\|u_0\|_{L^2(\mathbb{T}^2)}^p \right] + \int_0^t E \left[\sup_{0 \leq \tau \leq s} \|\nabla w_\epsilon(\tau)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^p \right] ds. \end{aligned}$$

We additionally employed Jensen's and the Poincaré inequality here. Since $u_0 \in L^p(\Omega, H^1(\mathbb{T}^2))$, the monotone function

$$t \mapsto E \left[\sup_{0 \leq s \leq t} \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^p \right]$$

takes finite values by (3.3) and Theorem 2.3(i), (ii) and therefore an application of Grönwall's inequality yields

$$(4.9) \quad E \left[\sup_{0 \leq s \leq T} \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^p \right] \lesssim_{\Lambda, p, T} E \left[\|u_0\|_{H^1(\mathbb{T}^2)}^p \right]$$

In order to obtain the above inequality also for $p \in (0, 2)$ we observe that $\mathbb{1}_R u_\epsilon$ coincides with the process $u_{\epsilon, R}$ obtained by constructing the splitting scheme with initial value $\mathbb{1}_R u_0$ for $R \in \mathfrak{F}_0$. Indeed, from the properties in Theorem 2.3(i) we conclude that $\mathcal{S}_{\alpha, q, \delta}$ maps 0 to the solution which is 0 for all times. Consequently, we have

$$v_{\epsilon, R}|_{[0, \delta)} = \mathbb{1}_R v_\epsilon|_{[0, \delta)} \quad \text{and} \quad w_{\epsilon, R}(0) = \mathbb{1}_R w_\epsilon(0).$$

Therefore $w_{\epsilon, R}|_{[0, \delta)}$ and $\mathbb{1}_R w_\epsilon|_{[0, \delta)}$ are both solutions to (3.2) and have the same initial value such that we can conclude $w_{\epsilon}^{(R)}|_{[0, \delta)} = \mathbb{1}_R w_\epsilon|_{[0, \delta)}$. It is left to apply the uniqueness statement from Theorem 3.1 and repeat these arguments on $[j\delta, (j+1)\delta)$ for $j = 1, \dots, N$. Hence, applying (4.9) to $w_{\epsilon, R}$ with exponent $p = 2$ yields

$$E \left[\mathbb{1}_R \sup_{0 \leq s \leq T} \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 \right] \lesssim_{\Lambda, T} E \left[\mathbb{1}_R \|u_0\|_{H^1(\mathbb{T}^2)}^2 \right].$$

Since $R \in \mathfrak{F}_0$ was arbitrary, it follows that

$$E \left[\sup_{0 \leq s \leq T} \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 \middle| \mathfrak{F}_0 \right] \lesssim_{\Lambda, T} \|u_0\|_{H^1(\mathbb{T}^2)}^2.$$

For $p \in (0, 2)$ we can use Jensen's inequality to deduce that

$$E \left[\sup_{0 \leq s \leq T} \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^p \middle| \mathfrak{F}_0 \right] \leq E \left[\sup_{0 \leq s \leq T} \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 \middle| \mathfrak{F}_0 \right]^{\frac{p}{2}} \lesssim_{\Lambda, T} \|u_0\|_{H^1(\mathbb{T}^2)}^p$$

and it is left to take the expectation. Finally, we use Theorem 2.3(ii) to obtain (4.9) with w_ϵ replaced by u_ϵ which together with (4.2) implies (4.3). \square

Lemma 4.4. *We have*

$$E\left[\|J_\epsilon\|_{L^2(0,T;L^{q'}(\mathbb{T}^2))}^{\frac{p}{2}}\right] \lesssim_{\Lambda,p,q,T} E\left[\|u_0\|_{H^1(\mathbb{T}^2)}^p\right]$$

for $p \in (0, \infty)$.

Proof. We observe that as a consequence of Theorem 2.3(iii) and (4.2)

$$\begin{aligned} \|J_\epsilon\|_{L^2(j\delta,(j+1)\delta;L^{q'}(\mathbb{T}^2))}^2 &\lesssim_q \|\nabla v_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^4 - \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^4 \\ &\quad + \left(\int_{\mathbb{T}^2} u_0 dx\right)^2 \left(\|\nabla v_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 - \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2\right). \end{aligned}$$

Using that

$$\|J_\epsilon\|_{L^2(0,T;L^{q'}(\mathbb{T}^2))}^2 = \sum_{j=0}^N \|J_\epsilon\|_{L^2(j\delta,(j+1)\delta;L^{q'}(\mathbb{T}^2))}^2$$

we obtain the bound

$$\begin{aligned} \|J_\epsilon\|_{L^2(0,T;L^{q'}(\mathbb{T}^2))}^2 &\lesssim_q \sum_{j=0}^N \|\nabla v_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^4 - \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^4 \\ &\quad + \sum_{j=0}^N \left(\int_{\mathbb{T}^2} u_0 dx\right)^2 \left(\|\nabla v_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 - \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2\right). \end{aligned}$$

Applying the $\frac{p}{4}$ -th power and using that $(a_1 + \dots + a_4)^{\frac{p}{4}} \lesssim_p a_1^{\frac{p}{4}} + \dots + a_4^{\frac{p}{4}}$ we conclude

$$\begin{aligned} \|J_\epsilon\|_{L^2(0,T;L^{q'}(\mathbb{T}^2))}^{\frac{p}{2}} &\lesssim_{p,q} \left|0 \vee \left(\|\nabla u_0\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^4 - \|\nabla w_\epsilon(T-\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^4\right)\right|^{\frac{p}{4}} \\ &\quad + \left|\sum_{j=0}^{N-1} \|\nabla v_\epsilon((j+1)\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^4 - \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^4\right|^{\frac{p}{4}} \\ &\quad + \left(\int_{\mathbb{T}^2} u_0 dx\right)^{\frac{p}{2}} \left|0 \vee \left(\|\nabla u_0\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 - \|\nabla w_\epsilon(T-\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2\right)\right|^{\frac{p}{4}} \\ (4.10) \quad &\quad + \left|\left(\int_{\mathbb{T}^2} u_0 dx\right)^2 \sum_{j=0}^{N-1} \|\nabla v_\epsilon((j+1)\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 - \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2\right|^{\frac{p}{4}}. \end{aligned}$$

The expectation of the first and the third summand of the right-hand side of (4.10) can be each estimated by $E[\|u_0\|_{H^1(\mathbb{T}^2)}^p]$. To control also the second term we apply Itô's formula, see e.g. [27, Theorem 15.19], to the composition of $(\cdot)^2$ with the real-valued semimartingale (4.4) and obtain that

$$\begin{aligned} \|\nabla w_\epsilon(t)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^4 &= \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^4 + 2 \int_{j\delta}^t \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 dM_{2,j}(s) \\ &\quad + 4 \int_{j\delta}^t \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 \langle \nabla w_\epsilon(s), \nabla A^\epsilon(w_\epsilon(s)) \rangle ds \\ &\quad + 2 \int_{j\delta}^t \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 \sum_{l=1}^\infty \lambda_l^2 \int_{j\delta}^t \|\nabla \operatorname{div}(w_\epsilon(s)\psi_l)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 ds \end{aligned}$$

$$+ 4 \sum_{l=1}^{\infty} \lambda_l^2 \int_{j\delta}^t \langle \nabla \operatorname{div}(w_\epsilon(s)\psi_l), \nabla w_\epsilon(s) \rangle^2 ds.$$

Using (3.8), (4.8) we conclude for the endpoint $t = (j + 1)\delta$ that

$$\begin{aligned} & \|\nabla v_\epsilon((j + 1)\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^4 - \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^4 - 2 \int_{j\delta}^{(j+1)\delta} \|\nabla w_\epsilon(s)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 dM_{2,j}(s) \\ & \lesssim_{\Lambda} \int_{j\delta}^{(j+1)\delta} \|w_\epsilon(s)\|_{H^1(\mathbb{T}^2)}^4 ds. \end{aligned}$$

Summing up over j , taking the $\frac{p}{4}$ -power, applying the BDG-inequalities, (4.8) and (4.3) yields that

$$\begin{aligned} & E \left[\left| \sum_{j=0}^{N-1} \|\nabla v_\epsilon((j + 1)\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^4 - \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^4 \right|^{\frac{p}{4}} \right] \\ & \lesssim_{\Lambda, p} E \left[\left(\int_0^{T-\delta} \|w_\epsilon(s)\|_{H^1(\mathbb{T}^2)}^8 ds \right)^{\frac{p}{8}} + \left(\int_0^{T-\delta} \|w_\epsilon(s)\|_{H^1(\mathbb{T}^2)}^4 ds \right)^{\frac{p}{4}} \right] \\ & \lesssim_{p, T} E \left[\sup_{0 \leq t \leq T} \|w_\epsilon(t)\|^p \right] \lesssim_{\Lambda, p, T} E[\|u_0\|_{H^1(\mathbb{T}^2)}^p]. \end{aligned}$$

Similarly, by summing up over j in (4.5) and taking the power $\frac{p}{4}$ we obtain

$$\begin{aligned} & \left| \left(\int_{\mathbb{T}^2} u_0 dx \right)^2 \sum_{j=0}^{N-1} \|\nabla v_\epsilon((j + 1)\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 - \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 \right|^{\frac{p}{4}} \\ & \lesssim_{\Lambda, p} \left| \left(\int_{\mathbb{T}^2} u_0 dx \right)^2 M_2(T - \delta) \right|^{\frac{p}{4}} + \left(\left(\int_{\mathbb{T}^2} u_0 dx \right)^2 \int_0^{T-\delta} \|w_\epsilon(s)\|_{H^1(\mathbb{T}^2)}^2 ds \right)^{\frac{p}{4}} \end{aligned}$$

Taking the expectation, using the Burkholder–Davis–Gundy inequality, (4.8) and (4.3) yields the estimate

$$\begin{aligned} & E \left[\left| \left(\int_{\mathbb{T}^2} u_0 dx \right)^2 \sum_{j=0}^{N-1} \|\nabla v_\epsilon((j + 1)\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 - \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 \right|^{\frac{p}{4}} \right] \\ & \lesssim_{\Lambda, p} E \left[\left(\left(\int_{\mathbb{T}^2} u_0 dx \right)^4 \int_0^{T-\delta} \|w_\epsilon(s)\|_{H^1(\mathbb{T}^2)}^4 ds \right)^{\frac{p}{8}} + \left(\left(\int_{\mathbb{T}^2} u_0 dx \right)^2 \int_0^{T-\delta} \|w_\epsilon(s)\|_{H^1(\mathbb{T}^2)}^2 ds \right)^{\frac{p}{4}} \right] \\ & \lesssim_{p, T} \sqrt{E \left[\left(\int_{\mathbb{T}^2} u_0 dx \right)^p \right]} E \left[\sup_{0 \leq t \leq T} \|w_\epsilon(t)\|_{H^1(\mathbb{T}^2)}^p \right] \lesssim_{\Lambda, p, T} E[\|u_0\|_{H^1(\mathbb{T}^2)}^p]. \end{aligned}$$

Finally, taking the expectation of (4.10) and using the estimates on the individual summands yields the claim. \square

We also show tail estimates of the powers of v_ϵ in their respective space. We note that the obtained bound depends on N and will therefore be improved to a bound, which is uniform in N after letting $\epsilon \searrow 0$.

Lemma 4.5. *We have for $K \in (1, \infty)$ the estimate*

$$P(\{\|v_\epsilon\|_{L^2(0, T; H^2(\mathbb{T}^2))}^{\frac{\alpha+3}{2}} + \|v_\epsilon\|_{L^4(0, T; W^{1,4}(\mathbb{T}^2))}^{\frac{\alpha+3}{4}} > K\}) \lesssim_{\Lambda, \alpha, T, N} \frac{E[\|u_0\|_{H^1(\mathbb{T}^2)}^{\alpha+1}] + E[\|u_0\|_{H^1(\mathbb{T}^2)}^2]}{K^{\frac{2}{\alpha+3}}}.$$

Proof. As a consequence of Theorem 2.3(iv), (1.15) and Hölder’s inequality we have

$$\int_{j\delta}^{(j+1)\delta} \int_{\mathbb{T}^2} |Hv_\epsilon^{\frac{\alpha+3}{2}}|^2 + |\nabla v_\epsilon^{\frac{\alpha+3}{4}}|^4 dx dt$$

$$\begin{aligned} &\lesssim \int_{\mathbb{T}^2} G_\alpha(v_\epsilon(j\delta)) - G_\alpha(w_\epsilon(j\delta)) dx \\ &\lesssim_\alpha \|v_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2)}^{\alpha+1} + \|v_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2)} + \|w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2)}^{\alpha+1} + \|w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2)}. \end{aligned}$$

Summing up over j and taking the expectation yields that

$$(4.11) \quad E \left[\int_0^T \int_{\mathbb{T}^2} |Hv_\epsilon^{\frac{\alpha+3}{2}}|^2 + |\nabla v_\epsilon^{\frac{\alpha+3}{4}}|^4 dx dt \right] \lesssim_{\alpha,N} E \left[\sup_{0 \leq t \leq T} \|u_\epsilon(t)\|_{H^1(\mathbb{T}^2)}^{\alpha+1} + \sup_{0 \leq t \leq T} \|u_\epsilon(t)\|_{H^1(\mathbb{T}^2)} \right].$$

Moreover, by the Sobolev embedding theorem we have the estimate

$$\int_0^T \int_{\mathbb{T}^2} |v_\epsilon^{\frac{\alpha+3}{2}}(t)|^2 dx dt = \int_0^T \int_{\mathbb{T}^2} |v_\epsilon^{\frac{\alpha+3}{4}}(t)|^4 dx dt \lesssim_\alpha \int_0^T \|v_\epsilon\|_{H^1(\mathbb{T}^2)}^{\alpha+3} dt,$$

which implies by taking the $\frac{2}{\alpha+3}$ -th power and the expectation that

$$(4.12) \quad E \left[\left(\int_0^T \int_{\mathbb{T}^2} |v_\epsilon^{\frac{\alpha+3}{2}}(t)|^2 + |v_\epsilon^{\frac{\alpha+3}{4}}(t)|^4 dx dt \right)^{\frac{2}{\alpha+3}} \right] \lesssim_{\alpha,T} E \left[\sup_{0 \leq t \leq T} \|u_\epsilon(t)\|_{H^1(\mathbb{T}^2)}^2 \right].$$

Combining (4.11) and (4.12) with Chebyshev's inequality yields respectively that

$$\begin{aligned} P \left(\left\{ \int_0^T \int_{\mathbb{T}^2} |Hv_\epsilon^{\frac{\alpha+3}{2}}|^2 + |\nabla v_\epsilon^{\frac{\alpha+3}{4}}|^4 dx dt > K \right\} \right) &\lesssim_{\alpha,N} \frac{1}{K} E \left[\sup_{0 \leq t \leq T} \|u_\epsilon(t)\|_{H^1(\mathbb{T}^2)}^{\alpha+1} + \sup_{0 \leq t \leq T} \|u_\epsilon(t)\|_{H^1(\mathbb{T}^2)} \right], \\ P \left(\left\{ \int_0^T \int_{\mathbb{T}^2} |v_\epsilon^{\frac{\alpha+3}{2}}(t)|^2 + |v_\epsilon^{\frac{\alpha+3}{4}}(t)|^4 dx dt > K \right\} \right) &\lesssim_{\alpha,T} \frac{1}{K^{\frac{2}{\alpha+3}}} E \left[\sup_{0 \leq t \leq T} \|u_\epsilon(t)\|_{H^1(\mathbb{T}^2)}^2 \right]. \end{aligned}$$

Combining these estimates, the assumption $K \in (1, \infty)$ and the interpolation inequality

$$(4.13) \quad \|f\|_{H^2(\mathbb{T}^2)}^2 \lesssim \int_{\mathbb{T}^2} |f|^2 + |Hf|^2 dx, \quad f \in H^2(\mathbb{T}^2),$$

we obtain that

$$\begin{aligned} &P(\{ \|v_\epsilon^{\frac{\alpha+3}{2}}\|_{L^2(0,T;H^2(\mathbb{T}^2))}^2 + \|v_\epsilon^{\frac{\alpha+3}{4}}\|_{L^4(0,T;W^{1,4}(\mathbb{T}^2))}^4 > K \}) \\ &\lesssim_{\alpha,T,N} \frac{1}{K^{\frac{2}{\alpha+3}}} E \left[\sup_{0 \leq t \leq T} \|w_\epsilon(t)\|_{H^1(\mathbb{T}^2)}^{\alpha+1} + \sup_{0 \leq t \leq T} \|w_\epsilon(t)\|_{H^1(\mathbb{T}^2)}^2 \right]. \end{aligned}$$

It is left to apply (4.3) to conclude the claim. □

In the final part of the analysis of the approximate scheme, we show Hölder regularity in time of u_ϵ .

Lemma 4.6. *Let $\gamma \in (0, \frac{1}{2})$ and $K \in (1, \infty)$, then*

$$(4.14) \quad P(\{ \|u_\epsilon\|_{\gamma, W^{-1,q'}(\mathbb{T}^2)} > K \}) \lesssim_{\Lambda, q, \gamma, T} \frac{1 + E[\|u_0\|_{H^1(\mathbb{T}^2)}^2]}{K}.$$

Proof. We divide the proof into three steps.

Step 1 (Deterministic integrals). By Hölder's inequality we have

$$(4.15) \quad \left\| \int_s^t \operatorname{div} J_\epsilon(\tau) d\tau \right\|_{W^{-1,q'}(\mathbb{T}^2)} \leq \int_s^t \|\operatorname{div} J_\epsilon(\tau)\|_{W^{-1,q'}(\mathbb{T}^2)} d\tau \lesssim |t-s|^{\frac{1}{2}} \|J_\epsilon\|_{L^2(0,T,L^{q'}(\mathbb{T}^2))}$$

for any $s, t \in [0, T]$. Analogously, using that A^ϵ maps $H^1(\mathbb{T}^2)$ continuously to $H^{-1}(\mathbb{T}^2)$ due to (3.5) (with operator norm depending solely on Λ) we obtain that

$$(4.16) \quad \left\| \int_s^t A^\epsilon w_\epsilon(\tau) d\tau \right\|_{H^{-1}(\mathbb{T}^2)} \leq \int_s^t \|A^\epsilon w_\epsilon(\tau)\|_{H^{-1}(\mathbb{T}^2)} d\tau \lesssim_\Lambda |t-s| \sup_{0 \leq \tau \leq T} \|w_\epsilon(\tau)\|_{H^1(\mathbb{T}^2)}.$$

From the above inequalities, Lemma 4.4, and (4.3), we deduce that

$$(4.17) \quad \begin{aligned} P\left(\left[\int_0^\cdot \operatorname{div} J_\epsilon(s) ds\right]_{\frac{1}{2}, W^{-1, q'}(\mathbb{T}^2)} > K\right) &\lesssim_{\Lambda, q, T} \frac{E[\|u_0\|_{H^1(\mathbb{T}^2)}^2]}{K}, \\ P\left(\left[\int_0^\cdot A^\epsilon w_\epsilon(s) ds\right]_{1, H^{-1}(\mathbb{T}^2)} > K\right) &\lesssim_{\Lambda, T} \frac{E[\|u_0\|_{H^1(\mathbb{T}^2)}^2]}{K^2}. \end{aligned}$$

Step 2 (Stochastic integral). By [32, Theorem 3.2] we conclude that

$$P\left(\left[\sum_{l=1}^\infty \lambda_l \int_0^\cdot \operatorname{div}(w_\epsilon(s)\psi_l) d\beta_s^{(l)}\right]_{\gamma, L^2(\mathbb{T}^2)} > K \wedge \sup_{0 \leq t \leq T} \|B(w_\epsilon(t))\|_{L_2(H^2(\mathbb{T}^2, \mathbb{R}^2), L^2(\mathbb{T}^2))} \leq \sqrt{K}\right)$$

is dominated by $2e^{-C_\gamma T K}$, where $C_{\gamma, T} \in (0, \infty)$ is a suitable constant. We observe that due to (3.5), B maps continuously from $H^1(\mathbb{T}^2)$ to $L_2(H^2(\mathbb{T}^2, \mathbb{R}^2), L^2(\mathbb{T}^2))$ with operator norm only depending on Λ . Using additionally (4.3), we obtain the estimate

$$(4.18) \quad \begin{aligned} &P\left(\left[\sum_{l=1}^\infty \lambda_l \int_0^\cdot \operatorname{div}(w_\epsilon(s)\psi_l) d\beta_s^{(l)}\right]_{\gamma, L^2(\mathbb{T}^2)} > K\right) \\ &\leq 2e^{-C_\gamma T K} + \frac{E[\sup_{0 \leq t \leq T} \|B(w_\epsilon(t))\|_{H^1(\mathbb{T}^2)}^2]}{K} \lesssim_{\Lambda, \gamma, T} \frac{1 + E[\|u_0\|_{H^1(\mathbb{T}^2)}^2]}{K}. \end{aligned}$$

Step 3 (Combination of the estimates). Let $0 \leq s < t \leq T$. Splitting the process u_ϵ in its increments corresponding to v_ϵ and w_ϵ we obtain that

$$u_\epsilon(t) - u_\epsilon(s) = \int_{s'}^{t'} \operatorname{div} J_\epsilon(\tau) d\tau + \int_{s''}^{t''} A^\epsilon(w_\epsilon(\tau)) d\tau + \sum_{l=1}^\infty \lambda_l \int_{s''}^{t''} \operatorname{div}(w(\tau)\psi_l) d\beta_\tau^{(l)}$$

with an appropriate choice of $s', s'', t', t'' \in [0, T]$ satisfying in particular $|t' - s'|, |t'' - s''| < 2|t - s|$. Therefore, we can estimate $[u_\epsilon(t) - u_\epsilon(s)]_{\gamma, W^{-1, q'}(\mathbb{T}^2)}$ by

$$C_{q, T} \left(\left[\int_0^\cdot \operatorname{div} J_\epsilon(s) ds\right]_{\frac{1}{2}, W^{-1, q'}(\mathbb{T}^2)} + \left[\int_0^\cdot A^\epsilon w_\epsilon(s) ds\right]_{1, H^{-1}(\mathbb{T}^2)} + \left[\sum_{l=1}^\infty \lambda_l \int_0^\cdot \operatorname{div}(w_\epsilon(s)\psi_l) d\beta_s^{(l)}\right]_{\gamma, L^2(\mathbb{T}^2)} \right),$$

where $C_{q, T} < \infty$ is an appropriate constant. Invoking the estimates (4.17), (4.18) as well as the assumption $K \in (1, \infty)$ we conclude (4.14). \square

4.2. The vanishing viscosity limit

In this subsection we let T, q, α, u_0 and $R^{(k)}$ as in Theorem 4.1 and assume, as in the previous subsection, that u_0 is an \mathfrak{F}_0 -measurable random variable on a filtered probability space subject to the usual conditions with a family of independent Brownian motions $(\beta^{(l)})_{l \in \mathbb{N}}$. We let \mathcal{I} be a sequence converging to zero and apply for every $k \in \mathbb{N}$ and $\epsilon \in \mathcal{I}$ the construction from the previous subsection to the initial value $u_0^{(k)} = \mathbb{1}_{R^{(k)}} u_0$ and obtain a regularized splitting scheme consisting of $u_\epsilon^{(k)}, v_\epsilon^{(k)}, w_\epsilon^{(k)}, J_\epsilon^{(k)}$. We consider the sequence

$$(4.19) \quad ((\mathbb{1}_{R^{(l)}}, \beta^{(l)}, u_\epsilon^{(l)}, v_\epsilon^{(l)}, w_\epsilon^{(l)}, J_\epsilon^{(l)}, (v_\epsilon^{(l)})^{\frac{\alpha+3}{2}}, (v_\epsilon^{(l)})^{\frac{\alpha+3}{4}})_{l \in \mathbb{N}})_{\epsilon \in \mathcal{I}}$$

in the topological product space

$$(4.20) \quad \prod_{l=1}^\infty \mathbb{R} \times C([0, T]) \times C(0, T; L^2(\mathbb{T}^2)) \times L_{w^*}^\infty(0, T; H^1(\mathbb{T}^2)) \times L_{w^*}^\infty(0, T; H^1(\mathbb{T}^2)) \\ \times L_w^2(0, T; L^{q'}(\mathbb{T}^2)) \times L_w^2(0, T; H^2(\mathbb{T}^2)) \times L_w^4(0, T; W^{1,4}(\mathbb{T}^2)).$$

Proposition 4.7. *The sequence (4.19) is tight on (4.20)*

Proof. By Tychonoff’s theorem it is sufficient to show tightness of every component of (4.19) separately, so we fix an $l \in \mathbb{N}$. The distribution of $\mathbb{1}_{R^{(l)}}$ and $\beta^{(l)}$ is independent of ϵ and since the corresponding space is a Radon space, the sequences $(\mathbb{1}_{R^{(l)}})_{\epsilon \in \mathcal{I}}, (\beta^{(l)})_{\epsilon \in \mathcal{I}}$ are tight. Using (4.3) we deduce that

$$P(\{\|v_\epsilon^{(l)}\|_{L^\infty(0,T;H^1(\mathbb{T}^2))} > K\}) \lesssim_{\Lambda,T} \frac{E[\|u_0^{(l)}\|_{H^1(\mathbb{T}^2)}^2]}{K^2} \rightarrow 0$$

as $K \rightarrow \infty$ uniformly in ϵ such that tightness of $v_\epsilon^{(l)}$ is a consequence of the Banach–Alaoglu theorem. The components $w_\epsilon^{(l)}, J_\epsilon^{(l)}, (v_\epsilon^{(l)})^{\frac{\alpha+3}{2}}, (v_\epsilon^{(l)})^{\frac{\alpha+3}{4}}$ can be treated analogously using (4.3) and Lemmas 4.4 and 4.5. Lastly, we obtain from (4.3) and (4.14) that

$$P(\{\max\{\|u_\epsilon^{(l)}\|_{L^\infty(0,T;H^1(\mathbb{T}^2))}, [u_\epsilon^{(l)}]_{\gamma,W^{-1,q'}(\mathbb{T}^2)}\} > K\}) \lesssim_{\Lambda,q,\gamma,T} \frac{1 + E[\|u_0^{(l)}\|_{H^1(\mathbb{T}^2)}^2]}{K}$$

for $K \in (1, \infty)$ such that tightness of $u_\epsilon^{(l)}$ follows by [35, Theorem 5]. □

An application of [26, Theorem 2] yields that there exists for a subsequence, which we index again by \mathcal{I} , a complete probability space $(\tilde{\Omega}, \tilde{\mathfrak{A}}, \tilde{P})$ and a sequence of \mathfrak{B} -measurable random variables

$$((\mathbb{1}_{\tilde{R}^{(l)}}, \tilde{\beta}_\epsilon^{(l)}, \tilde{u}_\epsilon^{(l)}, \tilde{v}_\epsilon^{(l)}, \tilde{w}_\epsilon^{(l)}, \tilde{J}_\epsilon^{(l)}, \tilde{f}_\epsilon^{(l)}, \tilde{g}_\epsilon^{(l)})_{l \in \mathbb{N}})_{\epsilon \in \mathcal{I}}$$

with values in (4.20) such that

$$(4.21) \quad (\mathbb{1}_{\tilde{R}^{(l)}}, \tilde{\beta}_\epsilon^{(l)}, \tilde{u}_\epsilon^{(l)}, \tilde{v}_\epsilon^{(l)}, \tilde{w}_\epsilon^{(l)}, \tilde{J}_\epsilon^{(l)}, \tilde{f}_\epsilon^{(l)}, \tilde{g}_\epsilon^{(l)})_{l \in \mathbb{N}}$$

has the same distribution as

$$(\mathbb{1}_{R^{(l)}}, \beta^{(l)}, u_\epsilon^{(l)}, v_\epsilon^{(l)}, w_\epsilon^{(l)}, J_\epsilon^{(l)}, (v_\epsilon^{(l)})^{\frac{\alpha+3}{2}}, (v_\epsilon^{(l)})^{\frac{\alpha+3}{4}})_{l \in \mathbb{N}}$$

for every $\epsilon \in \mathcal{I}$. Moreover, as $\epsilon \searrow 0$, (4.21) converges to a \mathfrak{B} -measurable random variable

$$(\mathbb{1}_{\tilde{R}^{(l)}}, \tilde{\beta}^{(l)}, \tilde{u}^{(l)}, \tilde{v}^{(l)}, \tilde{w}^{(l)}, \tilde{J}^{(l)}, \tilde{f}^{(l)}, \tilde{g}^{(l)})_{l \in \mathbb{N}}$$

in (4.20).

Remark 4.8. In order to apply [26, Theorem 2] one needs to check that there exists a countable sequence of $[-1, 1]$ -valued continuous functions on (4.20) which separate the points. Such a sequence is straightforward to construct using point-evaluations for the spaces of continuous functions and separability of the respective (pre-) dual for the spaces equipped with weak (weak-*) topology, see [25, Proposition 1.2.29; Corollary 1.3.22].

Lemma 4.9. *The sets $(\tilde{R}^{(k)})_{k \in \mathbb{N}}$ form up to \tilde{P} -null sets a disjoint partition of $\tilde{\Omega}$. Moreover, the following holds \tilde{P} -almost surely for every $k \in \mathbb{N}$.*

(i) *The random variables*

$$\tilde{u}^{(k)}, \tilde{v}^{(k)}, \tilde{w}^{(k)}, \tilde{J}^{(k)}, \tilde{f}^{(k)} \text{ and } \tilde{g}^{(k)}$$

vanish outside of $\tilde{R}^{(k)}$.

(ii) *$\tilde{u}^{(k)}(t) \geq 0$ for all $t \in [0, T]$.*

(iii) *For almost all $t \in [0, T]$ we have*

$$(4.22) \quad \tilde{u}^{(k)}(t) = \begin{cases} \tilde{v}^{(k)}(2(t - j\delta) + j\delta), & j\delta \leq t < (j + \frac{1}{2})\delta, \\ \tilde{w}^{(k)}(2(t - (j + \frac{1}{2})\delta) + j\delta), & (j + \frac{1}{2})\delta \leq t < (j + 1)\delta. \end{cases}$$

(iv) *The tuples $(\tilde{v}^{(k)}, \tilde{J}^{(k)})$ are solutions to the deterministic thin-film equation on $[j\delta, (j + 1)\delta)$ satisfying property (iv) from Theorem 2.3 with initial value $\tilde{u}^{(k)}(j\delta)$ for every $j = 0, \dots, N$.*

$$(v) \quad \tilde{f}^{(k)} = (\tilde{v}^{(k)})^{\frac{\alpha+3}{2}} \text{ and } \tilde{g}^{(k)} = (\tilde{v}^{(k)})^{\frac{\alpha+3}{4}}.$$

Proof. For every $\epsilon \in \mathcal{I}$ we have

$$\tilde{E}[\mathbb{1}_{\tilde{R}_\epsilon^{(k_1)}} \mathbb{1}_{\tilde{R}_\epsilon^{(k_2)}}] = \delta_{k_1, k_2} P(R^{(k_1)}),$$

such that by letting $\epsilon \searrow 0$ we conclude the first part of the claim. Part (i) follows by letting $\epsilon \searrow 0$ in $\mathbb{1}_{\tilde{R}_\epsilon^{(k)}} \|\tilde{u}_\epsilon^{(k)}\|_{C(0, T; L^2(\mathbb{T}^2))} = 0$ and the same argument for the other random variables. Part (ii) is a consequence of $\tilde{u}_\epsilon^{(k)}(t) \geq 0$ together with conservation of this property under limits in $C(0, T; L^2(\mathbb{T}^2))$. Analogously, we deduce (iii) from the respective property in of $\tilde{u}_\epsilon^{(k)}$, $\tilde{v}_\epsilon^{(k)}$ and $\tilde{w}_\epsilon^{(k)}$. For (iv) and (v) we observe first that by measurability of $\mathcal{S}_{\alpha, q, \delta}$ we have

$$(4.23) \quad \mathcal{S}_{\alpha, q, \delta} \tilde{u}_\epsilon^{(k)}(j\delta) = (\tilde{v}_\epsilon^{(l)}|_{[j\delta, (j+1)\delta]}, \tilde{J}_\epsilon^{(k)}|_{[j\delta, (j+1)\delta]}, \tilde{f}_\epsilon^{(k)}|_{[j\delta, (j+1)\delta]}, \tilde{g}_\epsilon^{(k)}|_{[j\delta, (j+1)\delta]}).$$

In particular, we have $\tilde{f}_\epsilon^{(k)} = (\tilde{f}_\epsilon^{(k)})^{\frac{\alpha+3}{2}}$, $\tilde{g}_\epsilon^{(k)} = (\tilde{f}_\epsilon^{(k)})^{\frac{\alpha+3}{4}}$ and the right hand-side of (4.23) fulfills the properties stated in Theorem 2.3. If we let $\epsilon \searrow 0$ we deduce that the limit

$$(\tilde{v}^{(k)}|_{[j\delta, (j+1)\delta]}, \tilde{J}^{(k)}|_{[j\delta, (j+1)\delta]})$$

is a solution to the thin-film equation and that (v) holds true by Proposition 2.4. In light of 2.2 the initial value of (4.23) is indeed $\tilde{u}_\epsilon^{(k)}(j\delta)$. It is left to observe that property (iv) of Theorem 2.3 is preserved due to lower semi-continuity of the norm with respect to weak convergence. \square

By (4.22) we deduce that $\tilde{u}_\epsilon^{(k)}$ converges to $\tilde{u}^{(k)}$ also in $L^\infty_{w^*}(0, T; H^1(\mathbb{T}^2))$ and that $\tilde{u}^{(k)}$ is weakly continuous in $H^1(\mathbb{T}^2)$ again. Moreover, we identify in the following $\tilde{v}^{(k)}$ and $\tilde{w}^{(k)}$ with their versions such that (4.22) holds for all $t \in [0, T]$. We define $\tilde{\mathfrak{F}}$ as the augmentation of the filtration $\tilde{\mathfrak{G}}$ given by

$$\tilde{\mathfrak{G}}_t = \sigma(\{\mathbb{1}_{\tilde{R}^{(l)}}|_{[0, t]}, \tilde{J}^{(l)}|_{[0, t]} | l \in \mathbb{N}\} \cup \{\tilde{u}^{(l)}(s), \tilde{\beta}^{(l)}(s) | 0 \leq s \leq t, l \in \mathbb{N}\}),$$

where we consider $\tilde{J}^{(l)}|_{[0, t]}$ as a \mathfrak{B} -random variable in $L^2(0, t; L^{q'}(\mathbb{T}^2))$.

Lemma 4.10. *The processes $(\tilde{\beta}^{(l)})_{l \in \mathbb{N}}$ are a family of independent $\tilde{\mathfrak{F}}$ -Brownian motions. Moreover, we have for every $k \in \mathbb{N}$, $j \in \{0, \dots, N\}$ and $\varphi \in H^1(\mathbb{T}^2)$ that \tilde{P} -almost surely*

$$\langle \tilde{w}^{(k)}(t), \varphi \rangle - \langle \tilde{w}^{(k)}(j\delta), \varphi \rangle = \frac{1}{2} \sum_{l=1}^{\infty} \lambda_l^2 \int_{j\delta}^t \langle \operatorname{div}(\operatorname{div}(\tilde{w}^{(k)}(s)\psi_l)\psi_l), \varphi \rangle ds + \sum_{l=1}^{\infty} \lambda_l \int_{j\delta}^t \langle \operatorname{div}(\tilde{w}^{(k)}(s)\psi_l), \varphi \rangle d\beta_s^{(l)}$$

for all $t \in [j\delta, (j+1)\delta)$.

The proof of the lemma above is a simpler version of the proof of Theorem 5.12 and is therefore omitted. Finally, we observe that many of the estimates from the previous subsection carry over to their limit.

Proposition 4.11. *For every $k \in \mathbb{N}$ and $p \in (0, \infty)$ we have*

$$(4.24) \quad \begin{aligned} \tilde{E} \left[\sup_{0 \leq t \leq T} \|\tilde{u}^{(k)}\|_{H^1(\mathbb{T}^2)}^p \right] &\lesssim_{\Lambda, p, T} E \left[\|u_0^{(k)}\|_{H^1(\mathbb{T}^2)}^p \right], \\ \tilde{E} \left[\|\tilde{J}^{(k)}\|_{L^2(0, T; L^{q'}(\mathbb{T}^2))}^{\frac{p}{2}} \right] &\lesssim_{\Lambda, p, q, T} E \left[\|u_0^{(k)}\|_{H^1(\mathbb{T}^2)}^p \right]. \end{aligned}$$

Moreover, for any $\gamma \in (0, \frac{1}{2})$ and $K \in (1, \infty)$ it holds

$$\tilde{P}(\{[\tilde{u}^{(k)}]_{\gamma, W^{-1, q'}(\mathbb{T}^2)} > K\}) \lesssim_{\Lambda, q, \gamma, T} \frac{1 + E[\|u_0^{(k)}\|_{H^1(\mathbb{T}^2)}^2]}{K}.$$

Proof. The estimates (4.24) follow from lower semi-continuity of the norm with respect to weak (weak-*) convergence, (4.3), Lemma 4.4 and Fatou’s lemma. Moreover, we observe that

$$[\tilde{u}^{(k)}]_{\gamma, W^{-1, q'}(\mathbb{T}^2)} \leq \liminf_{\epsilon \searrow 0} [\tilde{u}_\epsilon^{(k)}]_{\gamma, W^{-1, q'}(\mathbb{T}^2)}$$

by convergence in $C(0, T; L^2(\mathbb{T}^2))$. Therefore, we have

$$\begin{aligned} P(\{[\tilde{u}^{(k)}]_{\gamma, W^{-1, q'}(\mathbb{T}^2)} > K\}) &\leq P\left(\left\{\liminf_{\epsilon \searrow 0} [\tilde{u}_\epsilon^{(k)}]_{\gamma, W^{-1, q'}(\mathbb{T}^2)} > K\right\}\right) \\ &\leq P\left(\liminf_{\epsilon \searrow 0} \{[\tilde{u}_\epsilon^{(k)}]_{\gamma, W^{-1, q'}(\mathbb{T}^2)} > K\}\right) \leq \liminf_{\epsilon \searrow 0} P(\{[\tilde{u}_\epsilon^{(k)}]_{\gamma, W^{-1, q'}(\mathbb{T}^2)} > K\}) \end{aligned}$$

and it is left to apply (4.14). □

Proof of Theorem 4.1. The limiting random variables $(\beta^{(l)})_{l \in \mathbb{N}}, \mathbb{1}_{\tilde{R}^{(k)}}, \tilde{u}^{(k)}$ and $\tilde{J}^{(k)}$ have the desired properties. Indeed, (i) is a consequence of

$$(\mathbb{1}_{\tilde{R}_\epsilon^{(k)}}, \tilde{u}_\epsilon^{(k)}(0))_{k \in \mathbb{N}} \sim (\mathbb{1}_{R^{(k)}}, u_0^{(k)})_{k \in \mathbb{N}}, \quad \epsilon \in \mathcal{I},$$

the convergence

$$(\mathbb{1}_{\tilde{R}_\epsilon^{(k)}}, \tilde{u}_\epsilon^{(k)}(0))_{k \in \mathbb{N}} \rightarrow (\mathbb{1}_{\tilde{R}^{(k)}}, \tilde{u}^{(k)}(0))_{k \in \mathbb{N}}$$

in $(\mathbb{R} \times L^2(\mathbb{T}^2))^\infty$ and Lemma 4.9(i). Part (ii) follows by the choice of $\tilde{\mathfrak{F}}$. Parts (iii), (iv), (v) and (vi) are the content of Lemma 4.9(iv), Lemma 4.10 and Proposition 4.11. □

5. Construction of solutions

Let finally μ, T, q, α as in Theorem 1.3. We apply Theorem 4.1 for every $N \in \mathbb{N}$ to a random variable which is distributed according to μ and obtain processes $u_N^{(k)}$, families of Brownian motions $\beta_N^{(l)}$ and random variables $\mathbb{1}_{R_N^{(k)}}, J_N^{(k)}$ for each $l, k \in \mathbb{N}$ satisfying the stated properties. We assume that these random variables are defined on the same probability space $(\Omega, \mathfrak{A}, P)$ with filtration \mathfrak{F} and moreover, that $\beta^{(l)} = \beta_N^{(l)}$ is independent of N . This does not influence the mathematical analysis since we analyze the solutions for each N separately and serves only notational convenience.

Remark 5.1. Alternatively, one can also apply the limiting procedure from Section 4.2 for all step numbers $N \in \mathbb{N}$ simultaneously to end up in the assumed situation. This, however, would lead to a notational mess in the previous section.

We remark also that we have dropped the \sim -notation since we want to pass to another probability space once more. For $k, N \in \mathbb{N}$ and we define $v_N^{(k)}$ and $w_N^{(k)}$ by

$$(5.1) \quad \begin{cases} u_N^{(k)}(t) = v_N^{(k)}(2(t - j\delta) + j\delta), & j\delta \leq t < (j + \frac{1}{2})\delta, \\ u_N^{(k)}(t) = w_N^{(k)}(2(t - (j + \frac{1}{2})\delta) + j\delta), & (j + \frac{1}{2})\delta \leq t < (j + 1)\delta, \end{cases}$$

where again $\delta = \delta(N) = \frac{T}{N+1}$.

5.1. Additional tightness properties

The approximate solutions $u_N^{(k)}, J_N^{(k)}$ satisfy the bounds from Theorem 4.1(v), (vi), which can as in Proposition 4.7 be used to derive tightness in suitable spaces.

Remark 5.2. In light of Theorem 4.1(i), the right-hand sides of the aforementioned bounds can be expressed in terms of the cut-off moments

$$(5.2) \quad v_{k, p} = \int \mathbb{1}_{\{k-1 \leq \|\cdot\|_{H^1(\mathbb{T}^2)} < k\}} \cdot \|\cdot\|_{H^1(\mathbb{T}^2)}^p d\mu$$

of the initial distribution μ . We remark that the notation (5.2) will be used during the remainder of this section.

In this subsection we provide an additional tightness property, which can be seen as the adaption of the compactness statement [33, Lemma 2.5] to our setting. Its proof relies on deriving a version of Lemma 4.5 with a uniform estimate in N and a simplified proof of [33, Lemma 2.5]. The former is based on a control of the entropy production along the stochastic dynamics. We point out that the simplification of the compactness proof is only possible due to the assumption $\alpha \in (-1, 0)$, which is less general than the situation in [33, Lemma 2.5].

Lemma 5.3. *It holds for every $k, N \in \mathbb{N}$ that*

$$E\left[\|(v_N^{(k)})^{\frac{\alpha+3}{2}}\|_{L^2(0,T;H^2(\mathbb{T}^2))}^2 + \|(v_N^{(k)})^{\frac{\alpha+3}{4}}\|_{L^4(0,T;W^{1,4}(\mathbb{T}^2))}^4\right] \lesssim_{\Lambda,\alpha,T} 1 + \nu_{k,\alpha+3}.$$

Proof. An application of Theorem 2.3(iv) yields the estimate

$$\begin{aligned} & \int_0^T \int_{\mathbb{T}^2} |H(v_N^{(k)})^{\frac{\alpha+3}{2}}|^2 + |\nabla(v_N^{(k)})^{\frac{\alpha+3}{4}}|^4 dx dt \\ & \lesssim_{\alpha} \sum_{j=0}^N \int_{\mathbb{T}^2} G_{\alpha}(v_N^{(k)}(j\delta)) - G_{\alpha}(w_N^{(k)}(j\delta)) dx \\ (5.3) \quad & = \int_{\mathbb{T}^2} G_{\alpha}(v_N^{(k)}(0)) - G_{\alpha}(w_N^{(k)}(N\delta)) dx + \sum_{j=0}^{N-1} \int_{\mathbb{T}^2} G_{\alpha}(v_N^{(k)}((j+1)\delta)) - G_{\alpha}(w_N^{(k)}(j\delta)) dx. \end{aligned}$$

The first summand can be estimated directly using the expression (1.15) by

$$C_{\alpha}(\|v_N^{(k)}(0)\|_{L^2(\mathbb{T}^2)}^{\alpha+1} + \|v_N^{(k)}(0)\|_{L^2(\mathbb{T}^2)} + \|w_N^{(k)}(0)\|_{L^2(\mathbb{T}^2)}^{\alpha+1} + \|w_N^{(k)}(0)\|_{L^2(\mathbb{T}^2)}) \lesssim_{\alpha} \sup_{0 \leq t \leq T} \|u_N^{(k)}(t)\|_{L^2(\mathbb{T}^2)} + 1.$$

Taking the expectation and employing Theorem 4.1(v) we obtain the estimate

$$(5.4) \quad E\left[\int_{\mathbb{T}^2} G_{\alpha}(v_N^{(k)}(0)) - G_{\alpha}(w_N^{(k)}(N\delta)) dx\right] \lesssim_{\Lambda,\alpha,T} 1 + \nu_{k,1}.$$

To estimate the second summand of the right-hand side of (5.3), we fix a function $\eta \in C^{\infty}(\mathbb{R})$ such that $0 \leq \eta \leq 1$, $\eta = 1$ on $[2, \infty)$ and $\eta = 0$ on $(-\infty, 1]$. We define smooth functions

$$\eta_{\kappa}(x) = \eta\left(\frac{x}{\kappa}\right), \quad G_{\alpha,\kappa}(x) = G_{\alpha}(x)\eta_{\kappa}(x).$$

for $\kappa > 0$. Correspondingly, we define the regularized functional

$$(5.5) \quad \phi_{\kappa} : L^2(\mathbb{T}^{\neq}) \rightarrow \mathbb{R}, \quad w \mapsto \int_{\mathbb{T}^2} G_{\alpha,\kappa}(w) dx$$

We observe that there is a constant $C_{\alpha,\kappa} < \infty$ such that

$$(5.6) \quad |G_{\alpha,\kappa}(x)| \leq C_{\alpha,\kappa}|x|^2, \quad |G'_{\alpha,\kappa}(x)| \leq C_{\alpha,\kappa}|x| \quad \text{and} \quad |G''_{\alpha,\kappa}(x)| \leq C_{\alpha,\kappa}$$

for each $x \in \mathbb{R}$. An application of Itô's formula to the composition of the functional ϕ_{κ} with the process $w_N^{(k)}$ satisfying the SPDE from Theorem 4.1(iv) yields that

$$\begin{aligned} (5.7) \quad \phi_{\kappa}(w_N^{(k)}(t)) &= \phi_{\kappa}(w_N^{(k)}(j\delta)) + \sum_{l=1}^{\infty} \lambda_l \int_{j\delta}^t \int_{\mathbb{T}^2} G'_{\alpha,\kappa}(w_N^{(k)}(s)) \operatorname{div}(w_N^{(k)}(s)\psi_l) dx d\beta^{(l)}(s) \\ &\quad - \frac{1}{2} \sum_{l=1}^{\infty} \int_{j\delta}^t \int_{\mathbb{T}^2} \lambda_l^2 G''_{\alpha,\kappa}(w_N^{(k)}(s)) [\operatorname{div}(w_N^{(k)}(s)\psi_l)] [\psi_l \cdot \nabla w_N^{(k)}(s)] dx ds \\ &\quad + \frac{1}{2} \sum_{l=1}^{\infty} \lambda_l^2 \int_{j\delta}^t \int_{\mathbb{T}^2} G''_{\alpha,\kappa}(w_N^{(k)}(s)) [\operatorname{div}(w_N^{(k)}(s)\psi_l)]^2 dx ds \end{aligned}$$

for $t \in [j\delta, (j + 1)\delta)$. For a justification of the applicability of Itô's formula see Appendix C. We note that as pointed out there, the above formula is also valid for the end-point $t = (j + 1)\delta$, but then the term on the left-hand side has to be replaced by $\phi_\kappa(v_N^{(k)}((j + 1)\delta))$. Due to the smoothness of ψ_l it holds

$$\begin{aligned} [\operatorname{div}(w\psi_l)][\psi_l \cdot \nabla w] &= [\operatorname{div}(w\psi_l)]^2 - [\operatorname{div}(w\psi_l)][w \operatorname{div} \psi_l] \\ &= [\operatorname{div}(w\psi_l)]^2 - [w \operatorname{div} \psi_l]^2 - w \nabla w \cdot [\psi_l \operatorname{div} \psi_l]. \end{aligned}$$

The derivative of $\zeta_\kappa(x) = \int_0^x y G''_{\alpha,\kappa}(y) dy$ is bounded such that an application of [9, Proposition 9.5] yields

$$\int_{\mathbb{T}^2} G''_{\alpha,\kappa}(w) w \nabla w \cdot [\psi_l \operatorname{div} \psi_l] dx = - \int_{\mathbb{T}^2} \zeta_\kappa(w) \operatorname{div}[\psi_l \operatorname{div} \psi_l] dx,$$

for $w \in H^1(\mathbb{T}^2)$. We also introduce the function $\theta_\kappa(x) = x^2 G''_{\alpha,\kappa}(x)$ and rewrite (5.7) using the previous identities as

$$\begin{aligned} \phi_\kappa(w_N^{(k)}(t)) &= \phi_\kappa(w_N^{(k)}(j\delta)) + \sum_{l=1}^\infty \lambda_l \int_{j\delta}^t \int_{\mathbb{T}^2} G'_{\alpha,\kappa}(w_N^{(k)}(s)) \operatorname{div}(w_N^{(k)}(s)\psi_l) dx d\beta^{(l)}(s) \\ &\quad + \frac{1}{2} \sum_{l=1}^\infty \lambda_l^2 \int_{j\delta}^t \int_{\mathbb{T}^2} \theta_\kappa(w_N^{(k)}(s)) (\operatorname{div} \psi_l)^2 dx ds \\ (5.8) \quad &\quad - \frac{1}{2} \sum_{l=1}^\infty \lambda_l^2 \int_{j\delta}^t \int_{\mathbb{T}^2} \zeta_\kappa(w_N^{(k)}(s)) \operatorname{div}[\psi_l \operatorname{div} \psi_l] dx ds. \end{aligned}$$

Using that

$$(5.9) \quad G''_{\alpha,\kappa}(x) = \frac{1}{\kappa^2} \eta''\left(\frac{x}{\kappa}\right) G_\alpha(x) + \frac{2}{\kappa} \eta'\left(\frac{x}{\kappa}\right) \left[\frac{x^\alpha}{\alpha} + r'_\alpha(x) \right] + \eta\left(\frac{x}{\kappa}\right) x^{\alpha-1}$$

and that η' and η'' vanish outside of $[1, 2]$ we deduce that $|\theta_\kappa(x)| \leq C_\alpha(1 + |x|)$. The same argument yields the bound $|\zeta_\kappa(x)| \leq C_\alpha(1 + |x|)$, indeed we can estimate for example

$$\left| \int_0^x \frac{y}{\kappa^2} \eta''\left(\frac{y}{\kappa}\right) G_\alpha(y) dy \right| \lesssim \frac{1}{\kappa} \int_0^{|x| \wedge 2\kappa} |G_\alpha(y)| dy \lesssim_\alpha \frac{|x| \wedge 2\kappa + (|x| \wedge 2\kappa)^2}{\kappa} \lesssim 1 + |x|.$$

Using (3.5) we obtain the estimates

$$\begin{aligned} \left| \frac{1}{2} \sum_{l=1}^\infty \lambda_l^2 \int_{j\delta}^{(j+1)\delta} \int_{\mathbb{T}^2} \theta_\kappa(w_N^{(k)}(s)) (\operatorname{div} \psi_l)^2 dx ds \right| &\lesssim_{\Lambda,\alpha} \delta \left(1 + \operatorname{ess\,sup}_{0 \leq t \leq T} \|w_N^{(k)}(t)\|_{L^2(\mathbb{T}^2)} \right), \\ \left| \frac{1}{2} \sum_{l=1}^\infty \lambda_l^2 \int_{j\delta}^{(j+1)\delta} \int_{\mathbb{T}^2} \zeta_\kappa(w_N^{(k)}(s)) \operatorname{div}[\psi_l \operatorname{div} \psi_l] dx ds \right| &\lesssim_{\Lambda,\alpha} \delta \left(1 + \operatorname{ess\,sup}_{0 \leq t \leq T} \|w_N^{(k)}(t)\|_{L^2(\mathbb{T}^2)} \right). \end{aligned}$$

For the series of stochastic integrals in (5.8) we observe that

$$\begin{aligned} E \left[\sum_{l=1}^\infty \lambda_l^2 \int_{j\delta}^{(j+1)\delta} \left(\int_{\mathbb{T}^2} G'_{\alpha,\kappa}(w_N^{(k)}(s)) \operatorname{div}(w_N^{(k)}(s)\psi_l) dx \right)^2 ds \right] \\ \lesssim_{\alpha,\kappa} E \left[\sum_{l=1}^\infty \lambda_l^2 \int_{j\delta}^{(j+1)\delta} (\|w_N^{(k)}(s)\|_{H^1(\mathbb{T}^2)})^2 ds \right] < \infty. \end{aligned}$$

We used (3.5) and that the function $G'_{\alpha,\kappa}$ is bounded in the first inequality and Theorem 4.1(v) for the second one. Hence, the series of stochastic integrals has integrable quadratic variation and is therefore a martingale. Therefore, taking the expectation of (5.8) with $t = (j + 1)\delta$, using the previous estimates, as well as Theorem 4.1(v) once more, yields that

$$E[\phi_\kappa(v_N^{(k)}((j + 1)\delta)) - \phi_\kappa(w_N^{(k)}(j\delta))] \lesssim_{\Lambda,\alpha} \delta \left(1 + E \left[\operatorname{ess\,sup}_{0 \leq t \leq T} \|w_N^{(k)}(t)\|_{L^2(\mathbb{T}^2)} \right] \right) \lesssim_{\Lambda,T} \delta(1 + \nu_{k,1}).$$

Finally, taking the expectation of (5.3) and using additionally (5.4), we end up with

$$E \left[\int_0^T \int_{\mathbb{T}^2} |H(v_N^{(k)})^{\frac{\alpha+3}{2}}|^2 + |\nabla(v_N^{(k)})^{\frac{\alpha+3}{4}}|^4 dx dt \right] \lesssim_{\Lambda, \alpha, T} 1 + \nu_{k,1}.$$

As in the proof of Lemma 4.5, we use that

$$E \left[\int_0^T \int_{\mathbb{T}^2} |(v_N^{(k)}(t))^{\frac{\alpha+3}{2}}|^2 + |(v_N^{(k)}(t))^{\frac{\alpha+3}{4}}|^4 dx dt \right] \lesssim_{\alpha, T} E \left[\sup_{0 \leq t \leq T} \|u_N^{(k)}(t)\|_{H^1(\mathbb{T}^2)}^{\alpha+3} \right] \\ \lesssim_{\Lambda, \alpha, T} \nu_{k, \alpha+3}$$

as a consequence of Theorem 4.1(v) and therefore

$$E \left[\|(v_N^{(k)})^{\frac{\alpha+3}{2}}\|_{L^2(0, T; H^2(\mathbb{T}^2))}^2 + \|(v_N^{(k)})^{\frac{\alpha+3}{4}}\|_{L^4(0, T; W^{1,4}(\mathbb{T}^2))}^4 \right] \lesssim_{\Lambda, \alpha, T} 1 + \nu_{k, \alpha+3}$$

by (4.13). □

Lemma 5.4. *For every $k \in \mathbb{N}$, the laws of $((v_N^{(k)})^{\frac{\alpha+3}{2}})_{N \in \mathbb{N}}$ are tight on $L^2(0, T; H^1(\mathbb{T}^2))$.*

Proof. We divide the proof into three steps.

Step 1 (Hölder regularity of $u_N^{(k)}$ in $L^2(\mathbb{T}^2)$). First, we observe that the paths of $u_N^{(k)}$ are weakly continuous in $H^1(\mathbb{T}^2)$ and in particular $\|u_N^{(k)}(t)\|_{H^1(\mathbb{T}^2)} \leq \|u_N^{(k)}\|_{L^\infty(0, T; H^1(\mathbb{T}^2))}$ for every $t \in [0, T]$. The Sobolev embedding theorem, see [34, Section 3.5.5], states that $W^{-1, q'}(\mathbb{T}^2) \hookrightarrow H^{\frac{-2}{q'}}(\mathbb{T}^2)$ and therefore we can estimate the Hölder seminorm $[u_N^{(k)}]_{\gamma, H^{\frac{-2}{q'}}(\mathbb{T}^2)}$ by $C_q [u_N^{(k)}]_{\gamma, W^{-1, q'}(\mathbb{T}^2)}$ for $\gamma \in (0, 1)$. The interpolation inequality

$$\|f\|_{L^2(\mathbb{T}^2)} \leq \|f\|_{H^1(\mathbb{T}^2)}^\theta \|f\|_{H^{\frac{-2}{q'}}(\mathbb{T}^2)}^{1-\theta}, \quad f \in H^1(\mathbb{T}^2),$$

with $\theta = \frac{\frac{2}{q'}}{1 + \frac{2}{q'}}$ can be derived using Fourier methods. We obtain the estimate

$$\|u_N^{(k)}(t) - u_N^{(k)}(s)\|_{L^2(\mathbb{T}^2)} \leq \|u_N^{(k)}(t) - u_N^{(k)}(s)\|_{H^1(\mathbb{T}^2)}^\theta \|u_N^{(k)}(t) - u_N^{(k)}(s)\|_{H^{\frac{-2}{q'}}(\mathbb{T}^2)}^{1-\theta} \\ \lesssim_q [u_N^{(k)}]_{\gamma, W^{-1, q'}(\mathbb{T}^2)}^{1-\theta} \|u_N^{(k)}\|_{L^\infty(0, T; H^1(\mathbb{T}^2))}^\theta |t - s|^{(1-\theta)\gamma}$$

on the increments, which yields that

$$(5.10) \quad [u_N^{(k)}]_{(1-\theta)\gamma, L^2(\mathbb{T}^2)} \lesssim_q [u_N^{(k)}]_{\gamma, W^{-1, q'}(\mathbb{T}^2)}^{1-\theta} \|u_N^{(k)}\|_{L^\infty(0, T; H^1(\mathbb{T}^2))}^\theta.$$

Step 2 (Integral estimate on the increments of $v_N^{(k)}$). In this part, we deduce from the first step an estimate on

$$\|\tau_h v_N^{(k)} - v_N^{(k)}\|_{L^4(0, T-h; L^2(\mathbb{T}^2))},$$

in similar terms, where τ_h denotes the translation operator by $h > 0$ in the time variable. To quantify the jumps in the paths of $v_N^{(k)}$ we introduce the function

$$\phi_{N, h} : [0, T] \rightarrow \mathbb{N}, \quad N \mapsto \lfloor t + h \rfloor_\delta - \lfloor t \rfloor_\delta,$$

which counts how many discretization points lie between t and $t + h$. The function $\lfloor \cdot \rfloor_\delta$ denotes here the biggest integer multiple of δ which is less or equal to its input value. Then we have

$$(5.11) \quad \|v_N^{(k)}(t + h) - v_N^{(k)}(t)\|_{L^2(\mathbb{T}^2)} \leq [u_N^{(k)}]_{(1-\theta)\gamma, L^2(\mathbb{T}^2)} (h + \phi_{N, h}(t)\delta)^{(1-\theta)\gamma}$$

for $t \in [0, T - h]$. We introduce the sets

$$C_{N,h} = \{t \in [0, T - h] | \phi_{N,h}(t) \neq 1\} \quad \text{and} \quad D_{N,h} = \{t \in [0, T - h] | \phi_{N,h}(t) = 1\},$$

distinguishing between points t , where one can estimate the right-hand side of (5.11) in terms of h or not. Indeed, if $t \in C_{N,h}$ it holds either $\phi_{N,h}(t) = 0$ or $\phi_{N,h}(t) \geq 2$ such that in any case $\phi_{N,h}(t)\delta \leq 2h$ and therefore

$$(h + \phi_{N,h}(t)\delta)^{(1-\theta)\gamma} \leq (3h)^{(1-\theta)\gamma}.$$

We deduce that

$$\begin{aligned} \|\tau_h v_N^{(k)} - v_N^{(k)}\|_{L^4(0, T-h; L^2(\mathbb{T}^2))}^4 &= \int_0^{T-h} \|v_N^{(k)}(t+h) - v_N^{(k)}(t)\|_{L^2(\mathbb{T}^2)}^4 dt \\ (5.12) \quad &\leq [u_N^{(k)}]_{(1-\theta)\gamma, L^2(\mathbb{T}^2)}^4 ((3h)^{4(1-\theta)\gamma} |C_{N,h}| + (h + \delta)^{4(1-\theta)\gamma} |D_{N,h}|). \end{aligned}$$

If $h \geq \delta$, we use the trivial estimate $|C_{N,h}| + |D_{N,h}| \leq T$ to conclude

$$(3h)^{4(1-\theta)\gamma} |C_{N,h}| + (h + \delta)^{4(1-\theta)\gamma} |D_{N,h}| \leq (3h)^{4(1-\theta)\gamma} T. \tag{5.13}$$

For $h < \delta$ we use instead that

$$t \in [j\delta, (j+1)\delta - h] \Rightarrow \phi_{N,h}(t) = 0 \Rightarrow t \in C_{N,h}$$

and consequently $|D_{N,h}| \leq (N+1)h$. We define the function

$$f_h(x) = \left(h + \frac{T}{x+1}\right)^{4(1-\theta)\gamma} (x+1)h, \quad x \in \left[1, \frac{T}{h} - 1\right]$$

such that

$$(h + \delta)^{4(1-\theta)\gamma} |D_{N,h}| \leq f_h(N)$$

and it suffices to estimate the maximum of f_h . Its derivative is given by

$$f'_h(x) = h \left(h + \frac{T}{x+1}\right)^{4(1-\theta)\gamma} - \frac{4(1-\theta)\gamma T \left(h + \frac{T}{x+1}\right)^{4(1-\theta)\gamma - 1} h(x+1)}{(x+1)^2},$$

which can vanish only if

$$\frac{4(1-\theta)\gamma T}{(x+1)\left(h + \frac{T}{x+1}\right)} = 1 \Rightarrow h(x+1) = (4(1-\theta)\gamma - 1)T.$$

We choose for the rest of the proof that $\gamma = \frac{1}{4}$ such that the above is not feasible for $x \in [1, \frac{T}{h} - 1]$. Hence f_h can attain its maximum only at the boundary points 1 and $\frac{T}{h} - 1$. Evaluating f_h gives

$$f_h(1) = \left(h + \frac{T}{2}\right)^{1-\theta} 2h \leq 2T^{1-\theta}h, \quad f_h\left(\frac{T}{h} - 1\right) = (2h)^{1-\theta}T.$$

We end up with the estimate

$$(3h)^{1-\theta} |C_{N,h}| + (h + \delta)^{1-\theta} |D_{N,h}| \leq 2T(3h)^{1-\theta} + 2T^{1-\theta}h.$$

We define the right-hand side as $g_{\theta,T}(h)$ and obtain from (5.10) and (5.12) that

$$\|\tau_h v_N^{(k)} - v_N^{(k)}\|_{L^4(0, T-h; L^2(\mathbb{T}^2))} \lesssim_q [u_N^{(k)}]_{\frac{1}{4}, W^{-1,q'}(\mathbb{T}^2)(\mathbb{T}^2)}^{1-\theta} \|u_N^{(k)}\|_{L^\infty(0, T; H^1(\mathbb{T}^2))}^\theta (g_{\theta,T}(h))^{\frac{1}{4}}. \tag{5.14}$$

By (5.13), this holds also if $h \geq \delta$.

Step 3 (Proof of tightness). Due to Theorem 4.1(v), (vi) and Lemma 5.3 we have

$$\begin{aligned} P\left(\left\{\sup_{0 \leq t \leq T} \|u_N^{(k)}\|_{H^1(\mathbb{T}^2)} > K\right\}\right) &\lesssim_{\Lambda, T} \frac{\nu_{k,2}}{K^2}, \\ \tilde{P}\left(\left\{[u_N^{(k)}]_{\frac{1}{4}, W^{-1, q'}(\mathbb{T}^2)} > K\right\}\right) &\lesssim_{\Lambda, q, T} \frac{1 + \nu_{k,2}}{K}, \\ P\left\{\|(v_N^{(k)})^{\frac{\alpha+3}{2}}\|_{L^2(0, T; H^2(\mathbb{T}^2))} > K\right\} &\lesssim_{\Lambda, \alpha, T} \frac{1 + \nu_{k, \alpha+3}}{K^2} \end{aligned}$$

for $K \in (1, \infty)$. In particular,

$$(5.15) \quad P((F_{N, K}^{(k)})^c) \lesssim_{\Lambda, \alpha, q, T} \frac{1 + \nu_{k, \alpha+3}}{K},$$

where we define

$$F_{N, K}^{(k)} = \left\{ \max \left\{ \sup_{0 \leq t \leq T} \|u_N^{(k)}\|_{H^1(\mathbb{T}^2)}, [u_N^{(k)}]_{\frac{1}{4}, W^{-1, q'}(\mathbb{T}^2)}, \|(v_N^{(k)})^{\frac{\alpha+3}{2}}\|_{L^2(0, T; H^2(\mathbb{T}^2))} \right\} \leq K \right\}.$$

Moreover, using that for $a, b \geq 0$ we have

$$\left| a^{\frac{\alpha+3}{2}} - b^{\frac{\alpha+3}{2}} \right| \leq \frac{\alpha+3}{2} |a-b| \max(a, b)^{\frac{\alpha+1}{2}} \lesssim |a-b| \left[a^{\frac{\alpha+1}{2}} + b^{\frac{\alpha+1}{2}} \right]$$

as a consequence of the fundamental theorem of calculus, we deduce that

$$\begin{aligned} \|\tau_h(v_N^{(k)})^{\frac{\alpha+3}{2}} - (v_N^{(k)})^{\frac{\alpha+3}{2}}\|_{L^2(0, T-h; L^{\frac{4}{\alpha+3}}(\mathbb{T}^2))} &\lesssim \|(\tau_h v_N^{(k)} - v_N^{(k)})(\tau_h(v_N^{(k)})^{\frac{\alpha+1}{2}} + (v_N^{(k)})^{\frac{\alpha+1}{2}})\|_{L^2(0, T-h; L^{\frac{4}{\alpha+3}}(\mathbb{T}^2))} \\ &\lesssim \|\tau_h v_N^{(k)} - \tilde{v}_N\|_{L^4(0, T-h; L^2(\mathbb{T}^2))} \|(v_N^{(k)})^{\frac{\alpha+1}{2}}\|_{L^4(0, T, L^{\frac{4}{\alpha+1}}(\mathbb{T}^2))} \end{aligned}$$

from Hölder's inequality. We estimate the latter term by

$$\|(v_N^{(k)})^{\frac{\alpha+1}{2}}\|_{L^4(0, T, L^{\frac{4}{\alpha+1}}(\mathbb{T}^2))} = \left(\int_0^T \left(\int_{\mathbb{T}^2} (v_N^{(k)})^2 dx \right)^{\alpha+1} dt \right)^{\frac{1}{4}} \lesssim_T \sup_{0 \leq t \leq T} \|v_N^{(k)}\|_{L^2(\mathbb{T}^2)}^{\frac{\alpha+1}{2}}.$$

We conclude by (5.14) that

$$\|\tau_h(v_N^{(k)})^{\frac{\alpha+3}{2}} - (v_N^{(k)})^{\frac{\alpha+3}{2}}\|_{L^2(0, T-h; L^{\frac{4}{\alpha+3}}(\mathbb{T}^2))} \lesssim_{q, T} [u_N^{(k)}]_{\frac{1}{4}, W^{-1, q'}(\mathbb{T}^2)}^{1-\theta} \|u_N^{(k)}\|_{L^\infty(0, T; H^1(\mathbb{T}^2))}^{\theta + \frac{\alpha+1}{2}} (g_{\theta, T}(h))^{\frac{1}{4}}.$$

Hence, for $\omega \in F_{N, K}^{(k)}$ we have that

$$\|\tau_h(v_N^{(k)}(\omega))^{\frac{\alpha+3}{2}} - (v_N^{(k)}(\omega))^{\frac{\alpha+3}{2}}\|_{L^2(0, T-h; L^{\frac{4}{\alpha+3}}(\mathbb{T}^2))} \lesssim_{q, T} K^{\frac{\alpha+3}{2}} (g_{\theta, T}(h))^{\frac{1}{4}}.$$

and

$$\|(v_N^{(k)}(\omega))^{\frac{\alpha+3}{2}}\|_{L^2(0, T; H^2(\mathbb{T}^2))} \leq K$$

and therefore $v_N^{(k)}(\omega)$ lies in a compact subset of $L^2(0, T; H^1(\mathbb{T}^2))$ by [35, Theorem 5, p.84], which we denote by $\chi_{q, \alpha, T, K}$. From (5.15) we deduce that

$$P(\{v_N^{(k)} \notin \chi_{q, \alpha, T, K}\}) \lesssim_{\Lambda, \alpha, q, T} \frac{1 + \nu_{k, \alpha+3}}{K}.$$

The tightness assertion follows since the right hand side goes uniformly in N to 0 as $K \rightarrow \infty$. \square

5.2. The time-step limit

In this last subsection, we finally let $N \rightarrow \infty$ and show that the limit satisfies the assertions of Theorem 1.3. This time we consider the sequence

$$(5.16) \quad \left((\mathbb{1}_{R_N^{(l)}}, \beta^{(l)}, u_N^{(l)}, v_N^{(l)}, w_N^{(l)}, J_N^{(l)}, (v_N^{(l)})^{\frac{\alpha+3}{2}}, (v_N^{(l)})^{\frac{\alpha+3}{4}}, (v_N^{(l)})^{\frac{\alpha+3}{2}})_{l \in \mathbb{N}} \right)_{N \in \mathbb{N}}$$

in the topological space

$$(5.17) \quad \prod_{l=1}^{\infty} \mathbb{R} \times C([0, T]) \times C(0, T; L^2(\mathbb{T}^2)) \times L_{w^*}^{\infty}(0, T; H^1(\mathbb{T}^2)) \times L_{w^*}^{\infty}(0, T; H^1(\mathbb{T}^2)) \\ \times L_w^2(0, T; L^{q'}(\mathbb{T}^2)) \times L_w^2(0, T; H^2(\mathbb{T}^2)) \times L_w^4(0, T; W^{1,4}(\mathbb{T}^2)) \times L^2(0, T; H^1(\mathbb{T}^2)).$$

Notice that this differs from (4.20) by the additional appearance of the space $L^2(0, T; H^1(\mathbb{T}^2))$.

Corollary 5.5. *The sequence (5.16) is tight on (5.17).*

Proof. This can be shown analogously to Proposition 4.7, using Theorem 4.1(v) and (vi) and invoking additionally the findings from Lemma 5.3 and Lemma 5.4. \square

As in Section 4.2, we obtain that for a subsequence indexed by $\mathcal{N} \subset \mathbb{N}$, there exists a sequence of \mathfrak{B} -measurable random variables

$$\left((\mathbb{1}_{\tilde{R}_N^{(l)}}, \tilde{\beta}_N^{(l)}, \tilde{u}_N^{(l)}, \tilde{v}_N^{(l)}, \tilde{w}_N^{(l)}, \tilde{J}_N^{(l)}, \tilde{f}_N^{(l)}, \tilde{g}_N^{(l)}, \tilde{h}_N^{(l)})_{l \in \mathbb{N}} \right)_{N \in \mathcal{N}}$$

defined on a complete probability space $(\tilde{\Omega}, \tilde{\mathfrak{A}}, \tilde{P})$, such that

$$(5.18) \quad \left(\mathbb{1}_{\tilde{R}_N^{(l)}}, \tilde{\beta}_N^{(l)}, \tilde{u}_N^{(l)}, \tilde{v}_N^{(l)}, \tilde{w}_N^{(l)}, \tilde{J}_N^{(l)}, \tilde{f}_N^{(l)}, \tilde{g}_N^{(l)}, \tilde{h}_N^{(l)} \right)_{l \in \mathbb{N}}$$

has the same distribution on (5.17) as

$$(5.19) \quad \left(\mathbb{1}_{R_N^{(l)}}, \beta^{(l)}, u_N^{(l)}, v_N^{(l)}, w_N^{(l)}, J_N^{(l)}, (v_N^{(l)})^{\frac{\alpha+3}{2}}, (v_N^{(l)})^{\frac{\alpha+3}{4}}, (v_N^{(l)})^{\frac{\alpha+3}{2}} \right)_{l \in \mathbb{N}}$$

for every $N \in \mathcal{N}$. Moreover, as $N \rightarrow \infty$, (5.18) converges to a \mathfrak{B} -measurable random variable

$$(5.20) \quad \left(\mathbb{1}_{\tilde{R}^{(l)}}, \tilde{\beta}^{(l)}, \tilde{u}^{(l)}, \tilde{v}^{(l)}, \tilde{w}^{(l)}, \tilde{J}^{(l)}, \tilde{f}^{(l)}, \tilde{g}^{(l)}, \tilde{h}^{(l)} \right)_{l \in \mathbb{N}}$$

in (5.17). The following properties are inherited from the approximating sequence.

Lemma 5.6. *The sets $(\tilde{R}^{(k)})_{k \in \mathbb{N}}$ form, up to \tilde{P} -null sets, a disjoint partition of $\tilde{\Omega}$. Moreover, the following holds \tilde{P} -almost surely for every $k \in \mathbb{N}$.*

(i) *The random variables*

$$\tilde{u}^{(k)}, \tilde{v}^{(k)}, \tilde{w}^{(k)}, \tilde{J}^{(k)}, \tilde{f}^{(k)}, \tilde{g}^{(k)} \text{ and } \tilde{h}^{(k)}$$

vanish outside of $\tilde{R}^{(k)}$.

(ii) *$\tilde{u}^{(k)}(t) \geq 0$ for all $t \in [0, T]$.*

(iii) *$\tilde{u}^{(k)} = \tilde{v}^{(k)} = \tilde{w}^{(k)}$.*

(iv) *$\tilde{f}^{(k)} = \tilde{h}^{(k)} = (\tilde{u}^{(k)})^{\frac{\alpha+3}{2}}$ and $\tilde{g}^{(k)} = (\tilde{u}^{(k)})^{\frac{\alpha+3}{4}}$.*

Proof. The claim regarding the sets $\tilde{R}^{(k)}$, as well as part (i) and (ii) follow as in the proof of Lemma 4.9. For part (iii) we conclude first from (5.1) that \tilde{P} -almost surely

$$(5.21) \quad \tilde{v}_N^{(k)}(t) = \tilde{u}_N^{(k)} \left(j\delta + \frac{t - j\delta}{2} \right), \quad t \in [j\delta, (j+1)\delta).$$

for almost all $t \in [0, T]$. Fixing such t that (5.21) holds for all $N \in \mathcal{N}$ and using that \hat{u}_N converges uniformly to an $L^2(\mathbb{T}^2)$ -continuous function we conclude that

$$(5.22) \quad \|\tilde{v}_N^{(k)}(t) - \tilde{u}^{(k)}(t)\|_{L^2(\mathbb{T}^2)} < \epsilon$$

for sufficiently large N . It follows that $\tilde{v}_N^{(k)} \rightarrow \tilde{u}^{(k)}$ in $L^\infty(0, T; L^2(\mathbb{T}^2))$ and therefore the limit has to coincide with $\tilde{v}^{(k)}$. The proof of $\tilde{u}^{(k)} = \tilde{w}^{(k)}$ works analogously. Since in contrast to the proof of Lemma 4.9 we cannot just rely on Proposition 2.4 for the identification of powers in (iv), we carry out the argument by hand. Since (5.18) and (5.19) have the same distribution, it holds

$$(5.23) \quad \tilde{f}_N^{(k)} = (\tilde{v}_N^{(k)})^{\frac{\alpha+3}{2}}$$

for every $N \in \mathcal{N}$. Due to the previously verified convergence $\tilde{v}_N^{(k)} \rightarrow \tilde{u}^{(k)}$ in $L^\infty(0, T; L^2(\mathbb{T}^2))$ it follows that the same convergence holds $[0, T] \times \mathbb{T}^2$ -almost everywhere up to taking a subsequence. Moreover, since $\tilde{v}_N^{(k)}$ is also weakly convergent in $L^\infty(0, T; H^1(\mathbb{T}^2))$, we conclude that it is uniformly in N bounded in $L^r([0, T] \times \mathbb{T}^2)$ for every $r > 0$ by the Sobolev embedding theorem. Vitali's convergence theorem yields that

$$(5.24) \quad (\tilde{v}_N^{(k)})^{\frac{\alpha+3}{2}} \rightarrow (\tilde{u}^{(k)})^{\frac{\alpha+3}{2}}$$

in $L^r([0, T] \times \mathbb{T}^2)$ for every $r > 0$. Invoking (5.23) and that $\tilde{f}_N^{(k)} \rightharpoonup f^{(k)}$ in $L^2(0, T; H^2(\mathbb{T}^2))$ we obtain the identification $\tilde{f}^{(k)} = (\tilde{u}^{(k)})^{\frac{\alpha+3}{2}}$. The remaining part of (iv) can be shown analogously. \square

Proposition 5.7. *For all $\eta \in L^\infty(0, T; W^{2,\infty}(\mathbb{T}^2))$ it holds*

$$\begin{aligned} \int_0^T \int_{\mathbb{T}^2} \tilde{J}^{(k)} \cdot \eta \, dx \, dt &= \int_0^T \int_{\{\tilde{u}^{(k)}(s) > 0\}} |\nabla \tilde{u}^{(k)}|^2 \nabla \tilde{u}^{(k)} \cdot \eta \, dx \, ds + \int_0^T \int_{\{\tilde{u}^{(k)}(s) > 0\}} \tilde{u}^{(k)} |\nabla \tilde{u}^{(k)}|^2 \operatorname{div} \eta \, dx \, ds \\ &+ 2 \int_0^T \int_{\{\tilde{u}^{(k)}(s) > 0\}} \tilde{u}^{(k)} \nabla^T \tilde{u}^{(k)} D\eta \nabla \tilde{u}^{(k)} \, dx \, ds + \int_0^T \int_{\mathbb{T}^2} (\tilde{u}^{(k)})^2 \nabla \tilde{u}^{(k)} \cdot \nabla \operatorname{div} \eta \, dx \, ds \end{aligned}$$

\tilde{P} -almost surely.

Proof. Since (5.18) and (5.19) have the same distribution, we conclude that

$$\begin{aligned} \int_{j\delta}^{(j+1)\delta} \int_{\mathbb{T}^2} \tilde{J}_N^{(k)} \cdot \eta \, dx \, dt &= \int_{j\delta}^{(j+1)\delta} \int_{\{\tilde{v}_N^{(k)}(s) > 0\}} |\nabla \tilde{v}_N^{(k)}|^2 \nabla \tilde{v}_N^{(k)} \cdot \eta \, dx \, ds \\ &+ \int_{j\delta}^{(j+1)\delta} \int_{\{\tilde{v}_N^{(k)}(s) > 0\}} \tilde{v}_N^{(k)} |\nabla \tilde{v}_N^{(k)}|^2 \operatorname{div} \eta \, dx \, ds \\ &+ 2 \int_{j\delta}^{(j+1)\delta} \int_{\{\tilde{v}_N^{(k)}(s) > 0\}} \tilde{v}_N^{(k)} \nabla^T \tilde{v}_N^{(k)} D\eta \nabla \tilde{v}_N^{(k)} \, dx \, ds \\ &+ \int_{j\delta}^{(j+1)\delta} \int_{\mathbb{T}^2} (\tilde{v}_N^{(k)})^2 \nabla \tilde{v}_N^{(k)} \cdot \nabla \operatorname{div} \eta \, dx \, ds \end{aligned}$$

by Theorem 4.1(iii) for every $N \in \mathcal{N}$ and $j \in \{0, \dots, N\}$. Summing up over j yields that

$$\begin{aligned} \int_0^T \int_{\mathbb{T}^2} \tilde{J}_N^{(k)} \cdot \eta \, dx \, dt &= \int_0^T \int_{\{\tilde{v}_N^{(k)}(s) > 0\}} |\nabla \tilde{v}_N^{(k)}|^2 \nabla \tilde{v}_N^{(k)} \cdot \eta \, dx \, ds + \int_0^T \int_{\{\tilde{v}_N^{(k)}(s) > 0\}} \tilde{v}_N^{(k)} |\nabla \tilde{v}_N^{(k)}|^2 \operatorname{div} \eta \, dx \, ds \\ &+ 2 \int_0^T \int_{\{\tilde{v}_N^{(k)}(s) > 0\}} \tilde{v}_N^{(k)} \nabla^T \tilde{v}_N^{(k)} D\eta \nabla \tilde{v}_N^{(k)} \, dx \, ds + \int_0^T \int_{\mathbb{T}^2} \tilde{v}_N^{(k)} \nabla \tilde{v}_N^{(k)} \cdot \nabla \operatorname{div} \eta \, dx \, ds. \end{aligned}$$

It is left to take the limit $N \rightarrow \infty$ in the above equality, which works exactly as in the deterministic case [33, Corollary 2.7, Theorem 3.2]. \square

Remark 5.8. We stress the importance of the additional convergence $(\tilde{v}_N^{(k)})^{\frac{\alpha+3}{2}} \rightarrow (\tilde{u}^{(k)})^{\frac{\alpha+3}{2}}$ in $L^2(0, T; H^1(\mathbb{T}^2))$ for the limiting argument [33, Corollary 2.7, Theorem 3.2].

The previous statement shows that the weak formulation of $\tilde{J}^{(k)} = (\tilde{u}^{(k)})^2 \nabla \Delta (\tilde{u}^{(k)})$ as in (1.7) is satisfied. We gather some more convergence and integrability results before we recover (1.8) as well. We note that we use again the convention to identify $\tilde{v}_N^{(k)} \in L^\infty(0, T; H^1(\mathbb{T}^2))$ with its version defined by (5.21) as well as the rounding function $\lfloor \cdot \rfloor_\delta$ from the proof of Lemma 5.4.

Lemma 5.9. For every $\varphi \in W^{1,q}(\mathbb{T}^2)$, $k \in \mathbb{N}$ and $t \in [0, T]$ it holds

$$(5.25) \quad \langle \tilde{v}_N^{(k)}(t), \varphi \rangle \rightarrow \langle \tilde{u}^{(k)}(t), \varphi \rangle,$$

$$(5.26) \quad \int_0^t \langle \operatorname{div}(\tilde{J}_N^{(k)}), \varphi \rangle ds \rightarrow \int_0^t \langle \operatorname{div}(\tilde{J}^{(k)}), \varphi \rangle ds,$$

$$(5.27) \quad \sum_{l=1}^{\infty} \int_0^{\lfloor t \rfloor_\delta} \lambda_l^2 \langle \operatorname{div}(\operatorname{div}(\tilde{w}_N^{(k)}(s) \psi_l) \psi_l), \varphi \rangle ds \rightarrow \sum_{l=1}^{\infty} \int_0^t \lambda_l^2 \langle \operatorname{div}(\operatorname{div}(\tilde{u}^{(k)}(s) \psi_l) \psi_l), \varphi \rangle ds,$$

$$(5.28) \quad \int_0^{\lfloor t \rfloor_\delta} \sum_{l=1}^{\infty} \lambda_l^2 \langle \operatorname{div}(\psi_l \tilde{w}_N^{(k)}), \varphi \rangle^2 ds \rightarrow \int_0^t \sum_{l=1}^{\infty} \lambda_l^2 \langle \operatorname{div}(\psi_l \tilde{u}^{(k)}), \varphi \rangle^2 ds,$$

$$(5.29) \quad \int_0^{\lfloor t \rfloor_\delta} \lambda_l \langle \operatorname{div}(\psi_l \tilde{w}_N^{(k)}), \varphi \rangle d\tau \rightarrow \int_0^t \lambda_l \langle \operatorname{div}(\psi_l \tilde{u}^{(k)}), \varphi \rangle d\tau,$$

$$(5.30) \quad \tilde{\beta}_N^{(l)}(\lfloor t \rfloor_\delta) \rightarrow \tilde{\beta}^{(l)}(t)$$

\tilde{P} -almost surely as $N \rightarrow \infty$.

Proof. The convergence (5.25) follows by (5.22). Part (5.26) is a direct consequence of $\tilde{J}_N^{(k)} \rightharpoonup \tilde{J}^{(k)}$ in $L^2(0, T; L^q(\mathbb{T}^2))$. Next, we observe that

$$\|\psi_l \cdot \nabla(\psi_l \cdot \nabla \varphi)\|_{H^{-1}(\mathbb{T}^2)} \lesssim \|\varphi\|_{H^1(\mathbb{T}^2)}$$

due to (3.5). Using $\tilde{w}_N^{(k)} \rightharpoonup^* \tilde{u}^{(k)}$ in $L^\infty(0, T; H^1(\mathbb{T}^2))$ we deduce that

$$(5.31) \quad \sum_{l=1}^{\infty} \int_0^t \lambda_l^2 \langle \tilde{w}_N^{(k)}(s), \psi_l \cdot \nabla(\psi_l \cdot \nabla \varphi) \rangle ds \rightarrow \sum_{l=1}^{\infty} \int_0^t \lambda_l^2 \langle \tilde{u}^{(k)}(s), \psi_l \cdot \nabla(\psi_l \cdot \nabla \varphi) \rangle ds.$$

Since weak-* convergent sequences are norm bounded, we obtain (5.27) by combining

$$\left| \sum_{l=1}^{\infty} \lambda_l^2 \int_{\lfloor t \rfloor_\delta}^t \langle \tilde{w}_N^{(k)}(s), \psi_l \cdot \nabla(\psi_l \cdot \nabla \varphi) \rangle ds \right| \leq \delta \sum_{l=1}^{\infty} \lambda_l^2 \|\tilde{w}_N^{(k)}\|_{L^\infty(0, T; H^1(\mathbb{T}^2))} \|\psi_l \cdot \nabla(\psi_l \cdot \nabla \varphi)\|_{H^{-1}}$$

with (5.31). For (5.28), we estimate

$$\begin{aligned} & \left| \int_0^t \sum_{l=1}^{\infty} \lambda_l^2 \langle \operatorname{div}(\psi_l \tilde{w}_N^{(k)}), \varphi \rangle^2 ds - \int_0^t \sum_{l=1}^{\infty} \lambda_l^2 \langle \operatorname{div}(\psi_l \tilde{u}^{(k)}), \varphi \rangle^2 ds \right| \\ & \leq \sum_{l=1}^{\infty} \lambda_l^2 \int_0^T |\langle \tilde{w}_N^{(k)}, \psi_l \cdot \nabla \varphi \rangle^2 - \langle \tilde{u}^{(k)}, \psi_l \cdot \nabla \varphi \rangle^2| ds \\ & \lesssim \sum_{l=1}^{\infty} \lambda_l^2 \int_0^T |\langle \tilde{w}_N^{(k)}, \psi_l \cdot \nabla \varphi \rangle - \langle \tilde{u}^{(k)}, \psi_l \cdot \nabla \varphi \rangle| (|\langle \tilde{w}_N^{(k)}, \psi_l \cdot \nabla \varphi \rangle| + |\langle \tilde{u}^{(k)}, \psi_l \cdot \nabla \varphi \rangle|) ds, \end{aligned}$$

where we employed that

$$a^2 - b^2 \leq 2|a - b| \max(|a|, |b|) \leq 2|a - b|(|a| + |b|)$$

for $a, b \in \mathbb{R}$. Since $\|\psi_l \cdot \nabla \varphi\|_{L^2(\mathbb{T}^2)}$ is uniformly in l bounded, we obtain further that

$$\begin{aligned}
& \left| \int_0^t \sum_{l=1}^{\infty} \lambda_l^2 \langle \operatorname{div}(\psi_l \tilde{w}_N^{(k)}), \varphi \rangle^2 ds - \int_0^t \sum_{l=1}^{\infty} \lambda_l^2 \langle \operatorname{div}(\psi_l \tilde{u}^{(k)}), \varphi \rangle^2 ds \right| \\
& \lesssim_{\Lambda, \varphi} \int_0^T \|\tilde{w}_N^{(k)} - \tilde{u}^{(k)}\|_{L^2(\mathbb{T}^2)} (\|\tilde{u}^{(k)}\|_{L^2(\mathbb{T}^2)} + \|\tilde{w}_N^{(k)}\|_{L^2(\mathbb{T}^2)}) dt \\
(5.32) \quad & \lesssim_T \|\tilde{w}_N^{(k)} - \tilde{u}^{(k)}\|_{L^\infty(0, T; L^2(\mathbb{T}^2))} (\|\tilde{u}^{(k)}\|_{L^\infty(0, T; L^2(\mathbb{T}^2))} + \sup_{N \in \mathcal{N}} \|\tilde{w}_N^{(k)}\|_{L^\infty(0, T; L^2(\mathbb{T}^2))})
\end{aligned}$$

As a consequence of the proof of Lemma 5.6(iii) we have $\tilde{w}_N^{(k)} \rightarrow \tilde{u}^{(k)}$ in $L^\infty(0, T; L^2(\mathbb{T}^2))$ as $N \rightarrow \infty$ and therefore the right-hand side of (5.32) tends to 0. Using that by the same arguments

$$\left| \int_{[t, \delta]}^t \sum_{l=1}^{\infty} \lambda_l^2 \langle \operatorname{div}(\psi_l \tilde{w}_N^{(k)}), \varphi \rangle^2 ds \right| \lesssim_{\Lambda, \varphi} \delta \|\tilde{w}_N^{(k)}\|_{L^\infty(0, T; L^2(\mathbb{T}^2))}^2$$

we obtain indeed (5.28). The convergence (5.29) can be derived analogously to (5.27). The last assertion (5.30) is a consequence of $\tilde{\beta}_N^{(l)} \rightarrow \tilde{\beta}^{(l)}$ in $C([0, T])$. \square

Lemma 5.10. *It holds for every $k \in \mathbb{N}$ that*

$$\begin{aligned}
& \tilde{E} \left[\sup_{0 \leq t \leq T} \|\tilde{u}^{(k)}\|_{H^1(\mathbb{T}^2)}^p \right] \lesssim_{\Lambda, p, T} \nu_{k, p}, \\
& \tilde{E} \left[\|\tilde{J}^{(k)}\|_{L^2(0, T; L^{q'}(\mathbb{T}^2))}^{\frac{p}{2}} \right] \lesssim_{\Lambda, p, q, T} \nu_{k, p}, \\
& \tilde{E} \left[\|\tilde{u}^{(k)}\|_{L^2(0, T; H^2(\mathbb{T}^2))}^{\frac{\alpha+3}{2}} + \|\tilde{u}^{(k)}\|_{L^4(0, T; W^{1,4}(\mathbb{T}^2))}^{\frac{\alpha+3}{4}} \right] \lesssim_{\Lambda, \alpha, T} 1 + \nu_{k, \alpha+3}.
\end{aligned}$$

Proof. This follows since (5.18) and (5.19) have the same distribution, lower semi-continuity of the norm with respect to weak and weak-* convergence, as well as the bounds from Theorem 4.1(v) and Lemma 5.3. \square

Finally, we define, as in Section 4.2, $\tilde{\mathfrak{F}}$ as the augmentation of the filtration $\tilde{\mathfrak{G}}$ given by

$$\tilde{\mathfrak{G}}_t = \sigma(\{\mathbb{1}_{\tilde{R}^{(l)}}, \tilde{J}^{(l)}|_{[0, t]} | l \in \mathbb{N}\} \cup \{\tilde{u}^{(l)}(s), \tilde{\beta}^{(l)}(s) | 0 \leq s \leq t, l \in \mathbb{N}\}),$$

where we consider $\tilde{J}^{(l)}|_{[0, t]}$ again as a \mathfrak{B} -random variable in $L^2(0, t; L^{q'}(\mathbb{T}^2))$.

Remark 5.11. The smallest σ -field $\tilde{\mathfrak{H}}_t$ on $\tilde{\Omega}$ such that $\phi(X)$ with

$$X = (\mathbb{1}_{\tilde{R}^{(l)}}, \tilde{\beta}^{(l)}|_{[0, t]}, \tilde{u}^{(l)}|_{[0, t]}, \tilde{J}^{(l)}|_{[0, t]})_{l \in \mathbb{N}}$$

is measurable for every bounded and continuous function

$$(5.33) \quad \phi: \prod_{l=1}^{\infty} \mathbb{R} \times C([0, t]) \times C(0, t; L^2(\mathbb{T}^2, \mathbb{R}^2)) \times L_w^2(0, t; L^{q'}(\mathbb{T}^2)) \rightarrow \mathbb{R}$$

coincides with $\tilde{\mathfrak{G}}_t$. Indeed, the inclusion $\tilde{\mathfrak{G}}_t \subset \tilde{\mathfrak{H}}_t$ follows since one can choose ϕ as a function depending only on one of the components of

$$\prod_{l=1}^{\infty} \mathbb{R} \times C([0, t]) \times C(0, t; L^2(\mathbb{T}^2, \mathbb{R}^2)) \times L_w^2(0, t; L^{q'}(\mathbb{T}^2)).$$

For the reverse inclusion $\tilde{\mathfrak{H}}_t \subset \tilde{\mathfrak{G}}_t$, we assume that ϕ as in (5.33) is bounded and continuous, such that it suffices to show that $\phi(X)$ is measurable with respect to $\tilde{\mathfrak{G}}_t$. In particular, ϕ is continuous as mapping from

$$(5.34) \quad \prod_{l=1}^{\infty} \mathbb{R} \times C([0, t]) \times C(0, t; L^2(\mathbb{T}^2, \mathbb{R}^2)) \times L^2(0, t; L^{q'}(\mathbb{T}^2))$$

into \mathbb{R} . But (5.34) is a complete separable metric space such that $\tilde{\mathfrak{G}}_t$ - \mathfrak{B} measurability of the (5.34)-valued random variable X can be checked using a suitable family of functions separating the points by [6, Theorem 6.8.9].

Theorem 5.12. *The processes $(\tilde{\beta}^{(l)})_{l \in \mathbb{N}}$ are a family of independent $\tilde{\mathfrak{F}}$ -Brownian motions. Moreover, we have for every $k \in \mathbb{N}$, $\varphi \in W^{1,q}(\mathbb{T}^2)$ and $t \in [0, T]$*

$$\begin{aligned} \langle \tilde{u}^{(k)}(t), \varphi \rangle - \langle \tilde{u}^{(k)}(0), \varphi \rangle &= - \int_0^t \langle \operatorname{div}(\tilde{J}^{(k)}), \varphi \rangle ds + \sum_{l=1}^{\infty} \int_0^t \lambda_l^2 \langle \operatorname{div}(\operatorname{div}(\tilde{u}^{(k)}(s)\psi_l)\psi_l), \varphi \rangle ds \\ &\quad + \sum_{l=1}^{\infty} \lambda_l \int_0^t \langle \operatorname{div}(\tilde{u}^{(k)}(s)\psi_l), \varphi \rangle d\tilde{\beta}_s^l \end{aligned}$$

\tilde{P} -almost surely.

Proof. For the claim regarding the family $(\tilde{\beta}^{(l)})_{l \in \mathbb{N}}$ we refer to [16]. For the remainder of the proof we fix $k \in \mathbb{N}$ and $\varphi \in W^{1,q}(\mathbb{T}^2)$ and define the $\tilde{\mathfrak{F}}$ -adapted process

$$M(t) = \langle \tilde{u}^{(k)}(t), \varphi \rangle - \langle \tilde{u}^{(k)}(0), \varphi \rangle + \int_0^t \langle \operatorname{div}(\tilde{J}^{(k)}), \varphi \rangle ds - \sum_{l=1}^{\infty} \int_0^t \lambda_l^2 \langle \operatorname{div}(\operatorname{div}(\tilde{u}^{(k)}(s)\psi_l)\psi_l), \varphi \rangle ds$$

and the approximating processes

$$M_N(t) = \langle \tilde{v}_N^{(k)}(t), \varphi \rangle - \langle \tilde{u}_N^{(k)}(0), \varphi \rangle + \int_0^t \langle \operatorname{div}(\tilde{J}^{(k)}), \varphi \rangle ds - \sum_{l=1}^{\infty} \int_0^{\lfloor t \rfloor \delta} \lambda_l^2 \langle \operatorname{div}(\operatorname{div}(\tilde{u}^{(k)}(s)\psi_l)\psi_l), \varphi \rangle ds.$$

As a consequence of (5.25), (5.26) and (5.27), $M_N(t)$ converges to $M(t)$ for every $t \in [0, T]$ as $N \rightarrow \infty$. Moreover, we let

$$(5.35) \quad \rho : \prod_{l=1}^{\infty} \mathbb{R} \times C([0, s]) \times C(0, s; L^2(\mathbb{T}^2, \mathbb{R}^2)) \times L_w^2(0, s; L^{q'}(\mathbb{T}^2)) \rightarrow \mathbb{R}$$

be bounded and continuous and consider the random variables

$$\begin{aligned} \rho &= \phi((\mathbb{1}_{\tilde{R}^{(l)}}, \tilde{\beta}^{(l)}|_{[0,s]}, \tilde{u}^{(l)}|_{[0,s]}, \tilde{J}^{(l)}|_{[0,s]})_{l \in \mathbb{N}}), \\ \rho_N &= \phi((\mathbb{1}_{\tilde{R}_N^{(l)}}, \tilde{\beta}_N^{(l)}|_{[0,s]}, \tilde{u}_N^{(l)}|_{[0,s]}, \tilde{J}_N^{(l)}|_{[0,s]})_{l \in \mathbb{N}}). \end{aligned}$$

As consequence of the convergence of (5.18) to (5.20) in (5.17) we have $\rho_N \rightarrow \rho$ as $N \rightarrow \infty$. Using that as a consequence of Theorem 4.1(iii) and (iv)

$$\begin{aligned} \langle v_N^{(k)}(t), \varphi \rangle - \langle u_N^{(k)}(0), \varphi \rangle + \int_0^t \langle \operatorname{div}(J_N^{(k)}), \varphi \rangle ds - \sum_{l=1}^{\infty} \int_0^{\lfloor t \rfloor \delta} \lambda_l^2 \langle \operatorname{div}(\operatorname{div}(w_N^{(k)}(s)\psi_l)\psi_l), \varphi \rangle ds \\ = \sum_{l=1}^{\infty} \lambda_l \int_0^{\lfloor t \rfloor \delta} \langle \operatorname{div}(w_N^{(k)}\psi_l), \varphi \rangle d\beta_s^l, \end{aligned}$$

we conclude by additionally invoking Theorem 4.1(ii) that

$$(5.36) \quad \begin{aligned} \tilde{E}[(M_N(t) - M_N(s + \kappa))\rho_N] &= 0, \\ \tilde{E}\left[\left(M_N^2(t) - M_N^2(s + \kappa) - \int_{\lfloor s + \kappa \rfloor \delta}^{\lfloor t \rfloor \delta} \sum_{l=1}^{\infty} \lambda_l^2 \langle \operatorname{div}(\psi_l \tilde{w}_N^{(k)}), \varphi \rangle^2 d\tau\right)\rho_N\right] &= 0, \\ \tilde{E}\left[\left(\tilde{\beta}_N^{(l)}(\lfloor t \rfloor \delta)M_N(t) - \tilde{\beta}_N^{(l)}(\lfloor s + \kappa \rfloor \delta)M_N(s + \kappa) - \int_{\lfloor s + \kappa \rfloor \delta}^{\lfloor t \rfloor \delta} \lambda_l \langle \operatorname{div}(\psi_l \tilde{w}_N^{(k)}), \varphi \rangle d\tau\right)\rho_N\right] &= 0. \end{aligned}$$

for $s, t \in [0, T]$, $\kappa > 0$ such that $s + \kappa \leq t$, and N large enough so that $\lfloor s + \kappa \rfloor_\delta \geq s$. Due to Theorem 4.1(v) and the Burkholder–Davis–Gundy inequality we have

$$(5.37) \quad \sup_{N \in \mathbb{N}} \tilde{E} \left[\|\tilde{w}_N^{(k)}\|_{L^\infty(0, T; H^1(\mathbb{T}^2))}^p + \sup_{\tau \in [0, T]} |M_N(\tau)|^p + |\tilde{\beta}_N^{(l)}(\tau)|^p \right] < \infty$$

for every $p \in (0, \infty)$ such that Vitali’s convergence theorem and (5.28), (5.29), (5.30) yield

$$(5.38) \quad \begin{aligned} \tilde{E}[(M(t) - M(s + \kappa))\rho] &= 0, \\ \tilde{E} \left[\left(M^2(t) - M^2(s + \kappa) - \int_{s+\kappa}^t \sum_{l=1}^\infty \lambda_l^2 \langle \operatorname{div}(\psi_l \tilde{u}^{(k)}), \varphi \rangle^2 d\tau \right) \rho \right] &= 0, \\ \tilde{E} \left[\left(\tilde{\beta}^{(l)}(t)M(t) - \tilde{\beta}^{(l)}(s + \kappa)M(s + \kappa) - \int_{s+\kappa}^t \lambda_l \langle \operatorname{div}(\psi_l \tilde{u}^{(k)}), \varphi \rangle d\tau \right) \rho \right] &= 0, \end{aligned}$$

by letting $N \rightarrow \infty$ in (5.36). Using that

$$\{ \{ \phi((\mathbb{1}_{\tilde{R}_N^{(l)}}, \tilde{\beta}_N^{(l)}|_{[0, s]}, \tilde{u}_N^{(l)}|_{[0, s]}, \tilde{J}_N^{(l)}|_{[0, s]})_{l \in \mathbb{N}}) \in B \} | \phi \text{ as in (5.35) cont. bdd., } B \in \mathfrak{B}(\mathbb{R}) \}$$

is an intersection stable generator of $\tilde{\mathfrak{G}}_t$ by Remark 5.11, we conclude that (5.38) holds for every $\tilde{\mathfrak{G}}_t$ -measurable and bounded random variable ρ . Finally, we let $0 \leq s' \leq t' \leq T$ and ρ be a $\tilde{\mathfrak{G}}_{s'}$ -measurable and bounded random variable. If we can show that

$$(5.39) \quad \begin{aligned} \tilde{E}[(M(t') - M(s'))\rho] &= 0, \\ \tilde{E} \left[\left(M^2(t') - M^2(s') - \int_{s'}^{t'} \sum_{l=1}^\infty \lambda_l^2 \langle \operatorname{div}(\psi_l \tilde{u}^{(k)}), \varphi \rangle^2 d\tau \right) \rho \right] &= 0, \\ \tilde{E} \left[\left(\tilde{\beta}^{(l)}(t)M(t') - \tilde{\beta}^{(l)}(s')M(s') - \int_{s'}^{t'} \lambda_l \langle \operatorname{div}(\psi_l \tilde{u}^{(k)}), \varphi \rangle d\tau \right) \rho \right] &= 0, \end{aligned}$$

the claim follows by [24, Proposition A.1], because M is $\tilde{\mathfrak{G}}$ -adapted and square-integrable due to Lemma 5.10. To this end, we let $\kappa' > 0$ and define $\kappa = \frac{\kappa'}{2}$, $s = s' + \kappa$ and $t = t' + 2\kappa$. By definition of the augmented filtration, there exists a \tilde{P} -version of ρ which is $\tilde{\mathfrak{G}}_s$ -measurable and moreover we have $s + \kappa \leq t$. Therefore, we can apply (5.38) and rephrase in terms of s', t', κ' to deduce that

$$\begin{aligned} \tilde{E}[(M(t' + \kappa') - M(s' + \kappa'))\rho] &= 0, \\ \tilde{E} \left[\left(M^2(t' + \kappa') - M^2(s' + \kappa') - \int_{s'+\kappa'}^{t'+\kappa'} \sum_{l=1}^\infty \lambda_l^2 \langle \operatorname{div}(\psi_l \tilde{u}^{(k)}), \varphi \rangle^2 d\tau \right) \rho \right] &= 0, \\ \tilde{E} \left[\left(\tilde{\beta}^{(l)}(t' + \kappa')M(t' + \kappa') - \tilde{\beta}^{(l)}(s' + \kappa')M(s' + \kappa') - \int_{s'+\kappa'}^{t'+\kappa'} \lambda_l \langle \operatorname{div}(\psi_l \tilde{u}^{(k)}), \varphi \rangle d\tau \right) \rho \right] &= 0. \end{aligned}$$

Since $\kappa' > 0$ was arbitrary, we can use continuity of $\tilde{\beta}^{(l)}$ and M , Vitalis’s theorem and the consequence

$$E \left[\sup_{\tau \in [0, T]} \|\tilde{u}^{(k)}(\tau)\|_{H^1(\mathbb{T}^2)}^p + |M(\tau)|^p + |\tilde{\beta}^{(l)}(\tau)|^p \right] < \infty$$

of Lemma 5.10 to let $\kappa' \searrow 0$ and obtain (5.39). □

Finally, we put

$$(5.40) \quad \tilde{u} = \sum_{k=0}^\infty \tilde{u}^{(k)}, \quad \tilde{J} = \sum_{k=0}^\infty \tilde{J}^{(k)},$$

which is in light of Lemma 5.6 equivalent to require

$$(5.41) \quad \tilde{u} = \tilde{u}^{(k)} \quad \text{and} \quad \tilde{J} = \tilde{J}^{(k)} \quad \text{on} \quad \tilde{R}^{(k)}.$$

Proof of Theorem 1.3. We first show that $(\tilde{\Omega}, \tilde{\mathfrak{A}}, \tilde{P}), \tilde{\mathfrak{F}}, (\tilde{\beta}^{(l)})_{l \in \mathbb{N}}, \tilde{u}$ together with \tilde{J} constitute a solution to the stochastic thin-film equation with q' -regular non linearity in the sense of Definition 1.1. By definition, $\tilde{\mathfrak{F}}$ fulfills the usual conditions and $(\tilde{\beta}^{(l)})_{l \in \mathbb{N}}$ is a family of independent Brownian motions by Theorem 5.12. Furthermore, \tilde{u} and \tilde{J} are, as each of their summands, an $H_w^1(\mathbb{T}^2)$ -continuous, $\tilde{\mathfrak{F}}$ -adapted process and a random variable in $L^2(0, T; L^{q'}(\mathbb{T}^2))$, respectively. Moreover, $\tilde{J}|_{[0,t]}$ is $\tilde{\mathfrak{F}}_t$ -measurable by definition of $\tilde{\mathfrak{F}}$ and we have $\sup_{0 \leq t \leq T} \|\tilde{u}(t)\|_{H^1(\mathbb{T}^2)} < \infty$ because of Lemma 5.10. Proposition 5.7 together with (5.40) yield that (1.7) is indeed fulfilled. Similarly, we obtain (1.8) from Theorem 5.12, (5.40) and the fact that one can pull the $\tilde{\mathfrak{F}}_0$ -measurable random variable $\mathbb{1}_{\tilde{R}^{(k)}}$ outside of the stochastic integrals in (1.8). For the initial condition, we observe that

$$(\mathbb{1}_{\tilde{R}_N^{(k)}}, \tilde{u}_N^{(k)}(0))_{k \in \mathbb{N}} \sim (\mathbb{1}_{R_N^{(k)}}, u_N^{(k)}(0))_{k \in \mathbb{N}}, \quad N \in \mathcal{N},$$

and

$$(\mathbb{1}_{\tilde{R}_N^{(k)}}, \tilde{u}_N^{(k)}(0))_{k \in \mathbb{N}} \rightarrow (\mathbb{1}_{\tilde{R}^{(k)}}, \tilde{u}^{(k)}(0))_{k \in \mathbb{N}}$$

as $N \rightarrow \infty$ in $(\mathbb{R} \times L^2(\mathbb{T}^2))^\infty$. Hence, we have

$$(\mathbb{1}_{\tilde{R}^{(k)}}, \tilde{u}^{(k)}(0))_{k \in \mathbb{N}} \sim (\mathbb{1}_{R_N^{(k)}}, u_N^{(k)}(0))_{k \in \mathbb{N}}$$

and therefore

$$\tilde{u}(0) = \sum_{k=1}^{\infty} \tilde{u}^{(k)}(0) \sim \sum_{k=1}^{\infty} u_N^{(k)}(0) \sim \mu$$

by Theorem 4.1(i). The non-negativity of $\tilde{u}(t)$ follows from Lemma 5.6(ii). From Lemma 5.10 together with the monotone convergence theorem we deduce the energy estimates (1.10) and (1.11). Finally due to Lemma 5.6(iv) we conclude that the additional spatial regularity property (1.12) is by construction fulfilled. \square

Appendix A: Properties of solutions to the deterministic thin-film equation

Proof of Theorem 2.3. Since $\alpha \in (-1, 0)$ and $v \in H^1(\mathbb{T}^2)$ we have due to (1.15) that

$$\int_{\mathbb{T}^2} G_\alpha(v_0) dx < \infty.$$

Therefore, [33, Theorem 3.2] applies and yields that a weak solution (v, J) with q' -regular non-linearity and initial value v_0 exists. The first part of (i) follows by testing the equation $\partial_t u = -\operatorname{div} J$ with $\varphi \otimes \mathbb{1}_{\mathbb{T}^2}$ for arbitrary $\varphi \in C_c^\infty((0, T))$. For the other properties we consider the approximation procedure in [33], which takes place in two steps. First problems of the form

$$(P_{\delta\epsilon}) \quad \begin{cases} \partial_t(v_{\delta\epsilon}) + \operatorname{div}(J_{\delta\epsilon}) = 0 & \text{in } L^2(0, T; H^{-1}(\mathbb{T}^2)), \\ J_{\delta\epsilon} = m_{\delta\epsilon}(v_{\delta\epsilon}) \nabla \Delta v_{\delta\epsilon} & \text{weakly,} \\ \operatorname{esslim}_{t \rightarrow 0} v_{\delta\epsilon}(t, \cdot) = v_0 + \delta + \epsilon^\theta & \text{in } H^1(\mathbb{T}^2), \end{cases}$$

are solved by [20, Theorem 1.1]. Letting $\epsilon \searrow 0$ yields solutions to

$$(P_\delta) \quad \begin{cases} \partial_t(u_\delta) + \operatorname{div}(J_{\delta\epsilon}) = 0, \\ J_\delta = m_\delta(v_\delta) \nabla \Delta v_\delta, \\ v_\delta(0, \cdot) = v_0 + \delta, \end{cases}$$

in the sense of [33, Definition 3.1] which again are used to construct v . The functions m_δ and $m_{\delta\epsilon}$ are auxiliary mobilities, which take the form $m_\delta(\tau) = \frac{\tau^2}{1+\delta\tau^2}$ and $m_{\delta\epsilon} = \frac{\tau^s m_\delta(\tau)}{\epsilon m_\delta(\tau) + \tau^s}$ for some $s > 4$, see [33, p.323, p.331], so we can choose for example $s = 5$. The number θ from $(P_{\delta\epsilon})$ is a sufficiently small constant. By [33, Lemma 2.1] it follows that

$$(A.1) \quad \|\nabla v_{\delta\epsilon}\|_{L^\infty(0,T;L^2(\mathbb{T}^2,\mathbb{R}^2))} \leq \|\nabla v_0\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}.$$

Since $v_{\delta\epsilon} \rightharpoonup^* v_\delta$ and $v_\delta \rightharpoonup^* v$ in $L^\infty(0, T; H^1(\mathbb{T}^2))$, see [33, Proposition 2.6], we conclude under additional consideration of Remark 2.2 that part (ii) holds true. Moreover, since also $v_{\delta\epsilon} \rightharpoonup v_\delta$ and $v_\delta \rightharpoonup v$ in $H^1(0, T; W^{-1,q'}(\mathbb{T}^2))$, it follows that strong convergence takes place in $C(0, T; L^2(\mathbb{T}^2))$ by Remark 2.2. Hence non negativity is preserved and the second part of (i) follows by the non negativity of $v_{\delta\epsilon}$, see [33, Lemma 2.1]. Furthermore, in [33, Equation (2.26)] one finds the estimate (iv). Finally, to convince ourselves also of part (iii), we conclude that as consequence of [33, Lemma 2.1] it holds

$$(A.2) \quad \operatorname{ess\,sup}_{T-\rho \leq t \leq T} \|\nabla v_{\delta\epsilon}(t)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 + 2 \int_0^{T-\rho} \|\sqrt{m_{\delta\epsilon}(v_{\delta\epsilon}(t))} \nabla \Delta v_{\delta\epsilon}(t)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 dt \leq \|\nabla v_0\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2$$

for any $\rho > 0$. Since by definition

$$m_{\delta\epsilon}(\tau) \leq m_\delta(\tau) \leq \tau^2$$

we obtain by Sobolev's inequality, see [1, Theorem 4.51], the periodic Poincaré inequality and (A.1) that

$$\begin{aligned} \|\sqrt{m_{\delta\epsilon}(v_{\delta\epsilon}(t))}\|_{L^r(\mathbb{T}^2)}^2 &\leq \|v_{\delta\epsilon}(t)\|_{L^r(\mathbb{T}^2)}^2 \lesssim_r \|v_{\delta\epsilon}(t)\|_{H^1(\mathbb{T}^2)}^2 \lesssim \|\nabla v_{\delta\epsilon}(t)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 + \left| \int_{\mathbb{T}^2} v_{\delta\epsilon}(t) dx \right|^2 \\ &\leq \|\nabla v_0\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 + \left| \int_{\mathbb{T}^2} v_0 + \delta + \epsilon^\theta dx \right|^2 \end{aligned}$$

for any $0 \leq t \leq T$ and $r \in [1, \infty)$. Because we have $J_{\delta\epsilon}(t) = m_{\delta\epsilon}(v_{\delta\epsilon}(t)) \nabla \Delta v_{\delta\epsilon}(t)$ for almost all $0 \leq t \leq T$, see [33, Lemma 2.1], we obtain by (A.1), (A.2), Hölder's inequality and the choice $\frac{1}{2} + \frac{1}{r} = \frac{1}{q'}$ that

$$\begin{aligned} C_q \operatorname{ess\,sup}_{T-\rho \leq t \leq T} \left[\|\nabla v_{\delta\epsilon}(t)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 \left(\|\nabla v_{\delta\epsilon}(t)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 + \left| \int_{\mathbb{T}^2} v_0 + \delta + \epsilon^\theta dx \right|^2 \right) \right] + \int_0^{T-\rho} \|J_{\delta\epsilon}(t)\|_{L^{q'}(\mathbb{T}^2,\mathbb{R}^2)}^2 dt \\ \leq C_q \|\nabla v_0\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 \left(\|\nabla v_0\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 + \left| \int_{\mathbb{T}^2} v_0 + \delta + \epsilon^\theta dx \right|^2 \right). \end{aligned}$$

Using that $J_{\delta\epsilon} \rightharpoonup J_\delta$ and $J_\delta \rightharpoonup J$ in $L^2(0, T; L^{q'}(\mathbb{T}^2, \mathbb{R}^2))$ as well as that $\nabla v_{\delta\epsilon} \rightharpoonup^* \nabla v_\delta$ and $\nabla v_\delta \rightharpoonup^* \nabla v$ in $L^\infty(0, T; L^2(\mathbb{T}^2, \mathbb{R}^2))$, see [33, Proposition 2.6], we infer that

$$\begin{aligned} C_q \operatorname{ess\,sup}_{T-\rho \leq t \leq T} \left[\|\nabla v(t)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 \left(\|\nabla v(t)\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 + \left| \int_{\mathbb{T}^2} v_0 dx \right|^2 \right) \right] + \int_0^{T-\rho} \|J(t)\|_{L^{q'}(\mathbb{T}^2,\mathbb{R}^2)}^2 dt \\ \leq C_q \|\nabla v_0\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 \left(\|\nabla v_0\|_{L^2(\mathbb{T}^2,\mathbb{R}^2)}^2 + \left| \int_{\mathbb{T}^2} v_0 dx \right|^2 \right) \end{aligned}$$

The claimed estimate follows by letting $\rho \searrow 0$ together with the weak continuity of v in $H^1(\mathbb{T}^2)$. \square

Appendix B: Gelfand triple of Bessel potential spaces

The purpose of this section is to verify that $H^2(\mathbb{T}^2) \subset H^1(\mathbb{T}^2) \subset L^2(\mathbb{T}^2)$ is a Gelfand triple, when equipping $H^2(\mathbb{T}^2)$ with the Bessel potential norm, as claimed in the proof of Theorem 3.1. We recall that the Bessel potential norm on $H^2(\mathbb{T}^2)$ is defined by

$$\|f\|_{H^2(\mathbb{T}^2)}^2 = \sum_{k \in \mathbb{Z}^2} (1 + |2\pi k|^2)^2 |\hat{f}(k)|^2,$$

where

$$(B.1) \quad \hat{f}(k) = \int_{\mathbb{T}^2} f(x) e^{-2\pi i k \cdot x} dx, \quad k \in \mathbb{Z}^2$$

is the k -th Fourier coefficient of a function $f \in L^2(\mathbb{T}^2)$. Moreover, by definition of the Bessel potential spaces

$$H^2(\mathbb{T}^2) = \{f \in L^2(\mathbb{T}^2) \mid \|f\|_{H^2(\mathbb{T}^2)} < \infty\}.$$

The pairing of two functions $f \in H^1(\mathbb{T}^2)$, $g \in H^2(\mathbb{T}^2)$ in $H^1(\mathbb{T}^2)$ can be rewritten by Parseval's relation [19, Proposition 3.2.7] as

$$(B.2) \quad (f, g)_{H^1(\mathbb{T}^2)} = (f, g)_{L^2(\mathbb{T}^2)} + (\nabla f, \nabla g)_{L^2(\mathbb{T}^2, \mathbb{R}^2)} = \sum_{k \in \mathbb{Z}^2} (1 + |2\pi k|^2) \hat{f}(k) \overline{\hat{g}(k)}$$

and therefore

$$|(f, g)_{H^1(\mathbb{T}^2)}| \leq \left(\sum_{k \in \mathbb{Z}^2} |\hat{f}(k)|^2 \right) \left(\sum_{k \in \mathbb{Z}^2} (1 + |2\pi k|^2)^2 |\hat{g}(k)|^2 \right) = \|f\|_{L^2(\mathbb{T}^2)} \|g\|_{H^2(\mathbb{T}^2)}$$

by the Cauchy–Schwarz inequality. Hence,

$$(B.3) \quad \|(f, \cdot)_{H^1(\mathbb{T}^2)}\|_{(H^2(\mathbb{T}^2))'} \leq \|f\|_{L^2(\mathbb{T}^2)}.$$

Moreover, since the coefficients are square summable, the series

$$\sum_{k \in \mathbb{Z}^2} \frac{\hat{f}(k)}{1 + |2\pi k|^2} e^{2\pi i k \cdot x}$$

converges to an element $g_f \in L^2(\mathbb{T}^2)$. Since

$$(B.4) \quad \|g_f\|_{H^2(\mathbb{T}^2)}^2 = \sum_{k \in \mathbb{Z}^2} (1 + |2\pi k|^2)^2 \left(\frac{|\hat{f}(k)|}{1 + |2\pi k|^2} \right)^2 = \|f\|_{L^2(\mathbb{T}^2)}^2 < \infty,$$

it satisfies $g_f \in H^2(\mathbb{T}^2)$. Using (B.2), we obtain that

$$(f, g_f)_{H^1(\mathbb{T}^2)} = \sum_{k \in \mathbb{Z}^2} (1 + |2\pi k|^2) \hat{f}(k) \frac{\overline{\hat{f}(k)}}{1 + |2\pi k|^2} = \|f\|_{L^2(\mathbb{T}^2)}^2,$$

such that

$$\|(f, \cdot)_{H^1(\mathbb{T}^2)}\|_{(H^2(\mathbb{T}^2))'} \geq \|f\|_{L^2(\mathbb{T}^2)}$$

by (B.4). Due to (B.3), the above inequality is an equality. Consequently, identifying $f \in H^1(\mathbb{T}^2)$ with $(f, \cdot)_{H^1(\mathbb{T}^2)}$ and taking the completion of these functions with respect to

$$\|(f, \cdot)_{H^1(\mathbb{T}^2)}\|_{(H^2(\mathbb{T}^2))'}$$

yields the space $L^2(\mathbb{T}^2)$. With this identification, the dual pairing between functions $g \in H^2(\mathbb{T}^2)$ and $f \in L^2(\mathbb{T}^2)$ is given by

$$\langle\langle f, g \rangle\rangle_{H^1(\mathbb{T}^2)} = \langle f, g \rangle + \langle \nabla f, \nabla g \rangle,$$

where we recall that $\langle \cdot, \cdot \rangle$ is the dual pairing in $L^2(\mathbb{T}^2)$. Indeed, for $f \in H^1(\mathbb{T}^2)$ this follows since it was identified with $(f, \cdot)_{H^1(\mathbb{T}^2)}$. For $f \in L^2(\mathbb{T}^2)$ we take a sequence $(f_n)_{n \in \mathbb{N}}$ from $H^1(\mathbb{T}^2)$ converging to f in $L^2(\mathbb{T}^2)$. Then also $\nabla f_n \rightarrow \nabla f$ in $H^{-1}(\mathbb{T}^2, \mathbb{R}^2)$ and hence

$$\langle\langle f, g \rangle\rangle_{H^1(\mathbb{T}^2)} \leftarrow \langle\langle f_n, g \rangle\rangle_{H^1(\mathbb{T}^2)} = \langle f_n, g \rangle + \langle \nabla f_n, \nabla g \rangle \rightarrow \langle f, g \rangle + \langle \nabla f, \nabla g \rangle$$

for all $g \in H^2(\mathbb{T}^2)$.

Appendix C: Justifications of Itô's formula

First, we justify the use of Itô's formula in the proof of Lemma 4.3. To this end, we introduce the equivalence relation

$$f \sim g \iff \exists c \in \mathbb{R} : f = g + c$$

for $f, g \in H^s(\mathbb{T}^2)$, $s \geq 0$ and write \dot{f} for the respective equivalence class in $H^s(\mathbb{T}^2)$. We recall that the Bessel potential space is given by

$$H^s(\mathbb{T}^2) = \{f \in L^2(\mathbb{T}^2) \mid \|f\|_{H^s(\mathbb{T}^2)} < \infty\},$$

where the appearing Bessel potential norm is defined as

$$\|f\|_{H^s(\mathbb{T}^2)}^2 = \sum_{k \in \mathbb{Z}^2} (1 + |2\pi k|^2)^s |\hat{f}(k)|^2$$

with $\hat{f}(k)$ being the k -th Fourier coefficient (B.1) of a function $f \in L^2(\mathbb{T}^2)$. Under this norm, the quotient space $H^s(\mathbb{T}^2)/\sim$ is equipped with

$$\begin{aligned} \|\dot{f}\|_{H^s(\mathbb{T}^2)/\sim}^2 &= \inf_{g \in \dot{f}} \sum_{k \in \mathbb{Z}^2} (1 + |2\pi k|^2)^s |\hat{g}(k)|^2 = \inf_{g \in \dot{f}} |\hat{g}(0)|^2 + \sum_{k \in \mathbb{Z}^2 \setminus \{(0,0)\}} (1 + |2\pi k|^2)^s |\hat{f}(k)|^2 \\ &= \sum_{k \in \mathbb{Z}^2 \setminus \{(0,0)\}} (1 + |2\pi k|^2)^s |\hat{f}(k)|^2. \end{aligned}$$

Here, we used in the second equality that

$$\int_{\mathbb{T}^2} e^{-2\pi k i \cdot x} dx = 0$$

for $k \in \mathbb{Z}^2 \setminus \{(0,0)\}$ and therefore

$$\hat{f}(k) = \hat{g}(k)$$

for all $g \in \dot{f}$. In the following, we write $\dot{H}^s(\mathbb{T}^2)$ for $H^s(\mathbb{T}^2)/\sim$ and equip it with the equivalent norm

$$\|\dot{f}\|_{\dot{H}^s(\mathbb{T}^2)}^2 = \sum_{k \in \mathbb{Z}^2 \setminus \{(0,0)\}} |2\pi k|^{2s} |\hat{f}(k)|^2.$$

Analogously to Appendix B, one verifies that $\dot{H}^0(\mathbb{T}^2)$ can be identified with the dual of $\dot{H}^2(\mathbb{T}^2)$ under the pairing in $\dot{H}^1(\mathbb{T}^2)$. Moreover, the dual pairing is given by

$$(C.1) \quad \langle \dot{f}, \dot{g} \rangle_{\dot{H}^1(\mathbb{T}^2)} = \sum_{k \in \mathbb{Z}^2 \setminus \{(0,0)\}} |2\pi k|^2 \hat{f}(k) \overline{\hat{g}(k)} = \langle \nabla f, \nabla g \rangle$$

for $\dot{f} \in \dot{H}^1(\mathbb{T}^2)$ and $\dot{g} \in \dot{H}^2(\mathbb{T}^2)$ by Parseval's relation [19, Proposition 3.2.7]. For general $\dot{f} \in \dot{H}^0(\mathbb{T}^2)$ the equality (C.1) holds by an approximation argument as in Appendix B.

Next, we denote by P_{hom} the operator mapping a function $f \in H^s(\mathbb{T}^2)$ to its equivalence class in $\dot{H}^s(\mathbb{T}^2)$, i.e. we set $P_{\text{hom}}f = \dot{f}$. Then

$$P_{\text{hom}} \in L(L^2(\mathbb{T}^2), \dot{H}^0(\mathbb{T}^2)) \cap L(H^1(\mathbb{T}^2), \dot{H}^1(\mathbb{T}^2))$$

and applying P_{hom} to equation (3.2) satisfied by w_ϵ on $[j\delta, (j+1)\delta)$ yields that

$$(C.2) \quad P_{\text{hom}}w_\epsilon(t) = P_{\text{hom}}w_\epsilon(j\delta) + \int_{j\delta}^t P_{\text{hom}}A^\epsilon(w_\epsilon(s)) ds + \int_{j\delta}^t P_{\text{hom}}B(w_\epsilon(s)) dV_t, \quad t \in [j\delta, (j+1)\delta),$$

where V is the cylindrical Wiener process in $H^2(\mathbb{T}^2, \mathbb{R}^2)$ given by

$$(C.3) \quad V_t = \sum_{l=1}^{\infty} \beta_l^{(l)} \psi_l.$$

Because of $P_{\text{hom}} \in L(H^2(\mathbb{T}^2), \dot{H}^2(\mathbb{T}^2))$, $w_\epsilon \in L^2([j\delta, (j+1)\delta] \times \Omega, H^2(\mathbb{T}^2))$ by Theorem 3.1, and the boundedness of the operators A^ϵ and B , it holds

$$(C.4) \quad P_{\text{hom}} w_\epsilon \in L^2([j\delta, (j+1)\delta] \times \Omega, \dot{H}^2(\mathbb{T}^2)),$$

$$(C.5) \quad P_{\text{hom}} A^\epsilon(w_\epsilon) \in L^2([j\delta, (j+1)\delta] \times \Omega, \dot{H}^0(\mathbb{T}^2)),$$

$$(C.6) \quad P_{\text{hom}} B(w_\epsilon) \in L^2([j\delta, (j+1)\delta] \times \Omega, L_2(H^2(\mathbb{T}^2, \mathbb{R}^2), \dot{H}^0(\mathbb{T}^2))).$$

Moreover, because of right-continuity in $H^1(\mathbb{T}^2)$, w_ϵ admits a progressively measurable, $H^2(\mathbb{T}^2)$ -valued $dt \otimes P$ -version by [30, Exercise 4.2.3]. Since later in the proof of Lemma 4.3 we integrate in time and take the expectation anyways, we denote this progressive version again by w_ϵ to ease the notation. By continuity of the involved operators, also the processes (C.5) and (C.6) are progressive when choosing this $dt \otimes P$ -version of w_ϵ , such that Itô's formula for the squared norm in $\dot{H}^1(\mathbb{T}^2)$ from [30, Theorem 4.2.5] is applicable to (C.2). Noting that by Parseval's relation, the norm in $\dot{H}^1(\mathbb{T}^2)$ can be written as

$$\|\dot{f}\|_{\dot{H}^1(\mathbb{T}^2)}^2 = \|\nabla f\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2,$$

we obtain that

$$\begin{aligned} \|\nabla w_\epsilon(t)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 &= \|\nabla w_\epsilon(j\delta)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 + 2 \int_{j\delta}^t \langle \nabla w_\epsilon(s), \nabla A^\epsilon(w_\epsilon(s)) \rangle ds \\ &\quad + 2 \int_{j\delta}^t \langle \nabla w_\epsilon(s), \nabla B(w_\epsilon(s)) dV_s \rangle + \sum_{l=1}^{\infty} \lambda_l^2 \int_{j\delta}^t \|\nabla \operatorname{div}(w_\epsilon(s) \psi_l)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)}^2 ds \end{aligned}$$

for $t \in [j\delta, (j+1)\delta)$. Writing the stochastic integral with respect to V as its series representation results in (4.4). Moreover, its quadratic variation is given by

$$(C.7) \quad 4 \int_{j\delta}^t \|\langle \nabla w_\epsilon(s), \nabla B(w_\epsilon(s)) \cdot \rangle\|_{L_2(H^2(\mathbb{T}^2, \mathbb{R}^2), \mathbb{R})}^2 ds = 4 \sum_{l=1}^{\infty} \lambda_l^2 \int_{j\delta}^t \langle \nabla \operatorname{div}(w_\epsilon(s) \psi_l), \nabla w_\epsilon(s) \rangle^2 ds.$$

Secondly, we justify the use of Itô's formula in the proof of Lemma 5.3, where we use instead [14, Proposition A.1]. Choosing $\psi = \mathbb{1}_{\mathbb{T}^2}$, $\varphi = G_{\alpha, \kappa}$ in the notation of this proposition, we see that the functional (5.5) is of the required form. Next, we observe that as a consequence of Theorem 4.1(iv), the process $w_N^{(k)}$ satisfies

$$dw_N^{(k)} = \operatorname{div}(G(t)) dt + H(t) dV_t$$

on $[j\delta, (j+1)\delta)$, where

$$\begin{aligned} G(t) &= \frac{1}{2} \sum_{l=1}^{\infty} \lambda_l^2 \operatorname{div}(w_N^{(k)}(t) \psi_l) \psi_l, \\ H(t)[v] &= \sum_{l=1}^{\infty} \lambda_l \langle v, \psi_l \rangle_{H^2(\mathbb{T}^2, \mathbb{R}^2)} \operatorname{div}(w_N^{(k)}(t) \psi_l), \quad v \in H^2(\mathbb{T}^2, \mathbb{R}^2) \end{aligned}$$

and V_t as in (C.3). By Theorem 4.1(v), we have

$$(C.8) \quad w_N^{(k)} \in L^2(\Omega, L^2(j\delta, (j+1)\delta; H^1(\mathbb{T}^2)))$$

and

$$w_N^{(k)} \in L^2(\Omega, C(j\delta, (j+1)\delta; L^2(\mathbb{T}^2))),$$

if we replace its terminal value $w_N^{(k)}((j+1)\delta)$ by $v_N^{(k)}((j+1)\delta)$. By (3.5) and $\Lambda \in l^2(\mathbb{N})$ we have that

$$\|G(t)\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)} \leq \frac{1}{2} \sum_{l=1}^{\infty} \lambda_l^2 \|\operatorname{div}(w_N^{(k)}(t)\psi_l)\psi_l\|_{L^2(\mathbb{T}^2, \mathbb{R}^2)} \lesssim \sum_{l=1}^{\infty} \lambda_l^2 \|w_N^{(k)}(t)\|_{H^1(\mathbb{T}^2)} \lesssim_{\Lambda} \|w_N^{(k)}(t)\|_{H^1(\mathbb{T}^2)}$$

and consequently (C.8) implies that

$$G \in L^2(\Omega, L^2(j\delta, (j+1)\delta; L^2(\mathbb{T}^2, \mathbb{R}^2))).$$

Similarly, we estimate

$$\|H(t)\|_{L^2(H^2(\mathbb{T}^2, \mathbb{R}^2), L^2(\mathbb{T}^2))}^2 = \sum_{l=1}^{\infty} \lambda_l^2 \|\operatorname{div}(w_N^{(k)}(t)\psi_l)\|_{L^2(\mathbb{T}^2)}^2 \lesssim \sum_{l=1}^{\infty} \lambda_l^2 \|w_N^{(k)}(t)\|_{H^1(\mathbb{T}^2)}^2 \lesssim_{\Lambda} \|w_N^{(k)}(t)\|_{H^1(\mathbb{T}^2)}^2,$$

such that

$$H \in L^2(\Omega, L^2(j\delta, (j+1)\delta; L_2(H^2(\mathbb{T}^2, \mathbb{R}^2), L^2(\mathbb{T}^2)))).$$

Hence, all the assumptions of [14, Proposition A.1] are satisfied, which results in (5.7).

Acknowledgements

The author thanks his doctoral advisor Manuel Gnann for many insightful discussions on this subject as well as the careful reading of this document. He thanks Mark Veraar for pointing out the possibility to relax integrability assumptions on the initial value. He also thanks the anonymous referees for the careful reading of this document and their valuable suggestions.

References

- [1] R. A. Adams and J. J. F. Fournier. *Sobolev spaces*. Elsevier/Academic Press, Amsterdam, 2003. MR2424078
- [2] A. Agresti and M. Veraar. Nonlinear parabolic stochastic evolution equations in critical spaces part II. *J. Evol. Equ.* **22** (2022) 1–96. MR4437443 <https://doi.org/10.1007/s00028-022-00786-7>
- [3] H. Amann. Compact embeddings of vector valued Sobolev and Besov spaces. *Glas. Mat.* **35** (2000) 161–177. MR1783238
- [4] F. Bernis and A. Friedman. Higher order nonlinear degenerate parabolic equations. *J. Differ. Equ.* **83** (1990) 179–206. MR1031383 [https://doi.org/10.1016/0022-0396\(90\)90074-Y](https://doi.org/10.1016/0022-0396(90)90074-Y)
- [5] D. Blömker. Nonhomogeneous noise and Q-Wiener processes on bounded domains. *Stoch. Anal. Appl.* **23** (2005) 255–273. MR2130349 <https://doi.org/10.1081/SAP-200050092>
- [6] V. I. Bogachev. *Measure theory. Vol. I, II*. Springer-Verlag, Berlin, 2007. MR2267655 <https://doi.org/10.1007/978-3-540-34514-5>
- [7] F. Boyer and P. Fabrie. *Mathematical tools for the study of the incompressible Navier–Stokes equations and related models*. Springer, New York, 2013. MR2986590 <https://doi.org/10.1007/978-1-4614-5975-0>
- [8] D. Breit, E. Feireisl and M. Hofmanová. *Stochastically forced compressible fluid flows*. De Gruyter, Berlin, 2018. MR3791804
- [9] H. Brezis. *Functional analysis, Sobolev spaces and partial differential equations*. Springer, New York, 2011. MR2759829
- [10] F. Cornalba A priori positivity of solutions to a non-conservative stochastic thin-film equation. Preprint, 2018. Available at arXiv:1811.07826.
- [11] K. Dareiotis, B. Gess, M. V. Gnann and G. Grün. Non-negative martingale solutions to the stochastic thin-film equation with nonlinear gradient noise. *Arch. Ration. Mech. Anal.* **242** (2021) 179–234. MR4302759 <https://doi.org/10.1007/s00205-021-01682-z>
- [12] B. Davidovitch, E. Moro and H. A. Stone. Spreading of viscous fluid drops on a solid substrate assisted by thermal fluctuations. *Phys. Rev. Lett.* **95** (2005) 244505.
- [13] N. De Nitti and J. Fischer. Sharp criteria for the waiting time phenomenon in solutions to the thin-film equation. *Comm. Partial Differential Equations* (2022) 1–41. MR4444309 <https://doi.org/10.1080/03605302.2022.2056702>
- [14] A. Debussche, M. Hofmanová and J. Vovelle. Degenerate parabolic stochastic partial differential equations: Quasilinear case. *Ann. Probab.* **44** (2016) 1916–1955. MR3502597 <https://doi.org/10.1214/15-AOP1013>
- [15] S. N. Ethier and T. G. Kurtz. *Markov processes: Characterization and convergence*. John Wiley & Sons, Inc., New York, 1986. MR0838085 <https://doi.org/10.1002/9780470316658>
- [16] J. Fischer and G. Grün. Existence of positive solutions to stochastic thin-film equations. *SIAM J. Math. Anal.* **50** (2018) 411–455. MR3755665 <https://doi.org/10.1137/16M1098796>
- [17] M. Gerencsér, I. Gyöngy and N. Krylov. On the solvability of degenerate stochastic partial differential equations in Sobolev spaces. *Stoch. Partial Differ. Equ. Anal. Comput.* **3** (2014) 52–83. MR3312592 <https://doi.org/10.1007/s40072-014-0042-6>
- [18] B. Gess and M. Gnann. The stochastic thin-film equation: Existence of nonnegative martingale solutions. *Stochastic Process. Appl.* **130** (2020) 7260–7302. MR4167206 <https://doi.org/10.1016/j.spa.2020.07.013>
- [19] L. Grafakos. *Classical Fourier analysis*. Springer, New York, 2014. MR3243734 <https://doi.org/10.1007/978-1-4939-1194-3>

- [20] G. Grün. Degenerate parabolic differential equations of fourth order and a plasticity model with non-local hardening. *Z. Anal. Anwend.* **14** (1995) 541–574. MR1362530 <https://doi.org/10.4171/ZAA/639>
- [21] G. Grün and L. Klein. Zero-contact angle solutions to stochastic thin-film equations. *J. Evol. Equ.* **22** (2022) 1–37. MR4455116 <https://doi.org/10.1007/s00028-022-00818-2>
- [22] G. Grün, K. Mecke and M. Rauscher. Thin-film flow influenced by thermal noise. *J. Stat. Phys.* **122** (2006) 1261–1291. MR2219535 <https://doi.org/10.1007/s10955-006-9028-8>
- [23] I. Gyöngy and N. Krylov. On the splitting-up method and stochastic partial differential equations. *Ann. Probab.* **31** (2003) 564–591. MR1964941 <https://doi.org/10.1214/aop/1048516528>
- [24] M. Hofmanová. Degenerate parabolic stochastic partial differential equations. *Stochastic Process. Appl.* **123** (2013) 4294–4336. MR3096355 <https://doi.org/10.1016/j.spa.2013.06.015>
- [25] T. Hytönen, J. van Neerven, M. Veraar and L. Weis. *Analysis in Banach spaces. Vol. I. Martingales and Littlewood–Paley theory*. Springer, Cham, 2016. MR3617205
- [26] A. Jakubowski. Short communication: The almost sure Skorokhod representation for subsequences in nonmetric spaces. *Theory Probab. Appl.* **42** (1998) 167–174. MR1453342 <https://doi.org/10.1137/S0040585X97976052>
- [27] O. Kallenberg. *Foundations of modern probability*. Springer-Verlag, New York, 1997. MR1464694
- [28] N. Krylov. A relatively short proof of Itô’s formula for spdes and its applications. *Stoch. Partial Differ. Equ. Anal. Comput.* **1** (2013) 152–174. MR3327504 <https://doi.org/10.1007/s40072-013-0003-5>
- [29] N. V. Krylov. *Introduction to the theory of random processes*. American Mathematical Society, Providence, RI, 2002. MR1885884 <https://doi.org/10.1090/gsm/043>
- [30] W. Liu and M. Röckner. *Stochastic partial differential equations: An introduction*. Springer, Cham, 2015. MR3410409 <https://doi.org/10.1007/978-3-319-22354-4>
- [31] S. Metzger and G. Grün Existence of nonnegative solutions to stochastic thin-film equations in two space dimensions. Preprint, 2021. Available at arXiv:2106.07973. MR4462584 <https://doi.org/10.4171/afb/476>
- [32] M. Ondreját and M. Veraar. On temporal regularity of stochastic convolutions in 2-smooth Banach spaces. *Ann. Inst. Henri Poincaré Probab. Stat.* **56** (2020) 1792–1808. MR4116708 <https://doi.org/10.1214/19-AIHP1017>
- [33] R. D. Passo, H. Garcke and G. Grün. On a fourth-order degenerate parabolic equation: Global entropy estimates, existence, and qualitative behaviour of solutions. *SIAM J. Math. Anal.* **29** (1998) 321–342. MR1616558 <https://doi.org/10.1137/S0036141096306170>
- [34] H.-J. Schmeisser and H. Triebel. *Topics in Fourier analysis and function spaces*. John Wiley & Sons, Ltd., Chichester, 1987. MR0891189
- [35] J. Simon. Compact sets in the space $L^p(0, T; B)$. *Ann. Mat. Pura Appl.* **146** (1986) 65–96. MR0916688 <https://doi.org/10.1007/BF01762360>