

**Delft University of Technology** 

## Deformation pattern in the thrust-related Parmelan Anticline (Bornes Massif, Subalpine Chains, Haute-Savoie, France): preliminary results

Berio, Luigi; Balsamo, Fabrizio; Mittempergher, Silvia; Mozafar, Mahtab; Storti, Fabrizio; Bistacchi, Andrea; Bruna, Pierre-Olivier; Bertotti, Giovanni

**Publication date** 2018 **Document Version** Final published version

## Citation (APA)

Berio, L., Balsamo, F., Mittempergher, S., Mozafar, M., Storti , F., Bistacchi, A., Bruna, P.-O., & Bertotti, G. (2018). *Deformation pattern in the thrust-related Parmelan Anticline (Bornes Massif, Subalpine Chains,* Hauté-Savoie, France): preliminary results. Abstract from EGU General Assembly 2018, Vienna, Austria.

#### Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

**Copyright** Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

#### Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

This work is downloaded from Delft University of Technology. For technical reasons the number of authors shown on this cover page is limited to a maximum of 10. Geophysical Research Abstracts Vol. 20, EGU2018-8923, 2018 EGU General Assembly 2018 © Author(s) 2018. CC Attribution 4.0 license.



# Deformation pattern in the thrust-related Parmelan Anticline (Bornes Massif, Subalpine Chains, Haute-Savoie, France): preliminary results

Luigi Berio (1), Fabrizio Balsamo (1), Silvia Mittempergher (2), Mahtab Mozafari (1), Fabrizio Storti (1), Andrea Bistacchi (2), Pierre-Olivier Bruna (3), and Giovanni Bertotti (3)

(1) NEXT—Natural and Experimental Tectonics Research Group, Department of Chemistry, Life Sciences and Environmental Sustainability, Parma University, Parco Area delle Scienze 157A, 43124 Parma, Italy, (2) Department of Earth and Environmental Sciences, University of Milano-Bicocca, Piazza della Scienza 4, 20126 Milano, Italy, (3) TU Delft, Department of Civil Engineering and Geosciences, Applied Geology Section, Stevinweg 51, 2628 CN Delft, The Netherlands

The Parmelan Anticline is located in the frontal part of the Bornes Massif where Upper Jurassic to Cretaceous sediments of the European passive margin overlain by Tertiary foredeep turbidites are exposed. The NE-SW trending Parmelan Anticline is characterized by steeply dipping limbs separated by a large flat plateau. The limbs and the crest represent discrete structural domains separated by narrow kinked hinge zones localized on inherited prefolding NE-SW trending extensional fault zones. In this contribution, we present a preliminary reconstruction of the kinematic evolution of this anticline by combining structural data obtained from detailed field mapping with microstructural and petrographic analyses. Relying on crosscutting relationships, a relative chronology between different calcite vein sets, bed-parallel and tectonic stylolites and conjugate systems of reverse and strike-slip faults is proposed. Their relative timing with respect to folding (i.e. pre- syn- or post-folding) is determined by comparing structural data collected in different structural domains after bedding dip removal. Upon unfolding, our data illustrate that pre-folding deformation structures related to layer-parallel shortening or even older, such as different sets of veins and tectonic stylolites, and conjugate systems of subsidiary reverse faults, are widespread. Because of their pre-folding origin, these structural elements are almost homogeneously distributed irrespectively of their position within the anticline. Accordingly, our results confirm the fundamental role played by structural inheritance, either related or unrelated to the stress field active during folding, to drive fold evolution and the associated deformation pattern.