

## Implications for seismic hazard assessment from the characterization of active fault zones near Ulaanbaatar with electrical resistivity models

Comeau, Matthew J.; Batmagnai, Erdenechimeg; Tserendug, Shoovdor; Bayartogtokh, Enkhzul; Kuvshinov, Alexey; Demberel, Sodnomsambuu

### Publication date

2024

### Document Version

Final published version

### Citation (APA)

Comeau, M. J., Batmagnai, E., Tserendug, S., Bayartogtokh, E., Kuvshinov, A., & Demberel, S. (2024). *Implications for seismic hazard assessment from the characterization of active fault zones near Ulaanbaatar with electrical resistivity models*. Abstract from 26th International Electromagnetic Induction Workshop 2024, Beppu, Japan. [https://www.emiw.org/emiw2024/abstracts/abstract-listing/pmfe-abstract-details?tx\\_powermail\\_pi2%5Baction%5D=show&tx\\_powermail\\_pi2%5Bcontroller%5D=Output&tx\\_powermail\\_pi2%5Bmail%5D=3906&cHash=9e4d07da9d7b4eb9b7dbc3b0d38d83c3](https://www.emiw.org/emiw2024/abstracts/abstract-listing/pmfe-abstract-details?tx_powermail_pi2%5Baction%5D=show&tx_powermail_pi2%5Bcontroller%5D=Output&tx_powermail_pi2%5Bmail%5D=3906&cHash=9e4d07da9d7b4eb9b7dbc3b0d38d83c3)

### Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

### Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

### Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

## Implications for seismic hazard assessment from the characterization of active fault zones near Ulaanbaatar with electrical resistivity models

Matthew J. Comeau<sup>1</sup>, Erdenechimeg Batmagnai<sup>2</sup>, Shoovdor Tserendug<sup>2</sup>, Enkhzul Bayartogtokh<sup>2</sup>, Alexey Kuvshinov<sup>3</sup>, and Sodnomsambuu Demberel<sup>2</sup>

<sup>1</sup> Delft University of Technology, The Netherlands, M.J.Comeau@tudelft.nl

<sup>2</sup> Institute of Astronomy and Geophysics Mongolian Academy of Sciences, Mongolia, Batmagnai@iag.ac.mn; Tserendug@iag.ac.mn; Enkhzul@iag.ac.mn; Demberel@iag.ac.mn

<sup>3</sup> ETH Zürich, Switzerland, Kuvshinov@erdw.ethz.ch

---

### SUMMARY

Within the Ulaanbaatar region, Mongolia, 120 earthquake events were recorded with a magnitude between 3.4 and 5.6 and 978 earthquake events had a magnitude between 2.5 and 3.4, for the period from 1994 to 2016. Several of these have been strongly felt by residents of Ulaanbaatar. Historical records, since 1905, show that Mongolia as a whole has experienced four major earthquakes with magnitudes larger than 8 and many moderate earthquakes with magnitudes larger than 5.5. The seismicity in Mongolia is concentrated along the Mongolian-Altai and Gobi-Altai, the Bulnay (north of the Khangai mountains), and around Mogod (east of the Khangai mountains).

In the west of the Ulaanbaatar region there are several prominent fault zones, some only identified very recently. The majority of the seismic events in this region are related to the Khustai, Sharkhai, and Avdar fault zones. Seismicity is typically detected in the upper ~16 km of the crust. These fault zones are 100+ km long and historical events are predicted to have produced vertical offsets of up to 10 m; some sections show a cumulative horizontal offset of up to 100 m. Based on paleo-seismic surveys, it is estimated that these fault zones could produce earthquakes up to magnitude 7. Therefore, these faults pose a serious threat and risk of damage to Ulaanbaatar.

In this presentation we aim to characterize the active fault zones near Ulaanbaatar with electrical resistivity models generated from magnetotelluric data. In mid-2024 we carried out measurements across the Khustai, Sharkhai, and Avdar fault zones and modeled the local features near the fault traces and the regional crustal features of the region. Preliminary models show several low-resistivity features (approximately <100  $\Omega$ m) in the near-surface. The upper crust (0-25 km depth) appears to have a generally high-resistivity (~10,000  $\Omega$ m), whereas the lower crust (25–50 km depth) appears to have a lower resistivity (approximately <100  $\Omega$ m).

We aim to give an integrated interpretation of the electrical conductivity structure of the subsurface with geomorphological and geological knowledge, geodetic measurements, paleo-seismic trenching, and near-surface ground-penetrating radar surveys. We also aim to discuss the relation with fault mechanical models and local fault damage zones, and the relevance of the low slip rate. Understanding the subsurface structure of the region and characterizing the active faults is an important step for assessing the seismic hazard.

**Keywords:** Electrical Resistivity; Fault; Hazard; Earthquake

---