

(Re-) Evaluating the Role of Microbes for Fluid Mud Rheology and Settling

Gebert, Julia; Kirichek, Alex ; de Lucas Pardo, Miguel; Amman, Baerbel ; Ohle, Nino

Publication date

2023

Document Version

Final published version

Citation (APA)

Gebert, J., Kirichek, A., de Lucas Pardo, M., Amman, B., & Ohle, N. (2023). *(Re-) Evaluating the Role of Microbes for Fluid Mud Rheology and Settling*. Abstract from 13th International SedNet Conference 2023, Lisbon, Portugal. <https://sednet.org/wp-content/uploads/2023/08/5.5F-Re-Evaluating-the-Role-of-Microbes-for-Fluid-Mud-Rheology-and-Settling.pdf>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

(Re-) Evaluating the Role of Microbes for Fluid Mud Rheology and Settling

Julia Gebert¹, Alex Kirichek², Miguel de Lucas Pardo³, Bärbel Amman⁴, Nino Ohle⁵

¹Department of Geoscience & Engineering & ²Department of Hydraulic Engineering, Delft University of Technology, Delft, the Netherlands

Phone: +31 152 782 798

E-mail: j.gebert@tudelft.nl

³Deltares, Delft, the Netherlands

⁴Niedersachsen Ports GmbH & Co. KG, branch Emden, Germany

⁵Hamburg Port Authority AöR, Hamburg, Germany

Conference theme number(s): 3

Introduction: Organic matter interacts with charged mineral surfaces and thereby influences sediment properties relevant for port navigability, such as rheological characteristics [1], [2], and settling and consolidation behavior [3], [4]. Microbially produced extracellular polymeric substances (EPS) have been thought to enhance suspension and thereby the desired low yield stresses of fluid mud (FM) [5]. This study investigates the relationship between concentration and composition of EPS, particle size distribution, mineralogy, settling rates and yield stresses in fluid mud from the ports of Emden and Hamburg.

Methods: Concentration and composition of EPS, microbial biomass, rheological parameters, settling rates, microbial community composition and standard solids properties were measured over several years in Port of Hamburg and over one year in Seaport Emden. Single analyses of clay mineralogical composition were carried out on samples from both ports.

Results: The fine-grained fraction ($< 63 \mu\text{m}$) in FM exceeded 90% in both ports. The microbial community at both sites was highly diverse and total EPS concentration significantly lower in the saline inner harbor of Seaport Emden than in the freshwater Port of Hamburg (Fig. 1, left). Here, FM at downstream site KH with the lowest ratio of biomass to clay (Fig. 1, right) featured hindered settling behaviour. Yield stress increased exponentially with density (Fig. 2, left), with a similar relationship for the two Emden locations and a more scattered one for the Port of Hamburg, in relation varying organic matter content [6]. Settling rates were inversely related to density, as shown by Emden data (Fig. 2, right).

Discussion: The favourable properties of fluid mud in Seaport Emden appear to result from the low density achieved by recirculation dredging and not by particularly high concentrations of EPS, which are also subject to strong seasonal variation. Comparative results from different sites within the Port of Hamburg show that a low ratio of biomass to clay experience promotes hindered settling, likely due to a lesser extent of floc formation. These results support the relevance of fresh, biomass-derived organic matter for particle-particle interactions [7], flocculation and settling behavior, but shed doubt on the assumption

that EPS contribute to keeping FM in suspension. The latter appears to be accomplished by the density window generated from recirculation dredging, with densities that are low enough to enable low yield stresses but high enough to allow for low settling rates.




Fig. 1: Left: EPS in Seaport Emden (sites GS and IH) and in Port of Hamburg. Right: Ratio of microbial biomass to clay at three sites in Port of Hamburg.




Fig. 2: Relationship between density and fluidic yield stress in fluid mud from Emden and Hamburg (left) and settling rate (Emden, right).

This study was funded by Niedersachsen Ports and Hamburg Port Authority and was carried out within the framework of the MUDNET academic network. <https://www.tudelft.nl/mudnet/>.

References: [1] Zanderet et al. (2022) *J Soils Sediments* 22: 2873-2882; [2] Shakeel et al. (2022) *J Soils Sediments* 22: 2883-2892; [3] Sills and Gonzalez (2001) *Géotechnique* 51: 629-639; [4] Jommi et al. (2019) *Géotechnique* 69: 753-766; [5] Wurpts et al. (2005) *Terra et Aqua* 99: 22-32; [6] Shakeel et al. (2019) *Geo Mar Lett* 39: 427-434 [7] Deng et al. (2019) *Marine Geology* 413: 71-84.