

## Formate Oxidase (FOx) from *Aspergillus oryzae*

### One Catalyst Enables Diverse H<sub>2</sub>O<sub>2</sub>-Dependent Biocatalytic Oxidation Reactions

Tieves, Florian; Willot, Sébastien Jean Paul; van Schie, Morten Martinus Cornelis Harald; Rauch, Marine Charène Renée; Younes, Sabry Hamdy Hamed; Zhang, Wuyuan; Dong, Jia Jia; Gomez de Santos, Patricia; Hollmann, Frank

**DOI**

[10.1002/anie.201902380](https://doi.org/10.1002/anie.201902380)

**Publication date**

2019

**Document Version**

Final published version

**Published in**

Angewandte Chemie - International Edition

**Citation (APA)**

Tieves, F., Willot, S. J. P., van Schie, M. M. C. H., Rauch, M. C. R., Younes, S. H. H., Zhang, W., Dong, J. J., Gomez de Santos, P., & Hollmann, F. (2019). Formate Oxidase (FOx) from *Aspergillus oryzae*: One Catalyst Enables Diverse H<sub>2</sub>O<sub>2</sub>-Dependent Biocatalytic Oxidation Reactions. *Angewandte Chemie - International Edition*, 58(23), 7873-7877. <https://doi.org/10.1002/anie.201902380>

**Important note**

To cite this publication, please use the final published version (if applicable).  
Please check the document version above.

**Copyright**

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

**Takedown policy**

Please contact us and provide details if you believe this document breaches copyrights.  
We will remove access to the work immediately and investigate your claim.

## Biocatalysis

International Edition: DOI: 10.1002/anie.201902380  
German Edition: DOI: 10.1002/ange.201902380**Formate Oxidase (FOx) from *Aspergillus oryzae*: One Catalyst Enables Diverse H<sub>2</sub>O<sub>2</sub>-Dependent Biocatalytic Oxidation Reactions**Florian Tieves<sup>†</sup>, Sébastien Jean-Paul Willot<sup>†</sup>, Morten Martinus Cornelis Harald van Schie, Marine Charlene Renée Rauch, Sabry Hamdy Hamed Younes, Wuyuan Zhang, JiaJia Dong, Patricia Gomez de Santos, John Mick Robbins, Bettina Bommarius, Miguel Alcalde, Andreas Sebastian Bommarius, and Frank Hollmann\*

**Abstract:** An increasing number of biocatalytic oxidation reactions rely on H<sub>2</sub>O<sub>2</sub> as a clean oxidant. The poor robustness of most enzymes towards H<sub>2</sub>O<sub>2</sub>, however, necessitates more efficient systems for in situ H<sub>2</sub>O<sub>2</sub> generation. In analogy to the well-known formate dehydrogenase to promote NADH-dependent reactions, we here propose employing formate oxidase (FOx) to promote H<sub>2</sub>O<sub>2</sub>-dependent enzymatic oxidation reactions. Even under non-optimised conditions, high turnover numbers for coupled FOx/peroxygenase catalysis were achieved.

**E**nzymatic oxidation and oxyfunctionalisation reactions are currently receiving tremendous interest in the context of preparative organic chemistry.<sup>[1]</sup> Especially if selectivity is desired, enzymatic reactions often excel over the chemical counterparts. Amongst available biocatalysts, monooxygenases are of particular interest.<sup>[2]</sup> Monooxygenases, however,

rely on molecular oxygen, which is reductively activated at the enzyme active site. The reducing equivalents required are mostly derived more or less directly from reduced nicotinamide cofactors (NAD(P)H). While issues regarding the in situ regeneration of NAD(P)H have largely been solved, so that it can be used catalytic amounts,<sup>[3]</sup> the so-called oxygen dilemma poses a more severe challenge:<sup>[4]</sup> many monooxygenases cannot utilise NAD(P)H directly but depend on single-electron mediators to transform the hydride transfer from NAD(P)H into two successive single-electron transport events. The reduced mediators, however, also directly interact with dissolved molecular oxygen and are re-oxidised (Scheme 1 a). As a consequence, reactive oxygen species are formed in a futile cycle that uncouples the regeneration reaction from the oxygenation reaction. In extreme cases, up to 95 % of the reducing equivalents provided by the co-substrate are wasted.<sup>[4]</sup>

A viable solution of the challenge outlined above is to make use of H<sub>2</sub>O<sub>2</sub>-dependent “Peroxisomes”.<sup>[5]</sup> By using H<sub>2</sub>O<sub>2</sub>, Peroxisomes actually make use of the oxygen dilemma instead of being hampered by it (Scheme 1 b).

A range of enzymes are able to use H<sub>2</sub>O<sub>2</sub>. Peroxidases for example, represent model enzymes for H<sub>2</sub>O<sub>2</sub>-dependent oxidation,<sup>[6]</sup> polymerization,<sup>[7]</sup> or halogenation reactions.<sup>[8]</sup> More recently, peroxygenases have received a lot of attention for selective oxyfunctionalisation reactions.<sup>[9]</sup> In addition to these reactions, hydrolase-catalysed formation of peracids for H<sub>2</sub>O<sub>2</sub>-driven epoxidation<sup>[10]</sup> and Baeyer–Villiger oxidations<sup>[11]</sup> are becoming popular.

Enzymes, however, are also prone to oxidative inactivation by H<sub>2</sub>O<sub>2</sub><sup>[12]</sup> which is why a broad range of in situ H<sub>2</sub>O<sub>2</sub> generation methods have been investigated in recent years (Table S2 compares some established systems with respect to efficiency and waste generation). The goal is to provide the production enzymes with H<sub>2</sub>O<sub>2</sub> at rates that allow high catalytic turnover while minimising the undesired oxidative inactivation by excess H<sub>2</sub>O<sub>2</sub>.<sup>[13]</sup> Today, glucose oxidase (GOx) is the catalyst of choice for in situ H<sub>2</sub>O<sub>2</sub> generation.<sup>[14]</sup> It couples the oxidation of glucose to the reductive activation of O<sub>2</sub> to form H<sub>2</sub>O<sub>2</sub> in a highly efficient and robust fashion. The GOx system, however, suffers from high levels of waste generation (196 g of gluconate waste per mol H<sub>2</sub>O<sub>2</sub> equivalent are generated).<sup>[15]</sup> Additionally, practical issues such as the high viscosity of the reaction medium have to be dealt with at larger reaction scales. Formate would be a more suitable reductant for the reductive activation of O<sub>2</sub> (generating only

[\*] Dr. F. Tieves,<sup>[†]</sup> S. J. Willot,<sup>[†]</sup> M. M. C. H. van Schie, M. C. R. Rauch, Dr. S. H. H. Younes, Dr. W. Zhang, Dr. J. J. Dong, Dr. F. Hollmann  
Department of Biotechnology, University of Technology Delft  
van der Massweg 9, 2629HZ Delft (The Netherlands)  
E-mail: f.hollmann@tudelft.nl

Dr. S. H. H. Younes  
Chemistry Department, Faculty of Science, Sohag University  
Sohag 82524 (Egypt)

P. Gomez de Santos, Prof. Dr. M. Alcalde  
Department of Biocatalysis, Institute of Catalysis, CSIC  
28049 Madrid (Spain)

Dr. J. M. Robbins, Dr. B. Bommarius, Prof. Dr. A. S. Bommarius  
School of Chemical and Biomolecular Engineering, Georgia Institute  
of Technology  
311 Ferst Drive, N.W., Atlanta, GA 30332 (USA)


Prof. Dr. A. S. Bommarius  
School of Chemistry and Biochemistry  
Georgia Institute of Technology  
901 Atlantic Drive, N.W., Atlanta, GA 30332 (USA)

[†] These authors contributed equally to this work.


Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:  
<https://doi.org/10.1002/anie.201902380>.

© 2019 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial, and no modifications or adaptations are made.

## a) Monooxygenase-catalysed oxyfunctionalisation reactions



## b) Peroxizyme-catalysed oxyfunctionalisation reactions




**Scheme 1.** Biocatalytic oxyfunctionalisations using monooxygenases (a) or peroxizymes (b). Monooxygenases often are prone to the oxygen dilemma while peroxizymes productively make use of the oxygen dilemma.

44 g of volatile and therefore not accumulating CO<sub>2</sub> waste per mol H<sub>2</sub>O<sub>2</sub> equivalent). The systems available today, however, either rely on bioincompatible transition-metal catalysts,<sup>[16]</sup> or are too complex<sup>[15,17]</sup> or too elaborate<sup>[18]</sup> to be practical.

Recently, a formic acid oxidase from *Aspergillus oryzae* (AoFOx) has been reported as the first member of the glucose-methanol-choline (GMC) oxidoreductase superfamily that oxidizes formic acid instead of simple alcohols.<sup>[19]</sup> This enzyme features an optimum pH range from 2.8–6.8 and a  $k_{\text{cat}}$  value of 82 s<sup>-1</sup> over that range. It contains an unusual 8-formyl flavin adenine dinucleotide (FAD) cofactor, which is formed in situ from FAD through self-oxidation. Its unique catalytic properties render AoFOx a promising candidate for H<sub>2</sub>O<sub>2</sub>-dependent enzymatic reactions. We therefore set out to evaluate the potential of AoFOx as a catalyst to promote H<sub>2</sub>O<sub>2</sub>-dependent biocatalytic oxidation reactions (Scheme 2).

AoFOx was prepared according to a previously published procedure.<sup>[19]</sup> In short, AoFOx was expressed in recombinant *Escherichia coli* and partially purified to remove catalase. Overall, from 1 L culture broth, 38 mg of purified enzyme were obtained within 1 day (Figure S1 in the Supporting Information).


Having AoFOx in hand, we decided to first apply this enzyme for some selective oxyfunctionalisation reactions catalysed by the recombinant evolved unspecific peroxygenase from *Agrocybe aegerita* (rAaeUPO) heterologously expressed in *Pichia pastoris*.<sup>[20]</sup> As model reaction, we first focused on the selective hydroxylation of ethyl benzene into (*R*)-1-phenylethanol. A preliminary optimisation of the reaction conditions (Figure 1) revealed that the bienzymatic



**Scheme 2.** The formate oxidase from *Aspergillus oryzae* (AoFOx) enables in situ H<sub>2</sub>O<sub>2</sub> generation from formate and ambient oxygen to promote a broad range of biocatalytic oxidation/oxyfunctionalisation reactions.

cascade operates optimally in slightly acidic reaction media (pH 6, Figure 1a, Figure S2), which is in line with the reported preferences of the enzymes.<sup>[21]</sup> An apparent optimal temperature of 25 °C was determined (Figure 1b, Figure S3). Between 20 and 35 °C, the initial rates of the overall system were largely temperature-independent (Figure S2) but the reaction ceased sooner at elevated temperatures. At 40 °C for example, no further product formation was observed after 2 h. In contrast, steady product accumulation occurred at 30 °C or lower. This behaviour can be attributed to the comparably poor thermal robustness of wt-AoFOx.

We determined an apparent optimal formate concentration of 200 mM (Figure 1d, Figure S4), which represents a compromise between the relatively high  $K_M$  value of wt-AoFOx at this pH<sup>[19d-f]</sup> and the decreasing peroxygenative activity of UPOs at higher formate concentrations.<sup>[22]</sup>



**Figure 1.** Characterisation of the reaction parameters that influence the efficiency of the bienzymatic hydroxylation of ethyl benzene. Individual reaction conditions are given in the captions of Figures S2–8.

The relative ratio of ( $\text{H}_2\text{O}_2$ -generating) *AoFOx* and ( $\text{H}_2\text{O}_2$ -consuming) *rAaeUPO* had a very pronounced effect on the efficiency of the overall reaction system (Figure 1c, Figure S5). The highest initial rate was observed at an equimolar ratio of the two enzymes, albeit at the expense of poor long-term stability of the overall system (after 5 h, no further product formation was observed; Figure S5). Lower ratios of *AoFOx* to *rAaeUPO* gave lower productivity but significantly greater robustness. At a ratio of 1:5, stable product formation for at least 24 h was observed.

The availability of molecular oxygen had a significant influence on the overall reaction (Figure 1e). Under ambient atmosphere without stirring, an  $\text{O}_2$  transfer rate of  $0.84 \pm 0.03 \text{ mmh}^{-1}$  was estimated (Figure S7), which limits the productivity of the overall system. Increasing the  $\text{O}_2$  availability by increasing the  $\text{O}_2$  partial pressure in the headspace of the reaction dramatically increased the productivity of the overall reaction more than ten-fold (Figure 1e, Figure S8).


It is worth mentioning that appropriate negative controls (i.e. reactions leaving out either one of the enzymes or reactions in the absence of formate) were performed for all of the reactions reported. With the sole exception of CytC-catalysed sulfoxidation, where traces of sulfoxide were also observed in the absence of CytC, the control reactions gave no product formation.

Next, we explored the enzyme and product scope of the *AoFOx*-catalysed  $\text{H}_2\text{O}_2$  generation system to promote various  $\text{H}_2\text{O}_2$ -dependent biocatalytic oxidation reactions (Figure 2).

First, we investigated some peroxygenase-catalysed hydroxylation and epoxidation reactions. The proposed  $\text{H}_2\text{O}_2$ -generation system enabled excellent catalytic performance of the peroxygenase used. Both product concentrations and *rAaeUPO*-turnover numbers were at least as high as for previous methods using more complicated  $\text{H}_2\text{O}_2$  generation systems.<sup>[15,18,20d,23]</sup>

The stereospecific hydroxylation of ethyl benzene was performed on a semi-preparative scale, yielding 434 mg of (*R*)-1-phenylethanol (>99% *ee*, see the Supporting Information for further details). A very satisfactory turnover number for the *AoFOx* of more than 300 000 was achieved, which suggests that this in situ  $\text{H}_2\text{O}_2$  generation system is economically feasible. It is also worth mentioning that up to  $31 \pm 3 \text{ mM}$  (*R*)-1-phenylethanol was produced (Figure S6), which is one of the highest numbers observed so far using *rAaeUPO*.<sup>[15]</sup> It should be mentioned here that in case of volatile reagents, imperfect mass balances were observed upon prolonged reaction times. We believe that this is a technical issue that will be overcome in future scale-up experiments.

Cytochrome C (CytC), another heme-containing protein capable of catalysing  $\text{H}_2\text{O}_2$ -driven oxygen transfer reactions, especially sulfoxidation,<sup>[16]</sup> was evaluated next. Compared to the turnover numbers observed with *rAaeUPO*, the numbers achieved with CytC appear rather low. However, these numbers are still significantly higher than those achieved previously using other  $\text{H}_2\text{O}_2$ -generation systems.<sup>[16]</sup> The lack of enantioselectivity in the sulfoxidation of thioanisole is in accordance with previous reports.<sup>[16]</sup> It should be kept in mind here that the natural role of CytC is not that of an enzyme but rather that of an electron-transport protein.



**Figure 2.** The scope of *AoFOx*-driven peroxzyme reactions: The peroxygenase from *Agroclybe aegerita* (*rAaeUPO*) enabled selective hydroxylations and epoxidations; lipase B from *Candida antarctica* (CalB) mediated chemoenzymatic epoxidation and Baeyer–Villiger oxidations; cytochrome C (CytC, a heme-containing electron-transport protein) was applied to the sulfoxidation of thioanisole; and V-dependent chloroperoxidase from *Curcuvaria inaequalis* (CVCPO)-initiated hydroxylation and halolactonisation reactions. For details about the reaction schemes and experimental results, please refer to the respective section in the Supporting Information. Yields shown are calculated from the product concentration divided by the initial starting material concentration.

Another important  $\text{H}_2\text{O}_2$ -driven reaction is the so-called perhydrolase reaction of lipases.<sup>[1b,24]</sup> In short, a lipase catalyses the perhydrolysis of carboxylic (esters) to yield a reactive peracid, which in turn can undergo Baeyer–Villiger oxidations of ketones or Prilezhaev oxidations of  $\text{C}=\text{C}$ -double bonds. Our proposed *AoFOx*  $\text{H}_2\text{O}_2$ -generation system proved to be applicable in principle to drive these reactions (Figure 2). Using the lipase B from *Candida antarctica*, CalB) together with octanoic acid as cocatalyst gave catalytic turnover in the chemoenzymatic Baeyer–Villiger oxidation of cyclohexanone as well as the chemoenzymatic epoxidation of styrene. However, compared to the other systems investigated here, rather low turnover numbers for the biocatalyst were observed. This can be attributed to the low affinity of CalB towards  $\text{H}_2\text{O}_2$  in aqueous systems<sup>[25]</sup> resulting in low CalB activity under the conditions chosen. Further investigations aiming at higher in situ  $\text{H}_2\text{O}_2$  concentrations are currently ongoing.

Finally, we evaluated *AoFOx* to promote halogenation reactions catalysed by the V-dependant haloperoxidase from

*Curvularia inaequalis*.<sup>[8b,c,26]</sup> The hydroxyhalogenation of styrene gave acceptable results in terms of product yield and catalyst performance. Again, the volatility of the reagents impaired the final product concentration and thereby the catalytic numbers. A completely different picture evolved, however, when using 4-pentenoic acid as starting material. Here, a perfect mass balance was observed and full conversion of the starting material into the desired bromolactone was observed. We also scaled up this reaction to the gram scale: Starting from 200 mm 4-pentenoic acid, 150 mm of the desired bromolactone was obtained, which could be separated from the reaction mixture by simple extraction (Figure S14). Thus, 1.6 g of the pure product was obtained.<sup>[27]</sup>

In conclusion, we present herein the proof-of-concept for a simple H<sub>2</sub>O<sub>2</sub> generation system based on formate oxidases such as the FOx from *Aspergillus oryzae* (AoFOx). This system stands out in terms of practical simplicity and excellent performance, even at this early stage of development. Furthermore, the turnover numbers achieved with AoFOx exceed those of established systems by orders of magnitude (Table S2).

Three decades ago, the introduction of formate dehydrogenases as NADH regeneration catalysts ushered in a new era in bioreduction catalysis.<sup>[28]</sup> We are convinced that formate oxidases will have a similar impact for biooxidation/function-alisation catalysis. Further developments in our laboratories will focus on further engineering AoFOx (in particular, a lower K<sub>M</sub> value towards formate is highly desirable) and further characterisation and optimisation of the synthetic schemes to fully explore its synthetic potential.

## Acknowledgements

The authors gratefully acknowledge funding by the European Research Commission (ERC consolidator grant, No. 648026), the European Union (H2020-BBI-PPP-2015-2-1-720297), the Netherlands Organisation for Scientific Research (VICI grant No. 724.014.003), the National Science Foundation (NSF) of the United States (grant IIP-1540017) and the Comunidad de Madrid Synergy CAM Project Y2018/BIO-4738-EVOCHIMERA-CM.

## Conflict of interest

The authors declare no conflict of interest.

**Keywords:** biocatalysis · formate oxidase · hydrogen peroxide · oxidation · oxyfunctionalisation

**How to cite:** *Angew. Chem. Int. Ed.* **2019**, *58*, 7873–7877  
*Angew. Chem.* **2019**, *131*, 7955–7959

- [1] a) Y. Liang, J. Wei, X. Qiu, N. Jiao, *Chem. Rev.* **2018**, *118*, 4912–4945; b) J. Dong, E. Fernández-Fueyo, F. Hollmann, C. Paul, M. Pesic, S. Schmidt, Y. Wang, S. Younes, W. Zhang, *Angew. Chem. Int. Ed.* **2018**, *57*, 9238–9261; *Angew. Chem.* **2018**, *130*, 9380–9404; c) N. D. Fessner, *ChemCatChem* **2019**, <https://doi.org/10.1002/cctc.201801829>.

- [2] a) E. Roduner, W. Kaim, B. Sarkar, V. B. Urlacher, J. Pleiss, R. Gläser, W.-D. Einicke, G. A. Sprenger, U. Beifuß, E. Klemm, C. Liebner, H. Hieronymus, S.-F. Hsu, B. Plietker, S. Laschat, *ChemCatChem* **2013**, *5*, 82–112; b) V. B. Urlacher, M. Girhard, *Trends Biotechnol.* **2012**, *30*, 26–36; c) B. M. Nestl, S. C. Hammer, B. A. Nebel, B. Hauer, *Angew. Chem. Int. Ed.* **2014**, *53*, 3070–3095; *Angew. Chem.* **2014**, *126*, 3132–3158; d) R. Bernhardt, V. B. Urlacher, *Appl. Microbiol. Biotechnol.* **2014**, *98*, 6185–6203; e) R. Fasan, *ACS Catal.* **2012**, *2*, 647–666; f) S. T. Jung, R. Lauchli, F. H. Arnold, *Curr. Opin. Biotechnol.* **2011**, *22*, 809–817.
- [3] a) S. Kara, J. H. Schrittwieser, F. Hollmann, M. B. Ansorge-Schumacher, *Appl. Microbiol. Biotechnol.* **2014**, *98*, 1517–1529; b) A. Weckbecker, H. Gröger, W. Hummel in *Biosystems Engineering I: Creating Superior Biocatalysts*, Vol. 120, Springer-Verlag Berlin, Berlin, **2010**, pp. 195–242.
- [4] D. Holtmann, F. Hollmann, *ChemBioChem* **2016**, *17*, 1391–1398.
- [5] As Peroxizymes, we define all enzymes that (in principle) can use H<sub>2</sub>O<sub>2</sub> as stoichiometric oxidant. Examples are the classical peroxidases and peroxygenases but also other heme-containing biocatalysts such as P450 monooxygenases and also hydrolases.
- [6] F. van Rantwijk, R. A. Sheldon, *Curr. Opin. Biotechnol.* **2000**, *11*, 554–564.
- [7] F. Hollmann, I. W. C. E. Arends, *Polymers* **2012**, *4*, 759–793.
- [8] a) C. J. Seel, A. Králík, M. Hacker, A. Frank, B. König, T. Gulder, *ChemCatChem* **2018**, *10*, 3960–3963; b) J. J. Dong, E. Fernández-Fueyo, J. Li, Z. Guo, R. Renirie, R. Wever, F. Hollmann, *Chem. Commun.* **2017**, *53*, 6207–6210; c) E. Fernández-Fueyo, M. van Wingerden, R. Renirie, R. Wever, Y. Ni, D. Holtmann, F. Hollmann, *ChemCatChem* **2015**, *7*, 4035–4038; d) D. Wischang, J. Hartung, *Tetrahedron* **2012**, *68*, 9456–9463.
- [9] a) M. Hofrichter, R. Ullrich, *Curr. Opin. Chem. Biol.* **2014**, *19*, 116–125; b) Y. Wang, D. Lan, R. Durrani, F. Hollmann, *Curr. Opin. Chem. Biol.* **2017**, *37*, 1–9.
- [10] a) J. Meyer-Wassewitz, D. Hohmann, M. B. Ansorge-Schumacher, M. Kraume, A. Drews, *Biochem. Eng. J.* **2017**, *126*, 68–77; b) P. Zhou, X. Wang, C. Zeng, W. Wang, B. Yang, F. Hollmann, Y. Wang, *ChemCatChem* **2017**, *9*, 934–936.
- [11] a) M. Markiton, S. Boncel, D. Janas, A. Chrobok, *ACS Sustainable Chem. Eng.* **2017**, *5*, 1685–1691; b) A. R. S. Teixeira, A. Flourat, A. M. Peru, F. Brunissen, F. Allais, *Front. Chem.* **2016**, *4*, 11; c) A. Drozd, A. Chrobok, *Chem. Commun.* **2016**, *52*, 1230–1233.
- [12] B. Valderrama, M. Ayala, R. Vazquez-Duhalt, *Chem. Biol.* **2002**, *9*, 555–565.
- [13] F. Sabuzi, E. Churakova, P. Galloni, R. Wever, F. Hollmann, B. Floris, V. Conte, *Eur. J. Inorg. Chem.* **2015**, 3519–3525.
- [14] R. A. Sheldon, P. C. Pereira, *Chem. Soc. Rev.* **2017**, *46*, 2678–2691.
- [15] Y. Ni, E. Fernández-Fueyo, A. G. Baraibar, R. Ullrich, M. Hofrichter, H. Yanase, M. Alcalde, W. J. H. van Berkel, F. Hollmann, *Angew. Chem. Int. Ed.* **2016**, *55*, 798–801; *Angew. Chem.* **2016**, *128*, 809–812.
- [16] F. Hollmann, A. Schmid, *J. Inorg. Biochem.* **2009**, *103*, 313–315.
- [17] J. Rocha-Martin, S. Velasco-Lozano, J. M. Guisan, F. Lopez-Gallego, *Green Chem.* **2014**, *16*, 303–311.
- [18] a) W. Zhang, E. Fernández-Fueyo, Y. Ni, M. van Schie, J. Gacs, R. Renirie, R. Wever, F. G. Mutti, D. Rother, M. Alcalde, F. Hollmann, *Nat. Catal.* **2018**, *1*, 55–62; b) W. Zhang, B. O. Burek, E. Fernández-Fueyo, M. Alcalde, J. Z. Bloh, F. Hollmann, *Angew. Chem. Int. Ed.* **2017**, *56*, 15451–15455; *Angew. Chem.* **2017**, *129*, 15654–15658.
- [19] a) D. Doubayashi, T. Ootake, Y. Maeda, M. Oki, Y. Tokunaga, A. Sakurai, Y. Nagaosa, B. Mikami, H. Uchida, *Biosci. Biotechnol. Biochem.* **2011**, *75*, 1662–1667; b) Y. Maeda, D. Doubayashi, M. Oki, H. Nose, A. Sakurai, K. Isa, Y. Fujii, H.

- Uchida, *Biosci. Biotechnol. Biochem.* **2009**, *73*, 2645–2649; c) Y. Maeda, D. Doubayashi, M. Oki, H. Nose, Y. Fujii, H. Uchida, *J. Biosci. Bioeng.* **2009**, *108*, S106–S106; d) J. M. Robbins, A. S. Bommarius, G. Gadda, *Arch. Biochem. Biophys.* **2018**, *643*, 24–31; e) J. M. Robbins, M. G. Souffrant, D. Hamelberg, G. Gadda, A. S. Bommarius, *Biochemistry* **2017**, *56*, 3800–3807; f) J. M. Robbins, J. Geng, B. A. Barry, G. Gadda, A. S. Bommarius, *Biochemistry* **2018**, *57*, 5818–5826.
- [20] a) P. Molina-Espeja, E. Garcia-Ruiz, D. Gonzalez-Perez, R. Ullrich, M. Hofrichter, M. Alcalde, *Appl. Environ. Microbiol.* **2014**, *80*, 3496–3507; b) P. Molina-Espeja, P. G. de Santos, M. Alcalde in *Directed Enzyme Evolution: Advances and Applications* (Ed.: M. Alcalde), Springer International Publishing, Cham, **2017**, pp. 127–143; c) P. Molina-Espeja, S. Ma, D. M. Mate, R. Ludwig, M. Alcalde, *Enzyme Microb. Technol.* **2015**, *73–74*, 29–33; d) P. Gomez de Santos, M. Canellas, F. Tieves, S. H. H. Younes, P. Molina-Espeja, M. Hofrichter, F. Hollmann, V. Guallar, M. Alcalde, *ACS Catal.* **2018**, *8*, 4789–4799.
- [21] R. Ullrich, J. Nüske, K. Scheibner, J. Spantzel, M. Hofrichter, *Appl. Environ. Microbiol.* **2004**, *70*, 4575–4581.
- [22] D. I. Perez, M. Mifsud Grau, I. W. C. E. Arends, F. Hollmann, *Chem. Commun.* **2009**, 6848–6850.
- [23] a) S. Peter, M. Kinne, R. Ullrich, G. Kayser, M. Hofrichter, *Enzyme Microb. Technol.* **2013**, *52*, 370–376; b) M. Kluge, R. Ullrich, K. Scheibner, M. Hofrichter, *Green Chem.* **2012**, *14*, 440–446.
- [24] a) F. Björkling, H. Frykman, S. E. Godtfredsen, O. Kirk, *Tetrahedron* **1992**, *48*, 4587–4592; b) F. Björkling, S. E. Godtfredsen, O. Kirk, *J. Chem. Soc. Chem. Commun.* **1990**, 1301–1303.
- [25] a) Y. Ma, P. Li, S. J.-P. Willot, W. Zhang, D. Ribitsch, Y. H. Choi, T. Zhang, R. Verpoort, F. Hollmann, Y. Wang, *ChemSusChem* **2019**, <https://doi.org/10.1002/cssc.201900043>; b) P. F. Zhou, D. M. Lan, G. M. Popowicz, X. P. Wang, B. Yang, Y. H. Wang, *Appl. Microbiol. Biotechnol.* **2017**, *101*, 5689–5697; c) P. Zhou, X. Wang, B. Yang, F. Hollmann, Y. Wang, *RSC Adv.* **2017**, *7*, 12518–12523.
- [26] a) R. Wever, M. A. van der Horst, *Dalton Trans.* **2013**, *42*, 11778–11786; b) R. Wever, R. Renirie, *Peroxidases and Catalases: Biochemistry Biophysics, Biotechnology, and Physiology*, Wiley, Hoboken, **2010**, pp. 363–385.
- [27] This corresponds to a 45 % yield of isolated product. However, upon performing a second extraction, more product (albeit contaminated with the starting material) was obtained.
- [28] a) Z. Shaked, G. M. Whitesides, *J. Am. Chem. Soc.* **1980**, *102*, 7104–7105; b) V. O. Popov, V. S. Lamzin, *Biochem. J.* **1994**, *301*, 625–643; c) V. I. Tishkov, V. O. Popov, *Biochemistry* **2004**, *69*, 1252–1253; d) A. S. Bommarius, M. Schwarm, K. Stingl, M. Kottenhahn, K. Huthmacher, K. Drauz, *Tetrahedron: Asymmetry* **1995**, *6*, 2851–2888.

Manuscript received: February 22, 2019

Accepted manuscript online: April 4, 2019

Version of record online: April 30, 2019