

Crustal-scale interferometric imaging in the Malargüe region, Argentina, with P-wave coda and multidimensional deconvolution

Nishitsuji, Yohei; Minato, Shohei; Hartstra, Iris; Boullenger, Boris; Wapenaar, Kees; Gomez, Martín; Draganov, Deyan; Hadziioannou, G

Publication date

2016

Document Version

Final published version

Published in

Geophysical Research Abstracts (online)

Citation (APA)

Nishitsuji, Y., Minato, S., Hartstra, I., Boullenger, B., Wapenaar, K., Gomez, M., Draganov, D., & Hadziioannou, G. (Ed.) (2016). Crustal-scale interferometric imaging in the Malargüe region, Argentina, with P-wave coda and multidimensional deconvolution. *Geophysical Research Abstracts (online)*, 18, 1-1. Article EGU2016-5798.

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.



Crustal-scale interferometric imaging in the Malargüe region, Argentina, with P-wave coda and multidimensional deconvolution

Yohei Nishitsuji (1), Shohei Minato (1), Iris Hartstra (1), Boris Boullenger (1), Kees Wapenaar (1), Martín Gomez (2), and Deyan Draganov (1)

(1) TU Delft, Department of Geoscience and Engineering, Delft, The Netherlands (y.nishitsuji@tudelft.nl), (2) Comisión Nacional de Energía Atómica, International Center for Earth Sciences, Buenos Aires, Argentina

Several seismic investigations - using receiver-function methods as well as tomographic approaches - have been carried out in the Malargüe region (Argentina) for various purposes over a few decades. We use a body-wave seismic interferometry (SI) approach to retrieve reflections later used for the consecutive imaging of the subsurface. We investigate the applicability of the body-wave SI using P-wave coda from local earthquakes with the aim to retrieve reflection responses from a part of the Andean crust below the seismic array we use. We called our technique local-earthquake P-wave coda (LEPC) SI. In this presentation, we show three different LEPC SI results based on three different SI theories: crosscorrelation, crosscoherence, and multidimensional deconvolution. We find that, from a structural-interpretation point of view, multidimensional deconvolution based on the truncated singular-value decomposition scheme provides us with a better structural imaging than the other SI approaches. We interpret deep thrust faults in the imaging results from LEPC SI, whose presence in this region has previously been indicated from interpretation of active seismic-survey data and exploration-well data. We also interpret dimmed-amplitude parts in the reflection image as possible melting zones that have been previously indicated by magnetotelluric methods. The LEPC SI method we propose could be used as a low-cost alternative to active-source seismic surveys for imaging and monitoring purposes of deeper geothermal reservoirs, e.g. in enhanced geothermal systems where the target structures are down to 10 km depth.