

Whole-Cell Photoenzymatic Cascades to Synthesize Long-Chain Aliphatic Amines and Esters from Renewable Fatty Acids

Cha, Hee Jeong; Hwang, Se Yeun; Lee, Da Som; Kumar, Akula Ravi; Kwon, Yong Uk; Voß, Moritz; Schuiten, Eva; Bornscheuer, Uwe T.; Hollmann, Frank; More Authors

DOI

[10.1002/anie.201915108](https://doi.org/10.1002/anie.201915108)

Publication date

2020

Document Version

Final published version

Published in

Angewandte Chemie - International Edition

Citation (APA)

Cha, H. J., Hwang, S. Y., Lee, D. S., Kumar, A. R., Kwon, Y. U., Voß, M., Schuiten, E., Bornscheuer, U. T., Hollmann, F., & More Authors (2020). Whole-Cell Photoenzymatic Cascades to Synthesize Long-Chain Aliphatic Amines and Esters from Renewable Fatty Acids. *Angewandte Chemie - International Edition*, 59(18), 7024-7028. <https://doi.org/10.1002/anie.201915108>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Green Open Access added to TU Delft Institutional Repository

'You share, we take care!' – Taverne project

<https://www.openaccess.nl/en/you-share-we-take-care>

Otherwise as indicated in the copyright section: the publisher is the copyright holder of this work and the author uses the Dutch legislation to make this work public.

Biocatalysis

International Edition: DOI: 10.1002/anie.201915108
German Edition: DOI: 10.1002/ange.201915108

Whole-Cell Photoenzymatic Cascades to Synthesize Long-Chain Aliphatic Amines and Esters from Renewable Fatty Acids

Hee-Jeong Cha, Se-Yeun Hwang, Da-Som Lee, Akula Ravi Kumar, Yong-Uk Kwon, Moritz Voß, Eva Schuitem, Uwe T. Bornscheuer, Frank Hollmann,* Deok-Kun Oh,* and Jin-Byung Park*

Abstract: Long-chain aliphatic amines such as (*S,Z*)-heptadec-9-en-7-amine and 9-aminoheptadecane were synthesized from ricinoleic acid and oleic acid, respectively, by whole-cell cascade reactions using the combination of an alcohol dehydrogenase (ADH) from *Micrococcus luteus*, an engineered amine transaminase from *Vibrio fluvialis* (Vf-ATA), and a photoactivated decarboxylase from *Chlorella variabilis* NC64A (Cv-FAP) in a one-pot process. In addition, long chain aliphatic esters such as 10-(heptanoyloxy)dec-8-ene and octyl-nonanoate were prepared from ricinoleic acid and oleic acid, respectively, by using the combination of the ADH, a Baeyer–Villiger monooxygenase variant from *Pseudomonas putida* KT2440, and the Cv-FAP. The target compounds were produced at rates of up to 37 U g⁻¹ dry cells with conversions up to 90%. Therefore, this study contributes to the preparation of industrially relevant long-chain aliphatic chiral amines and esters from renewable fatty acid resources.

Introduction

There is an increasing demand for the preparation of chemical building blocks from biomass-derived starting materials. Renewable oils and fatty acids are particularly interesting because they are abundant in nature and can be derived from non-edible sources. Oils and fatty acids are

already used for the synthesis of a variety of chemicals ranging from cosmetic esters to performance additives.^[1] For instance, hydroxy fatty acids serve as starting materials to manufacture flavors, resins, waxes, lubricants, and polymers, and as additives in coatings and paints.^[1b,2] More recently, hydroxy fatty acids have been used as starting materials in multi-enzyme cascades combining alcohol dehydrogenases (ADH) and amine transaminases (ATA; Scheme S1 in the Supporting Information).^[3] The hydroxy groups can also be converted in a sequence of ADH-catalyzed oxidation followed by a Baeyer–Villiger monooxygenase (BVMO) step; the resulting esters are hydrolyzed into alkanolic acids and ω-hydroxy fatty acids by an esterase.^[4] The ω-hydroxy fatty acids can be oxidized into α,ω-dicarboxylic acids by an alcohol/aldehyde dehydrogenase^[5] or converted into ω-amino fatty acids by an ADH and ω-transaminase.^[5d,6] Further synthetic possibilities comprise transformation into α,ω-diols by combined reduction through carboxylic acid reductases and ADHs,^[7] or into α,ω-diamines using carboxylic acid reductases, ADHs, and ω-transaminase.^[6]

Another promising transformation of fatty acids is decarboxylation to the corresponding C-1-shortened alkanes (Scheme S1). Established chemical methods, however, rely on rather harsh reaction conditions^[8] under which additional functional groups are unlikely to remain unaltered. A very promising solution to this challenge was reported recently by Beisson and co-workers.^[9] The photodecarboxylase from the algae *Chlorella variabilis* NC64A (Cv-FAP) catalyzes the selective light-driven decarboxylation of (fatty) acids^[10] and thereby possibly expands the scope of products derived from renewable oleochemicals significantly.

In this study, a novel fatty acid biotransformation pathway was investigated to produce long-chain aliphatic amines and esters from renewable fatty acids, which can be used as active ingredients in cosmetic formulations and performance additives in oleochemicals.^[1b,c,2b] For instance, the biotransformation of ricinoleic acid (**1**) into (*S,Z*)-heptadec-9-en-7-amine (**5**) and 10-(heptanoyloxy)dec-8-ene (**7**; Scheme 1) and the biotransformation of oleic acid [(*Z*)-octadec-9-enoic acid (**10**)] into 9-aminoheptadecane (**14**) and octylnonanoate (**16**; Scheme 2) were investigated by using Cv-FAP as a key enzyme.

Results and Discussion

In a first set of experiments, we created an *Escherichia coli* based whole-cell catalyst expressing Cv-FAP and tested it for

[*] H.-J. Cha, S.-Y. Hwang, D.-S. Lee, Dr. A. R. Kumar, Prof. Dr. J.-B. Park
Department of Food Science & Engineering
Ewha Womans University, Seoul 03760 (Republic of Korea)
E-mail: jbpark06@ewha.ac.kr


Dr. A. R. Kumar, Prof. Dr. Y.-U. Kwon
Department of Chemistry and Nanoscience
Ewha Womans University, Seoul 03760 (Republic of Korea)

M. Voß, E. Schuitem, Prof. Dr. U. T. Bornscheuer
Institute of Biochemistry
Department of Biotechnology & Enzyme Catalysis
Greifswald University, 17487 Greifswald (Germany)


Prof. Dr. F. Hollmann
Department of Biotechnology, Delft University of Technology
Van der Maasweg 9, 2629HZ Delft (The Netherlands)
E-mail: f.hollmann@tudelft.nl

Prof. Dr. D.-K. Oh
Department of Bioscience and Biotechnology, Konkuk University
Seoul 05029 (Republic of Korea)
E-mail: deokkun@konkuk.ac.kr

Supporting information (including experimental procedures, analytical details, and additional experimental data) and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.201915108>.



Scheme 1. Cascades for the transformation of ricinoleic acid (1) into secondary fatty alcohols (2) or amines (5), fatty acid esters (7), or the corresponding acids (8) and alcohol (9).^[3,5b,11] Details about the catalysts used can be found in Table 2.



Scheme 2. Cascades for the transformation of oleic acid (10) into fatty amines (14), fatty acid esters (16), or the corresponding acids (17) and alcohol (18).^[1f,3,5b] Details about the catalysts used can be found in Table 2.

the transformation of ricinoleic acid (1) into (*Z*)-heptadec-9-en-7-ol (2; Scheme 1).

Since the expression level of a long-chain fatty acid transporter (FadL) in *E. coli* had previously been shown to accelerate whole-cell biotransformation rates of long-chain fatty acids,^[11,12] two types of *E. coli* based biocatalysts (Cat-1 and Cat-2; Table 1) were constructed. Cat-1 is the recombinant *E. coli* expressing the Cv-FAP in the cytoplasm while Cat-2 is the recombinant *E. coli* expressing Cv-FAP in the cytoplasm and FadL in the outer membrane.

Cat-1 almost completely converted ricinoleic acid (1) into (*Z*)-heptadec-9-en-7-ol (2) with in only 15 min (Figure 1, and Table 2, and Figure S1 in the Supporting Information). The transformation took significantly longer (30 min) using Cat-2 under otherwise identical conditions (Figure 1). Similar observations were also made for the transformations of oleic acid and linoleic acid (Figure S2). The lower rates might be due to a negative impact of the expression of FadL on Cv-FAP expression (Figure S3). Therefore, we used Cat-1 as a chassis for all further catalyst designs.

Next, we aimed at the conversion of ricinoleic acid (1) into (*S,Z*)-heptadec-9-en-7-amine (5; Cascade 1, Figure 2). For this, we used a previously reported recombinant *E. coli* system (Cat-3; Table 1)^[3] comprising a secondary alcohol dehydrogenase (sADH) with a transaminase (Vf-ATA_{H3-RA}) and the long-chain fatty acid transporter (FadL) with the aforementioned Cat-1. Although the transformation proceeded smoothly to full conversion (Figure S4a), the enantiomeric purity of the desired (*S,Z*)-heptadec-9-en-7-amine (5) was disappointingly low (27% *ee*, Scheme S3 and Figure S5). Therefore, Vf-ATA_{H3-RA} was subjected to structure-based rational engineering to improve its enantioselectivity. Substrate docking revealed that Lys163 may play a crucial role (Figure S6). Indeed, substitution of this amino acid with an aspartic acid (Vf-ATA_{H3-RA K163D}) resulted in a more active and more enantio-

selective (71% *ee*) catalyst (Cat-4; Table 1, Table S2, and Figure S4b). Overall, using the combination of Cat-1 and Cat-4 gave 78% conversion of ricinoleic acid (1) into (*S,Z*)-heptadec-9-en-7-amine (5; 71% *ee*, Table 2).

It is worth noting that at the time the illumination was initiated, not all starting material had been converted into (*S,Z*)-12-aminooctadec-9-enoic acid (4), therefore, trace amounts of the decarboxylation products of the non-reacted starting material and intermediate were also observed (Figure 2).

Table 1: List of the recombinant biocatalysts used in this study.

Name	Description	Remarks
Cat-1	<i>E. coli</i> BL21 (DE3) pET28a-Cv-FAP	Photodecarboxylase (Cv-FAP)
Cat-2	<i>E. coli</i> BL21 (DE3) pACYC-FadL/pET28a-Cv-FAP	Photodecarboxylase (Cv-FAP) + Fatty acid transporter (FadL)
Cat-3	<i>E. coli</i> BL21 (DE3) pACYC-ADH-FadL/pET24b-Vf-ATA _{H3-RA}	Secondary alcohol dehydrogenase (sADH) + Fatty acid transporter (FadL) + Amine transaminase (Vf-ATA _{H3-RA})
Cat-4	<i>E. coli</i> BL21 (DE3) pACYC-ADH-FadL/pET24b-Vf-ATA _{H3-RA} RA K163D	Secondary alcohol dehydrogenase (sADH) + Fatty acid transporter (FadL) + Amine transaminase (Vf-ATA _{H3-RA} K163D)
Cat-5	<i>E. coli</i> BL21 (DE3) pAPTm-E6BVMO _{-C302L} -ADH/pCon-FadL	Secondary alcohol dehydrogenase (sADH) + Fatty acid transporter (FadL) + Baeyer–Villiger monooxygenase (E6BVMO _{-C302L})
Cat-6	<i>E. coli</i> BL21 (DE3) pACYC-ADH-FadL-OhyA/pET24b-Vf-ATA _{H3-RA}	Secondary alcohol dehydrogenase (sADH) + Fatty acid transporter (FadL) + Fatty acid hydratase (OhyA) + Amine transaminase (Vf-ATA _{H3-RA})
Cat-7	<i>E. coli</i> BL21 (DE3) $\Delta infA$ pACYC-ADH-FadL-OhyA/ pSTAPL_E6BVMO _{-C302L} _M340L_Lv3	Secondary alcohol dehydrogenase (sADH) + Fatty acid transporter (FadL) + Fatty acid hydratase (OhyA) + Baeyer–Villiger monooxygenase (E6BVMO _{-C302L} _M340L)

Cv-FAP = photodecarboxylase from *Chlorella variabilis* NC64A. FadL = long chain fatty acid transporter. sADH = secondary alcohol dehydrogenase from *Micrococcus luteus*. Vf-ATA_{H3-RA} = amine transaminase variant from *Vibrio fluvialis*. Vf-ATA_{H3-RA}K163D = the engineered Vf-ATA_{H3-RA} to improve enantioselectivity. E6BVMO_{-C302L} = engineered variant of the Baeyer–Villiger monooxygenase from *Pseudomonas putida* KT2440. OhyA = fatty acid double-bond hydratase from *Stenotrophomonas maltophilia*. See the Table S1 for details of the plasmids.

In another example we evaluated the conversion of ricinoleic acid (**1**) into (*Z*)-10-(heptanoyloxy)dec-8-ene (**7**; Cascade 2, Figure 3). For this Cat-1 and Cat-5 were combined. The latter comprised recombinant *E. coli* expressing the




Figure 1. Biotransformation of ricinoleic acid (**1**, black circles) into (*Z*)-heptadec-9-en-7-ol (**2**, blue squares) using Cat-1 (closed symbols) or Cat-2 (open symbols). Conditions: $c(\mathbf{1}) = 10 \text{ mM}$, $c(\text{Cat-1})$ or $c(\text{Cat-2}) = 3.6 \text{ g}_{\text{CDW}} \text{ L}^{-1}$, illumination with blue light ($\lambda = 450 \text{ nm}$).

FadL,^[4,12,13] sADH and a Baeyer–Villiger monooxygenase variant from *Pseudomonas putida* KT2440 (E6BVMO_{-C302L}). The biotransformation was initiated by adding ricinoleic acid (**1**, 15 mM) to the culture broth of Cat-5 (Figure 3). After 8 h, the transformation of ricinoleic acid into 11-(heptanoyloxy)undec-9-enoic acid (**6**) was nearly complete and Cat-1 was added and illumination was initiated for another 4 h. 9.7 mM of the desired product (*Z*)-10-(heptanoyloxy)dec-8-ene (**7**) was obtained (Figure 3 and Figure S8a). The overall isolated yield from ricinoleic acid was 52% with a purity of 92% (Table 2 and Figure S8b). Upon addition of the esterase from *Pseudomonas fluorescens* SIK WI (*Pf*-Esterase), the ester product (**7**) could also be hydrolyzed into *n*-heptanoic acid (**8**) and 2-decen-1-ol (**9**) from the ester compound (**7**; Scheme 1 and Figure S8c).

A key element of the transformations shown above is the hydroxy functionality present in ricinoleic acid. To broaden the substrate scope, we decided to enlarge the cascades with a C=C-double bond hydration step (Scheme 2). To achieve this, Cat-3 and Cat-5 were supplemented with a fatty acid double bond hydratase (OhyA) from *Stenotrophomonas maltophilia* generating Cat-6 and Cat-7 (Table 1).

Combining Cat-6 with Cat-1 enabled us to transform oleic acid (**10**, 5.2 mM) into 9-aminoheptadecane (**14**; Cascade 3, Figure 4a). In contrast to Cascade 1, the amine donor for the reductive amination step could not be added from the beginning of the reaction since it severely inhibited OhyA.^[3] Nevertheless, the target product 9-aminoheptadecane (**14**) was obtained in an overall 58% conversion (at 3 mM; Table 2).

Finally, we performed the transformation of oleic acid (**10**, 10.5 mM) into octylnonanoate (**16**; Cascade 4, Figure 4b). Using a combination of Cat-7 and Cat-1, octylnonanoate (**16**)

Table 2: Summary of the products obtained through the presented photoenzymatic cascades.

Cascade	Product	Yield [mM]	Conversion [%]
–	(<i>R,Z</i>)-heptadec-9-en-7-ol (2)	9.1	91
1	(<i>S,Z</i>)-heptadec-9-en-7-amine (5)	3.9	78
2	10-(heptanoyloxy)dec-8-ene (7)	9.7	65
3	9-aminoheptadecane (14)	3	58
4	octylnonanoate (16)	7.2	69




Figure 2. Cascade-1: Biotransformation of ricinoleic acid (1, black ●) into (S,Z)-heptadec-9-en-7-amine (5, blue ▲) via (Z)-12-oxooctadec-9-enoic acid (3, green ▲) and (S,Z)-12-aminooctadec-9-enoic acid (4, red ■). Conditions: $c(1) = 5 \text{ mM}$, $c(\text{Cat-1}) = 3.6 \text{ g}_{\text{CDW}} \text{ L}^{-1}$, $c(\text{Cat-4}) = 7.2 \text{ g}_{\text{CDW}} \text{ L}^{-1}$, $c(\text{methyl benzyl amine}) = 10 \text{ mM}$, illumination with blue light ($\lambda = 450 \text{ nm}$) was initiated at $t = 2 \text{ h}$. Minor amounts of the decarboxylation products of ○ and ▲ were observed as well.




Figure 3. Cascade-2: Biotransformation of ricinoleic acid (1, black ●) into 10-(heptanoyloxy)dec-8-ene (7, blue ◆) via (Z)-12-oxooctadec-9-enoic acid (3, green ▲) and 11-heptanoyloxyundec-9-enoic acid (6, red ■). Conditions: $c(1) = 15 \text{ mM}$, $c(\text{Cat-1}) = 3.6 \text{ g}_{\text{CDW}} \text{ L}^{-1}$, $c(\text{Cat-5}) = 3.6 \text{ g}_{\text{CDW}} \text{ L}^{-1}$, illumination with blue light ($\lambda = 450 \text{ nm}$) was initiated at $t = 8 \text{ h}$. Minor amounts of the decarboxylation products of 1 (○) and 3 (▲) were observed as well.

was obtained at an overall conversion of 69% (7.2 mM) from oleic acid (Table 2). The ester (16) was isolated in an overall yield of 48% with a purity of 90% (Figure S10c). Addition of *Pf*-esterase into the reaction medium resulted in complete hydrolysis of 16 into *n*-nonanoic acid (17) and *n*-octanol (18).

Conclusion

These results demonstrate that long-chain aliphatic amines (e.g., (S,Z)-heptadec-9-en-7-amine and 9-aminoheptadecane) can be produced from renewable fatty acids (e.g., ricinoleic acid and oleic acid) by whole-cell biocatalytic





Figure 4. a) Cascade-3: Biotransformation of oleic acid (10, black ●) into 9-aminoheptadecane (14, blue ▲) via (R)-10-hydroxyoctadecanoic acid (11, grey ▲), 10-oxooctadecanoic acid (12, green ▼), and 10-aminooctadecanoic acid (13, red ■). Conditions: $c(10) = 5.2 \text{ mM}$, $c(\text{Cat-1}) = 3.6 \text{ g}_{\text{CDW}} \text{ L}^{-1}$, $c(\text{Cat-6}) = 7.2 \text{ g}_{\text{CDW}} \text{ L}^{-1}$, illumination with blue light ($\lambda = 450 \text{ nm}$) was initiated at $t = 6 \text{ h}$. b) Cascade-4: Biotransformation of oleic acid (10, black ●) into octylnonanoate (16, blue ◆) via (R)-10-hydroxyoctadecanoic acid (11, grey ▲), 10-oxooctadecanoic acid (12, green ▼), and 9-(nonanoyloxy)nonanoic acid (15, red ■). Conditions: $c(10) = 10.5 \text{ mM}$, $c(\text{Cat-1}) = 3.6 \text{ g}_{\text{CDW}} \text{ L}^{-1}$, $c(\text{Cat-7}) = 3.6 \text{ g}_{\text{CDW}} \text{ L}^{-1}$, illumination with blue light ($\lambda = 450 \text{ nm}$) was initiated at $t = 7 \text{ h}$.

casades using the combination of the enzymes sADH, Vf-ATA, and Cv-FAP in a one-pot process. In addition, long-chain aliphatic esters (e.g., 10-(heptanoyloxy)dec-8-ene and octylnonanoate) were prepared from renewable fatty acids by using a combination of sADH, E6BVMO_{C302L}, and Cv-FAP. The target products were synthesized at a rather high rate with high conversion. This study will contribute to the preparation of industrially relevant long-chain aliphatic amines and esters from renewable oils and fatty acids.

Acknowledgements

This work was supported by C1 Gas Refinery Research Center (NRF Grant Number: 2018M3D3A1A01055735-3) of the National Research Foundation (NRF) of Korea funded by Ministry of Science and ICT and by the Marine Biomaterials

Research Center grant from the Marine Biotechnology Program [No. D11013214H480000100] funded by the Ministry of Oceans and Fisheries, Korea. F.H. gratefully acknowledges support by The Netherlands Organisation for Scientific Research through a VICI grant (No. 724.014.003) and the European Research Commission (ERC consolidator grant No. 648026). U.T.B. gratefully acknowledges support by the European Union's Horizon 2020 Programme (Marie-Curie Actions, grant No. 722610) and the German Research Foundation, DFG (grant No.: BO 1862/16-1).

Conflict of interest

The authors declare no conflict of interest.

Keywords: Biocatalysis · fatty acids · long-chain aliphatic amines · photodecarboxylase · whole-cell biocatalysis

How to cite: *Angew. Chem. Int. Ed.* **2020**, *59*, 7024–7028
Angew. Chem. **2020**, *132*, 7090–7094

- [1] a) M. A. Meier, J. O. Metzger, U. S. Schubert, *Chem. Soc. Rev.* **2007**, *36*, 1788–1802; b) U. Biermann, U. Bornscheuer, M. A. Meier, J. O. Metzger, H. J. Schäfer, *Angew. Chem. Int. Ed.* **2011**, *50*, 3854–3871; *Angew. Chem.* **2011**, *123*, 3938–3956; c) U. T. Bornscheuer, *Lipid Modification by Enzymes and Engineered Microbes*, Elsevier, Amsterdam, **2018**; d) M. Schrewe, M. K. Julsing, K. Lange, E. Czarnotta, A. Schmid, B. Bühler, *Biotechnol. Bioeng.* **2014**, *111*, 1820–1830; e) S. Koppireddi, J. H. Seo, E. Y. Jeon, P. S. Chowdhury, H. Y. Jang, J. B. Park, Y. U. Kwon, *Adv. Synth. Catal.* **2016**, *358*, 3084–3092; f) E.-Y. Jeon, J.-H. Seo, W.-R. Kang, M.-J. Kim, J.-H. Lee, D.-K. Oh, J.-B. Park, *ACS Catal.* **2016**, *6*, 7547–7553; g) H.-Y. Jang, K. Singha, H.-H. Kim, Y.-U. Kwon, J.-B. Park, *Green Chem.* **2016**, *18*, 1089–1095; h) M. B. Ansorge-Schumacher, O. Thum, *Chem. Soc. Rev.* **2013**, *42*, 6475–6490; i) S. Kern, A. Himmelspach, K. Grammann, O. Thum, A. Liese, *Org. Process Res. Dev.* **2016**, *20*, 1930–1936; j) J. Y. Park, S. H. Lee, K. R. Kim, J.-B. Park, D.-K. Oh, *J. Biotechnol.* **2015**, *208*, 1–10.
- [2] a) C. T. Hou, *Handbook of industrial biocatalysis*, CRC, Boca Raton, **2005**; b) U. Schörken, P. Kempers, *Eur. J. Lipid Sci. Technol.* **2009**, *111*, 627–645.
- [3] a) D. S. Lee, J. W. Song, M. Voß, E. Schuiten, R. K. Akula, Y. U. Kwon, U. Bornscheuer, J. B. Park, *Adv. Synth. Catal.* **2019**, *361*, 1359–1367; b) J.-W. Song, J.-H. Seo, D.-K. Oh, U. T. Bornscheuer, J.-B. Park, *Catal. Sci. Technol.* **2020**, *10*, 46–64.
- [4] E.-J. Seo, H.-J. Kim, M.-J. Kim, J.-S. Kim, J.-B. Park, *Chem. Commun.* **2019**, *55*, 14462–14465.
- [5] a) H. J. Cha, E. J. Seo, J. W. Song, H. J. Jo, A. R. Kumar, J. B. Park, *Adv. Synth. Catal.* **2018**, *360*, 696–703; b) J. W. Song, E. Y. Jeon, D. H. Song, H. Y. Jang, U. T. Bornscheuer, D. K. Oh, J. B. Park, *Angew. Chem. Int. Ed.* **2013**, *52*, 2534–2537; *Angew. Chem.* **2013**, *125*, 2594–2597; c) E.-J. Seo, Y. J. Yeon, J.-H. Seo, J.-H. Lee, J. P. Boñgol, Y. Oh, J. M. Park, S.-M. Lim, C.-G. Lee, J.-B. Park, *Bioresour. Technol.* **2018**, *251*, 288–294; d) J. W. Song, J. H. Lee, U. T. Bornscheuer, J. B. Park, *Adv. Synth. Catal.* **2014**, *356*, 1782–1788.
- [6] S. Sung, H. Jeon, S. Sarak, M. M. Ahsan, M. D. Patil, W. Kroutil, B.-G. Kim, H. Yun, *Green Chem.* **2018**, *20*, 4591–4595.
- [7] M. Ahsan, S. Sung, H. Jeon, M. Patil, T. Chung, H. Yun, *Catalysts* **2017**, *8*, 4.
- [8] G. J. S. Dawes, E. L. Scott, J. Le Notre, J. P. Sanders, J. H. Bitter, *Green Chem.* **2015**, *17*, 3231–3250.
- [9] a) D. Sorigué, B. Légeret, S. Cuiné, S. Blangy, S. Moulin, E. Billon, P. Richaud, S. Brugière, Y. Couté, D. Nurizzo, P. Müller, K. Brettel, D. Pignol, P. Arnoux, Y. Li-Beisson, G. Peltier, F. Beisson, *Science* **2017**, *357*, 903–907; b) D. Sorigué, B. Légeret, S. Cuiné, P. Morales, B. Mirabella, G. Guédeney, Y. Li-Beisson, R. Jetter, G. Peltier, F. Beisson, *Plant Physiol.* **2016**, *171*, 2393–2405.
- [10] a) W. Zhang, M. Ma, M. M. Huijbers, G. A. Filonenko, E. A. Pidko, M. van Schie, S. de Boer, B. O. Burek, J. Z. Bloh, W. J. van Berkel, *J. Am. Chem. Soc.* **2019**, *141*, 3116–3120; b) M. M. Huijbers, W. Zhang, F. Tonin, F. Hollmann, *Angew. Chem. Int. Ed.* **2018**, *57*, 13648–13651; *Angew. Chem.* **2018**, *130*, 13836–13839; c) Y. Ma, X. Zhang, W. Zhang, P. Li, Y. Li, F. Hollmann, Y. Wang, *ChemPhotoChem* **2020**, <https://doi.org/10.1002/cptc.201900205>.
- [11] H.-Y. Jang, E.-Y. Jeon, A.-H. Baek, S.-M. Lee, J.-B. Park, *Proc. Biochem.* **2014**, *49*, 617–622.
- [12] E.-Y. Jeon, J.-W. Song, H.-J. Cha, S.-M. Lee, J. Lee, J.-B. Park, *J. Biotechnol.* **2018**, *281*, 161–167.
- [13] a) J.-H. Seo, H.-H. Kim, E.-Y. Jeon, Y.-H. Song, C.-S. Shin, J.-B. Park, *Sci. Rep.* **2016**, *6*, 28223; b) J.-M. Woo, E.-Y. Jeon, E.-J. Seo, J.-H. Seo, D.-Y. Lee, Y. J. Yeon, J.-B. Park, *Sci. Rep.* **2018**, *8*, 10280.

Manuscript received: November 26, 2019
Accepted manuscript online: January 19, 2020
Version of record online: February 25, 2020