

**Droplet motion with contact-line friction
long-time asymptotics in complete wetting**

Giacomelli, Lorenzo; Gnann, Manuel V.; Peschka, Dirk

DOI

[10.1098/rspa.2023.0090](https://doi.org/10.1098/rspa.2023.0090)

Publication date

2023

Document Version

Final published version

Published in

Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences

Citation (APA)

Giacomelli, L., Gnann, M. V., & Peschka, D. (2023). Droplet motion with contact-line friction: long-time asymptotics in complete wetting. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 479(2274), Article 20230090. <https://doi.org/10.1098/rspa.2023.0090>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Green Open Access added to TU Delft Institutional Repository

'You share, we take care!' - Taverne project

<https://www.openaccess.nl/en/you-share-we-take-care>

Otherwise as indicated in the copyright section: the publisher is the copyright holder of this work and the author uses the Dutch legislation to make this work public.

DROPLET MOTION WITH CONTACT-LINE FRICTION: LONG-TIME ASYMPTOTICS IN COMPLETE WETTING

LORENZO GIACOMELLI, MANUEL V. GNANN, AND DIRK PESCHKA

ABSTRACT. We consider the thin-film equation for a class of free boundary conditions modelling friction at the contact line, as introduced by E and Ren. Our analysis focuses on formal long-time asymptotics of solutions in the perfect wetting regime. In particular, through the analysis of quasi-self-similar solutions, we characterize the profile and the spreading rate of solutions depending on the strength of friction at the contact line, as well as their (global or local) corrections, which are due to the dynamical nature of the free boundary conditions. These results are complemented with full transient numerical solutions of the free boundary problem.

1. INTRODUCTION

1.1. Thin-film equations. Thin-film equations are a class of fourth-order degenerate-parabolic equations whose prototype, in one space dimension, is

$$\partial_t h + \partial_y(m(h)\partial_y^3 h) = 0 \quad \text{on } \{h > 0\} := \{(t, y) \in (0, \infty) \times \mathbb{R} : h(t, y) > 0\}, \quad (1.1)$$

where m is a mobility which degenerates at $h = 0$.

Thin-film equations are formally derived as leading-order approximations of the Navier-Stokes equations in a suitable regime, which is known as the *lubrication approximation* [1–3]. In this case, h represents the height of a thin layer or droplet of a Newtonian fluid on a flat solid substrate and the mobility m often has the form $m(h) = h^3 + b^{3-n}h^n$, the parameters b and n being related to a slip condition imposed at the liquid-solid interface: in particular, $n = 1$ corresponds to Greenspan’s slip condition [4], $n = 2$ corresponds to (linear) Navier slip and $n = 3$ (or $b = 0$) corresponds to no slip at the substrate. The case $m(h) = h$ may also be seen as the lubrication approximation of the two-dimensional Hele-Shaw flow in the half-space [5], and in this case the lubrication approximation has been given rigorous justifications [5–9]. Nonlinear free boundary problems with $m(h) = 1$ are discussed in the context of surface diffusion [10]. General traveling wave solutions for $m(h) = h^n$ with $0 \leq n < 3$ and beyond are discussed in [11–13].

We are interested in situations in which an interface exists which separates a dry region of the substrate from a wetted one:

$$\{h(t, \cdot) > 0\} = (s_-(t), s_+(t)). \quad (1.2)$$

In this case, (1.1) is complemented by two obvious boundary conditions at $s_{\pm}(t)$, respectively: the defining condition

$$h(t, s_{\pm}(t)) = 0, \quad (1.3)$$

and the kinematic condition

$$(h^{-1}m(h)\partial_y^3 h)|_{y=s_{\pm}(t)} = \dot{s}_{\pm}(t), \quad (1.4)$$

where here and after we agree that

$$f|_{y=s_+} = \lim_{\substack{y \rightarrow s_+ \\ y < s_+}} f(y), \quad f|_{y=s_-} = \lim_{\substack{y \rightarrow s_- \\ y > s_-}} f(y).$$

We are interested in situations in which the support $\{h(t, \cdot) > 0\}$ is allowed to evolve in time, leading to a genuine free boundary problem for which a third condition is required. Appropriate choices of such a third boundary condition are being debated since decades by now. Starting with the work of Bernis and Friedman [14], most of the analytical theory concentrated on the condition

LG acknowledges discussions with Maria Chiricotto. MVG appreciates discussions with Jochen Denzler, Robert McCann, and Christian Seis regarding self-similar asymptotics preceding the preparation of this work. MVG was partially supported by the Deutsche Forschungsgemeinschaft (DFG) under project # 334362478. DP thanks Luca Heltai and Marita Thomas for fruitful discussions and acknowledges the financial support within the DFG-Priority Programme 2171 by project # 422792530.

of constantly zero contact angle, focusing on existence of weak solutions [15–19], on their qualitative properties [20–32], and on well-posedness in weighted spaces [33–42]. For the constant, non-zero contact angle case we refer to [43–51]. For quasistatic models of droplet evolution we refer to [52] for the Stokes flow and to [4, 53, 54] for thin-film models.

We focus on a class of contact-line conditions, first considered in [4, 55] in special cases, that relate the contact-line velocity, \dot{s} , to the dynamic contact angle. Analytical works for this class are limited to a few contributions [56–58]. The class has been motivated and generalized by E and Ren building up on a simple, basic principle: consistency with the second law of thermodynamics in the isothermal case [59–61]. We now introduce this class directly at the level of lubrication theory.

1.2. Contact-line frictional laws in complete wetting. At leading order in lubrication approximation, and after a normalization, the surface energy reads

$$\mathcal{E}[h(t)] = \int_{s_-(t)}^{s_+(t)} \left(\frac{1}{2} (\partial_y h)^2 - S \right) dy, \quad (1.5)$$

where $S \in \mathbb{R}$ is (proportional to) the so-called spreading coefficient [6]. We are interested in a regime where the spreading coefficient of the solid/liquid/vapor system vanishes, i.e., the complete wetting regime $S = 0$. This is a generic situation in the so-called “moist” case, which concerns for instance a surface which has been pre-exposed to vapor [62]: thus

$$\mathcal{E}[h(t)] = \frac{1}{2} \int_{s_-(t)}^{s_+(t)} (\partial_y h)^2 dy, \quad (1.6)$$

in what follows (for the case $S \neq 0$, see e.g. the discussions in [45, 63–65]). After integrations by parts (see [58, formula (1.13)]), (1.1)–(1.4) formally yield

$$\begin{aligned} \frac{d}{dt} \mathcal{E}[h(t)] &= \dot{s}_+(t) \frac{1}{2} (\partial_y h)^2|_{y=s_+} - \dot{s}_-(t) \frac{1}{2} (\partial_y h)^2|_{y=s_-} + \int_{s_-(t)}^{s_+(t)} \partial_y h (\partial_t \partial_y h) dy \\ &= -\dot{s}_+(t) \left(\frac{1}{2} (\partial_y h)^2 - h \partial_y^2 h \right)|_{y=s_+} + \dot{s}_-(t) \left(\frac{1}{2} (\partial_y h)^2 - h \partial_y^2 h \right)|_{y=s_-} \\ &\quad - \int_{s_-(t)}^{s_+(t)} m(h) (\partial_y^3 h)^2 dy. \end{aligned} \quad (1.7)$$

If by contradiction $h(\partial_y^2 h)|_y = s_{\pm}(t) =: k_{\pm} \neq 0$ then one would have

$$\frac{1}{2} \partial_y ((\partial_y h)^2) = (\partial_y h) (\partial_y^2 h) \sim k_{\pm} h^{-1} \partial_y h = k_{\pm} \partial_y \log h \quad \text{as } y \rightarrow s_{\pm}(t)^{\mp},$$

whence $(\partial_y h)^2$ would become unbounded as $y \rightarrow s_{\pm}(t)^{\mp}$. Hence, $h(\partial_y^2 h)|_y = s_{\pm}(t) = 0$ for solutions with finite slope at the contact line, which is the class we are interested in. Therefore, (1.7) reads as

$$\frac{d}{dt} \mathcal{E}[h(t)] = -\frac{1}{2} \dot{s}_+(t) (\partial_y h)^2|_{y=s_+(t)} + \frac{1}{2} \dot{s}_-(t) (\partial_y h)^2|_{y=s_-(t)} - \int_{s_-(t)}^{s_+(t)} m(h) (\partial_y^3 h)^2 dy. \quad (1.8)$$

Consistency with the second law of thermodynamics implies that the first two terms on the right-hand side of (1.8) must be non-positive. A simple form of constitutive relations which enforces non-positivity is

$$(\partial_y h)^2|_x = s_{\pm}(t) = f_{\ell}(\pm \dot{s}_{\pm}) \quad \text{with } f_{\ell}(\sigma) \sigma \geq 0 \quad \text{for all } \sigma \in \mathbb{R}. \quad (1.9)$$

According to (1.9), receding fronts with speed $\sigma < 0$ only exist if $f_{\ell}(\sigma) = 0$. Hence, in complete wetting, receding fronts (if any) have zero contact angle. Furthermore, note that the more standard zero contact-angle condition $(\partial_y h)(t, s_{\pm}(t)) = 0$ corresponds to the limit of vanishing contact-line dissipation, $f_{\ell} \equiv 0$. For the analogue of (1.9) in partial wetting ($S < 0$) we refer to [58, 66].

The simple argument above can in fact be embedded into a more general formal gradient-flow structure of the system, based on a separation of dual forces into a bulk and a contact-line dissipation. This structure, which we elaborate in Appendix A, is also at the basis of the discretization that we adopt in numerical simulations.

1.3. Goals. Intermediate asymptotics for (1.1) with the Ren-E boundary condition (1.9), such as the Voinov-Cox-Hocking law (see the discussions in [13, 67–70]), has been formally worked out in [57]. Instead, here we will focus on the long-time dynamics. In complete wetting, it is expected that generic solutions spread indefinitely, covering the whole real line with a layer of zero-thickness in the limit as $t \rightarrow +\infty$. Therefore, the long-time dynamics can be equivalently captured by considering a power-law form $m(h) = h^n$ of the mobility, instead of its full form $m(h) = h^3 + b^{3-n}h^n$. Then (1.1) reads

$$\partial_t h + \partial_y (h^n \partial_y^3 h) = 0 \quad \text{on } \{h > 0\}, \quad n \in [1, 3), \quad (1.10)$$

and we are led to consider the following free boundary problem:

$$\partial_t h + \partial_y (h^n \partial_y^3 h) = 0 \quad \text{for } y \in (s_-(t), s_+(t)), \quad (1.11a)$$

$$h = 0 \quad \text{at } y = s_{\pm}(t), \quad (1.11b)$$

$$(\partial_y h)^2 = f_{\ell}(\pm \dot{s}_{\pm}(t)) \quad \text{at } y = s_{\pm}(t), \quad (1.11c)$$

$$h^{n-1}(\partial_y^3 h) = \dot{s}_{\pm}(t) \quad \text{at } y = s_{\pm}(t). \quad (1.11d)$$

In the absence of contact-line friction ($f_{\ell} \equiv 0$), (1.11) admits self-similar solutions [71], which are expected to describe the long-time dynamics of generic solutions (however, rigorous results are available for $n = 1$ only [36, 39, 72–75]). If $f_{\ell} \neq 0$, (1.11c) breaks the self-similar structure of (1.11a). Nevertheless, long-time dynamics may be inferred from the analysis of (quasi-)self-similar solutions, where the non-self-similar part of the operator is seen as a small modulation in time. This method has already been applied to thin-film equations for related asymptotic studies [67, 69].

1.4. The model problem. We assume prototypical power-law forms of f_{ℓ} , i.e.,

$$f_{\ell}(\sigma) = d g_{\alpha}(\sigma), \quad \text{where } g_{\alpha}(\sigma) = \max\{0, |\sigma|^{\alpha-1}\sigma\} \quad \text{or } g_{\alpha}(\sigma) = |\sigma|^{\alpha-1}\sigma, \quad (1.12)$$

and $\alpha > 0$ and $d > 0$ are constants encoding the strength of friction at the contact line. For $S = 0$, the former g_{α} allows for receding fronts, while the latter alternative does not.

Equation (1.11a) and the kinematic condition (1.11d) have two scaling invariances,

$$(t, y, h) \mapsto (T_* \hat{t}, Y_* \hat{y}, H_* \hat{h}), \quad \text{where } T_* = Y_*^4 H_*^{-n}. \quad (1.13)$$

We use one invariance to normalize the droplet's mass to be 2, i.e.

$$M := \int_{s_-(t)}^{s_+(t)} h(t, y) dy = 2H_* Y_*. \quad (1.14)$$

With the choice in (1.12), (1.11c) reads as

$$(\partial_{\hat{y}} \hat{h}|_{\hat{y}=\hat{s}_{\pm}(\hat{t})})^2 = D^2 g_{\alpha}(\pm \frac{d\hat{s}_{\pm}}{d\hat{t}}(\hat{t})) \quad \text{where } D^2 := d \frac{Y_*^2 Y_*^{\alpha}}{H_*^2 T_*^{\alpha}}.$$

If $\alpha \neq \frac{4}{n+3}$, we may use the second scaling invariance to fix the constant $D > 0$ to be 1:

$$1 = D^2 = \frac{dY_*^{2+\alpha}}{H_*^2 T_*^{\alpha}} \stackrel{(1.13)}{=} \frac{dY_*^{2-3\alpha}}{H_*^{2-\alpha n}} \stackrel{(1.14)}{=} \frac{d(\frac{M}{2})^{2-3\alpha}}{H_*^{4-\alpha(n+3)}} \Leftrightarrow H_*^{4-\alpha(n+3)} = d(\frac{M}{2})^{2-3\alpha}. \quad (1.15)$$

Therefore, removing hats, we will consider the following free boundary problem with initial mass $M = 2$ and with $D = 1$ if $\alpha \neq \frac{4}{n+3}$:

$$\partial_t h + \partial_y (h^n \partial_y^3 h) = 0 \quad \text{for } y \in (s_-(t), s_+(t)), \quad (1.16a)$$

$$h = 0 \quad \text{at } y = s_{\pm}(t), \quad (1.16b)$$

$$(\frac{1}{D} \partial_y h)^2 = g_{\alpha}(\pm \dot{s}_{\pm}(t)) \quad \text{at } y = s_{\pm}(t), \quad (1.16c)$$

$$h^{n-1} \partial_y^3 h = \dot{s}_{\pm}(t) \quad \text{at } y = s_{\pm}(t), \quad (1.16d)$$

$$\int_{s_-(t)}^{s_+(t)} h(t, y) dy = 2. \quad (1.16e)$$

We seek even solutions, with $s = s(t) = s_+(t) = -s_-(t)$ denoting the position of the right free boundary. Furthermore, since we are interested in the long-time dynamics, we seek solutions with

advancing contact lines $\dot{s} > 0$, whence (1.12) reduces to $g_\alpha(\dot{s}) = (\dot{s})^\alpha$. It is convenient to pass to a fixed domain by the change of variables

$$h(t, y) = s^{-1}H(t, x), \quad x = s^{-1}y. \quad (1.17)$$

Then, taking symmetry into account, (1.16) reads as

$$s^{n+4}\partial_t H - s^{n+3}\dot{s}\partial_x(xH) + \partial_x(H^n\partial_x^3 H) = 0 \quad \text{for } x \in (0, 1), \quad (1.18a)$$

$$H = 0 \quad \text{at } x = 1, \quad (1.18b)$$

$$s^{-4}(D^{-1}\partial_x H)^2 = (\dot{s})^\alpha \quad \text{at } x = 1, \quad (1.18c)$$

$$s^{-(n+3)}H^{n-1}\partial_x^3 H = \dot{s} \quad \text{at } x = 1, \quad (1.18d)$$

$$\partial_x H = \partial_x^3 H = 0 \quad \text{at } x = 0, \quad (1.18e)$$

$$\int_0^1 H(t, x)dx = 1, \quad (1.18f)$$

which we will consider in what follows. Note that the combination of (1.18c) and (1.18d) yields

$$(D^{-1}\partial_x H)^2 = s^{4-\alpha(n+3)}(H^{n-1}\partial_x^3 H)^\alpha \quad \text{at } x = 1. \quad (1.19)$$

A generic symmetric solution of problem (1.16) is shown in Fig. 1, where we choose initial conditions highlighting transient behavior and convergence to self-similar solutions for $h(t, y)$.

1.5. Outline. It is apparent from (1.19) that (1.18) has a self-similar structure if and only if

$$\alpha = \frac{4}{n+3}. \quad (1.20)$$

In §2 we discuss the case (1.20), identifying for any $D \geq 0$ a unique self-similar profile (cf. Theorem 2.2); we also discuss its behavior with respect to D and its stability. In §3 we investigate quasi-self-similar solutions. Our analysis shows that the long-time dynamics is dominated by:

- contact-line dissipation if $\alpha < \frac{4}{n+3}$ (strong contact-line friction, §3.2): in particular, the long-time scaling law depends on α ;
- bulk dissipation if $\alpha > \frac{4}{n+3}$ (weak contact-line friction, §3.3), leading to the same long-time scaling law as in the case of null contact-line dissipation.

For (1.20), both dissipations contribute equally and we speak of the balanced case. A summary of the results obtained is given in §4, which also contains their discussion and indicates further directions. Finally, in Appendix A we detail the gradient-flow formulation of the problem and we discuss how this formulation drives our numerical scheme.

2. SELF-SIMILAR SOLUTIONS

We seek symmetric and mass-preserving self-similar solutions of (1.16) in the case when the balance condition (1.20) holds. In terms of H , this translates into the ansatz

$$H(t, x) = H(x) \quad \text{and} \quad s(t) = (\gamma^{-1}B^2t)^\gamma, \quad \gamma = \frac{1}{n+4}, \quad (2.1)$$




FIGURE 1. Solution $h(t, y)$ of the transient problem (1.16) with $n = \alpha = 1$ and $D = 1$ with (symmetric) initial data $h(t = 0, y) = \frac{15}{4}((1+y)(1-y) - \frac{2}{5}(1 + \cos(\pi y)))$ for $s_\pm(t = 0) = \pm 1$.

where $B > 0$ is an unknown constant. Inserting (2.1) into (1.18), integrating once using (1.18e) and recalling (1.19), we find

$$H^{n-1} \frac{d^3 H}{dx^3} = B^2 x \quad \text{in } (0, 1), \quad (2.2a)$$

$$\frac{dH}{dx} = 0 \quad \text{at } x = 0, \quad (2.2b)$$

$$H = 0 \quad \text{at } x = 1, \quad (2.2c)$$

$$\left(\frac{dH}{dx}\right)^2 = D^2 (H^{n-1} \frac{d^3 H}{dx^3})^\alpha \stackrel{(2.2a)}{=} D^2 B^{2\alpha} \quad \text{at } x = 1, \quad (2.2d)$$

$$\int_0^1 H(x) dx = 1, \quad (2.2e)$$

where we recall that in this case D cannot be set to 1 by scaling. For $n = 1$, we have $\alpha \stackrel{(1.20)}{=} 1$ and (2.2) can be integrated explicitly to the unique solution

$$H_D(x) = \frac{B_D D}{2} (1 - x^2) + \frac{B_D^2}{24} (1 - x^2)^2, \quad B = B_D := \frac{15}{2} \left(-D + \sqrt{\frac{4}{5} + D^2} \right). \quad (2.3)$$

In particular,

$$B_D^2 \begin{cases} = 45 & \text{for } D = 0, \\ \sim 9/D^2 & \text{as } D \rightarrow \infty, \end{cases} \quad \text{hence } H_D(x) \begin{cases} = \frac{15}{8} (1 - x^2)^2 & \text{for } D = 0 \\ = \frac{3}{2} (1 - x^2) & \text{as } D \rightarrow \infty. \end{cases} \quad (2.4)$$

Remark 2.1. We remark that H_D for $D = 0$ coincides with the *Smyth-Hill solution* [76]. Further note that the profile (2.3) was formulated already in [72, Eq. (6.1)] and later on in [37], without a justification of the contact-angle dynamics, that is, instead of (2.2d) the condition $\frac{dH}{dx} = \text{constant}$ at $x = 1$ was assumed (cf. [37, (2.3)]).

For $n > 1$, the solution of (2.2) is not explicit. However, the situation is analogous:

Theorem 2.2. *Let $n \in [1, 3]$. For any $D \geq 0$ there exists a unique solution (B_D, H_D) to (2.2). In addition, $B_{D=0} > 0$, $(B_{D=0}, H_{D=0})$ is the unique solution to (2.2) with $D = 0$, and*

$$DB_D^\alpha \rightarrow -3 \quad \text{and} \quad H_D(x) \rightarrow \frac{3}{2} (1 - x^2) \quad \text{in } C([0, 1]) \quad \text{as } D \rightarrow +\infty.$$

Remark 2.3. Theorem 2.2 shows that for large, respectively small, contact-line frictional coefficients the evolution is controlled by the contact-line frictional law, respectively the complete wetting regime. In addition, since $s(t) \sim (\gamma^{-1} B_D^2 t)^\gamma$ and $B_D \rightarrow 0$ as $D \rightarrow +\infty$, as $D \rightarrow +\infty$ solutions approach a quasi-stationary interface shape with respect to the time-scale t^γ . The behavior of H_D for varying D is shown in Fig. 2.




FIGURE 2. Exact self-similar solutions $H_D(x) \equiv H(x)$ of the ODE system (2.2) and first derivative $\frac{dH_D}{dx}$ for $n = 1$ (first two panels) and $n = 2$ (last two panels) with $\alpha = 4/(n+3)$ shown for various friction coefficients $0 \leq D \leq \infty$ ($D = 0$ with $\frac{dH_D}{dx} = 0$ at $x = \pm 1$; $D \rightarrow +\infty$ with linear $\frac{dH_D}{dx}$). Numerical solutions are obtained by shooting method.

We conclude the section with the proof of Theorem 2.2.

Proof of Theorem 2.2. We rescale (2.2) as follows:

$$\hat{x} = \hat{x}_B x \quad \text{and} \quad \hat{H} = \hat{x}_B^{-1} H, \quad \hat{x}_B := B \frac{2}{n+4}, \quad (2.5)$$

so that (B, H) solves (2.2) if and only if (B, \hat{H}) solves

$$\hat{H}^{n-1} \frac{d^3 \hat{H}}{d\hat{x}^3} = \hat{x} \quad \text{for } \hat{x} \in (0, \hat{x}_B), \quad (2.6a)$$

$$\frac{d\hat{H}}{d\hat{x}} = 0 \quad \text{at } \hat{x} = 0, \quad (2.6b)$$

$$\left(\hat{H}, \frac{d\hat{H}}{d\hat{x}}\right) = (0, -DB^{\alpha - \frac{4}{n+4}}) \stackrel{(1.20)}{=} (0, -DB^{\frac{4}{(n+3)(n+4)}}) \quad \text{at } \hat{x} = \hat{x}_B, \quad (2.6c)$$

$$\int_0^{\hat{x}_B} u \, d\hat{x} = 1. \quad (2.6d)$$

In [67, Theorem 2.1], and in [71, Theorem 1.2] for $D = 0$ (up to a rescaling), it is shown that the system

$$u^{n-1} \frac{d^3 u}{dy^3} = y \quad \text{for } y \in (0, y_\theta), \quad (2.7a)$$

$$\frac{du}{dy} = 0 \quad \text{at } y = 0, \quad (2.7b)$$

$$\left(u, \frac{du}{dy}\right) = (0, -\theta) \quad \text{at } y = y_\theta, \quad (2.7c)$$

$$\int_0^{y_\theta} u \, dy = 1. \quad (2.7d)$$

has for any $\theta \geq 0$ a unique solution $(y, u) = (y_\theta, u_\theta)$ (see [37, Theorem 3.2 and Theorem 3.3] for an alternative proof of existence and uniqueness). Furthermore, y_θ is a decreasing function of θ with $0 < y_0 < \infty$ and $y_\infty = 0$. Note that the proof in [67] applies to inhomogeneous mobilities, but carries over to our situation without any change of the reasoning.

If $D = 0$, (2.6) coincides with (2.7) with $\theta = 0$, hence Theorem 2.2 holds with $(\hat{x}_B, \hat{H}) = (y_0, u)$. If $D > 0$, for any $B > 0$ let

$$\theta_B := DB^{\frac{4}{(n+3)(n+4)}}$$

and let (y_B, u_B) be the unique solution to (2.7) with $\theta = \theta_B$. Since $B \mapsto \theta_B$ is increasing, in view of the above, $B \mapsto y_B$ is decreasing from y_0 to $y_\infty = 0$: hence there exists a unique B (whence a unique θ) such that $y_B = \hat{x}_B = B^{\frac{2}{n+4}}$. This proves the well-posedness of (2.2).

It remains to consider the limit as $D \rightarrow +\infty$. Let θ_D , B_D , and $y_D = \hat{x}_D$ be the unique constants identified above. By construction, they are related by

$$\theta_D = DB_D^{\frac{4}{(n+3)(n+4)}} \quad \text{and} \quad y_D = B_D^{\frac{2}{n+4}}.$$

If by contradiction $\theta_D \rightarrow \bar{\theta} \in [0, +\infty)$ for a subsequence $D \rightarrow +\infty$ (not relabeled), then on one hand $B_D \rightarrow 0$, hence $y_D \rightarrow 0$; on the other hand, (2.7) would imply that $y_D \rightarrow \bar{y} \in (0, y_0]$, a contradiction. Therefore $\theta_D \rightarrow +\infty$, hence $y_D = B_D^{\frac{2}{n+4}} \rightarrow 0$ as $D \rightarrow +\infty$. Hence, in the limit $D \rightarrow +\infty$, it follows from (2.2) because of $B_D \rightarrow 0$ that $H_D(x)$ converges uniformly in $[0, 1]$ to the unique solution of

$$\frac{d^3 H}{dx^3} = 0 \quad \text{in } (0, 1), \quad H(1) = \frac{dH}{dx}(0) = 0, \quad \int_0^1 H(x) dx = 1.$$

Hence, $DB_D^\alpha \rightarrow -3$ and $H_D(x) \rightarrow \frac{3}{2}(1 - x^2)$ as $D \rightarrow \infty$. \square

3. QUASI-SELF-SIMILAR SOLUTIONS

3.1. Scaling and ansatz. As we have just seen, only the case $\alpha = \frac{4}{n+3}$ yields exact self-similar solutions to (1.18). For $\alpha \neq \frac{4}{n+3}$, we may nevertheless consider quasi-self-similar solutions, which are expected to describe the large-time asymptotics of generic ones. As mentioned in §1.3, this method has been applied to thin-film equations for related asymptotic studies [67, 69]. Quasi-self-similar solutions are characterized by ignoring the explicit dependence of H on time in (1.18), that is, the term $\partial_t H$ is dropped while the dependence on time through s is retained. We may then integrate (1.18a) using the boundary conditions (1.18b), (1.18d), and $|\dot{s}| < \infty$. Recalling also

(1.19), this yields

$$H^{n-1} \partial_x^3 H = s^{n+3} \dot{s} x \quad \text{for } x \in (0, 1), \quad (3.1a)$$

$$H = 0 \quad \text{at } x = 1, \quad (3.1b)$$

$$(\partial_x H)^2 = s^{4-\alpha(n+3)} (H^{n-1} \partial_x^3 H)^\alpha \quad \text{at } x = 1, \quad (3.1c)$$

$$\partial_x H = 0 \quad \text{at } x = 0, \quad (3.1d)$$

$$\int_0^1 H dx = 1, \quad (3.1e)$$

where in view of (1.15) we have assumed without loss of generality $D = 1$. The free boundary condition (3.1c) suggests that for $t \gg 1$, i.e. $s \gg 1$, the solution is

- dominated by contact-line friction if $\alpha < \frac{4}{n+3}$ (strong contact-line friction);
- a perturbation of the complete wetting solution if $\alpha > \frac{4}{n+3}$ (weak contact-line friction).

In the next two sections we will show that this is indeed the case. To this aim, we will use the asymptotic expansion

$$s^{n+4}(t) = s_0^{n+4} + s_1^{n+4}(1 + o(1)), \quad H(t, x) = H_0(x) + \omega H_1(x) + O(\omega^2), \quad (3.2)$$

with $s_0(t) \gg 1$ and $s_1(t) \ll s_0(t)$ for $t \gg 1$, and $\omega = \omega(s) \ll 1$ as $s \rightarrow \infty$. Under the expansion (3.2), we obviously have from (3.1b) and (3.1d) that

$$H_0(1) = 0, \quad \frac{dH_0}{dx}(0) = 0, \quad H_1(1) = 0, \quad \frac{dH_1}{dx}(0) = 0, \quad (3.3)$$

and from (3.1e) that

$$\int_0^1 H_0 dx = 1, \quad \int_0^1 H_1 dx = 0. \quad (3.4)$$

On the other hand, the leading order term in the contact-line condition (3.1c) depends on the sign of $\alpha - \frac{4}{n+3}$. This motivates distinguishing the two cases.

3.2. The case $\alpha < \frac{4}{n+3}$: strong contact-line friction. Since $\alpha < \frac{4}{n+3}$ and $s_0 \gg 1$, using (3.2), at leading order the bulk equation (3.1a) and the contact-line condition (3.1c) are

$$H_0^{n-1} \frac{d^3 H_0}{dx^3} = s_0^{n+3} \dot{s}_0 x \quad \text{for } x \in (0, 1) \quad \text{and} \quad H_0^{n-1} \frac{d^3 H_0}{dx^3} = 0 \quad \text{at } x = 1, \quad (3.5)$$

respectively. The two equations are obviously incompatible, pointing to the necessity of considering the correction ωH_1 . Therefore,

$$\frac{d^3 H_0}{dx^3} = 0 \quad \text{in } (0, 1) \quad (3.6)$$

and the leading-order terms in (3.1a) and (3.1c) are, respectively,

$$\omega \left(H_0^{n-1} \frac{d^3 H_1}{dx^3} + (n-1) H_0^{n-2} \frac{d^3 H_0}{dx^3} H_1 \right) = s_0^{n+3} \dot{s}_0 x \quad \text{for } x \in (0, 1), \quad (3.7)$$

$$\left(\frac{dH_0}{dx} \right)^2 = s_0^{4-\alpha(n+3)} \omega^\alpha \left(H_0^{n-1} \frac{d^3 H_1}{dx^3} + (n-1) H_0^{n-2} \frac{d^3 H_0}{dx^3} H_1 \right)^\alpha \quad \text{at } x = 1. \quad (3.8)$$

Combining (3.3), (3.4), and (3.6) yields n -independently the unique solution

$$H_0 = \frac{3}{2}(1-x^2) \quad \text{for } x \in [-1, 1]. \quad (3.9)$$

On the other hand, separation of variables in (3.7) yields with (3.6) and (3.9),

$$\frac{d^3 H_1}{dx^3} = x(1-x^2)^{1-n} \quad \text{in } (0, 1), \quad (3.10)$$

$$\omega = \left(\frac{2}{3} \right)^{n-1} s_0^{n+3} \dot{s}_0, \quad (3.11)$$

where, since the expansion (3.2) only depends on the product ωH_1 , we were free to choose the normalization factor of ω . Now, we use

$$\begin{aligned}
0 &\stackrel{(3.4)}{=} \int_0^1 \left(\frac{dx}{dx}\right) H_1 dx = 1 \cdot H_1(1) - 0 \cdot H_1(0) - \int_0^1 x \frac{dH_1}{dx} dx \\
&\stackrel{(3.3)}{=} -\frac{1}{2} \int_0^1 \left(\frac{d}{dx}(x^2 - 1)\right) \frac{dH_1}{dx} dx = -0 \cdot \frac{dH_1}{dx}(1) + \frac{1}{2} \frac{dH_1}{dx}(0) + \frac{1}{2} \int_0^1 (x^2 - 1) \frac{d^2 H_1}{dx^2} dx \\
&\stackrel{(3.3)}{=} \frac{1}{6} \int_0^1 \left(\frac{d}{dx}(x^3 - 3x + 2)\right) \frac{d^2 H_1}{dx^2} dx \\
&= 0 \cdot \frac{d^2 H_1}{dx^2}(1) - \frac{1}{3} \frac{d^2 H_1}{dx^2}(0) - \frac{1}{6} \int_0^1 (x^3 - 3x + 2) \frac{d^3 H_1}{dx^3} dx,
\end{aligned}$$

giving with (3.10),

$$\frac{d^2 H_1}{dx^2}(0) = -C_1, \quad C_1 := \frac{1}{2} \int_0^1 x(x^3 - 3x + 2)(1 - x^2)^{1-n} dx. \quad (3.12)$$

Thus we can integrate (3.10) as follows:

$$\begin{aligned}
\frac{d^2 H_1}{dx^2}(x) &\stackrel{(3.10),(3.12)}{=} -C_1 + \int_0^x x_1 (1 - x_1^2)^{1-n} dx_1, \\
\frac{dH_1}{dx}(x) &\stackrel{(3.3)}{=} -C_1 x + \int_0^x x_1 (x - x_1) (1 - x_1^2)^{1-n} dx_1,
\end{aligned}$$

so that with (3.3),

$$H_1(x) = \frac{C_1}{2}(1 - x^2) - \int_x^1 \int_0^{x_1} x_2 (x_1 - x_2) (1 - x_2^2)^{1-n} dx_2 dx_1 \quad \text{for } x \in [-1, 1]. \quad (3.13)$$

For $n = 1$ we have $C_1 = \frac{1}{10}$ and this solution reads

$$H_1(x) = \frac{1}{20}(1 - x^2) - \frac{1}{24}(1 - x^4) = \frac{1}{120}(1 - x^2)(1 - 5x^2) \quad \text{for } x \in [-1, 1]. \quad (3.14)$$

For $n = 2$ we get $C_1 = \frac{1}{6}(5 - \log 2)$ and the correction

$$H_1(x) = \frac{C_1}{2}(1 - x^2) - \frac{1}{4}(3 - 3x^2 - 2 \log 4 + (x - 1)^2 \log(1 - x) + (x + 1)^2 \log(1 + x)). \quad (3.15)$$

Next, we compute s_0 and ω . With help of (3.9) and (3.10), (3.8) takes the form

$$3^2 = s_0^{4-\alpha(n+3)} \omega^\alpha \left(\frac{3}{2}\right)^{\alpha(n-1)} \iff \omega = 3^{\frac{2}{\alpha}} \left(\frac{2}{3}\right)^{n-1} s_0^{n+3-\frac{4}{\alpha}}. \quad (3.16)$$

Thus we obtain

$$\dot{s}_0 \stackrel{(3.11)}{=} \left(\frac{3}{2}\right)^{n-1} \omega s_0^{-n-3} \stackrel{(3.16)}{=} 3^{\frac{2}{\alpha}} s_0^{-\frac{4}{\alpha}},$$

so that by normalizing $s_0(0) = 0$ through a time shift,

$$s_0(t) = 3^{\frac{1-\gamma}{2}} \gamma^{-\gamma} t^\gamma, \quad \gamma = \frac{\alpha}{\alpha+4} < \frac{1}{n+4} \quad (3.17)$$

because $\alpha < \frac{4}{n+3}$, and consequently

$$\omega(t) \stackrel{(3.16)}{=} 3^{\frac{1-\gamma}{2\gamma}} \left(\frac{2}{3}\right)^{n-1} (s_0(t))^{-\frac{1-\gamma(n+4)}{\gamma}} = \gamma^{1-\gamma(n+4)} 2^{n-1} 3^{\frac{6-n-(n+4)\gamma}{2}} t^{-(1-\gamma(n+4))}. \quad (3.18)$$

Finally, the correction s_1 is obtained from the next-to-leading order terms in equation (3.1a):

$$s_1^{n+3} \dot{s}_1 = O(\omega^2) \stackrel{(3.18)}{=} O(t^{-2(1-\gamma(n+4))}) \iff s_1^{n+4} = \begin{cases} O(t^{2(n+4)\gamma-1}) & \text{if } \gamma \neq \frac{1}{2(n+4)}, \\ O(\log t) & \text{if } \gamma = \frac{1}{2(n+4)}. \end{cases} \quad (3.19)$$

Since

$$s = (s_0^{n+4} + s_1^{n+4}(1 + o(1)))^{\frac{1}{n+4}} = s_0(1 + O(s_1^{n+4} s_0^{-n-4})), \quad (3.20)$$

we have

$$s(t) \stackrel{(3.17),(3.19)}{=} 3^{\frac{1-\gamma}{2}} \gamma^{-\gamma} t^\gamma \begin{cases} (1 + O(t^{-(1-\gamma(n+4))})) & \text{if } \gamma \neq \frac{1}{2(n+4)}, \\ (1 + O(t^{-\frac{1}{2}} \log t)) & \text{if } \gamma = \frac{1}{2(n+4)}. \end{cases} \quad (3.21)$$

The combination of (3.9), (3.13), and (3.18) in (3.2) yields at leading order as $s \rightarrow \infty$ (or equivalently $t \rightarrow \infty$) the quasi-self-similar profile H according to

$$\begin{aligned} H(t, x) &= \frac{3}{2}(1 - x^2) + C_2 t^{-(1-\gamma(n+4))} H_1(x) + O(t^{-2(1-\gamma(n+4))}) \quad \text{for } x \in [-1, 1], \\ C_2 &= \gamma^{1-\gamma(n+4)} 2^{n-1} 3^{\frac{6-n-(n+4)\gamma}{2}} \end{aligned} \quad (3.22)$$

where $\gamma = \frac{\alpha}{\alpha+4}$ and H_1 is defined in (3.12)-(3.13). For $n = 1$ this equation reduces to

$$H(t, x) = \frac{3}{2}(1 - x^2) + \frac{3^{\frac{5(1-\gamma)}{2}} \gamma^{1-5\gamma}}{120} t^{-(1-5\gamma)} (1 - x^2)(1 - 5x^2) + O(t^{-2(1-5\gamma)}) \quad (3.23)$$

for $x \in [-1, 1]$.

3.3. The case $\alpha > \frac{4}{n+3}$: weak contact-line friction. On assuming $\alpha > \frac{4}{n+3}$, separation of variables in (3.1a) yields, at leading order for $s_0 \gg 1$,

$$\dot{s}_0 = B_0^2 s_0^{-(n+3)} \quad (3.24)$$

and

$$H_0^{n-1} \frac{d^3 H_0}{dx^3} = B_0^2 x \quad \text{in } (0, 1) \quad (3.25)$$

for some unknown constant $B_0 > 0$, whilst the condition (3.1c) at the contact line yields

$$\frac{dH_0}{dx} = 0 \quad \text{at } x = 1. \quad (3.26)$$

The system for H_0 is complemented with the boundary conditions (3.3) and the mass constraint (3.4), hence it coincides with (2.2) with $D = 0$. Therefore

$$(B_0, H_0) \text{ is the unique solution to (2.2) with } D = 0, \quad (3.27)$$

(cf. Theorem 2.2 and [71, Theorem 1.2]), i.e. $H_0 = H_{D=0}$ is the unique exact self-similar solution of (1.10) with zero contact angle and unit mass. Integrating (3.24) with normalization $s_0(0) = 0$ then yields

$$s_0(t) = ((n+4)B_0^2 t)^{\frac{1}{n+4}}. \quad (3.28)$$

Remark 3.1. It is to be noted that the leading-order expansion given by s_0 and H_0 is self-consistent. This is in contrast to the case of strong contact-line friction, where the leading-order expansions in the bulk and at the contact line were incompatible with each other (cf. (3.5)). However, since we are interested in quantifying the correction coming from the contact-line frictional law, we shall be looking at ωH_1 and s_1 in this case, too.

At next-to-leading order for $s_0 \gg 1$, the contact-line condition (3.1c) implies

$$\frac{dH_1}{dx} = -\left(H_0^{n-1} \frac{d^3 H_0}{dx^3}\right)^{\frac{\alpha}{2}} \stackrel{(3.25)}{=} -B_0^{-\alpha} \quad \text{at } x = 1, \quad (3.29)$$

$$\omega = s_0^{-\frac{\alpha(n+3)-4}{2}} \stackrel{(3.28)}{=} ((n+4)B_0^2 t)^{-\frac{\alpha(n+3)-4}{2(n+4)}}, \quad (3.30)$$

where we have normalized ω conveniently since only the product ωH_1 enters the expansion in (3.2). Furthermore, we have used that $\omega \frac{dH_1}{dx}$ has to have (strictly) negative sign at $x = 1$ for the expansion to make sense around $x = 1$ and be nontrivial. In the bulk, the next-to-leading order terms in (3.1a),

$$\omega \left((n-1)H_0^{n-2} \frac{d^3 H_0}{dx^3} H_1 + H_0^{n-1} \frac{d^3 H_1}{dx^3} \right) = s_1^{n+3} \dot{s}_1 x \quad \text{for } x \in (0, 1),$$

yield by separation of variables with (3.25)

$$\omega = -C_2^{-1} B_0^{\frac{2(1-n)}{n}} s_1^{n+3} \dot{s}_1 \quad (3.31)$$

and

$$(n-1)xH_1 + f^n \frac{d^3 H_1}{dx^3} = -C_2 x f \quad \text{in } (0, 1) \quad (3.32)$$

for some $C_2 > 0$ without loss of generality (since the expansion only depends on ωH_1), where we have defined

$$f = B_0^{-\frac{2}{n}} H_0. \quad (3.33)$$

Integrating (3.31) using (3.30) we obtain

$$\frac{ds_1^{n+4}}{dt} = O(\omega) = O\left(t^{\frac{4-\alpha(n+3)}{2(n+4)}}\right) \Rightarrow s_1^{n+4} = \begin{cases} O\left(t^{\frac{2(n+6)-\alpha(n+3)}{2(n+4)}}\right) & \text{if } \alpha \neq 2\frac{n+6}{n+3}, \\ O(\log t) & \text{if } \alpha = 2\frac{n+6}{n+3}, \end{cases}$$

and with (3.20) and (3.28) this produces

$$s(t) = ((n+4)B_0^2 t)^{\frac{1}{n+4}} \begin{cases} (1 + O(t^{-\frac{\alpha(n+3)-4}{2(n+4)}})) & \text{if } \alpha \neq 2\frac{n+6}{n+3}, \\ (1 + O(t^{-1} \log t)) & \text{if } \alpha = 2\frac{n+6}{n+3}. \end{cases} \quad (3.34)$$

Next, we turn our attention to determining H_1 and C_2 . We distinguish three cases.

$n = 1$. For $n = 1$, an explicit integration gives (cf. (2.3), (2.4), and (3.33))

$$f(x) = \frac{1}{24}(1-x^2)^2, \quad B_0^2 = 45 \quad \text{and} \quad H_0 = \frac{15}{8}(1-x^2)^2 \quad \text{for } x \in [-1, 1], \quad (3.35)$$

whence $s_0(t) \stackrel{(3.28)}{=} (225t)^{\frac{1}{5}}$, and (3.30) and (3.34) reduce to

$$\omega = (225t)^{-\frac{2(\alpha-1)}{5}}, \quad s(t) = (225t)^{\frac{1}{5}} \begin{cases} (1 + O(t^{-\frac{2(\alpha-1)}{5}})) & \text{if } \alpha \neq \frac{7}{2}, \\ (1 + O(t^{-1} \log t)) & \text{if } \alpha = \frac{7}{2}. \end{cases} \quad (3.36)$$

We integrate (3.32) using the four conditions (3.3), (3.4), and (3.29), leading to $C_2 = (45)^{\frac{\alpha}{2}}$ and

$$H_1 = \frac{(45)^{\frac{\alpha}{2}}}{6} \left(\frac{1}{5}(1-x^2) - \frac{1}{4}(1-x^2)^2 \right) = -\frac{(45)^{\frac{\alpha}{2}}}{120} (1-x^2)(1-5x^2). \quad (3.37)$$

Collecting (3.35), (3.36) and (3.37) in (3.2), at leading order as $s \rightarrow \infty$ (or equivalently $t \rightarrow \infty$) the approximate position of the free boundary is given by (3.36), and the quasi-self-similar profile H is given by

$$H(t, x) = \frac{15}{8}(1-x^2)^2 - \frac{1}{8} 3^{\frac{\alpha-1}{5}} 5^{-\frac{3\alpha+2}{10}} t^{-\frac{2(\alpha-1)}{5}} (1-x^2)(1-5x^2) + O\left(t^{-\frac{4(\alpha-1)}{5}}\right).$$

$n \in (1, \frac{3}{2})$. We now consider $n > 1$. Defining g as

$$H_1 = -\frac{C_2}{n} f + g, \quad (3.38)$$

we get from (3.32)

$$(n-1)xg + f^n \frac{d^3 g}{dx^3} = 0 \quad \text{for } x \in (0, 1), \quad (3.39a)$$

and the boundary conditions (3.3) and (3.29) transform into

$$g = 0 \quad \text{at } x = 1, \quad (3.39b)$$

$$\frac{dg}{dx} = -B_0^\alpha \quad \text{at } x = 1, \quad (3.39c)$$

$$\frac{dg}{dx} = 0 \quad \text{at } x = 0. \quad (3.39d)$$

Once g is determined, the mass constraint (3.4) leads to

$$C_2 = -\frac{n \int_0^1 g(x) dx}{\int_0^1 f(x) dx}. \quad (3.40)$$

In order to determine solvability of (3.39), we use [71, Theorem 1.3], that is,

$$f = \begin{cases} C_3(1-x)^2(1+o(1)) & \text{for } 0 < n < \frac{3}{2}, \\ C_4(1-x)^2(-\log(1-x))^{\frac{2}{3}}(1+o(1)) & \text{for } n = \frac{3}{2}, \\ C_5(1-x)^{\frac{3}{n}}(1+o(1)) & \text{for } \frac{3}{2} < n < 3, \end{cases} \quad \text{as } x \nearrow 1, \quad (3.41)$$

where $C_3, C_4, C_5 > 0$ only depend on n . The asymptotics (3.41) imply that (3.39) has no solution for $n \in [\frac{3}{2}, 3)$: indeed, for $n \in (\frac{3}{2}, 3)$ from (3.39a)-(3.39c) and (3.41) we infer

$$\frac{d^3 g}{dx^3} = \begin{cases} -\frac{1}{2} B_0^\alpha C_4^{-\frac{3}{2}} (1-x)^{-2} (-\log(1-x))^{-1} & \text{for } n = \frac{3}{2}, \\ -(n-1) B_0^\alpha C_5^{-n} (1-x)^{-2} (1+o(1)) & \text{for } n \in (\frac{3}{2}, 3), \end{cases} \quad \text{as } x \nearrow 1,$$

in contradiction with (3.39c).

For $n \in [1, \frac{3}{2})$, we apply a further splitting according to

$$g = \frac{1}{2} B_0^\alpha (1 - x^2) + v + w, \quad (3.42)$$

where

$$f^n \frac{d^3 v}{dx^3} = \frac{1-n}{2} B_0^\alpha x (1 - x^2) \quad \text{for } x \in (0, 1), \quad (3.43a)$$

$$v = \frac{dv}{dx} = 0 \quad \text{at } x = 1, \quad (3.43b)$$

$$\frac{dv}{dx} = 0 \quad \text{at } x = 0, \quad (3.43c)$$

and

$$\mathcal{L}w := (n-1)xw + f^n \frac{d^3 w}{dx^3} = (1-n)xv \quad \text{for } x \in (0, 1), \quad (3.44a)$$

$$w = \frac{dw}{dx} = 0 \quad \text{at } x = 1, \quad (3.44b)$$

$$\frac{dw}{dx} = 0 \quad \text{at } x = 0. \quad (3.44c)$$

We first construct a solution to (3.43). We have

$$\begin{aligned} \frac{d^2 v}{dx^2} &\stackrel{(3.43a)}{=} -\frac{n-1}{2} B_0^\alpha \int_0^x \frac{x_1(1-x_1^2)}{(f(x_1))^n} dx_1 + C_6, \\ \frac{dv}{dx} &\stackrel{(3.43c)}{=} -\frac{n-1}{2} B_0^\alpha \int_0^x \int_0^{x_1} \frac{x_2(1-x_2^2)}{(f(x_2))^n} dx_2 dx_1 + C_6 x \\ &= -\frac{n-1}{2} B_0^\alpha \int_0^x \frac{x_2(x-x_2)(1-x_1^2)}{(f(x_2))^n} dx_2 + C_6 x, \\ C_6 &\stackrel{(3.43b)}{=} \frac{n-1}{2} B_0^\alpha \int_0^1 \frac{x(1-x)^2(1+x)}{(f(x))^n} dx, \end{aligned}$$

so that

$$v \stackrel{(3.43b)}{=} \frac{n-1}{2} B_0^\alpha \int_x^1 \int_0^{x_1} \frac{x_2(x-x_2)(1-x_1^2)}{(f(x_2))^n} dx_2 dx_1 - \frac{C_6}{2} (1-x^2). \quad (3.45)$$

Notably, in view of (3.41), (3.45) only yields a well-defined solution for $n \in [1, \frac{3}{2})$. Lastly, in order to find the solution of (3.44), we use

$$\begin{aligned} \int_0^1 x f^{-n} w (\mathcal{L}w) dx &= (n-1) \int_0^1 x^2 f^{-n} w^2 dx + \int_0^1 x w \frac{d^3 w}{dx^3} dx \\ &\stackrel{(3.44b)}{=} (n-1) \int_0^1 x^2 f^{-n} w^2 dx - \int_0^1 w \frac{d^2 w}{dx^2} dx - \frac{1}{2} \int_0^1 x \frac{d}{dx} \left(\frac{dw}{dx} \right)^2 dx \\ &\stackrel{(3.44b), (3.44c)}{=} (n-1) \int_0^1 x^2 f^{-n} w^2 dx + \frac{3}{2} \int_0^1 \left(\frac{dw}{dx} \right)^2 dx > 0 \quad \text{for all } w \neq 0. \end{aligned}$$

This proves coercivity for $n \in [1, \frac{3}{2})$, so that the Lax-Milgram theorem yields existence of a unique solution to (3.44). Together with (3.45), this yields existence of a unique solution g to (3.39), hence of a unique H_1 .

$n \in (\frac{3}{2}, 3)$. The afore-mentioned non-existence of solutions to (3.39) for mobility exponents $n \in [\frac{3}{2}, 3)$ entails that the ansatz (3.2) breaks down for this range of mobilities in the regime of strong contact-line friction $\alpha > \frac{4}{n+3}$. This necessitates a matched-asymptotics approach in which we distinguish between outer and inner region. Since the case $n = \frac{3}{2}$ contains a logarithmic resonance (see (3.41)), we only concentrate on the range $n \in (\frac{3}{2}, 3)$ in what follows.

With H_0 and s_0 given by (3.27) and (3.28), respectively, it follows from the previous analysis that $s = s_0(1 + o(1))$ and $H(t, x) = H_0(x)$ in the outer region (whose extent is to be determined). Near $x = 1$, from (3.33) and (3.41), we infer for $n \in (\frac{3}{2}, 3)$ that

$$H_0 = C_5 B_0^{\frac{2}{n}} (1-x)^{\frac{3}{n}} (1 + o(1)) \quad \text{for } 0 < 1-x \ll 1, \quad (3.46)$$

where more precisely

$$C_5 = \left(\frac{3}{n} \left(\frac{3}{n} - 1 \right) \left(2 - \frac{3}{n} \right) \right)^{-\frac{1}{n}}. \quad (3.47)$$

In the inner region $x \nearrow 1$, we use the traveling-wave ansatz

$$H(t, x) = sF_{\text{in}}(\xi) \quad \xi = s(1 - x). \quad (3.48)$$

Inserted into (1.18a) with $s = s_0(1 + o(1))$, this gives, after an integration using (1.18b) and (1.18d),

$$F_{\text{in}}^{n-1} \frac{d^3 F_{\text{in}}}{d\xi^3} = -\dot{s}_0 \quad \text{for } \xi > 0. \quad (3.49a)$$

The boundary condition (1.18d) is then automatically satisfied; the conditions (1.18b) and (1.18c) translate into

$$F_{\text{in}} = 0 \quad \text{and} \quad \left(\frac{dF_{\text{in}}}{d\xi}\right)^2 = (\dot{s}_0)^\alpha \quad \text{at } \xi = 0, \quad (3.49b)$$

and are complemented with the condition

$$\frac{d^2 F_{\text{in}}}{d\xi^2} \rightarrow 0 \quad \text{as } \xi \rightarrow \infty \quad \text{for } \frac{3}{2} < n < 3, \quad (3.49c)$$

which in view of (3.46) is required for the matching to the outer solution H_0 . Well-posedness of (3.49) may be obtained using the strategies in [56] or [13], where related problems were considered. Here we sketch the application of the argument in [13] to the present case. For $\xi \ll 1$, a one-parametric solution family to (3.49a) and (3.49b) is given by

$$F_{\text{in}}(\xi) = \begin{cases} \left(\dot{s}_0^{\frac{\alpha}{2}} \xi - \frac{\dot{s}_0^{1+\frac{\alpha}{2}(1-n)}}{(4-n)(3-n)(2-n)} \xi^{4-n} + a_{\text{in}} \xi^2 + \text{h.o.t.}\right) & \text{for } n \neq 2 \\ \dot{s}_0^{\frac{\alpha}{2}} \left(\xi + \frac{\dot{s}_0^{1-\alpha}}{2} \xi^2 \log \xi + a_{\text{in}} \xi^2 + \text{h.o.t.}\right) & \text{for } n = 2 \end{cases} \quad \text{as } \xi \searrow 0, \quad a_{\text{in}} \in \mathbb{R} \quad (3.50)$$

where h.o.t. denotes higher-order terms. For $\xi \gg 1$, a one-parametric (plus translation) solution family to (3.49a) and (3.49c) is given by

$$F_{\text{in}} = \dot{s}_0^{\frac{1}{n}} \left(C_5 \xi^{\frac{3}{n}} + b_{\text{in}} \xi + \text{h.o.t.}\right) \quad \text{for } \frac{3}{2} < n < 3 \quad \text{as } \xi \rightarrow \infty, \quad b_{\text{in}} \in \mathbb{R}. \quad (3.51)$$

Therefore, for $\frac{3}{2} < n < 3$, (3.50) and (3.51) define two two-dimensional solution manifolds (with parameters (ξ, a_{in}) and (ξ, b_{in}) , respectively) of the three-dimensional dynamical system $(F_{\text{in}}, \frac{dF_{\text{in}}}{d\xi}, \frac{d^2 F_{\text{in}}}{d\xi^2})$: their intersection is the one-dimensional solution curve determining a_{in} and b_{in} and defining the solution to (3.49).

In terms of x , expansion (3.51) translates at leading order into

$$H(t, x) \stackrel{(3.48)}{=} s_0 F_{\text{in}} = \dot{s}_0^{\frac{1}{n}} s_0^{\frac{n+3}{n}} C_5 (1-x)^{\frac{3}{n}} (1+o(1)) \quad \text{for } s^{-1} \ll 1-x \ll 1. \quad (3.52)$$

For $s \gg 1$, expansions (3.46) and (3.52) thus have an overlapping region, $s^{-1} \ll 1-x \ll 1$: matching them yields

$$\dot{s}_0^{\frac{1}{n}} s_0^{\frac{n+3}{n}} C_5 = B_0^{\frac{2}{n}} C_5,$$

which is fulfilled because of (3.24). Since

$$\begin{aligned} s_0 F_{\text{in}} &\stackrel{(3.50)}{=} s_0^2 \dot{s}_0^{\frac{\alpha}{2}} (1-x)(1+o(1)) \stackrel{(3.24)}{=} B_0^\alpha s_0^{\frac{4-\alpha(n+3)}{2}} (1-x)(1+o(1)) \\ &\stackrel{(3.28)}{=} B_0^\alpha ((n+4)B_0^2 t)^{\frac{4-\alpha(n+3)}{2(n+4)}} (1-x)(1+o(1)) \end{aligned} \quad \text{for } 0 \leq 1-x \ll s^{-1},$$

we conclude that $H(t, x) = \tilde{H}_0(t, x)(1+o(1))$, where

$$\tilde{H}_0(t, x) \sim \begin{cases} H_0(x) & \text{for } 1-x \gg s^{-1} \\ B_0^\alpha ((n+4)B_0^2 t)^{-\beta} (1-x) & \text{for } 1-x \ll s^{-1}, \end{cases} \quad s \gg 1, \quad \beta = \frac{\alpha(n+3) - 4}{2(n+4)} > 0. \quad (3.53)$$

4. CONCLUSIONS AND OUTLOOK

We have obtained an asymptotic description of the long-time dynamics of solutions (H, s) to (1.18). This translates back to the original height via $h(t, y) = s^{-1}H(t, s^{-1}y)$ with support $(-s, s)$.

In the balanced case, $\alpha = \frac{4}{n+3}$, we have shown that (1.18) admits for any $D \geq 0$ a unique self-similar profile, determined by (2.2), and s scales like $t^{\frac{1}{n+4}}$. Both profile and speed depend on D , the profile ranging from the zero contact-angle one for $D = 0$ to a parabolic shape as $D \rightarrow +\infty$. Numerical solutions to (2.2) have been provided in Fig. 2.

A very interesting question concerns stability, i.e., the convergence of solutions H of (1.18) to the self-similar profile (2.3). This issue has already been raised in [72, §6] and [36, §8] in




FIGURE 3. For $D = 1$ and $n = 1$ (left) or $n = 2$ (right), the solution $H(t, x)$ to (1.18) for increasing times $0 \leq t < \infty$ (blue lines, time increases along arrows) with initial datum $H(t = 0, x) = \frac{15}{4}((1+x)(1-x) - \frac{2}{5}(1 + \cos(\pi x)))$ (red dashed line), compared with the exact self-similar solutions $H_D(x)$ to (2.2) approached as $t \rightarrow \infty$ (red dotted line).

a similar context and could be faced, starting from $n = 1$, either by energy-entropy methods [72–75], by studying global-in-time classical solutions for perturbations of special solutions like a self-similar profile [35, 36, 39], a traveling wave [40, 42], or an equilibrium-stationary solution [33, 34, 46, 47, 49–51]. In the three latter cases, the difficulty lies in finding suitable estimates for the linearized evolution of perturbations. Unlike in the case of the Smyth-Hill profile [76], this linearization does not carry an apparent symmetric structure, which is why the linear analysis is presumably more involved. Fig. 3 provides numerical simulations supporting convergence in the balanced case for $n = 1$ and $n = 2$.

For strong contact-line friction, $\alpha < \frac{4}{n+3}$, H and s obey the asymptotic

$$H(t, x) = \frac{3}{2}(1 - x^2) + C_2 t^{-(1-\gamma(n+4))} H_1(x) + O(t^{-2(1-\gamma(n+4))}) \quad \text{for } x \in [-1, 1], \quad (4.1a)$$

$$s(t) = 3^{\frac{1-\gamma}{2}} \gamma^{-\gamma} t^\gamma \begin{cases} (1 + O(t^{-(1-\gamma(n+4))})) & \text{if } \gamma \neq \frac{1}{2(n+4)}, \\ (1 + O(t^{-\frac{1}{2}} \log t)) & \text{if } \gamma = \frac{1}{2(n+4)}, \end{cases} \quad (4.1b)$$

where H_1 and C_2 are defined in §3.2 (see (3.12)-(3.13) and (3.22)), and

$$H_1(x) = \frac{1}{120}(1 - x^2)(1 - 5x^2) \quad \text{if } n = 1.$$

The leading-order profile is a parabolic one, with finite non-zero contact angle, and coincides with the exact self-similar profile in the limiting case $D = \infty$ (§2). In other words, $H_D \rightarrow H_0$ as $D \rightarrow \infty$. In addition, the evolution of the contact line, as given by (4.1b), is slower than the standard one and is dominated by the contact-line frictional exponent α . Therefore, for $\alpha < \frac{4}{n+3}$ contact-line friction dominates the long-time dynamics uniformly in space and time (Fig. 4, first column). Notably, the correction H_1 , which is dictated by attaining the dynamical contact-line condition, is not localized near the contact line, but rather propagates throughout the solution's support (Fig. 5, middle and right).

For weak contact-line friction, $\alpha > \frac{4}{n+3}$, the leading-order dynamics instead coincide with the zero-contact-angle ones in terms of both profile and speed, in the sense that

$$H(t, x) = H_0(x) \quad \text{and} \quad s(t) = ((n+4)B_0^2 t)^{\frac{1}{n+4}}, \quad (4.2)$$

where (B_0, H_0) is the unique solution to (2.2) with $D = 0$. In other words, $H_0 = H_{D=0}$ is the unique self-similar profile of (1.10) with zero contact angle and normalized mass (Fig. 4, second column). Our results on the corrections turn out to depend on the mobility exponent n : if $n \in [1, 3/2)$ we find a global estimate as above, in the sense that

$$H(t, x) = H_0(x) + ((n+4)B_0^2 t)^{-\beta} H_1(x) + O(t^{-2\beta}), \quad \beta = \frac{\alpha(n+3) - 4}{2(n+4)}, \quad (4.3a)$$

$$s(t) = ((n+4)B_0^2 t)^{\frac{1}{n+4}} \begin{cases} (1 + O(t^{-\beta})) & \text{if } \alpha \neq \frac{2n+6}{n+3}, \\ (1 + O(t^{-1} \log t)) & \text{if } \alpha = \frac{2n+6}{n+3} \end{cases} \quad (4.3b)$$




FIGURE 4. For $D = 1$ and $n = 1$ (top) or $n = 2$ (bottom), the solution $H(t, x)$ to (1.18) for different t (blue lines) with initial datum $H(t = 0, x) = \frac{15}{4}((1+x)(1-x) - \frac{2}{5}(1 + \cos(\pi x)))$ (red dashed line) for strong contact line friction $\alpha = 1/2$ (left) and weak contact line friction $\alpha = 2$ (right) compared with the exact self-similar solutions $H_{\tilde{D}}(x)$ to (2.2) approached as $t \rightarrow \infty$ (red dotted lines, with $\tilde{D} \rightarrow \infty$ for $\alpha = 1/2$, cf. Theorem 2.2, and $\tilde{D} = 0$ for $\alpha = 2$). Note that for $\alpha = \frac{1}{2}$ solutions go slightly above the graph of $H_{\tilde{D}}$ before relaxing (a manifestation of the lack of a comparison principle). The time evolution is indicated by dark grey arrows.




FIGURE 5. Convergence of the correction to the self-similar solution, where (left) we show the L^1 norm of the difference between $C(t, x) = (H(t, x) - H_0(x))/|H(t, \cdot) - H_0(\cdot)|_\infty$ and the theoretically predicted, normalized correction $\pm H_1(x)/|H_1(\cdot)|_\infty$. For comparison, in (middle) and (right) we show $C(t, x)$ at $t = 200$ for $n = 1$, respectively $n = 2$. Corrections C (colored) are compared with theoretically predicted ones (dashed/dotted), when the latter are available.

where $H_1(x)$ is uniquely determined in §3.3. On the other hand, if $n \in (3/2, 3)$ we are only able to qualify a *local* correction:

$$H(t, x) \sim \begin{cases} H_0(x) & \text{for } 1 - x \gg t^{-\frac{1}{n+4}} \\ B_0^\alpha ((n+4)B_0^2 t)^{-\beta} (1-x) & \text{for } 1 - x \ll t^{-\frac{1}{n+4}}. \end{cases}$$

It is apparent from our numerical simulation (Fig. 5, right) that the correction should consist, also in this case, of a globally defined function H_1 and s_1 as in (4.3), with the same time exponent β ; however, at the moment this is left as an open question.




FIGURE 6. Convergence to self-similar solutions observed in the long-time behavior $s(t) \sim t^\gamma$ of numerical solutions to (1.18) for mobility exponents $n = 1, 2$, where (left) we show the logarithmic time-derivative $\frac{d \log s}{d \log t}$ approaching the predicted γ (black dotted line) for $n = 1$ and (right) we show the evaluation of the logarithmic time-derivative at $t = 10^7$ for various α and for $n = 1, 2$ compared to the theoretical prediction.




FIGURE 7. Time correction, in the form $\frac{d}{dt}(t^{-\gamma}s(t))$ versus t , for numerical solutions to (1.18), shown using full lines for strong contact line friction $n = 1, \alpha < 1$ (left), weak contact line friction $n = 1, \alpha > 1$ (center), and strong contact line friction $n = 2, \alpha < 4/5$ (right). Dashed lines are the corresponding theoretical predictions.

As in the balanced case, a rigorous stability result is expected to hold for $\alpha \neq \frac{4}{n+3}$ and would be interesting to be pursued, showing convergence of solutions H of (1.18) to the corresponding H_0 . In this respect, Fig. 4 and 6 provide numerical simulations supporting convergence, and Fig. 5 and 7 numerically validate the next-to-leading-order corrections to the macroscopic profile H_0 .

A final remark concerns dependence on the contact-line frictional coefficient d (cf. (1.12)), which we could scale out. Let s_* be the asymptotic position of the free boundary of the solution to (1.16) with $D = 1$ and $M = 2$, as given by (4.1b) and (4.2). Keeping mass equal to two, but for a generic $d > 0$, the asymptotic position of the contact line in the original equation (1.11) is given by

$$s(t) = Y_* s_*(T_*^{-1}t), \quad \text{where } H_* \stackrel{(1.14)}{=} Y_*^{-1} \stackrel{(1.13)}{=} T_*^{-\frac{1}{n+4}} \stackrel{(1.15)}{=} d^{\frac{1}{4-\alpha(n+3)}}, \quad \alpha \neq \frac{4}{n+3},$$

yielding

$$s(t) \sim \begin{cases} 3^{\frac{1-\gamma}{2}} \gamma^{-\gamma} d^{-\frac{1}{\alpha+4}} t^\gamma & \text{if } \alpha < \frac{4}{n+3} \\ ((n+4)B_0^2 t)^{\frac{1}{n+4}} & \text{if } \alpha > \frac{4}{n+3} \end{cases} \quad \text{as } t \rightarrow +\infty.$$

This shows that in the strong case, as expected, larger frictional coefficients d yield slower speeds; notably, however, in the weak case the leading-order asymptotic speed in (1.11) is instead oblivious of both α and d , hence universal.

APPENDIX A. THE GRADIENT-FLOW FORMULATION AND ITS DISCRETIZATION

A.1. Gradient-flow formulation. Problem (1.11) and its discretization are based on a gradient flow formulation for the height h and the wetted area $\{h > 0\} = (s_-, s_+)$ as in (1.2). The gradient

flow

$$\dot{h} = -\partial_\eta \Psi^*(h, \mathcal{DE}[h]). \quad (\text{A.1})$$

is formally defined in terms of the energy \mathcal{E} in (1.5) and the dual dissipation potential

$$\Psi^*(h, \eta) = \frac{1}{2} \int_{\{h>0\}} m(h) (\partial_y \pi)^2 dy + \frac{1}{2} \int_{\partial\{h>0\}} m_{\text{CL}}(\partial_y h) \zeta^2 ds, \quad \eta = (\pi, \zeta), \quad (\text{A.2})$$

where $\Psi^*(h, 0) \equiv 0$ and Ψ^* is convex in the second argument. The bulk mobility m and the contact-line mobility m_{CL} are non-negative functions, which in the case of (1.11) are given by

$$m(h) = h^n \quad \text{and} \quad m_{\text{CL}}(z) = \frac{2}{d^\frac{1}{\alpha}} |z|^\frac{2}{\alpha}. \quad (\text{A.3})$$

The first term in Ψ^* encodes the standard dissipation of the viscous fluid with nontrivial slip boundary conditions, whereas the second term encodes the extra dissipation at the contact line $y = s_\pm$. This formulation relies on the formal assumption that for fixed time there exists a representation of the dual force $\eta = (\pi : \{h > 0\} \rightarrow \mathbb{R}, \zeta : \partial\{h > 0\} \rightarrow \mathbb{R})$ such that

$$\langle \eta, \dot{h} \rangle = \int_{\{h>0\}} \pi \dot{h} dy + \int_{\partial\{h>0\}} \zeta \dot{h} ds \quad \text{for any rate } \dot{h}(t) : \{h > 0\} \rightarrow \mathbb{R}. \quad (\text{A.4})$$

Using this representation¹, one identifies η with $\langle \eta, \bar{v} \rangle = \langle \mathcal{DE}[h], \bar{v} \rangle$ in a weak formulation, i.e.,

$$\int_{\{h>0\}} \pi \bar{v} dy + \int_{\partial\{h>0\}} \zeta \bar{v} ds = \int_{\{h>0\}} \partial_y h \partial_y \bar{v} dy + \int_{\partial\{h>0\}} \frac{1}{2|\partial_y h|} ((\partial_y h)^2 + (-2S)) \bar{v} ds. \quad (\text{A.5a})$$

Testing the gradient flow (A.1) with $\bar{\eta} = (\bar{\pi}, \bar{\zeta})$ and using (A.2) gives $\langle \bar{\eta}, \dot{h} \rangle = \langle \bar{\eta}, -\partial_\eta \Psi^*(h, \mathcal{DE}[h]) \rangle$, or in full detail

$$\int_{\{h>0\}} \dot{h} \bar{\pi} dy + \int_{\partial\{h>0\}} \dot{h} \bar{\zeta} ds = - \int_{\{h>0\}} m(h) \partial_y \pi \partial_y \bar{\pi} dy - \int_{\partial\{h>0\}} m_{\text{CL}}(\partial_y h) \zeta \bar{\zeta} ds. \quad (\text{A.5b})$$

We seek (\dot{h}, π, ζ) that satisfy (A.5) for all test functions $(\bar{v}, \bar{\pi}, \bar{\zeta})$. By testing this weak formulation with $(\bar{v}, \bar{\pi}, \bar{\zeta}) = (\dot{h}, \pi, \zeta)$, we can deduce the energy descent

$$\frac{d}{dt} \mathcal{E}[h(t)] = \langle \eta, \dot{h} \rangle = - \left(\int_{\{h>0\}} m(h) (\partial_y \pi)^2 dy + \int_{\partial\{h>0\}} m_{\text{CL}}(\partial_y h) \zeta^2 ds \right) \leq 0. \quad (\text{A.6})$$

Additionally, the solution $h(t, y)$ and the contact line $s_\pm(t)$ satisfy the kinematic condition

$$\frac{d}{dt} h(t, s_\pm(t)) = \dot{h}(t, s_\pm(t)) + \dot{s}_\pm(t) \partial_y h(t, s_\pm(t)) = 0, \quad (\text{A.7})$$

which we can use to reconstruct the boundary velocity and evolve the domain $\{h > 0\}$ and the solution h . Using integration by parts and assuming the solution is smooth enough, from (A.5a) we identify $\pi = -\partial_y^2 h$ and $\zeta = \frac{1}{2|\partial_y h|} ((-2S) - (\partial_y h)^2)$. Using (A.5b) we can recover the general evolution of h and $\{h > 0\}$. We will now focus on the complete wetting case, $S = 0$. Then the thin-film dynamics are governed by

$$\dot{h} = \partial_y (m(h) \partial_y \pi), \quad \pi = -\partial_y^2 h \quad \text{in } \{h > 0\}, \quad (\text{A.8a})$$

$$\dot{h} = -\dot{s}_\pm \partial_y h = -m_{\text{CL}}(\partial_y h) \zeta, \quad \zeta = -\frac{1}{2} |\partial_y h| \quad \text{on } \partial\{h > 0\}. \quad (\text{A.8b})$$

with natural boundary conditions for π in (A.8a). Assuming (A.3), at the contact line $\partial\{h > 0\}$ we get for positive speeds (which are of interest to us)

$$\dot{h} = 2d^{-\frac{1}{\alpha}} |\partial_y h|^\frac{2}{\alpha} \left(\frac{1}{2} |h_y| \right), \quad \text{or equivalently} \quad \dot{s}_\pm = \pm d^{-\frac{1}{\alpha}} |\partial_y h|^\frac{2}{\alpha} \quad \text{at } \partial\{h > 0\}, \quad (\text{A.9})$$

which coincides with (1.12). The Ren-E model with quadratic dissipation has $\alpha = 1$; for a derivation from the Stokes problem see [66].

¹For dual forces η that do not admit such a representation, we formally set $\Psi^*(h, \eta) = \infty$.

A.2. Discretization. The weak formulation (A.5) is discretized using a standard finite element discretization in space. In the derivative of the energy, we replace $\partial_y h$ by $\partial_y h + \tau \partial_y \dot{h}$ in order to achieve a semi-implicit treatment of the highest-order derivative in the time-discretization, a standard method in higher-order parabolic equations. For given h , we seek (\dot{h}, π, ζ) using P_1 finite elements defined in $\{h > 0\}$ and on $\partial\{h > 0\}$, respectively, and solve

$$\int_{\{h>0\}} \pi v - \tau \partial_y \dot{h} \partial_y \bar{v} \, dy + \int_{\partial\{h>0\}} \zeta \bar{v} \, ds = \int_{\{h>0\}} \partial_y h \partial_y \bar{v} \, dy + \int_{\partial\{h>0\}} \frac{1}{2} |\partial_y h| \bar{v} \, ds, \quad (\text{A.10a})$$

$$\int_{\{h>0\}} \dot{h} \bar{\pi} \, dy + \int_{\partial\{h>0\}} \dot{h} \bar{\zeta} \, ds = - \int_{\{h>0\}} m(h) \partial_y \pi \partial_y \bar{\pi} \, dy - \int_{\partial\{h>0\}} m_{\text{CL}}(\partial_y h) \zeta \bar{\zeta} \, ds. \quad (\text{A.10b})$$

where in the numerical scheme $m = m(h)$ and $m_{\text{CL}} = m_{\text{CL}}(\partial_y h)$ are evaluated explicitly from the previous time step. Now, we introduce an arbitrary Lagrangian-Eulerian method by constructing a mapping $\xi : \{h > 0\}(t_0) \rightarrow \{h > 0\}(t)$ using the linear construction

$$\xi(t, y) = \frac{s_+(t) - s_-(t)}{s_+(t_0) - s_-(t_0)} (y - s_-(t_0)) + s_-(t), \quad (\text{A.11})$$

which allows to define the function $H(t) : \{h > 0\}(t_0) \rightarrow \mathbb{R}$ via $H(t, y) = h(t, \xi(t, y))$. By construction we have $H(t, s_{\pm}(t_0)) \equiv 0$ and correspondingly $\partial_t H = 0$ at the fixed contact line $y = s_{\pm}(t_0)$. For the time derivatives we have $\dot{H}(t, y) = \dot{h}(t, \xi) + \partial_y h(t, \xi) \dot{\xi}$. The knowledge of \dot{h} from the solution of (A.10) and the boundary condition $\dot{H} = 0$ entirely determine the time derivatives \dot{H} and the mapping $\dot{\xi}$. Note that, for a moving front, $\partial_y h = 0$ at $\partial\{h > 0\}$ is not an issue for the application of P_1 finite elements, since on the last element connected to s_{\pm} , (A.9) yields a possibly small but nonzero value of $|\partial_y h|$. In the reference domain we can update both the height function $H(t + \tau, y) = H(t, y) + \tau \dot{H}(t, y)$ and the map $\xi(t + \tau, y) = \xi(t, y) + \tau \dot{\xi}(t, y)$, which uniquely determines $h(t, y)$ on a moving domain. A similar approach for partial wetting is studied in [66] and higher-dimensional extensions are discussed in [77].

REFERENCES

- [1] Alexander Oron, Stephen H. Davis, and S. George Bankoff. Long-scale evolution of thin liquid films. *Rev. Mod. Phys.*, 69:931–980, Jul 1997.
- [2] Daniel Bonn, Jens Eggers, Joseph Indekeu, Jacques Meunier, and Etienne Rolley. Wetting and spreading. *Rev. Mod. Phys.*, 81:739–805, May 2009.
- [3] M. Günther and G. Prokert. A justification for the thin film approximation of Stokes flow with surface tension. *J. Differential Equations*, 245(10):2802–2845, 2008.
- [4] H. P. Greenspan. On the motion of a small viscous droplet that wets a surface. *J. Fluid Mech.*, 84(1):125–143, 1978.
- [5] L. Giacomelli and F. Otto. Rigorous lubrication approximation. *Interfaces Free Bound.*, 5(4):483–529, 2003.
- [6] L. Giacomelli and F. Otto. Variational formulation for the lubrication approximation of the Hele-Shaw flow. *Calc. Var. Partial Differential Equations*, 13(3):377–403, 2001.
- [7] Hans Knüpfer and Nader Masmoudi. Well-posedness and uniform bounds for a nonlocal third order evolution operator on an infinite wedge. *Comm. Math. Phys.*, 320(2):395–424, 2013.
- [8] H. Knüpfer and N. Masmoudi. Darcy’s flow with prescribed contact angle: Well-posedness and lubrication approximation. *Archive for Rational Mechanics and Analysis*, 218(2):589–646, 2015.
- [9] Bogdan-Vasile Matioc and Georg Prokert. Hele-Shaw flow in thin threads: a rigorous limit result. *Interfaces Free Bound.*, 14(2):205–230, 2012.
- [10] Marion Dziwnik, Maciek Korzec, Andreas Münch, and Barbara Wagner. Stability analysis of unsteady, nonuniform base states in thin film equations. *Multiscale Modeling & Simulation*, 12(2):755–780, 2014.
- [11] S. Boatto, L. P. Kadanoff, and P. Olla. Traveling-wave solutions to thin-film equations. *Phys. Rev. E* (3), 48(6):4423–4431, 1993.
- [12] JR King and M Bowen. Moving boundary problems and non-uniqueness for the thin film equation. *European Journal of Applied Mathematics*, 12(3):321–356, 2001.
- [13] Lorenzo Giacomelli, Manuel V. Gnann, and Felix Otto. Rigorous asymptotics of traveling-wave solutions to the thin-film equation and Tanner’s law. *Nonlinearity*, 29(9):2497–2536, 2016.
- [14] Francisco Bernis and Avner Friedman. Higher order nonlinear degenerate parabolic equations. *J. Differential Equations*, 83(1):179–206, 1990.
- [15] E. Beretta, M. Bertsch, and R. Dal Passo. Nonnegative solutions of a fourth-order nonlinear degenerate parabolic equation. *Arch. Rational Mech. Anal.*, 129(2):175–200, 1995.
- [16] A. L. Bertozzi and M. Pugh. The lubrication approximation for thin viscous films: regularity and long-time behavior of weak solutions. *Comm. Pure Appl. Math.*, 49(2):85–123, 1996.
- [17] M. Bertsch, R. Dal Passo, H. Garcke, and G. Grün. The thin viscous flow equation in higher space dimensions. *Adv. Differential Equations*, 3(3):417–440, 1998.

- [18] Günther Grün. Droplet spreading under weak slippage—existence for the Cauchy problem. *Comm. Partial Differential Equations*, 29(11-12):1697–1744, 2004.
- [19] Lidia Ansini and Lorenzo Giacomelli. Doubly nonlinear thin-film equations in one space dimension. *Arch. Ration. Mech. Anal.*, 173(1):89–131, 2004.
- [20] Francisco Bernis. Finite speed of propagation and continuity of the interface for thin viscous flows. *Adv. Differential Equations*, 1(3):337–368, 1996.
- [21] Francisco Bernis. Finite speed of propagation for thin viscous flows when $2 \leq n < 3$. *C. R. Acad. Sci. Paris Sér. I Math.*, 322(12):1169–1174, 1996.
- [22] J. Hulshof and A. E. Shishkov. The thin film equation with $2 \leq n < 3$: finite speed of propagation in terms of the L^1 -norm. *Adv. Differential Equations*, 3(5):625–642, 1998.
- [23] R. Dal Passo, L. Giacomelli, and G. Grün. A waiting time phenomenon for thin film equations. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 30(2):437–463, 2001.
- [24] R. Dal Passo, L. Giacomelli, and A. E. Shishkov. The thin film equation with nonlinear diffusion. *Comm. Partial Differential Equations*, 26(9-10):1509–1557, 2001.
- [25] Günther Grün. Droplet spreading under weak slippage: a basic result on finite speed of propagation. *SIAM J. Math. Anal.*, 34(4):992–1006, 2003.
- [26] G. Grün. Droplet spreading under weak slippage: the waiting time phenomenon. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 21(2):255–269, 2004.
- [27] L. Giacomelli and A. E. Shishkov. Propagation of support in one-dimensional convected thin-film flow. *Indiana Univ. Math. J.*, 54(4):1181–1215, 2005.
- [28] Lorenzo Giacomelli and Günther Grün. Lower bounds on waiting times for degenerate parabolic equations and systems. *Interfaces Free Bound.*, 8(1):111–129, 2006.
- [29] J. Fischer. Optimal lower bounds on asymptotic support propagation rates for the thin-film equation. *J. Differential Equations*, 255(10):3127–3149, 2013.
- [30] J. Fischer. Upper Bounds on Waiting Times for the Thin-Film Equation: The Case of Weak Slippage. *Arch. Rational Mech. Anal.*, 211(3):771–818, 2014.
- [31] Julian Fischer. Behaviour of free boundaries in thin-film flow: the regime of strong slippage and the regime of very weak slippage. *Ann. Inst. H. Poincaré C Anal. Non Linéaire*, 33(5):1301–1327, 2016.
- [32] Nicola De Nitti and Julian Fischer. Sharp criteria for the waiting time phenomenon in solutions to the thin-film equation. *Comm. Partial Differential Equations*, 47(7):1394–1434, 2022.
- [33] L. Giacomelli, H. Knüpfer, and F. Otto. Smooth zero-contact-angle solutions to a thin-film equation around the steady state. *J. Differential Equations*, 245(6):1454–1506, 2008.
- [34] Björn Bringmann, Lorenzo Giacomelli, Hans Knüpfer, and Felix Otto. Corrigendum to “Smooth zero-contact-angle solutions to a thin-film equation around the steady state” [J. Differential Equations 245 (2008) 1454–1506]. *J. Differential Equations*, 261(2):1622–1635, 2016.
- [35] L. Giacomelli, M. Gnann, and F. Otto. Regularity of source-type solutions to the thin-film equation with zero contact angle and mobility exponent between $3/2$ and 3 . *European Journal of Applied Mathematics*, 24:735–760, 10 2013.
- [36] Manuel V. Gnann. Well-posedness and self-similar asymptotics for a thin-film equation. *SIAM J. Math. Anal.*, 47(4):2868–2902, 2015.
- [37] Fethi Ben Belgacem, Manuel V. Gnann, and Christian Kuehn. A dynamical systems approach for the contact-line singularity in thin-film flows. *Nonlinear Anal.*, 144:204–235, 2016.
- [38] Manuel V. Gnann. On the regularity for the Navier-slip thin-film equation in the perfect wetting regime. *Arch. Ration. Mech. Anal.*, 222(3):1285–1337, 2016.
- [39] Christian Seis. The thin-film equation close to self-similarity. *Anal. PDE*, 11(5):1303–1342, 2018.
- [40] L. Giacomelli, M.V. Gnann, H. Knüpfer, and F. Otto. Well-posedness for the Navier-slip thin-film equation in the case of complete wetting. *J. Differential Equations*, 257(1):15–81, 2014.
- [41] Manuel V. Gnann and Mircea Petrache. The Navier-slip thin-film equation for 3D fluid films: existence and uniqueness. *J. Differential Equations*, 265(11):5832–5958, 2018.
- [42] Manuel V. Gnann, Slim Ibrahim, and Nader Masmoudi. Stability of receding traveling waves for a fourth order degenerate parabolic free boundary problem. *Adv. Math.*, 347:1173–1243, 2019.
- [43] Felix Otto. Lubrication approximation with prescribed nonzero contact angle. *Comm. Partial Differential Equations*, 23(11-12):2077–2164, 1998.
- [44] M. Bertsch, L. Giacomelli, and G. Karali. Thin-film equations with “partial wetting” energy: existence of weak solutions. *Phys. D*, 209(1-4):17–27, 2005.
- [45] A. Mellet. The thin film equation with non-zero contact angle: a singular perturbation approach. *Comm. Partial Differential Equations*, 40(1):1–39, 2015.
- [46] Hans Knüpfer. Well-posedness for the Navier slip thin-film equation in the case of partial wetting. *Comm. Pure Appl. Math.*, 64(9):1263–1296, 2011.
- [47] Hans Knüpfer. Well-posedness for a class of thin-film equations with general mobility in the regime of partial wetting. *Arch. Ration. Mech. Anal.*, 218(2):1083–1130, 2015.
- [48] Sergey Degtyarev. Classical solvability of the multidimensional free boundary problem for the thin film equation with quadratic mobility in the case of partial wetting. *Discrete Contin. Dyn. Syst.*, 37(7):3625–3699, 2017.
- [49] Hans Knüpfer. Erratum to: Well-posedness for a class of thin-film equations with general mobility in the regime of partial wetting. *Arch. Ration. Mech. Anal.*, 2023. in preparation.
- [50] Elias Esselborn. Relaxation rates for a perturbation of a stationary solution to the thin-film equation. *SIAM J. Math. Anal.*, 48(1):349–396, 2016.

- [51] Mohamed Majdoub, Nader Masmoudi, and Slim Tayachi. Relaxation to equilibrium in the one-dimensional thin-film equation with partial wetting and linear mobility. *Comm. Math. Phys.*, 385(2):837–857, 2021.
- [52] Ciro Semprebon and Martin Brinkmann. On the onset of motion of sliding drops. *Soft Matter*, 10(18):3325–3334, 2014.
- [53] Natalie Grunewald and Inwon Kim. A variational approach to a quasi-static droplet model. *Calc. Var. Partial Differential Equations*, 41(1-2):1–19, 2011.
- [54] M. G. Delgadino and A. Mellet. On the relationship between the thin film equation and tanner’s law. *Communications on Pure and Applied Mathematics*, 74(3):507–543, 2021.
- [55] P. Ehrhard and S. H. Davis. Nonisothermal spreading of liquid drops on horizontal plates. *J. Fluid Mech.*, 229:365–388, 1991.
- [56] M. Chiricotto and L. Giacomelli. Droplets spreading with contact-line friction: lubrication approximation and traveling wave solutions. *Communications in Applied and Industrial Mathematics*, 2(2), 2011.
- [57] M. Chiricotto and L. Giacomelli. Scaling laws for droplets spreading under contact-line friction. *Commun. Math. Sci.*, 11(2):361–383, 2013.
- [58] Maria Chiricotto and Lorenzo Giacomelli. Weak solutions to thin-film equations with contact-line friction. *Interfaces Free Bound.*, 19(2):243–271, 2017.
- [59] Weiqing Ren and Weinan E. Boundary conditions for the moving contact line problem. *Physics of Fluids*, 19(2):022101, 2007.
- [60] Weiqing Ren, Dan Hu, and Weinan E. Continuum models for the contact line problem. *Physics of Fluids*, 22(10):102103, 2010.
- [61] W. Ren and W. E. Derivation of continuum models for the moving contact line problem based on thermodynamic principles. *Comm. Math. Sci.*, 9(2):597–606, 2011.
- [62] P. G. de Gennes. Wetting - Statics and dynamics. *Rev. Mod. Phys.*, 57(3):827–863, 1985.
- [63] F. Otto. Lubrication approximation with prescribed nonzero contact angle. *Comm. Partial Differential Equations*, 23(11-12):2077–2164, 1998.
- [64] M. Bertsch, L. Giacomelli, and G. Karali. Thin-film equations with “partial wetting” energy: existence of weak solutions. *Phys. D*, 209(1-4):17–27, 2005.
- [65] R. Durastanti and L. Giacomelli. Spreading equilibria under mildly singular potentials: pancakes versus droplets. *Journal of Nonlinear Science*, 32(5), 2022.
- [66] Dirk Peschka. Variational approach to dynamic contact angles for thin films. *Physics of Fluids*, 30(8):082115, 2018.
- [67] M. Bertsch, R. Dal Passo, S. H. Davis, and L. Giacomelli. Effective and microscopic contact angles in thin film dynamics. *European J. Appl. Math.*, 11(2):181–201, 2000.
- [68] L. Giacomelli and F. Otto. Droplet spreading: intermediate scaling law by PDE methods. *Comm. Pure Appl. Math.*, 55(2):217–254, 2002.
- [69] L. Ansini and L. Giacomelli. Shear-thinning liquid films: macroscopic and asymptotic behaviour by quasi-self-similar solutions. *Nonlinearity*, 15(6):2147–2164, 2002.
- [70] Manuel V. Gnann and Anouk C. Wisse. The Cox-Voinov law for traveling waves in the partial wetting regime. *Nonlinearity*, 35(7):3560–3592, 2022.
- [71] F. Bernis, L. A. Peletier, and S. M. Williams. Source type solutions of a fourth order nonlinear degenerate parabolic equation. *Nonlinear Anal.*, 18(3):217–234, 1992.
- [72] J. A. Carrillo and G. Toscani. Long-time asymptotics for strong solutions of the thin film equation. *Comm. Math. Phys.*, 225(3):551–571, 2002.
- [73] Eric A. Carlen and Süleyman Ulusoy. Asymptotic equipartition and long time behavior of solutions of a thin-film equation. *J. Differential Equations*, 241(2):279–292, 2007.
- [74] Daniel Matthes, Robert J. McCann, and Giuseppe Savaré. A family of nonlinear fourth order equations of gradient flow type. *Comm. Partial Differential Equations*, 34(10-12):1352–1397, 2009.
- [75] Eric A. Carlen and Süleyman Ulusoy. Localization, smoothness, and convergence to equilibrium for a thin film equation. *Discrete Contin. Dyn. Syst.*, 34(11):4537–4553, 2014.
- [76] N. F. Smyth and J. M. Hill. High-order nonlinear diffusion. *IMA J. Appl. Math.*, 40(2):73–86, 1988.
- [77] Dirk Peschka and Luca Heltai. Model hierarchies and higher-order discretisation of time-dependent thin-film free boundary problems with dynamic contact angle. *Journal of Computational Physics*, 464:111325, 2022.

(Lorenzo Giacomelli) SBAI DEPARTMENT, SAPIENZA UNIVERSITY OF ROME - VIA A. SCARPA 16, 00161 ROMA, ITALY - LORENZO.GIACOMELLI@UNIROMA1.IT

(Manuel V. Gnann) DELFT INSTITUTE OF APPLIED MATHEMATICS, FACULTY OF ELECTRICAL ENGINEERING, MATHEMATICS AND COMPUTER SCIENCE, DELFT UNIVERSITY OF TECHNOLOGY, MEKELWEG 4, 2628 CD DELFT, THE NETHERLANDS

(Dirk Peschka) WEIERSTRASS INSTITUTE - MOHRENSTRASSE 39, 10117 BERLIN, GERMANY - DIRK.PESCHKA@WIAS-BERLIN.DE