



Delft University of Technology

Document Version

Final published version

Licence

CC BY

Citation (APA)

Hermans, T., Malagón-Santos, V., Katsman, C., Jane, R., Rasmussen, D., Haasnoot, M., Garner, G., Kopp, R., Oppenheimer, M., & Slangen, A. (2023). *The Timing of Decreasing Coastal Flood Protection Due to Sea-Level Rise*. Abstract from EGU General Assembly 2023, Vienna, Austria. <https://doi.org/10.5194/egusphere-egu23-4834>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

In case the licence states "Dutch Copyright Act (Article 25fa)", this publication was made available Green Open Access via the TU Delft Institutional Repository pursuant to Dutch Copyright Act (Article 25fa, the Taverne amendment). This provision does not affect copyright ownership.
Unless copyright is transferred by contract or statute, it remains with the copyright holder.

Sharing and reuse

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

This work is downloaded from Delft University of Technology.

Abstract EGU23-4834

EGU23-4834

<https://doi.org/10.5194/egusphere-egu23-4834>

EGU General Assembly 2023

© Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.



The Timing of Decreasing Coastal Flood Protection Due to Sea-Level Rise

Tim Hermans^{1,2,3}, Victor Malagón-Santos¹, Caroline Katsman⁴, Robert Jane⁵, Dj Rasmussen⁶, Marjolijn Haasnoot^{7,8}, Gregory Garner⁹, Robert Kopp^{10,11}, Michael Oppenheimer^{6,12,13}, and Aimée Slangen¹¹NIOZ Royal Netherlands Institute for Sea Research, Department of Estuarine & Delta Systems, PO Box 140, 4400 AC Yerseke, The Netherlands²Department of Geosciences and Remote Sensing, Delft University of Technology, Stevinweg 1, 2628 CN Delft, The Netherlands³Institute for Marine and Atmospheric Research Utrecht, Utrecht University, Utrecht, The Netherlands (t.h.j.hermans@uu.nl)⁴Department of Hydraulic Engineering, Delft University of Technology, Stevinweg 1, 2628 CN Delft, The Netherlands⁵Department of Civil, Construction and Environmental Engineering, University of Central Florida, Orlando, FL, USA⁶Princeton School of Public & International Affairs, Princeton University, Princeton, NJ, USA⁷Deltares, Delft, Boussinesqweg 1, 2629 HV Delft, the Netherlands⁸Utrecht University, Department of Geosciences, Princetonlaan 8a, Utrecht, the Netherlands⁹GRO Intelligence, 505 Park Ave 12th Floor, New York, NY 10022, USA¹⁰Department of Earth and Planetary Sciences, Rutgers University, Piscataway, NJ, USA¹¹Rutgers Institute of Earth, Ocean, and Atmospheric Sciences, Rutgers University, New Brunswick, NJ, US¹²Department of Geosciences, Princeton University, Princeton, NJ, USA¹³High Meadows Institute, Princeton University, Princeton, NJ, USA

Sea-level rise (SLR) amplifies the frequency of extreme sea levels as it raises their baseline height. Projections of the frequency amplification of extremes are often computed for arbitrary future years and relative to the historical centennial event, which is not necessarily meaningful locally. Consequently, such projections may not provide salient information to adaptation planners, as they do not indicate when certain flood risk thresholds will be crossed given the current degree of local coastal flood protection.

To better support adaptation planning, we introduce a framework that extends the emerging timing perspective on sea-level rise to the frequency amplification of extreme sea levels. Moreover, by relating amplification factors to local flood protection standards estimated with the FLOPROS modelling approach, we project the timing of decreases in the local degree of protection. The sea-level rise required for such decreases is derived from extreme sea-level distributions inferred from GESLA3 observations and combined with the relative sea-level projections of the Sixth Assessment Report of the IPCC until 2150 to compute the timing of these decreases at tide gauges globally.

Our central estimates indicate that the estimated degrees of protection will be exceeded 10 times as frequently within the next 30 years (the lead time that large adaptation measures may take) at 26 & 32% of the tide gauges considered, and annually at 4 & 8%, for respectively a low & high emissions scenario (SSP1-2.6 & SSP3-7.0). Even though our results are based on estimated degrees of protection, they highlight that at several locations substantial decreases in the degree of protection may occur before large adaptation measures can be completed. Furthermore, we find that under SSP3-7.0, the same decreases in the degree of coastal protection will occur substantially faster in the future as sea-level rise accelerates. Our projection framework adds a new perspective on the frequency amplifications of extremes that may help adaptation planners to assess the available lead time and useful lifetime of protective infrastructure, given unacceptable decreases in the degree of coastal protection.

How to cite: Hermans, T., Malagón-Santos, V., Katsman, C., Jane, R., Rasmussen, D., Haasnoot, M., Garner, G., Kopp, R., Oppenheimer, M., and Slangen, A.: The Timing of Decreasing Coastal Flood Protection Due to Sea-Level Rise, EGU General Assembly 2023, Vienna, Austria, 24–28 Apr 2023, EGU23-4834, <https://doi.org/10.5194/egusphere-egu23-4834>, 2023.