

Finding the essential

Improving conservation monitoring across scales

Guerra, C.A.; Pendleton, L.; Drakou, E.G.; Proença, V.; Appeltans, W.; Domingos, T.; Geller, G.; Giamberini, S.; Ziemba, A.; More Authors

DOI

[10.1016/j.gecco.2019.e00601](https://doi.org/10.1016/j.gecco.2019.e00601)

Publication date

2019

Document Version

Final published version

Published in

Global Ecology and Conservation

Citation (APA)

Guerra, C. A., Pendleton, L., Drakou, E. G., Proença, V., Appeltans, W., Domingos, T., Geller, G., Giamberini, S., Ziemba, A., & More Authors (2019). Finding the essential: Improving conservation monitoring across scales. *Global Ecology and Conservation*, 18, Article e00601. <https://doi.org/10.1016/j.gecco.2019.e00601>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.



ELSEVIER

Contents lists available at ScienceDirect

Global Ecology and Conservation

journal homepage: <http://www.elsevier.com/locate/gecco>

Review Paper

Finding the essential: Improving conservation monitoring across scales



C.A. Guerra ^{a, b, *}, L. Pendleton ^{c, d}, E.G. Drakou ^{c, e}, V. Proença ^f, W. Appeltans ^g, T. Domingos ^f, G. Geller ^{h, i}, S. Giamberini ^j, M. Gill ^k, H. Hummel ^l, S. Imperio ^{j, u}, M. McGeoch ^m, A. Provenzale ^j, I. Serral ⁿ, A. Stritih ^o, E. Turak ^{p, q}, P. Vihervaara ^r, A. Ziemba ^{s, t}, H.M. Pereira ^{a, b, v}

^a Institute of Biology, Martin Luther University Halle-Wittenberg, Am Kirchtor 1, 06108, Halle (Saale), Germany

^b German Centre for Integrative Biodiversity Research (iDiv), Halle-Jena-Leipzig, Deutscher Platz 5e, 04103, Leipzig, Germany

^c International Chair of Excellence, University of Brest, Ifremer, CNRS, UMR 6308, AMURE, IUEM, 29280, Plouzané, France

^d Duke Nicholas Institute for Environmental Policy Solutions, USA

^e Faculty of Geo-Information Science and Earth Observation (ITC), University of Twente, P.O. Box 6, 7500 AA, Enschede, the Netherlands

^f MARETEC, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1, 1040-001, Lisboa, Portugal

^g UNESCO Intergovernmental Oceanographic Commission, IOC Project Office for IODE, Wandelaarkaai 7, 8400, Oostende, Belgium

^h Group on Earth Observations, Geneva, Switzerland

ⁱ NASA Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

^j Institute of Geosciences and Earth Resources, National Research Council, Via Moruzzi 1, 56124, Pisa, Italy

^k NatureServe, Arlington, Virginia, USA

^l Royal Netherlands Institute for Sea Research, Yerseke and Utrecht University, Yerseke, the Netherlands

^m School of Biological Sciences, Monash University, Clayton 3800, Victoria, Australia

ⁿ Grumets, CREAM, Cerdanyola del Vallès, Barcelona, Spain

^o Planning of Landscape and Urban Systems, Swiss Federal Institute of Technology (ETH), Stefano-Francini-Platz 5, 8093, Zurich, Switzerland

^p NSW Office of Environment and Heritage, 10 Valentine Avenue Parramatta, NSW, 2150, Australia

^q Australian Museum, 6 College Street, Sydney, NSW 2000, Australia

^r Finnish Environment Institute, Biodiversity Center, Ecosystem services Unit, Latokartanonkaari 11, FI-00790, Helsinki, Finland

^s Deltares, Department of Information, Resilience and Planning, Boussinesqweg 1, 2629 HV, Delft, Netherlands

^t Delft University of Technology, Applied Mathematics, Mekelweg 2, 2628 CD, Delft, Netherlands

^u Italian Institute for Environmental Protection and Research (ISPRA), via Ca' Fornacetta 9, 40064 Ozzano Emilia, BO, Italy

^v Center for Latin America Studies, Stanford University, Stanford, CA 94305, USA

ARTICLE INFO

Article history:

Received 18 January 2019

Received in revised form 23 March 2019

Accepted 24 March 2019

Keywords:

Essential variables

Global monitoring

Biodiversity

Ecosystem services

Systems approach

ABSTRACT

To account for progress towards conservation targets, monitoring systems should capture not only information on biodiversity but also knowledge on the dynamics of ecological processes and the related effects on human well-being. Protected areas represent complex social-ecological systems with strong human-nature interactions. They are able to provide relevant information about how global and local scale drivers (e.g., climate change, land use change) impact biodiversity and ecosystem services. Here we develop a framework that uses an ecosystem-focused approach to support managers in identifying essential variables in an integrated and scalable approach. We advocate that this approach can complement current essential variable developments, by allowing conservation managers to draw on system-level knowledge and theory of biodiversity and ecosystems to identify locally important variables that meet the local or sub-global needs for conservation data. This requires the development of system narratives and causal diagrams that pinpoints the

* Corresponding author. Institute of Biology, Martin Luther University Halle-Wittenberg, Am Kirchtor 1, 06108, Halle (Saale), Germany.
E-mail address: carlos.guerra@div.de (C.A. Guerra).

social-ecological variables that represent the state and drivers of the different components, and their relationships. We describe a scalable framework that builds on system based narratives to describe all system components, the models used to represent them and the data needed. Considering the global distribution of protected areas, with an investment in standards, transparency, and on active data mobilisation strategies for essential variables, these have the potential to be the backbone of global biodiversity monitoring, benefiting countries, biodiversity observation networks and the global biodiversity community.

© 2019 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Conservation management mediates the interactions between nature, human development and environmental impacts (Johnson et al., 2017; Pecl et al., 2017). It needs to be guided by clear conservation goals, and underpinned by information on the state and trends of social-ecological interactions (Ostrom, 2009). Protected areas are managed for a variety of conservation goals, from the protection of keystone species to the conservation and maintenance of entire ecosystems (Naughton-Treves et al., 2005). Despite their current aims, future expectations for their role may change due to changing environmental, demographic, social and political conditions, including new conservation objectives that were not previously considered (Rands et al., 2010).

To account for progress towards conservation targets, monitoring systems should capture information on biodiversity change but also knowledge on the dynamics of ecological processes and their effects on human well-being (Defries and Nagendra, 2017; Eisenhauer et al., 2019). These, integrated data needs will require the implementation of new monitoring approaches that include a wider characterization and quantification of social-ecological systems, and the optimization of available operational resources for long-term monitoring (Carpenter et al., 2009). Effective monitoring systems must also capture information on pressures and threats to allow for the identification of causal mechanisms of change (Maron et al., 2017). Without such integration, monitoring systems would be blind to the mechanisms driving ecological trends and thus, unable to guide effective policy responses (Nichols and Williams, 2006).

Protected areas constitute, in many parts of the planet, one of the main sources of in situ biodiversity information (Martin et al., 2012). These areas represent complex social-ecological systems with strong human-nature interactions, being able to provide relevant information about how global and local scale drivers impact biodiversity and ecosystem services. Nevertheless, effective monitoring systems must be scalable as many drivers of biodiversity change occur at scales beyond individual conservation areas, thereby, requiring integrated monitoring approaches that allow for the detection of drivers operating at broader scales (Navarro et al., 2018; Eisenhauer et al., 2019).

Recently, Pereira et al. (2013) argued that, to overcome the current gap in biodiversity data and to improve the scalability of observation systems, the scientific and practice communities should focus on identifying and capturing essential variables, i.e. critical measurements required for the study, report, and management of global biodiversity and ecosystem change, as well as other social-ecological dimensions e.g. (Bojinski et al., 2014). Here we develop a framework that uses an ecosystem-focused approach to support managers in identifying essential variables in an integrated and scalable approach.

2. From essential variables to conservation management

Monitoring biodiversity, entire ecosystems, and the services they provide often involves the implementation of costly monitoring programs that seek to combine targeted, hypothesis-driven and/or surveillance monitoring from different sources (Lindenmayer et al., 2018; Pereira et al., 2017). When establishing a monitoring program, conservation managers face the challenge of balancing available resources between monitoring activities, operational costs, research and conservation action (Gill et al., 2017). This often leaves monitoring programs with limited financial and logistic resources, hampering their usefulness and encompassing nature. Since the purpose of monitoring activities in this context is to facilitate more effective conservation actions, it is imperative that monitoring systems be as targeted and efficient as possible. If they fail to meet this objective, they risk misinforming conservation actions and draining resources (Legg and Nagy, 2006; Cameron et al., in press).

The process of identifying essential variables is not an exclusion process but rather a priority setting process, where central elements of monitoring effort are identified to improve the understanding of social-ecological dynamics. It facilitates the prioritization and optimization of resources and allows for a transparent and direct way to communicate monitoring needs through different levels of knowledge and decision-making. The identification of essential variables often uses a top-down approach, where experts define at the global level what variables should be monitored. Here we advocate that this approach can be complemented with a bottom-up approach, where conservation managers draw on system-level knowledge and theory (Liu et al., 2015) to identify locally important variables that meet local or sub-global needs for conservation data (Turak et al., 2016a, Fig. 1).

Our framework proceeds in four sequential steps, that we introduce briefly below:

1. **Narratives** are used to describe the complexity of ecological and social interactions that influence a specific conservation or management goal. These narratives should be harnessed to identify the major **system components, functions and processes**, and the underlying causal relationships that affect them. Using existing ecological knowledge, ecosystems supporting similar communities and processes, or facing comparable anthropogenic pressures can be grouped and systematically described (Turak et al., 2016b). These elements collectively summarize the monitoring needs to assess a given conservation goal. By examining the causal relations that link the social-ecological system, it is also possible to represent the focal system elements that are conditioned or condition the entire social-ecological system.
2. To quantitatively address these system elements and their changes through time, **models** that can describe the spatial and temporal interactions are often necessary. These models seek to describe critical ecosystem dynamics and can only be operationalized contingent upon the availability of interpretable information and technical capacity to implement them. Some of these models only use static variables, describing the state of a given system component (e.g. topography), but others rely on dynamic variables (e.g. species abundance) that can be analyzed to evaluate trends and make scenarios of future conservation trajectories.
3. Independent of the type of information used, modelling approaches rely on a set of **variables** that condense and summarize observations. These strongly depend on source datasets, making their identification a critical step to operationalize the models and create the foundation for the design and implementation of monitoring systems. Within these, conservation managers are able to identify concrete groups of locally important variables that together describe the social-ecological system.
4. **Observations** constitute the source datasets that are the basic building blocks of any given monitoring system and of which the variables are derived. These are usually dynamic measures that require the consideration of both in situ and remote (e.g. satellite data) monitoring activities (Vihervaara et al., 2017), but can also include information with smaller temporal variability that is still critical to understand system dynamics (e.g. soil type). Conservation managers must prioritize which observations must be collected in situ. Other data requirements can be met through the establishment of institutional collaborations that allows to maintain a constant flow of information, recognizing the fact that many important ecological processes and drivers of change are occurring at scales beyond the extent of the protected area in question.

3. Creating narratives to capture the complexity of social-ecological systems

The use of narratives can contribute to the identification of monitoring priorities targeting specific system elements that are fundamental to understand a social-ecological system. Such narratives have been widely used to facilitate communication between stakeholders engaged in biodiversity conservation and beyond (Metz et al., 2007; Moon et al., 2019). At the same




Fig. 1. Conceptual representation of a system approach to identify essential variables and monitoring priorities in conservation areas. The arrows in the Figure differentiate between direct data dependencies (full lines) and the expected causal relations between system components and essential variables (dashed lines). Illustrative implementation examples are given in Fig. 2a and Fig. 2b.

However, matching locally important variables with global essential variables is needed to foster the scalability of the data collected by protected areas. For instance, collection, mobilisation, and publishing of data regarding species distribution and population structure can use the standards, methods and tools being developed for essential biodiversity variables (Kissling et al., 2018). In doing so, automated data flows can be established and feed the development of global datasets critical for biodiversity monitoring and research (Navarro et al., 2018).

Still, there are consistency and scalability issues when several conservation areas consider the same essential variable. As an example, different conservation areas can identify species distribution as a monitoring variable without the necessary thematic consistency (e.g. in Fig. 2a species distribution corresponds to forest species while in Fig. 2b it corresponds to cetaceans). Addressing such thematic, and eventually temporal, inconsistencies will be critical when considering interoperability across conservation areas. A solution would imply direct coordination of monitoring activities (e.g. at the national level) that allows information to move across scales and ecosystems, as part of a multi-level conservation strategy.

Ultimately, the global selection of essential biodiversity variables needs also to be informed by these bottom-up variable selection efforts. Considering the relevance of the information collected by conservation areas, developing global monitoring schemes for essential biodiversity variables needs to draw on those efforts. Therefore, locally important variables which are identified across multiple sub-global regions or protected areas are likely to be those most viable to be monitored globally. This happens not only from a practical data availability point of view but also because variables that are systematically identified as locally important are probably also globally important for international conventions and countries.

5. Future implications

One of the challenges faced by conservation managers is to establish, maintain and highlight the need for long-term repeated measurements of data relevant for conservation purposes. It is critical that any monitoring system first clearly considers the policy and conservation objectives that the system needs to serve, ensuring a clear link to decision-making. These links typically involve a co-development process where decision-makers, managers, and researchers work together to identify key goals and objectives providing a clear roadmap for the development and operation of the monitoring system (Craine et al., 2007). With clear objectives in hand, the identification of essential variables is a useful way to prioritize data collection efforts needed for operational and policy-relevant conservation monitoring systems. It allows to focus resources, identify capacity building plus data needs, and to mobilize knowledge including the resources necessary to sustain local and cross-scale assessments.

Establishing a group of essential variables that can commonly be assessed across conservation goals and conservation areas allows monitoring systems to contribute directly to high level (e.g. national and international) monitoring efforts. At the same time, the benefits generated by such a coordinated monitoring program can be multiplied across spatial scales, increasing the effectiveness and usefulness of conservation management and monitoring for a variety of objectives. This harmonized approach requires specific guidelines that structure and standardize monitoring efforts but has the potential to maximize the outcomes from conservation efforts, allowing monitoring efforts to go beyond borders, ecosystem types and realms. It also requires that, in the context of conservation monitoring, essential biodiversity variables cannot be seen as a closed box and EBVs have to be seen in the context of others (e.g., ECVs, EOVs, etc.).

Considering their global distribution, covering all ecoregions (Dinerstein et al., 2017), current targets (e.g., Aichi target 11), and the essential variable based approach here described, protected areas have the potential to be the backbone of global biodiversity monitoring. Even considering that only a fraction of these globally distributed areas would be able or willing to participate in such initiative, thousands of sites would be able to report data. A step forward would be to attach to global conservation targets the global monitoring of these conservation areas and to establish a global biodiversity monitoring backbone. With strong political support, an investment in standards, transparent methods, and on active data mobilisation strategies (Navarro et al., 2018), in the face of climate and land use change, these areas could be pivotal as early warning systems to signal major regional and global nature shifts. Such an approach would benefit from being written into the post-2020 CBD targets, giving the International Union for Conservation of Nature the necessary political support to make this a reality and provide a global baseline for nature status and conservation.

Acknowledgement

This work was supported by the European Union's Horizon 2020 research and innovation programme under grant agreement No 641762—ECOPOTENTIAL project ("Improving future ecosystem benefits through earth observations"). The work of Linwood Pendleton and Evangelia Drakou was supported by the "Laboratoire d'Excellence" LabexMER (ANR-10-LABX-19) at the European Institute of Marine Sciences (IUEM).

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.gecco.2019.e00601>.

References

- Bojinski, S., Verstraete, M., Peterson, T.C., Richter, C., Simmons, A., Zemp, M., 2014. The concept of essential climate variables in support of climate research, applications, and policy. *Bull. Am. Meteorol. Soc.* 95, 1431–1443. <https://doi.org/10.1175/BAMS-D-13-00047.1>.
- Cameron, E. K., I. S. Martins, P. Lavelle, J. Mathieu, L. Tedersoo, M. Bahram, F. Gottschall, et al. In press. Global mismatches in aboveground and belowground biodiversity. *Conservation Biology*. <https://doi.org/10.1111/cobi.13311>.
- Carpenter, S.R., Mooney, H. A., Agard, J., Capistrano, D., Defries, R.S., Díaz, S., Dietz, T., Duraiappah, A.K., Oteng-Yeboah, A., Pereira, H.M., Perrings, C., Reid, W.V., Sarukhan, J., Scholes, R.J., Whyte, A., 2009. Science for managing ecosystem services: beyond the Millennium Ecosystem Assessment. *Proc. Natl. Acad. Sci. U.S.A.* 106, 1305–1312. <https://doi.org/10.1073/pnas.0808772106>.
- Craine, J.M., Battersby, J., Elmore, A.J., Jones, A.W., 2007. Building EDENs: the rise of environmentally distributed ecological networks. *Bioscience* 57, 45. <https://doi.org/10.1641/B570108>.
- Defries, R., Nagendra, H., 2017. Ecosystem management as a wicked problem. *Science* 270, 265–270. <https://doi.org/10.1126/science.aal1950>.
- Dinerstein, E., Olson, D., Joshi, A., Vynne, C., Burgess, N.D., Wikramanayake, E., Hahn, N., Palminteri, S., Hedao, P., Noss, R., Hansen, M., Locke, H., Ellis, E.C., Jones, B., Barber, C.V., Hayes, R., Kormos, C., Martin, V., Crist, E., Sechrest, W., Price, L., Baillie, J.E.M., Weeden, D., Suckling, K., Davis, C., Sizer, N., Moore, R., Thau, D., Birch, T., Potapov, P., Turubanova, S., Tyukavina, A., De Souza, N., Pinte, L., Brito, J.C., Llewellyn, O.A., Miller, A.G., Patzelt, A., Ghazanfar, S.A., Timberlake, J., Klöser, H., Shennan-Farpon, Y., Kindt, R., Lillesø, J.P.B., Van Breugel, P., Graudal, L., Voge, M., Al-Shammari, K.F., Saleem, M., 2017. An ecoregion-based approach to protecting half the terrestrial realm. *Bioscience* 67, 534–545. <https://doi.org/10.1093/biosci/bix014>.
- Eisenhauer, N., Bonn, A., Guerra, C., 2019. Recognizing the quiet extinction of invertebrates. *Nature Communications* 10, 50. <https://doi.org/10.1038/s41467-018-07916-1>.
- Gill, D.A., Mascia, M.B., Ahmadi, G.N., Glew, L., Lester, S.E., Barnes, M., Craigie, I., Darling, E.S., Free, C.M., Geldmann, J., Holst, S., Jensen, O.P., White, A.T., Basurto, X., Coad, L., Gates, R.D., Guannel, G., Mumby, P.J., Thomas, H., Whitmee, S., Woodley, S., Fox, H.E., 2017. Capacity shortfalls hinder the performance of marine protected areas globally. *Nature* 543, 665–669. <https://doi.org/10.1038/nature21708>.
- Giling, D.P., Beaumelle, L., Phillips, H.R.P., Cesarz, S., Eisenhauer, N., Ferlian, O., Gottschall, F., Guerra, C., Hines, J., Sendek, A., Siebert, J., Thakur, M.P., Barnes, A. D., 2018. A niche for ecosystem multifunctionality in global change research. *Global Change Biology* 1–12. <https://doi.org/10.1111/gcb.14528>.
- Hayes, K.R., Dambacher, J.M., Hosack, G.R., Bax, N.J., Dunstan, P.K., Fulton, E.A., Thompson, P.A., Hartog, J.R., Hobday, A.J., Bradford, R., Foster, S.D., Hedge, P., Smith, D.C., Marshall, C.J., 2015. Identifying indicators and essential variables for marine ecosystems. *Ecol. Indic.* 57, 409–419. <https://doi.org/10.1016/j.ecolind.2015.05.006>.
- Johnson, C.N., Balmford, A., Brook, B.W., Buettel, J.C., Galetti, M., Guangchun, L., Wilmshurst, J.M., 2017. Biodiversity losses and conservation responses in the Anthropocene. *Science* 356. <https://doi.org/10.1126/science.aam9317>.
- Kissling, W.D., Ahumada, J.A., Bowser, A., Fernandez, M., Fernández, N., García, E.A., Guralnick, R.P., Isaac, N.J.B., Kelling, S., Los, W., Mcrae, L., Mihoub, J.B., Obst, M., Santamaria, M., Skidmore, A.K., Williams, K.J., Agosti, D., Amariles, D., Arvanitidis, C., Bastin, L., De Leo, F., Egloff, W., Elith, J., Hobern, D., Martin, D., Pereira, H.M., Pesole, G., Peterseil, J., Saarenmaa, H., Schigel, D., Schmeller, D.S., Segata, N., Turak, E., Uhlir, P.F., Wee, B., Hardisty, A.R., 2018. Building essential biodiversity variables (EBVs) of species distribution and abundance at a global scale. *Biol. Rev.* 93, 600–625. <https://doi.org/10.1111/brv.12359>.
- Legg, C.J., Nagy, L., 2006. Why most conservation monitoring is, but need not be, a waste of time. *J. Environ. Manag.* 78, 194–199. <https://doi.org/10.1016/j.jenvman.2005.04.016>.
- Lindenmayer, D.B., Likens, G.E., 2010. Direct measurement versus surrogate indicator species for evaluating environmental change and biodiversity loss. *Ecosystems* 14, 47–59. <https://doi.org/10.1007/s10021-010-9394-6>.
- Lindenmayer, D.B., Likens, G.E., Franklin, J.F., 2018. Earth observation networks (EONs): finding the right balance. *Trends Ecol. Evol.* 33, 1–3. <https://doi.org/10.1016/j.tree.2017.10.008>.
- Liu, J., Mooney, H., Hull, V., Davis, S.J., Gaskell, J., Hertel, T., Lubchenco, J., Seto, K.C., Gleick, P., Kremen, C., Li, S., 2015. Systems integration for global sustainability. *Science* 347, 1258832. <https://doi.org/10.1126/science.1258832>.
- Maron, M., Mitchell, M.G.E., Runtz, R.K., Rhodes, J.R., Mace, G.M., Keith, D.A., Watson, J.E.M., 2017. Towards a threat assessment framework for ecosystem services. *Trends Ecol. Evol.* 32, 240–248. <https://doi.org/10.1016/j.tree.2016.12.011>.
- Martin, L.J., Blossey, B., Ellis, E., 2012. Mapping where ecologists work: biases in the global distribution of terrestrial ecological observations. *Front. Ecol. Environ.* 10, 195–201. <https://doi.org/10.1890/110154>.
- Masó, J., Serrall, I., Domingo-marimón, C., Zabala, A., Masó, J., Serrall, I., Domingo-marimón, C., Zabala, A., 2019. Earth observations for sustainable development goals monitoring based on essential variables and driver-pressure-state-impact-response indicators. *Int. J. Digit. Earth* 0, 1–19. <https://doi.org/10.1080/17538947.2019.1576787>.
- Metz, D., Klassen, S., McMillan, B., Clough, M., Olson, J., 2007. Building a foundation for the use of historical narratives. *Sci. Educ.* 16, 313–334. <https://doi.org/10.1007/s11191-006-9024-z>.
- Moon, K., Guerrero, A.M., Adams, V.M., Biggs, D., Blackman, D.A., Craven, L., Dickinson, H., Ross, H., 2019. Mental models for conservation research and practice. *Conserv. Lett.* 1–11. <https://doi.org/10.1111/conl.12642>.
- Naughton-Treves, L., Holland, M.B., Brandon, K., 2005. The role of protected areas in conserving biodiversity and sustaining local livelihoods. *Annu. Rev. Environ. Resour.* 30, 219–252. <https://doi.org/10.1146/annurev.energy.30.050504.164507>.
- Navarro, L.M., Fernández, N., Guerra, C., Guralnick, R., Kissling, W.D., Londoño, M.C., Karger, F.M., Turak, E., Balvanera, P., Costello, M.J., Delavaud, A., Serafy, G. El, Ferrier, S., Geijzendorffer, I., Geller, G.N., Jetz, W., Kim, E.-S., Kim, H., Martin, C.S., McGeoch, M.A., Mwampamba, T.H., Nel, J.L., Nicholson, E., Pettorelli, N., Schaeppman, M.E., Skidmore, A., Pinto, I.S., Vergara, S., Vihervaara, P., Xu, H., Yahara, T., Gill, M., Pereira, H.M., 2018. Monitoring biodiversity change through effective global coordination. *Curr. Opin. Environ. Sustain.* <https://doi.org/10.1016/j.cosust.2018.02.005>.
- Nichols, J.D., Williams, B.K., 2006. Monitoring for conservation. *Trends Ecol. Evol.* 21, 668–673. <https://doi.org/10.1016/j.tree.2006.08.007>.
- Ostrom, E., 2009. A general framework for analyzing sustainability of social-ecological systems. *Science* 325, 419–422. [https://doi.org/10.1126/S0169-1317\(03\)00213-8](https://doi.org/10.1126/S0169-1317(03)00213-8).
- Pecl, G.T., Araújo, M.B., Bell, J.D., Blanchard, J., Bonebrake, T.C., Chen, I.-C., Clark, T.D., Colwell, R.K., Danielsen, F., Evengård, B., Falconi, L., Ferrier, S., Frusher, S., García, R.A., Griffith, R.B., Hobday, A.J., Janion-Scheepers, C., Jarzyna, M.A., Jennings, S., Lenoir, J., Linnetved, H.I., Martin, V.Y., McCormack, P.C., McDonald, J., Mitchell, N.J., Mustonen, T., Pandolfi, J.M., Pettorelli, N., Popova, E., Robinson, S.A., Scheffers, B.R., Shaw, J.D., Sorte, C.J.B., Strugnelli, J.M., Sunday, J.M., Tuanmu, M.-N., Vergés, A., Villanueva, C., Wernberg, T., Wapstra, E., Williams, S.E., 2017. Biodiversity redistribution under climate change: impacts on ecosystems and human well-being. *Science* (New York, N.Y.) 355. <https://doi.org/10.1126/science.aai9214>.
- Pereira, H.M., Belnap, J., Böhm, M., Brummitt, N., García-Moreno, J., Gregory, R., Martin, L., Peng, C., Proença, V., Schmeller, D., van Swaay, C., 2017. Monitoring essential biodiversity variables at the species level. In: Walters, M., Scholes, R.J. (Eds.), *The GEO Handbook on Biodiversity Observation Networks*. Springer International Publishing, Cham, pp. 79–105. https://doi.org/10.1007/978-3-319-27288-7_4.
- Pereira, H.M., Ferrier, S., Walters, M., Geller, G.N., Jongman, R.H.G., Scholes, R.J., Bruford, M.W., Brummitt, N., Butchart, S.H.M., Cardoso, A.C., Coops, N.C., Dulloo, E., Faith, D.P., Freyhof, J., Gregory, R.D., Heip, C., Höft, R., Hurr, G., Jetz, W., Karp, D.S., McGeoch, M.A., Obura, D., Onoda, Y., Pettorelli, N., Reyers, B., Sayre, R., Scharlemann, J.P.W., Stuart, S.N., Turak, E., Walpole, M., Wegmann, M., 2013. Essential biodiversity variables. *Science* 339, 277–278. <https://doi.org/10.1126/science.1229931>.
- Rands, M.R.W., Adams, W.M., Bunnell, L., Butchart, S.H.M., Clements, A., Coomes, D., Entwistle, A., Hodge, I., Kapos, V., Scharlemann, J.P.W., Sutherland, W.J., Vira, B., 2010. Biodiversity conservation: challenges beyond 2010. *Science* 329, 1298–1303. <https://doi.org/10.1126/science.1189138>.
- Schmeller, D.S., Weatherdon, L.V., Loyau, A., Bondeau, A., Brotons, L., Brummitt, N., Geijzendorffer, I.R., Haase, P., Kuemmerlen, M., Martin, C.S., Mihoub, J.-B., Rocchini, D., Saarenmaa, H., Stoll, S., Regan, E.C., 2017. A suite of essential biodiversity variables for detecting critical biodiversity change. *Biol. Rev.* <https://doi.org/10.1111/brv.12332>.

- Spruijt, P., Knol, A.B., Vasileiadou, E., Devilee, J., Lebrecht, E., Petersen, A.C., 2014. Roles of scientists as policy advisers on complex issues: a literature review. *Environ. Sci. Policy* 40, 16–25. <https://doi.org/10.1016/j.envsci.2014.03.002>.
- Turak, E., Brazill-Boast, J., Cooney, T., Drielsma, M., Delacruz, J., Dunkerley, G., Fernandez, M., Ferrier, S., Gill, M., Jones, H., Koen, T., Leys, J., McGeoch, M., Mihoub, J.B., Scanes, P., Schmeller, D., Williams, K., 2016a. Using the essential biodiversity variables framework to measure biodiversity change at national scale. *Biol. Conserv.* <https://doi.org/10.1016/j.biocon.2016.08.019>.
- Turak, E., Harrison, I., Dudgeon, D., Abell, R., Bush, A., Darwall, W., Finlayson, C.M., Ferrier, S., Freyhof, J., Hermoso, V., Juffe-Bignoli, D., Linke, S., Nel, J., Patricio, H.C., Pittock, J., Raghavan, R., Revenga, C., Simaika, J.P., De Wever, A., 2016b. Essential Biodiversity Variables for measuring change in global freshwater biodiversity. *Biol. Conserv.* <https://doi.org/10.1016/j.biocon.2016.09.005>.
- Vihervaara, P., Auvinen, A.-P., Mononen, L., Törmä, M., Ahlroth, P., Anttila, S., Böttcher, K., Forsius, M., Heino, J., Heliölä, J., Koskelainen, M., Kuussaari, M., Meissner, K., Ojala, O., Tuominen, S., Viitasalo, M., Virkkala, R., 2017. How Essential Biodiversity Variables and remote sensing can help national biodiversity monitoring. *Glob. Ecol. Conserv.* 10, 43–59. <https://doi.org/10.1016/j.gecco.2017.01.007>.