

The minimum degree of minimal Ramsey graphs for cliques

Bamberg, John; Bishnoi, Anurag; Lesgourgues, Thomas

DOI

[10.1112/blms.12658](https://doi.org/10.1112/blms.12658)

Publication date

2022

Document Version

Final published version

Published in

Bulletin of the London Mathematical Society

Citation (APA)

Bamberg, J., Bishnoi, A., & Lesgourgues, T. (2022). The minimum degree of minimal Ramsey graphs for cliques. *Bulletin of the London Mathematical Society*, 54(5), 1827-1838. <https://doi.org/10.1112/blms.12658>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

RESEARCH ARTICLE

The minimum degree of minimal Ramsey graphs for cliques

John Bamberg¹ | Anurag Bishnoi² | Thomas Lesgourgues³

¹Department of Mathematics and Statistics, The University of Western Australia, Crawley, Australia

²Delft Institute of Applied Mathematics, TU Delft, Delft, The Netherlands

³School of Mathematics and Statistics, UNSW Sydney, Sydney, Australia

Correspondence

Anurag Bishnoi, Delft Institute of Applied Mathematics, TU Delft, Building 36, Mekelweg 4, CD Delft 2628, Netherlands.
Email: anurag.2357@gmail.com

Funding information

Australian Research Council, Grant/Award Number: DE190100666; Australian Government Research Training Program Scholarship; UNSW

Abstract

We prove that $s_r(K_k) = O(k^5 r^{5/2})$, where $s_r(K_k)$ is the Ramsey parameter introduced by Burr, Erdős and Lovász in 1976, which is defined as the smallest minimum degree of a graph G such that any r -colouring of the edges of G contains a monochromatic K_k , whereas no proper subgraph of G has this property. The construction used in our proof relies on a group theoretic model of generalised quadrangles introduced by Kantor in 1980.

MSC (2020)

05C55, 05D10, 51E12 (primary)

1 | INTRODUCTION

A graph G is called r -Ramsey for another graph H , denoted by $G \rightarrow (H)_r$, if every r -colouring of the edges of G contains a monochromatic copy of H . Observe that if $G \rightarrow (H)_r$, then every graph containing G as a subgraph is also r -Ramsey for H . Some very interesting questions arise when we study graphs G which are minimal with respect to $G \rightarrow (H)_r$, that is, $G \rightarrow (H)_r$ but there is no proper subgraph G' of G such that $G' \rightarrow (H)_r$. We call such graphs r -Ramsey minimal for H and we denote the set of all r -Ramsey minimal graphs for H by $\mathcal{M}_r(H)$. The classical result of Ramsey [21] implies that for any finite graph H and positive integer r , there exists a graph G that is r -Ramsey for H , that is, $\mathcal{M}_r(H)$ is non-empty.

Some of the central problems in graph Ramsey theory are concerned with the case where H is a clique K_k . For example, the most well-studied parameter is the Ramsey number $R_r(k)$, that denotes the smallest number of vertices of any graph in $\mathcal{M}_r(K_k)$. The classical work of Erdős

[8] and Erdős and Szekeres [9] shows that $2^{k/2} \leq R_2(k) \leq 2^{2k}$. While these bounds have been improved since then, most recently by Sah [23] (also see [24] and [5]), the constants in the exponent have stayed the same. We refer the reader to the survey of Conlon, Fox and Sudakov [6] for more on this and other graph Ramsey problems.

Several other questions on $\mathcal{M}_r(H)$ have also been explored; for example, the well-studied size-Ramsey number $\hat{R}_r(H)$ which is the minimum number of edges of a graph in $\mathcal{M}_r(H)$. We refer the reader to [1, 3, 18, 22] for various results on minimal Ramsey problems. In this paper, we will be interested in the *smallest minimum degree of an r -Ramsey minimal graph*, which is defined by

$$s_r(H) := \min_{G \in \mathcal{M}_r(H)} \delta(G),$$

for a finite graph H and positive integer r , where $\delta(G)$ denotes the minimum degree of G . Trivially, we have $s_r(H) \leq R_r(H) - 1$, since the complete graph on $R_r(H)$ vertices is r -Ramsey for H and has minimum degree $R_r(H) - 1$. The study of this parameter was initiated by Burr, Erdős and Lovász [2] in 1976. They were able to show the rather surprising exact result, $s_2(K_k) = (k - 1)^2$, which is far away from the trivial exponential bound of $s_2(K_k) \leq R_r(k) - 1$. The behaviour of this function is still not so well understood for $r > 2$ colours. Fox et al. [10] determined this function asymptotically for every fixed k up to a polylogarithmic factor, and for $k = 3$ their result was further improved by Guo and Warnke [12] who managed to obtain matching logarithmic factors.

Theorem 1.1 (Fox, Grinshpun, Liebenau, Person, Szabó).

(i) *There exist constants $c, C > 0$ such that for all $r \geq 2$, we have*

$$cr^2 \ln r \leq s_r(K_3) \leq Cr^2 \ln^2 r.$$

(ii) *For all $k \geq 4$, there exist constants $c_k, C_k > 0$ such that for all $r \geq 3$, we have*

$$c_k r^2 \frac{\ln r}{\ln \ln r} \leq s_r(K_k) \leq C_k r^2 (\ln r)^{8(k-1)^2}.$$

Theorem 1.2 (Guo, Warnke).

$$s_r(K_3) = \Theta(r^2 \ln r).$$

The constant in the upper bound of Theorem 1.1(ii) is rather large ($C_k \sim k^2 e^{4k^2 \ln 2}$), and in particular not polynomial in k . To remedy this, they proved the following general upper bound which is polynomial in both k and r .

Theorem 1.3 (Fox, Grinshpun, Liebenau, Person, Szabó). *For all $k, r \geq 3$,*

$$s_r(K_k) \leq 8(k - 1)^6 r^3.$$

For a fixed r and $k \rightarrow \infty$, Hàn, Rödl and Szabó [13] determined this function up-to polylogarithmic factors by proving the following.

Theorem 1.4 (Hàn, Rödl, Szabó). *There exists a constant k_0 such that for every $k > k_0$ and $r < k^2$*

$$s_r(K_k) \leq 80^3 (r \ln r)^3 (k \ln k)^2.$$

We prove the following general upper bound that improves Theorem 1.3, and thus provides the best known upper bound on $s_r(K_k)$ outside the special ranges covered by Theorems 1.1 and 1.4.

Theorem 1.5. *There exists an absolute constant C such that for all $r \geq 2, k \geq 3$,*

$$s_r(K_k) \leq C(k-1)^5 r^{5/2}.$$

Our proof uses the equivalence between $s_r(K_k)$ and another extremal function, called the *r -colour k -clique packing number* [10], defined as follows. Let $P_r(k)$ denote the minimum n for which there exist K_{k+1} -free pairwise edge disjoint graphs G_1, \dots, G_r on a common vertex set V of size n such that for any r -colouring of V , there exists an i such that G_i contains a copy of K_k all of whose vertices are coloured in the i^{th} colour.

Lemma 1.6 (see [10, Theorem 1.5]). *For all integers $r, k \geq 2$ we have $s_r(K_{k+1}) = P_r(k)$.*

Our graphs G_i in the packing would come from certain point-line geometries known as generalised quadrangles that we define in the next section. In Section 3, we show that any packing of ‘triangle-free’ point-line geometries implies an upper bound on $P_r(k)$, assuming certain conditions on the parameters of the geometry. In Section 4 we give a packing of certain subgeometries of the so-called Hermitian generalised quadrangles using a group theoretic model given by Kantor in the 1980s [15], and deduce that this packing implies our main result.

2 | BACKGROUND

A (finite) generalised quadrangle \mathcal{Q} of order (s, t) is an incidence structure of points \mathcal{P} , lines \mathcal{L} , together with a symmetric point-line incidence relation satisfying the following axioms.

- (i) Each point lies on $t + 1$ lines ($t \geq 1$) and two distinct points are incident with at most one line.
- (ii) Each line lies on $s + 1$ points ($s \geq 1$) and two distinct lines are incident with at most one point.
- (iii) If P is a point and ℓ is a line not incident with P , then there is a unique point on ℓ collinear with P .

Note that the third axiom above ensures that there are no triangles (that is, three distinct lines pairwise meeting in three distinct points) in \mathcal{Q} . The standard reference on finite generalised quadrangles is the book by Payne and Thas [20]. The *collinearity graph* of a generalised quadrangle is the graph on the set of points with two points adjacent when they are both incident with a common line. A collineation θ of \mathcal{Q} , that is, an automorphism of its collinearity graph, is an *elation* about the point P if it is either the identity collineation, or it fixes each line incident with P and fixes no point not collinear with P . If there is a group E of elations of \mathcal{Q} about the point P such that E acts regularly on the points not collinear with P , then we say that \mathcal{Q} is an *elation generalised quadrangle* with elation group E and *base point* P . Necessarily, E has order $s^2 t$, as there are $s^2 t$ points not collinear to a given point in any generalised quadrangle.

Now suppose we have a finite group E of order $s^2 t$ where $s, t > 1$. A *Kantor family* of E is a set $\mathcal{A} := \{A_i : i = 0, \dots, t\}$ of subgroups of order s , and a set $\mathcal{A}^* := \{A_i^* : i = 0, \dots, t\}$ of subgroups of order st , such that the following are satisfied.

TABLE 1 The points and lines of the elation generalised quadrangle arising from a Kantor family (n.b., $A_i \in \mathcal{A}, A_i^* \in \mathcal{A}^*, g \in E$)

Points	Lines
elements g of E	the right cosets $A_i g$
right cosets $A_i^* g$	symbols $[A_i]$
a symbol ∞ .	
Incidence:	$g \sim A_i g$ $A_i^* h \sim A_i g$, where $A_i g \subseteq A_i^* h$ $A_i^* h \sim [A_i]$ $\infty \sim [A_i]$

(K0) $A_i \leq A_i^*$ for all $i \in \{0, \dots, t\}$.

(K1) $A_i \cap A_j^* = \{1\}$ whenever $i \neq j$.

(K2) $A_i A_j \cap A_k = \{1\}$ whenever i, j, k are distinct.

Due to a theorem of Kantor (cf. [15, Theorem A.3.1]), a Kantor family as described above, gives rise to an *elation* generalised quadrangle of order (s, t) , which we briefly describe in Table 1.

We will simply be needing to use the Kantor family for a well-known family of generalised quadrangles, where the Heisenberg groups appear as the group E in the description above. We remark that the main property we will need is (K2), since it ensures that lines of the form $A_i g$ never form a triangle. For self-containment, we give a proof here. Suppose f, g, h are three elements of E forming the vertices of a triangle. Then there are three elements $A, B, C \in \mathcal{A}$ such that $Af = Ag$, $Bg = Bh$, $Ch = Cf$. Therefore, $fg^{-1} \in A$, $gh^{-1} \in B$, $fh^{-1} \in C$, from which it follows that $fh^{-1} = (fg^{-1})(gh^{-1}) \in AB \cap C$. Since $f \neq h$, we have $AB \cap C \neq \{1\}$. So the condition $AB \cap C = \{1\}$ given by (K2) ensures that there are no triangles.

In this paper, we will also need two commonly used maps on finite fields: the *trace* and *norm* maps. For a finite field, they are the analogues of ‘real part’ and ‘square-modulus’ for the complex numbers. We will not be needing the general theory of traces and norms, just two functions in particular. The *relative trace map* from \mathbb{F}_{q^2} to \mathbb{F}_q is defined by $\text{Tr}(x) = x + x^q$. If ϕ is a field-automorphism of \mathbb{F}_{q^2} , then $\text{Tr}(\phi(x)) = \text{Tr}(x)$ for all $x \in \mathbb{F}_{q^2}$. Also, Tr is additive; that is, $\text{Tr}(x + y) = \text{Tr}(x) + \text{Tr}(y)$ for \mathbb{F}_{q^2} . The *relative norm map* \mathbb{F}_{q^2} to \mathbb{F}_q is defined by $\text{N}(x) = x^{q+1}$. It is also invariant under field-automorphisms, and is surjective. We will make use of the following property: if q is odd, then every element of \mathbb{F}_q is a square in \mathbb{F}_{q^2} . To see this, let $t \in \mathbb{F}_q$. Since N is surjective, there exists $x \in \mathbb{F}_{q^2}$ such that $t = x^{q+1}$. Since q is odd, the element $y = x^{(q+1)/2}$ is well defined, and then $t = y^2$.

3 | PACKING GENERALISED QUADRANGLES

A partial linear space is a point-line incidence structure with the property that any two distinct points are incident to at most one common line. A triangle-free partial linear space of order (s, t) is an incidence structure satisfying Axioms (i) and (ii) of a generalised quadrangle, and (iii) there are no three distinct lines pairwise meeting each other in three distinct points. Clearly, any sub-geometry of a generalised quadrangle where the number of points on a line and the number of

lines through a point are constants is a triangle-free partial linear space. We now prove the main lemma that will imply Theorem 1.5 once we have the construction outlined in Section 4. Our proof follows the same idea as in Dudek and Rödl [7], and Fox et al. [10].

Lemma 3.1. *Let r, k, s, t be positive integers. Say there exists a family $(\mathcal{I}_i)_{i=1}^r$ of triangle-free partial linear spaces of order (s, t) , on the same point set \mathcal{P} and pairwise disjoint line-sets $\mathcal{L}_1, \dots, \mathcal{L}_r$, such that the point-line geometry $(\mathcal{P}, \bigcup_{i=1}^r \mathcal{L}_i)$ is also a partial linear space. If $s \geq 3rk \ln k$ and $t \geq 3k(1 + \ln r)$, then $P_r(k) \leq |\mathcal{P}|$.*

Proof. In order to show that $P_r(k) \leq |\mathcal{P}|$, we will exhibit K_{k+1} -free pairwise edge disjoint graphs G_1, \dots, G_r on the common vertex set $V = \mathcal{P}$, such that for any r -colouring of V , there exists an i such that G_i contains a copy of K_k all of whose vertices are coloured in the i th colour. We start by recalling the following properties about each partial linear space $\mathcal{I}_i, i \in \{1, \dots, r\}$.

(P1) Every point $p \in \mathcal{P}$ is incident with $t + 1$ lines of \mathcal{L}_i .

(P2) Every line $\ell \in \mathcal{L}_i$ contains $s + 1$ points from \mathcal{P} .

(P3) Any two points of \mathcal{P} lie on at most one line of \mathcal{L}_i .

(P4) \mathcal{I}_i is triangle-free.

Furthermore, given that $(\mathcal{P}, \bigcup_{i=1}^r \mathcal{L}_i)$ is a partial linear space and the line-sets $\mathcal{L}_1, \dots, \mathcal{L}_r$ are disjoint,

(P5) For any $i \neq j$, and any $\ell \in \mathcal{L}_i, m \in \mathcal{L}_j, \ell$ and m are incident with at most one common point.

Let $i \in \{1, \dots, r\}$, $\ell_1 = \lfloor \frac{s+1}{k} \rfloor$ and $\ell_2 = \lceil \frac{s+1}{k} \rceil$. For each line $\ell \in \mathcal{L}_i$, we select uniformly at random one partition of ℓ among all $\ell = \bigcup_{j=1}^k L_j^{(\ell)}$, where $L_j^{(\ell)}$ denotes the j th component of the partition, such that for some k' , $|L_1^{(\ell)}|, \dots, |L_{k'}^{(\ell)}| = \ell_1$ and $|L_{k'+1}^{(\ell)}|, \dots, |L_k^{(\ell)}| = \ell_2$. Choices for distinct lines in \mathcal{L}_i are independent.

We define a graph $G_i = (V, E_i)$ on the vertex set $V = \mathcal{P}$ as follows. For every $\ell \in \mathcal{L}_i$, we include the edges of a complete k -partite graph between the vertex sets $L_j^{(\ell)}$ for $j \in \{1, \dots, k\}$. Note that the graph G_i is a collection of Turán graphs on $(s + 1)$ vertices with k parts. Each Turán graph comes from one line $\ell \in \mathcal{L}_i$. By property (P3), any two points are incident with at most one line, therefore the different Turán graphs are edge-disjoint. Furthermore, by property (P4), G_i is K_{k+1} -free. Finally, by property (P5), for any $i \neq j \in \{1, \dots, r\}$, G_i and G_j are edge disjoint.

In order to conclude, we need to show that with positive probability, for any r -colouring of V , there exists an i such that G_i contains a copy of K_k all of whose vertices are coloured in the i th colour. Note that given G_1, \dots, G_r on the vertex set $V = \mathcal{P}$, in any r -colouring of V , at least one of the colours occurs at least $|\mathcal{P}|/r$ times. Therefore if for every G_i , every set of at least $|\mathcal{P}|/r$ vertices contains a copy of K_k , then we get the desired property. The choices of partitions being done independently, to conclude our proof it suffices to show that for each $i \in \{1, \dots, r\}$, with positive probability every set of at least $|\mathcal{P}|/r$ vertices contains a copy of K_k in G_i .

Fix $i \in \{1, \dots, r\}$. For a subset $W \subseteq \mathcal{P}$, let $\mathcal{A}(W)$ denote the event that the induced subgraph $G_i[W]$ contains no copy of K_k . Let $U \subset \mathcal{P}$ with $|U| = \lfloor \frac{|\mathcal{P}|}{r} \rfloor$. By property (P4), any copy of K_k can only appear from one line $\ell \in \mathcal{L}_i$, that is,

$$\mathcal{A}(U) \subseteq \bigcap_{\ell \in \mathcal{L}_i} \mathcal{A}(U \cap \ell).$$

All the events $\mathcal{A}(U \cap \ell)$ are independent, therefore

$$\mathbb{P}(\mathcal{A}(U)) \leq \prod_{\ell \in \mathcal{L}_i} \mathbb{P}(\mathcal{A}(U \cap \ell)).$$

For a given line $\ell \in \mathcal{L}_i$, let $u_\ell = |U \cap \ell|$, and let $\ell = \bigcup_{j=1}^k L_j^{(\ell)}$ be the random partition of ℓ . Note that $U \cap \ell$ contains no copy of K_k if and only if there exists $j \in \{1, \dots, k\}$ such that $U \cap L_j^{(\ell)} = \emptyset$. For a fixed $j \in \{1, \dots, k\}$,

$$\mathbb{P}(U \cap L_j^{(\ell)} = \emptyset) = \frac{\binom{s+1-u_\ell}{|L_j^{(\ell)}|}}{\binom{s+1}{|L_j^{(\ell)}|}} \leq \left(1 - \frac{u_\ell}{s+1}\right)^{|L_j^{(\ell)}|} \leq \exp\left(-\frac{\ell_1 u_\ell}{s+1}\right).$$

Therefore

$$\begin{aligned} \mathbb{P}(\mathcal{A}(U)) &\leq \prod_{\ell \in \mathcal{L}_i} \mathbb{P}(\exists j \in \{1, \dots, k\}, U \cap L_j^{(\ell)} = \emptyset) \\ &\leq k^{|\mathcal{L}_i|} \exp\left(-\sum_{\ell \in \mathcal{L}_i} \frac{\ell_1 u_\ell}{s+1}\right). \end{aligned}$$

Because every point of U is incident with $t+1$ lines from \mathcal{L}_i (by property (P1)), $\sum_{\ell \in \mathcal{L}_i} u_\ell = \sum_{\ell \in \mathcal{L}_i} |U \cap \ell| = (t+1)|U|$, and thus

$$\mathbb{P}(\mathcal{A}(U)) \leq k^{|\mathcal{L}_i|} \exp\left(-\frac{\ell_1(t+1)|U|}{s+1}\right).$$

Finally,

$$\begin{aligned} \mathbb{P}\left(\exists U \in \binom{\mathcal{P}}{\lfloor \frac{|\mathcal{P}|}{r} \rfloor} : \mathcal{A}(U)\right) &\leq \binom{|\mathcal{P}|}{\lfloor \frac{|\mathcal{P}|}{r} \rfloor} k^{|\mathcal{L}_i|} \exp\left(-\frac{t+1}{s+1} \ell_1 \lfloor \frac{|\mathcal{P}|}{r} \rfloor\right) \\ &\leq (re)^{|\mathcal{P}|/r} k^{|\mathcal{L}_i|} \exp\left(-\frac{t+1}{s+1} \lfloor \frac{s+1}{k} \rfloor \lfloor \frac{|\mathcal{P}|}{r} \rfloor\right) \end{aligned}$$

Given that $|\mathcal{P}| \geq s \geq 3rk \ln k$, $r \geq 2$ and $k \geq 3$, we have

$$\left\lfloor \frac{s+1}{k} \right\rfloor \geq \frac{5s+1}{6k}, \quad \left\lfloor \frac{|\mathcal{P}|}{r} \right\rfloor \geq \frac{5|\mathcal{P}|}{6r}.$$

Therefore,

$$\begin{aligned} \mathbb{P}\left(\exists U \in \binom{\mathcal{P}}{\lfloor \frac{|\mathcal{P}|}{r} \rfloor} : \mathcal{A}(U)\right) &\leq (re)^{|\mathcal{P}|/r} k^{|\mathcal{L}_i|} \exp\left(-\frac{t+1}{s+1} \frac{5(s+1)}{6k} \frac{5|\mathcal{P}|}{6r}\right) \\ &\leq \exp\left[|\mathcal{P}| \left(\frac{1+\ln r}{r} + \frac{|\mathcal{L}_i|}{|\mathcal{P}|} \ln k - \frac{25}{36} \frac{t+1}{rk}\right)\right]. \end{aligned}$$

By double counting (using properties (P1) and (P2)), we know that

$$|\mathcal{L}_i|(s+1) = |\mathcal{P}|(t+1),$$

and therefore

$$\mathbb{P}\left(\exists U \in \binom{\mathcal{P}}{\lfloor \frac{|\mathcal{P}|}{r} \rfloor} : \mathcal{A}(U)\right) \leq \exp\left[|\mathcal{P}|\left(\frac{1+\ln r}{r} + \frac{t+1}{s+1} \ln k - \frac{25}{36} \frac{t+1}{rk}\right)\right]. \quad (3.1)$$

Note that since $s \geq 3rk \ln k$, we have

$$\frac{25}{36} \frac{t+1}{rk} > \frac{2(t+1)}{s+1} \ln k,$$

and since $t \geq 3k(1 + \ln r)$, we have

$$\frac{25}{36} \frac{t+1}{rk} > \frac{2(1 + \ln r)}{r}.$$

Therefore,

$$\mathbb{P}\left(\exists U \in \binom{\mathcal{P}}{\lfloor \frac{|\mathcal{P}|}{r} \rfloor} : \mathcal{A}(U)\right) < 1.$$

Then there exists an instance of G_i such that every subset of \mathcal{P} with at least $\lfloor \frac{|\mathcal{P}|}{r} \rfloor$ vertices contains a copy of K_k in G_i . \square

4 | THE CONSTRUCTION

Let q be a prime power, and denote the finite field of order q^2 by \mathbb{F}_{q^2} . We will use the model of the Hermitian generalised quadrangle $H(3, q^2)$ that appears in [14, Section 3] (see Example 3). For a definition of $H(3, q^2)$ see [20, Chapter 3].

Let E be the group defined on $\mathbb{F}_{q^2} \times \mathbb{F}_q \times \mathbb{F}_{q^2}$ by the following operation:

$$(a, \gamma, b) \circ (a', \gamma', b') = (a + a', \gamma + \gamma' + \text{Tr}(b^q a'), b + b'),$$

where $\text{Tr}(x) = x + x^q$ is the relative trace map from \mathbb{F}_{q^2} to \mathbb{F}_q . It turns out that E is the *Heisenberg group* of order q^5 with centre of order q .

We can construct a generalised quadrangle by constructing a Kantor family of E . Define

$$\begin{aligned} A_\infty^* &= \{(0, \gamma, a) : a \in \mathbb{F}_{q^2}, \gamma \in \mathbb{F}_q\}, \\ A_t^* &= \{(a, \gamma, at) : a \in \mathbb{F}_{q^2}, \gamma \in \mathbb{F}_q\}, \quad t \in \mathbb{F}_q, \\ A_\infty &= \{(0, 0, a) : a \in \mathbb{F}_{q^2}\}, \\ A_t &= \{(a, a^{q+1}t, at) : a \in \mathbb{F}_{q^2}\}, \quad t \in \mathbb{F}_q. \end{aligned}$$

Then $\mathcal{A} := \{A_\infty\} \cup \{A_t : t \in \mathbb{F}_q\}$ and $\mathcal{A}^* := \{A_\infty^*\} \cup \{A_b^* : b \in \mathbb{F}_q\}$ form a Kantor family of E giving rise to a generalised quadrangle isomorphic to $H(3, q^2)$.[†]

From now on, we will assume that q is odd. Let κ be an element of \mathbb{F}_{q^2} . For each $\lambda \in \mathbb{F}_{q^2}$, define $\tau_\lambda : E \rightarrow E$ as follows. Let

$$\begin{aligned}\tau_\lambda : (a, 0, 0) &\mapsto \left(a, \text{Tr}(\lambda a + \kappa \lambda a^q + \frac{1}{2} \lambda^q a^2), \lambda a^q \right), \\ \tau_\lambda : (0, \gamma, b) &\mapsto (0, \gamma, b).\end{aligned}$$

By Axiom (K1), we have $|A_0 A_\infty^*| = |A_0| |A_\infty^*| / |A_0 \cap A_\infty^*| = s(st)/1 = |E|$. So $E = A_0 A_\infty^*$. Therefore, we can write every element $g \in E$ as $g = g_0 g_\infty^*$, with $g_0 \in A_0$ and $g_\infty^* \in A_\infty^*$. Define $\tau_\lambda(g) := \tau_\lambda(g_0) \circ \tau_\lambda(g_\infty^*)$.

Lemma 4.1. *For every $\lambda \in \mathbb{F}_{q^2}$, τ_λ is an automorphism of E .*

Proof. Let $\lambda \in \mathbb{F}_{q^2}$. It suffices to show that τ_λ is a homomorphism from A_0 to E , since τ_λ is clearly bijective. Let $a_1, a_2 \in \mathbb{F}_{q^2}$. Then

$$\tau_\lambda(a_1 + a_2, 0, 0) = \left(a_1 + a_2, \text{Tr}(\lambda(a_1 + a_2) + \kappa \lambda(a_1 + a_2)^q + \frac{1}{2} \lambda^q (a_1 + a_2)^2), \lambda(a_1 + a_2)^q \right)$$

Using $(x + y)^q = x^q + y^q$ in \mathbb{F}_{q^2} , we get,

$$\tau_\lambda(a_1 + a_2, 0, 0) = (a_1 + a_2, \text{Tr}(\lambda a_1 + \kappa \lambda a_1^q + \lambda^q a_1^2 + \lambda a_2 + \kappa \lambda a_2^q + \lambda^q a_2^2 + \lambda^q a_1 a_2), \lambda a_1^q + \lambda a_2^q).$$

Using the fact that $\text{Tr}(\cdot)$ is additive, we obtain

$$\begin{aligned}\tau_\lambda(a_1 + a_2, 0, 0) &= \left(a_1 + a_2, \text{Tr}(\lambda a_1 + \kappa \lambda a_1^q + \frac{1}{2} \lambda^q a_1^2) \right. \\ &\quad \left. + \text{Tr}(\lambda a_2 + \kappa \lambda a_2^q + \frac{1}{2} \lambda^q a_2^2) + \text{Tr}(\lambda^q a_1 a_2), \lambda a_1^q + \lambda a_2^q \right) \\ &= \left(a_1, \text{Tr}(\lambda a_1 + \kappa \lambda a_1^q + \frac{1}{2} \lambda^q a_1^2), \lambda a_1^q \right) \circ \left(a_2, \text{Tr}(\lambda a_2 + \kappa \lambda a_2^q + \frac{1}{2} \lambda^q a_2^2), \lambda a_2^q \right) \\ &= \tau_\lambda(a_1, 0, 0) \circ \tau_\lambda(a_2, 0, 0).\end{aligned}$$

Therefore, τ_λ is an automorphism of E . □

Lemma 4.2. *For every odd prime power q , there exists $\kappa \in \mathbb{F}_{q^2}$ such that $\text{Tr}(\kappa a + a^{2q-1}) \neq 0$ for all non-zero $a \in \mathbb{F}_{q^2}$.*

Proof. Suppose that $\text{Tr}(\kappa a + a^{2q-1}) = \kappa a + \kappa^q a^q + a^{2q-1} + a^{2-q} = 0$ for some non-zero $a \in \mathbb{F}_{q^2}$. Multiplying by a^{q-2} and defining $y = -a^{q-1}$ we get the following cubic equation

$$y^3 - \kappa^q y^2 + \kappa y - 1 = 0.$$

[†] In [14], the dual of this generalised quadrangle is defined, denoted by $O^-(6, q)$, but it is well known that $H(3, q^2)$ is isomorphic to the dual of the elliptic generalised quadrangle $O^-(6, q)$ (see [20, Chapter 3], where $O^-(6, q)$ is denoted by $Q(5, q)$).

Therefore if κ is such that the cubic is irreducible, then we get a contradiction. We show that such a choice of κ exists. Let t be a non-square in \mathbb{F}_q and suppose that α is an element of \mathbb{F}_{q^2} such that $\alpha^2 = t$. By the Hansen–Mullen Irreducibility Conjecture (which is true, see [4, Theorem 2.7]), there exists a monic cubic of the form $x^3 + ux^2 - tx + v$ irreducible in $\mathbb{F}_q[x]$ for some $u, v \in \mathbb{F}_q$. Let

$$\kappa = \left(\frac{-tu + 3v}{tu + v} + \frac{4t}{tu + v} \alpha \right)^q.$$

By [16, Theorem 3], $y^3 - \kappa y^2 + \kappa y - 1$ is irreducible in $\mathbb{F}_{q^2}[y]$. \square

Theorem 4.3. *Let κ be an element of \mathbb{F}_{q^2} such that $\text{Tr}(\kappa a + a^{2q-1}) \neq 0$ for all non-zero $a \in \mathbb{F}_{q^2}$. For each $\lambda \in \mathbb{F}_{q^2}$ and $t \in \mathbb{F}_q$, let*

$$A_t^\lambda = \{(a, a^{q+1}t + \text{Tr}(\lambda a + \kappa \lambda a^q + \frac{1}{2}\lambda^q a^2), at + \lambda a^q) : a \in \mathbb{F}_{q^2}\},$$

and let

$$S := \{\{A_t^\lambda : t \in \mathbb{F}_q\} : \lambda \in \mathbb{F}_{q^2}\}.$$

Then:

- (i) every element of S is a set of subgroups and any two cosets from subgroups of different such sets intersect each other in at most one element;
- (ii) if we let \mathcal{P} be the underlying set of E , then for every $\lambda \in \mathbb{F}_{q^2}$ the set of lines $\mathcal{L}_\lambda = \{A_t^\lambda : g \in E, t \in \mathbb{F}_q\}$ gives rise to a triangle-free partial linear space $(\mathcal{P}, \mathcal{L}_\lambda)$ of order $(q^2 - 1, q - 1)$.

Proof. First, for each $\lambda \in \mathbb{F}_{q^2}$ and $t \in \mathbb{F}_q$, we have

$$A_t^\lambda = \tau_\lambda(A_t).$$

Since τ_λ is an automorphism of the underlying group (as per Lemma 4.1), it follows that $\{\tau_\lambda(A_\infty)\} \cup \{\tau_\lambda(A_t) : t \in \mathbb{F}_q\}$ and $\{\tau_\lambda(A_\infty^*)\} \cup \{\tau_\lambda(A_b^*) : b \in \mathbb{F}_q\}$ also form a Kantor family, and give rise to an isomorphic generalised quadrangle. Therefore, by taking cosets of the subgroups A_t^λ , $t \in \mathbb{F}_q$, as lines we get a subgeometry of this generalised quadrangle which is a triangle-free partial linear space of order $(q^2 - 1, q - 1)$. We now prove the first part.

First we show that $A_{t_1}^{\lambda_1} \cap A_{t_2}^{\lambda_2} = \{(0, 0, 0)\}$ whenever $\lambda_1 \neq \lambda_2$. An element of $A_{t_1}^{\lambda_1} \cap A_{t_2}^{\lambda_2}$ is of the form $(a, a^{q+1}t_1 + \text{Tr}(\lambda_1 a + \kappa \lambda_1 a^q + \frac{1}{2}\lambda_1^q a^2), at_1 + \lambda_1 a^q)$ for some $a \in \mathbb{F}_{q^2}$, but it is also $(b, b^{q+1}t_2 + \text{Tr}(\lambda_2 b + \kappa \lambda_2 b^q + \frac{1}{2}\lambda_2^q b^2), bt_2 + \lambda_2 b^q)$ for some $b \in \mathbb{F}_{q^2}$. Therefore, $a = b$ and hence

$$a^{q+1}(t_1 - t_2) = \text{Tr} \left((\lambda_2 - \lambda_1)(a + \kappa a^q + \frac{1}{2}a^{2q}) \right)$$

$$a(t_1 - t_2) = (\lambda_2 - \lambda_1)a^q.$$

Now $a^{q+1}(t_1 - t_2) = a^q a(t_1 - t_2)$ and so

$$\text{Tr} \left((\lambda_2 - \lambda_1)(a + \kappa a^q + \frac{1}{2}a^{2q}) \right) = a^{2q}(\lambda_2 - \lambda_1).$$

Expanding this equation gives us

$$(\lambda_2 - \lambda_1)(a + \kappa a^q - \frac{1}{2}a^{2q}) + (\lambda_2 - \lambda_1)^q(a + \kappa a^q + \frac{1}{2}a^{2q})^q = 0.$$

Suppose, by way of contradiction, that $a \neq 0$. Since $\lambda_2 \neq \lambda_1$, we can rewrite the equation as

$$(\lambda_2 - \lambda_1)^{q-1} = -\frac{a + \kappa a^q - \frac{1}{2}a^{2q}}{(a + \kappa a^q + \frac{1}{2}a^{2q})^q}.$$

Let $N : \mathbb{F}_{q^2} \rightarrow \mathbb{F}_q$ be the relative norm function, defined by $N(x) = x^{q+1}$. The left-hand side has norm 1 and hence

$$N(-a^q + \kappa^q a - \frac{1}{2}a^2) = N(a^q + \kappa^q a + \frac{1}{2}a^2).$$

Now $N(-1) = 1$ and we can factor out $N(a)$, so

$$N(a^{q-1} + \kappa^q + \frac{1}{2}a) - N(a^{q-1} + \kappa^q - \frac{1}{2}a) = 0.$$

Note that by definition of the relative norm map,

$$\begin{aligned} N(x + y + z) &= (x + y + z)(x + y + z)^q \\ &= (x + y + z)(x^q + y^q + z^q) \\ &= xx^q + yy^q + zz^q + xy^q + yx^q + xz^q + zx^q + yz^q + zy^q \end{aligned}$$

Using that in \mathbb{F}_{q^2} , $yx^q = (y^q x)^q$, we obtain

$$N(x + y + z) = N(x) + N(y) + N(z) + \text{Tr}(xy^q + xz^q + y^q z).$$

Therefore,

$$\begin{aligned} 0 &= \left(N(a^{q-1}) + N(\kappa^q) + N(\frac{1}{2}a) + \text{Tr}(a^{q-1}\kappa + \frac{1}{2}a^{2q-1} + \frac{1}{2}\kappa a) \right) \\ &\quad - \left(N(a^{q-1}) + N(\kappa^q) + N(\frac{1}{2}a) + \text{Tr}(a^{q-1}\kappa - \frac{1}{2}a^{2q-1} - \frac{1}{2}\kappa a) \right) \\ &= \text{Tr}(\kappa a + a^{2q-1}), \end{aligned}$$

a contradiction. So $a = 0$ and $A_{t_1}^{\lambda_1} \cap A_{t_2}^{\lambda_2} = \{(0, 0, 0)\}$.

Now for any two subgroups H, K of a group, the intersection of two cosets of H and K is either empty, or a coset of $H \cap K$, which proves our claim. \square

Corollary 4.4. *There exists an absolute constant C such that for all $r \geq 2, k \geq 3$, we have $s_r(K_k) \leq C(k-1)^5 r^{5/2}$.*

Proof. Let $r \geq 2, k \geq 3, c = \frac{10+9\ln 2}{3\sqrt{2}}$ and let q be the smallest prime such that $q \geq ck\sqrt{r}$. By Lemma 4.2 and Theorem 4.3(ii), there exists a family of $r \leq q^2$ triangle-free partial linear spaces

of order $(q^2 - 1, q - 1)$, on the same point set \mathcal{P} and pairwise disjoint line-sets $\mathcal{L}_1, \dots, \mathcal{L}_r$, and by Theorem 4.3(i), the point-line geometry $(\mathcal{P}, \bigcup_{i=1}^r \mathcal{L}_i)$ is also a partial linear space. Note that $q^2 - 1 \geq 3rk \ln k$ and $q - 1 \geq 3k(1 + \ln r)$. Combining Lemmas 1.6 and 3.1, $s_r(K_{k+1}) = P_r(k) \leq |\mathcal{P}|$. By Bertrand's postulate, $q \leq 2ck\sqrt{r}$, and using $|\mathcal{P}| = q^5$ yields the desired bound, with $C = \lceil (2c)^5 \rceil = 26,282$. Note that, in light of Conjecture 5.2, we did not try to further optimise this constant. \square

5 | CONCLUDING REMARKS

While generalised quadrangles have been used extensively in extremal combinatorics, and particularly Ramsey theory (for example, [7, 11, 17, 19, 25]), our result appears to be the first instance in Ramsey theory where the group theoretic structure of these geometries is exploited. We are hopeful that Kantor's model of generalised quadrangles will lead to new results in other Ramsey problems as well.

In the probabilistic argument of Section 3, note that if we use $s + 1 = q^2$ and $t + 1 = q$, then from equation (3.1) it follows that we can solve the following quadratic inequality in q to ensure that the probability is < 1 :

$$\frac{25}{36} \frac{1}{rk} q^2 - \frac{1 + \ln r}{r} q - \ln k > 0.$$

One can check that this inequality is satisfied for all $q \geq \frac{36}{25}k(1 + \ln r) + \frac{6}{5}\sqrt{rk \ln k}$. Using that for any $a, b > 0$, $(a + b)^5 \leq 2^4(a^5 + b^5)$, we obtained the following more refined upper bound.

Theorem 5.1. *For all $r \geq 2, k \geq 2$,*

$$s_r(K_k) \leq 2^{12}[(k - 1)^5 \ln^5 r + (k - 1)^{5/2} r^{5/2} \ln^{5/2}(k - 1)].$$

For further improvements to our upper bound, we should perhaps explore triangle-free partial linear spaces that do not arise from generalised quadrangles. Moreover, if we could make the probabilistic argument of Section 3 deterministic, then this could also lead to an improvement in the bound. We would like to make the following conjecture.

Conjecture 5.2. *For all $r \geq 2, k \geq 2$*

$$s_r(K_k) \leq Ck^2r^2f(\ln k, \ln r)$$

for some constant $C > 0$ and a constant degree polynomial function f .

ACKNOWLEDGEMENTS

We are grateful to Anita Liebenau for helpful discussions. We also thank Simona Boyadzhyska for pointing out a significant gap in our first draft that was fixed later. We also want to thank the anonymous referee for a thorough proofreading and many helpful remarks.

Anurag Bishnoi's research supported by a Discovery Early Career Award of the [Australian Research Council](#) (No. DE190100666).

Thomas Lesgourgues' research supported by an Australian Government Research Training Program Scholarship and the School of Mathematics and Statistics, [UNSW](#).

JOURNAL INFORMATION

The *Bulletin of the London Mathematical Society* is wholly owned and managed by the London Mathematical Society, a not-for-profit Charity registered with the UK Charity Commission. All surplus income from its publishing programme is used to support mathematicians and mathematics research in the form of research grants, conference grants, prizes, initiatives for early career researchers and the promotion of mathematics.

REFERENCES

1. S. A. Burr, P. Erdős, R. J. Faudree, C. C. Rousseau, and R. H. Schelp, *Ramsey-minimal graphs for star-forests*, *Discrete Math.* **33** (1981), no. 3, 227–237.
2. S. A. Burr, P. Erdős, and L. Lovász, *On graphs of Ramsey type*, *Ars Combin.* **1** (1976), no. 1, 167–190.
3. S. A. Burr, J. Nešetřil, and V. Rödl, *On the use of senders in generalized Ramsey theory for graphs*, *Discrete Math.* **54** (1985), no. 1, 1–13.
4. S. D. Cohen, *Explicit theorems on generator polynomials*, *Finite Fields Appl.* **11** (2005), no. 3, 337–357.
5. D. Conlon, *A new upper bound for diagonal Ramsey numbers*, *Ann. of Math. (2)* **170** (2009), no. 2, 941–960.
6. D. Conlon, J. Fox, and B. Sudakov, *Recent developments in graph Ramsey theory*, *Surveys in Combinatorics 2015*, *London Math. Soc. Lecture Note Ser.*, vol. 424, Cambridge Univ. Press, Cambridge, 2015, pp. 49–118.
7. A. Dudek and V. Rödl, *On K_s -free subgraphs in K_{s+k} -free graphs and vertex Folkman numbers*, *Combinatorica* **31** (2011), no. 1, 39–53.
8. P. Erdős, *Some remarks on the theory of graphs*, *Bull. Amer. Math. Soc.* **53** (1947), 292–294.
9. P. Erdős and G. Szekeres, *A combinatorial problem in geometry*, *Compos. Math.* **2** (1935), 463–470.
10. J. Fox, A. Grinshpun, A. Liebenau, Y. Person, and T. Szabó, *On the minimum degree of minimal Ramsey graphs for multiple colours*, *J. Combin. Theory Ser. B* **120** (2016), 64–82.
11. Z. Füredi, A. Naor, and J. Verstraëte, *On the Turán number for the hexagon*, *Adv. Math.* **203** (2006), no. 2, 476–496 (en).
12. H. Guo and L. Warnke, *Packing nearly optimal Ramsey $R(3, t)$ graphs*, *Combinatorica* **40** (2020), no. 1, 63–103.
13. H. Hàn, V. Rödl, and T. Szabó, *Vertex Folkman numbers and the minimum degree of minimal Ramsey graphs*, *SIAM J. Discrete Math.* **32** (2018), no. 2, 826–838.
14. W. M. Kantor, *Generalized quadrangles associated with $G_2(q)^*$* , *J. Combin. Theory Ser. A* **29** (1980), 212–219.
15. W. M. Kantor, *Generalized polygons, SCABs and GABs*, *Buildings and the geometry of diagrams (Como, 1984)*, *Lecture Notes in Math.*, vol. 1181, Springer, Berlin, 1986, pp. 79–158.
16. H. D. Kim, J. Moon Kim, and I. Yie, *Certain cubic polynomials over finite fields*, *J. Korean Math. Soc.* **46** (2009), no. 1, 1–12.
17. A. Kostochka, D. Mubayi, and J. Verstraëte, *Hypergraph Ramsey numbers: triangles versus cliques*, *J. Combin. Theory Ser. A* **120** (2013), no. 7, 1491–1507.
18. T. Łuczak, *On Ramsey minimal graphs*, *Electron. J. Combin.* **1** (1994), 4.
19. D. Mubayi and J. Verstraëte, *A note on pseudorandom Ramsey graphs*, arXiv:1909.01461, 2019.
20. S. E. Payne and J. A. Thas, *Finite generalized quadrangles*, 2nd ed., *EMS Series of Lectures in Mathematics*, Eur. Math. Soc., Zürich, 2009.
21. F. P. Ramsey, *On a Problem of Formal Logic*, *Proc. Lond. Math. Soc. (2)* **30** (1929), no. 4, 264–286.
22. V. Rödl and M. Siggers, *On Ramsey minimal graphs*, *SIAM J. Discrete Math.* **22** (2008), no. 2, 467–488.
23. A. Sah, *Diagonal Ramsey via effective quasirandomness*, Preprint, arXiv:2005.09251, 2020.
24. J. Spencer, *Ramsey's theorem—a new lower bound*, *J. Combinatorial Theory Ser. A* **18** (1975), 108–115.
25. M. Tait, *Degree Ramsey numbers for even cycles*, *Discrete Mathematics* **341** (2018), no. 1, 104–108 (en).