



Delft University of Technology

Sobolev Algebras Through a 'Carré du Champ' Identity

Bernicot, Frédéric; Frey, Dorothee

DOI

[10.1017/S0013091518000214](https://doi.org/10.1017/S0013091518000214)

Publication date

2018

Document Version

Accepted author manuscript

Published in

Proceedings of the Edinburgh Mathematical Society

Citation (APA)

Bernicot, F., & Frey, D. (2018). Sobolev Algebras Through a 'Carré du Champ' Identity. *Proceedings of the Edinburgh Mathematical Society*, 61(4), 1041-1054. <https://doi.org/10.1017/S0013091518000214>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

SOBOLEV ALGEBRAS THROUGH A “CARRÉ DU CHAMP” IDENTITY

FRÉDÉRIC BERNICOT AND DOROTHEE FREY

ABSTRACT. We consider abstract Sobolev spaces of Bessel-type associated with an operator. In this work, we pursue the study of algebra properties of such functional spaces through the corresponding semigroup. As a follow-up of [4], we show that by making use of the property of a “carré du champ” identity, this algebra property holds in a wider range than previously shown.

1. INTRODUCTION

1.1. Setting. Let (M, d) be a locally compact separable metric space, equipped with a Borel measure μ , finite on compact sets and strictly positive on any non-empty open set. For Ω a measurable subset of M , we shall denote $\mu(\Omega)$ by $|\Omega|$. For all $x \in M$ and all $r > 0$, denote by $B(x, r)$ the open ball for the metric d with centre x and radius r , and by $V(x, r)$ its measure $|B(x, r)|$. For a ball B of radius r and a real $\lambda > 0$, denote by λB the ball concentric with B and with radius λr . We shall sometimes denote by $r(B)$ the radius of a ball B . We will use $u \lesssim v$ to say that there exists a constant C (independent of the important parameters) such that $u \leq Cv$, and $u \simeq v$ to say that $u \lesssim v$ and $v \lesssim u$. Moreover, for $\Omega \subset M$ a subset of finite and non-vanishing measure and $f \in L^1_{loc}(M, \mu)$, $\bar{f}_\Omega = \frac{1}{|\Omega|} \int_\Omega f d\mu$ denotes the average of f on Ω .

From now on, we assume that (M, d, μ) is a doubling metric measure space, which means that the measure μ satisfies the doubling property, that is

$$(VD) \quad V(x, 2r) \lesssim V(x, r), \quad \forall x \in M, r > 0.$$

As a consequence, there exists $\nu > 0$ such that

$$(VD_\nu) \quad V(x, r) \lesssim \left(\frac{r}{s}\right)^\nu V(x, s), \quad \forall r \geq s > 0, x \in M.$$

We then consider an unbounded operator L on $L^2(M, \mu)$ as well as an ‘abstract’ notion of gradient operator Γ under the following assumptions:

- Assumptions on L and Γ .**
- Assume that L is an injective, ω -accretive operator with dense domain $\mathcal{D} \subset L^2(M, \mu)$, where $0 \leq \omega < \pi/2$. Assume that there exists a bilinear operator Γ , with domain $\mathcal{F} \times \mathcal{F}$ for some subset \mathcal{F} of $L^2(M, \mu)$, with $\mathcal{D} \subset \mathcal{F}$.
 - For every $f \in \mathcal{F}$, we set $\Gamma(f) := |\Gamma(f, f)|^{1/2}$ and assume that Γ satisfies the inequality

$$(1.1) \quad |\Gamma(f, g)| \leq \Gamma(f)\Gamma(g), \quad \forall f, g \in \mathcal{F}.$$

Date: December 6, 2017.

FB’s research was supported by the ERC project FAnFARé no. 637510 and by the ANR project HAB no. ANR-12-BS01-0013.

Moreover, assume that

$$(R_2) \quad \|\Gamma f\|_2 \lesssim \|L^{1/2} f\|_2, \quad \forall f \in \mathcal{D}.$$

- Assume that the semigroup $(e^{-tL})_{t>0}$ admits a kernel representation with a kernel p_t satisfying the upper Gaussian pointwise estimates

$$(UE) \quad |p_t(x, y)| \lesssim \frac{1}{V(x, \sqrt{t})} \exp\left(-\frac{d^2(x, y)}{Ct}\right), \quad \forall t > 0, \text{ a.e. } x, y \in M.$$

- Assume that the semigroup $(e^{-tL})_{t>0}$ and its gradient satisfy L^2 Davies-Gaffney estimates, which means that for every $r > 0$ and all balls B_1, B_2 of radius r

$$(DG) \quad \|e^{-r^2 L}\|_{L^2(B_1) \rightarrow L^2(B_2)} + \|r\Gamma e^{-r^2 L}\|_{L^2(B_1) \rightarrow L^2(B_2)} \lesssim e^{-c \frac{d^2(B_1, B_2)}{r^2}}.$$

By our assumptions, $(e^{-tL})_{t>0}$ is bounded analytic on $L^p(M, \mu)$ for $p \in (1, \infty)$ and uniformly bounded on $L^p(M, \mu)$ for $p \in [1, \infty]$, see [5, Corollary 1.5]. Note that (DG) for the semigroup is a consequence of (UE). By analyticity of the semigroup, the property (UE), and thus also (DG), extends to the collections $((tL)^n e^{-tL})_{t>0}$ for every integer $n \geq 0$. The operator Γ is a sublinear operator, acting like the length of the gradient on a Riemannian manifold.

We also assume that Γ and L are related by a weak version of a “carré du champ” identity:

“Carré du champ” identity. Assume that Γ and L satisfy the following: for every $t > 0$ and all functions $f, g \in L^\infty(M, \mu) \cap \mathcal{D}$

$$(1.2) \quad e^{-tL} L(fg) = e^{-tL} [Lf \cdot g] + e^{-tL} [f \cdot Lg] - 2e^{-tL} \Gamma(f, g).$$

This equality can be viewed in $L^2_{\text{loc}}(M, \mu)$, since for functions f, g chosen as above, we know that $\Gamma(f, g) \in L^1(M, \mu)$ and so the LHS and RHS are both locally in $L^2(M, \mu)$ due to (UE).

Remark 1.1. • Note that the full “carré du champ” identity, which is

$$(1.3) \quad L(fg) = Lf \cdot g + f \cdot Lg - 2\Gamma(f, g),$$

is stronger than the previous assumption. It is not clear on which set of functions such an identity may be assumed.

- Let us emphasise that the proofs developed in the next sections do not really require the exact identity (1.2). It would be sufficient to only assume the following inequality: for every $t > 0$ and all functions $f, g \in L^\infty(M, \mu) \cap \mathcal{D}$

$$(1.4) \quad |e^{-tL} L(fg) - (e^{-tL} [Lf \cdot g] + e^{-tL} [f \cdot Lg])| \lesssim |e^{-tL} \Gamma(f, g)|.$$

We will assume the above throughout the paper. We abbreviate the setting with (M, μ, Γ, L) .

1.2. The algebra property. Following up on [4], we aim to prove that the (Bessel-type) Sobolev spaces satisfy an algebra property under our assumptions. Such property is very well understood in the Euclidean space and goes back to initial works by Strichartz [13], Kato and Ponce [9], and then Coifman and Meyer [6, 11] using the paraproduct decomposition. We refer the reader to [4] and references therein for a more complete review of the literature on this topic. This algebra

property and the corresponding Leibniz rule is crucial in order to study nonlinear PDEs.

In this current work, we are going to describe how the “carré du champ” property allows us to improve the main results of [4]. Note that the “carré du champ” identity combined with (1.1) encodes an algebra property at the order of regularity 1, since the operator Γ (resp. L) is implicitly an operator of order 1 (resp. 2). We will show how to deduce algebra properties of fractional order from this.

Let us first give a rigorous sense to what we mean by the Algebra property for Sobolev spaces. We follow the approach of [4]. Denote by $\mathcal{C}_0(M)$ the space of continuous functions on M which vanish at infinity, and $\mathfrak{C} := \mathcal{C}_0(M) \cap \mathcal{F}$. We define $\dot{L}_\alpha^p(M, L, \mu) \cap L^\infty(M, \mu)$ as the completion of

$$\{f \in \mathfrak{C}, L^{\alpha/2}f \in L^p(M, \mu)\}$$

with respect to the norm $\|L^{\alpha/2}f\|_p + \|f\|_\infty$. We denote in the sequel $\|L^{\alpha/2}f\|_p$ by $\|f\|_{p,\alpha}$.

Let us recall our definition of the algebra property $A(p, \alpha)$ from [4, Definition 1.1].

Definition 1.2. For $\alpha > 0$ and $p \in (1, \infty)$ we say that property $A(p, \alpha)$ holds if:

- the space $\dot{L}_\alpha^p(M, L, \mu) \cap L^\infty(M, \mu)$ is an algebra for the pointwise product;
- and the Leibniz rule inequality is valid:

$$\|fg\|_{p,\alpha} \lesssim \|f\|_{p,\alpha}\|g\|_\infty + \|f\|_\infty\|g\|_{p,\alpha}, \quad \forall f, g \in \dot{L}_\alpha^p(M, L, \mu) \cap L^\infty(M, \mu).$$

1.3. Main result. For $p \in [1, \infty]$, we say that the semigroup satisfies gradient bounds (G_p) if

$$(G_p) \quad \sup_{t>0} \|\sqrt{t}\Gamma e^{-tL}\|_{p \rightarrow p} < \infty.$$

Let us observe that by (R_2) and (UE) , it is classical that our previous assumptions already imply (G_p) for $p \in (1, 2]$.

Our main result reads as follows:

Theorem 1.3. Let (M, μ, Γ, L) as in Subsection 1.1 with a homogeneous dimension $\nu > 2$. Assume in addition (G_{p_0}) for some $p_0 \in [2, \nu)$. Then $A(p, \alpha)$ holds for every $p \in (1, p_0)$ with $\alpha \in (0, 1)$, and for every $p \in (p_0, \infty)$ with $0 < \alpha < \frac{p_0}{p}$.

The condition $p_0 < \nu$ is not relevant and not used, but for $p_0 > \nu$ the result was already obtained in [4] in a more general framework. That is why we restrict our attention here to the range $2 \leq p_0 < \nu$.

We use a slightly different decomposition of the product than in [4]. Indeed in [4], the product of two functions was decomposed into two paraproducts. Here, we decompose it into three terms (two paraproducts and a ‘resonant part’). The two paraproducts are completely uncritical, whereas the third one carries the most subtle information encoded in the resonances. The “carré du champ” identity now allows us to handle this third part in a better way. This allows to improve over [4] in the case $p > 2$.

Proof. The theorem will be proved in the following sections. The proof goes through the use of Stein’s complex interpolation between the two endpoints $(\alpha, p) = (1, p_0)$

and $(\alpha, p) = (0, \infty)$.

The case $p \in (1, p_0)$ is obtained as the combination of the paraproduct decompositions (3.1) and (3.2) with the boundedness results of Propositions 3.2, 3.7 and 3.8. The case $p \in [p_0, \infty)$ is shown by combining the paraproduct decompositions (3.1) and (3.2) with the boundedness results of Propositions 3.2 and 3.9. \square

1.4. Comparison to previous results and examples. Let us compare this result with what we have previously obtained in [4, Theorem 1.5]. First, let us mention that even if [4] was written in the setting of a Dirichlet form (which is a particular case of our current setting here), all of the results in [4] can be described in our present setting, without assuming the “carré du champ” identity, with identical proofs. The extra main property used in [4] (instead of (1.2)) is the following inequality

$$(1.5) \quad \left| \int Lf \cdot g \, d\mu \right| \lesssim \int \Gamma f \cdot \Gamma g \, d\mu$$

for all functions $f, g \in \mathcal{F}$.

Let us now compare our result with the one of [4]:

- The two approaches rely on the same framework given by a ‘gradient’ operator Γ satisfying a Leibniz rule and a semigroup $(e^{-tL})_{t>0}$. The main difference is that [4] requires (1.5), whereas here we assume (1.2) or in fact the weaker version (1.4). We first observe that in the case of a self-adjoint and conservative operator L , then by integrating (1.4) implies exactly (1.5). So our current assumption is stronger than the one used in [4] and corresponds to a pointwise version; it is therefore natural that we are able to obtain a wider range of exponents. To be more precise, for $p > p_0$ we improve the range $\alpha \in (0, 1 - \nu(\frac{1}{p_0} - \frac{1}{p}))$ (obtained in [4]) to $\alpha \in (0, \frac{p_0}{p})$.
- Moreover, we only detail the proofs of [4] and of the current work in the setting where the semigroup is supposed to satisfy (UE), which corresponds to pointwise (or L^1 - L^∞) local estimates. However, it is by now well-known that all the employed arguments can be extended to a more general framework where the semigroup is only assumed to have local L^{p_-} - L^{p_+} estimates for some $p_- < 2 < p_+$. In such a situation the condition on the exponents α, p such that $A(\alpha, p)$ can be proved will depend on p_-, p_+ . A careful examination reveals the following difference: in [4], we make appear only one Γ operator, evaluated on a product and then use a Leibniz property. In the current work, the “carré du champ” identity (1.2) makes appear the product of two Γ operators. So combining the Γ operator (on which we assume L^{p_-} - L^{p_0} local estimates through (G_{p_0})) and the local L^{p_-} - L^{p_+} estimates on the semigroup will then lead to more restrictions in the current setting than in [4]. Thus also from this point of view it is natural that we can obtain a wider range for the Sobolev algebra property, because of our stronger assumption.

As a conclusion of the comparison: our previous work [4] and this current one are both interesting in themselves and each of them brings results in its proper framework. If one can fit into the current framework, then it is better to follow the current approach, where we develop a simpler proof for the range $(1, p_0]$ and a wider range for $p > p_0$ by taking advantage of the “carré du champ” identity.

However [4] explains how we can still prove the Algebra property in a more general setting, yet with a smaller range.

Let us now describe some examples where the extra assumption in terms of “carré du champ” identity is satisfied:

- The Dirichlet form setting (as detailed in [4]) with a “carré du champ”. In such a case, the “carré du champ” operator Γ satisfies the ‘strong’ (pointwise) identity (1.3), as well as (1.1).
- In the Euclidean setting $M = \mathbb{R}^n$ (or more generally in a doubling Riemannian manifold), consider $A = A(x)$ a complex matrix-valued function with bounded measurable coefficients, satisfying the ellipticity (or accretivity) condition

$$(1.6) \quad \lambda|\xi|^2 \leq \Re\langle A(x)\xi, \xi \rangle \quad \text{and} \quad |\langle A(x)\xi, \zeta \rangle| \leq \Lambda|\xi||\zeta|,$$

for some constants $\lambda, \Lambda > 0$ and every $x \in \mathbb{R}^n$, $\xi, \zeta \in \mathbb{R}^n$.

For such a complex matrix-valued function A , we may define a second order divergence form operator

$$L = L_A f := -\operatorname{div}(A \nabla f),$$

which we first interpret in the sense of maximal accretive operators via a sesquilinear form. That is, $\mathcal{D}(L)$ is the largest subspace contained in $W^{1,2} := \mathcal{D}(\nabla)$ for which

$$\left| \int_M \langle A \nabla f, \nabla g \rangle d\mu \right| \leq C \|g\|_2 \quad \forall g \in W^{1,2},$$

and we define Lf by

$$\langle Lf, g \rangle = \int_M \langle A \nabla f, \nabla g \rangle d\mu$$

for $f \in \mathcal{D}(L)$ and $g \in W^{1,2}$. Thus defined, $L = L_A$ is a maximal-accretive operator on L^2 and $\mathcal{D}(L)$ is dense in $W^{1,2}$.

For such an operator we have the pointwise “carré du champ” identity (1.3) with the operator

$$\Gamma(f, g) := \Re\langle A \nabla f, \nabla g \rangle.$$

The ellipticity condition then implies (1.1).

- In the case of a non-selfadjoint operator L , we can also consider the following example: in the Euclidean space, associated with a rather singular function a , consider the operator $L(f) = -\Delta(af)$. It is non-selfadjoint and non conservative, but some of the arguments of [4] or those developed here can be used, if we can prove (UE) and (DG). We refer the reader to [10] (extended to a doubling setting in [7]), where it is proven that if the measurable function a has an accretive real part, then the semigroup e^{-tL^*} satisfies (UE) and by duality it is also true for e^{-tL} . Combining this with Riesz transform estimates in L^2 also gives L^2 Davies-Gaffney estimates (DG) for the operator L . For such an operator, it is interesting to observe that assumption (1.5) (used for [4]) relies on a Lipschitz condition on a although the assumption (1.4) (used here) will require a C^2 -condition on a .

2. TECHNICAL PRELIMINARIES

Let us give some notation and a few reminders about certain operators constructed from the functional calculus of L . We refer to [4] for more details. We first define *approximation operators*, which are the elementary objects to build a paraproduct associated with a semigroup.

Definition 2.1. Let $N \in \mathbb{N}$, $N > 0$, and set $c_N = \int_0^{+\infty} s^N e^{-s} \frac{ds}{s}$. For $t > 0$, define

$$(2.1) \quad Q_t^{(N)} := c_N^{-1} (tL)^N e^{-tL}$$

and

$$(2.2) \quad P_t^{(N)} := \phi_N(tL),$$

with $\phi_N(x) := c_N^{-1} \int_x^{+\infty} s^N e^{-s} \frac{ds}{s}$, $x \geq 0$.

Let us define some suitable sets of test functions. Let us recall that $\mathfrak{C} := \mathcal{C}_0(M) \cap \mathcal{F}$.

Definition 2.2. For $p \in (1, +\infty)$, we define the set of test functions

$$\mathcal{S}^p = \mathcal{S}^p(M, L) := \{f \in \mathfrak{C} \cap L^p : \exists g, h \in L^2 \cap L^p, f = Lg \text{ and } h = Lf\},$$

and

$$\mathcal{S} = \bigcup_{p \in (1, +\infty)} \mathcal{S}^p.$$

We recall from [4, Proposition 2.13] that (UE) implies square function estimates for $Q_t^{(N)}$ in L^p .

Lemma 2.3. Let $p \in (1, \infty)$, $N \in \mathbb{N}$ and $\alpha > 0$. Under (UE), one has

$$\left\| \left(\int_0^\infty |(tL)^\alpha P_t^{(N)} f|^2 \frac{dt}{t} \right)^{1/2} \right\|_p \lesssim \|f\|_p$$

for all $f \in L^p(M, \mu)$.

A direct consequence of the above is the following orthogonality lemma. See [4, Lemma 2.15] for a slightly less general version.

Lemma 2.4. Let $p \in (1, \infty)$, $N \in \mathbb{N}$ and $\alpha > 0$. Assume (UE). Then

$$\left\| \int_0^{+\infty} (tL)^\alpha P_t^{(N)} F_t \frac{dt}{t} \right\|_p \lesssim \left\| \left(\int_0^{+\infty} |F_t|^2 \frac{dt}{t} \right)^{1/2} \right\|_p,$$

where $F_t(x) := F(t, x)$, $F : (0, +\infty) \times M \rightarrow \mathbb{R}$ is a measurable function such that the RHS has a meaning and is finite.

Under the additional assumption (G_{p_0}) for some $p_0 > 2$, one also has square function estimates involving Γ .

Lemma 2.5. Let $N \in \mathbb{N}$ and $\alpha \in (0, 1)$. Assume (G_{p_0}) for some $p_0 \in (2, \infty)$. Then for every $p \in (1, p_0)$,

$$\left\| \left(\int_0^\infty |\sqrt{t}\Gamma(tL)^{-\alpha/2} P_t^{(N)} f|^2 \frac{dt}{t} \right)^{1/2} \right\|_p \lesssim \|f\|_p$$

for all $f \in L^p(M, \mu)$.

Proof. The proof of [4, Proposition 2.14] has to be adapted as follows. By writing

$$P_t^{(N)} f = \int_t^\infty Q_s^{(N)} f \frac{ds}{s},$$

one has the pointwise estimate

$$|\sqrt{t}\Gamma(tL)^{-\alpha/2} P_t^{(N)} f| \leq \int_t^\infty \left(\frac{t}{s}\right)^{\frac{1-\alpha}{2}} |\sqrt{s}\Gamma(sL)^{-\alpha/2} Q_s^{(N)} f| \frac{ds}{s}.$$

Since $\alpha \in (0, 1)$, Hardy's inequality yields

$$\left(\int_0^\infty |\sqrt{t}\Gamma(tL)^{-\alpha/2} P_t^{(N)} f|^2 \frac{dt}{t}\right)^{1/2} \lesssim \left(\int_0^\infty |\sqrt{t}\Gamma(tL)^{-\alpha/2} Q_t^{(N)} f|^2 \frac{dt}{t}\right)^{1/2}.$$

Having this pointwise inequality, one can proceed as before in [4, Proposition 2.14]. \square

3. MAIN RESULT

From now on, fix $D \in \mathbb{N}$ in the definition of $Q_t^{(D)}$ and $P_t^{(D)}$ sufficiently large ($D > 4\nu$ will suffice), and write $Q_t := Q_t^{(D)}$ and $P_t := P_t^{(D)}$.

We define paraproducts associated with the underlying operator L . Note however that the definitions differ from those in [4].

For $g \in L^\infty(M, \mu)$, we define the paraproduct Π_g on \mathcal{S} by

$$\Pi_g^{(D)}(f) = \Pi_g(f) := \int_0^\infty P_t(Q_t f \cdot P_t g) \frac{dt}{t}, \quad f \in \mathcal{S}.$$

For every $p \in (1, \infty)$ and every $f \in \mathcal{S}^p$, the integral is absolutely convergent in $L^p(M, \mu)$. We refer the reader to [4, Section 3] for the details, noting that $(P_t)_{t>0}$ is bounded uniformly in $L^p(M, \mu)$.

We define the resonant term Π on \mathcal{S} by

$$\Pi^{(D)}(f, g) = \Pi(f, g) := \int_0^\infty Q_t(P_t f \cdot P_t g) \frac{dt}{t}, \quad f, g \in \mathcal{S}.$$

We discuss the question of absolute convergence of the integral in $\Pi(f, g)$ after Proposition 3.2.

Lemma 3.1 (Product decomposition). *For every $p \in (1, \infty)$ and every $f, g \in \mathcal{S}^p$, we have the product decomposition*

$$(3.1) \quad fg = \Pi(f, g) + \Pi_g(f) + \Pi_f(g) \quad \text{in } L^p(M, \mu).$$

Proof. Since $\mathcal{S}^p \subseteq L^\infty(M, \mu)$, we have $f \cdot g, P_t f \cdot P_t g \in L^p(M, \mu)$. We recall from [4, Proposition 2.11, Lemma 3.1] that in the L^p sense, $f \cdot g = \lim_{t \rightarrow 0} P_t f \cdot P_t g$ and $0 = \lim_{t \rightarrow \infty} P_t f \cdot P_t g$, where the latter makes use of our assumption $N(L) = \{0\}$. The same arguments then also imply that

$$\begin{aligned} f \cdot g &= \lim_{t \rightarrow 0} P_t(P_t f \cdot P_t g), \\ 0 &= \lim_{t \rightarrow \infty} P_t(P_t f \cdot P_t g) \end{aligned}$$

in the L^p sense. Since P_t and Q_t are defined such that $Q_t = -t\partial_t P_t$, we obtain

$$\begin{aligned} fg &= \lim_{t \rightarrow 0} P_t(P_t f \cdot P_t g) - \lim_{t \rightarrow \infty} P_t(P_t f \cdot P_t g) = - \int_0^\infty \partial_t(P_t(P_t f \cdot P_t g)) dt \\ &= \int_0^\infty Q_t(P_t f \cdot P_t g) \frac{dt}{t} + \int_0^\infty P_t(Q_t f \cdot P_t g) \frac{dt}{t} + \int_0^\infty P_t(P_t f \cdot Q_t g) \frac{dt}{t}, \end{aligned}$$

which is the stated decomposition. \square

The critical term in the product decomposition is the resonant term $\Pi(f, g)$. We have shown already in [4, Proposition 3.3] that the paraproduct $\Pi_g(f)$ is bounded in \dot{L}_α^p for all $\alpha \in (0, 1)$, without other assumption than (UE). Let us mention that the result remains true for $\alpha \geq 1$.

Proposition 3.2. *Let $p \in (1, \infty)$, $\alpha \in (0, 1)$ and $g \in L^\infty(M, \mu)$. Then Π_g is well-defined on \mathcal{S}^p with for every $f \in \mathcal{S}^p$*

$$\|\Pi_g(f)\|_{p, \alpha} \lesssim \|f\|_{p, \alpha} \|g\|_\infty.$$

Let us now have a look at the resonant term $\Pi(f, g)$. We use the assumed “carré du champ” identity (1.2) to write, with $\tilde{Q}_t := (tL)^{-1}Q_t$,

$$\begin{aligned} \Pi(f, g) &= \int_0^\infty (tL)^{-1}Q_t tL(P_t f \cdot P_t g) \frac{dt}{t} \\ &= \int_0^\infty \tilde{Q}_t(tLP_t f \cdot P_t g) \frac{dt}{t} + \int_0^\infty \tilde{Q}_t(P_t f \cdot tLP_t g) \frac{dt}{t} \\ (3.2) \quad &\quad - 2 \int_0^\infty \tilde{Q}_t \Gamma(\sqrt{t}P_t f, \sqrt{t}P_t g) \frac{dt}{t}. \end{aligned}$$

For the first term one can use the same arguments as for $\Pi_g(f)$ to show that for $p \in (1, \infty)$, $g \in L^\infty(M, \mu)$ and $f \in \mathcal{S}^p$, the integral converges absolutely in $L^p(M, \mu)$. By interchanging the roles of f and g , the same holds true for the second term. In the third term, for every $0 < \varepsilon < R < \infty$, the finite integral \int_ε^R is well-defined. The results of Proposition 3.5 and Proposition 3.8 below in particular imply that the integral converges absolutely in $L^p(M, \mu)$.

Instead of showing the boundedness of $\Pi(f, g)$ in \dot{L}_α^p directly, we first show its boundedness in $L^q(M, \mu)$ for large $q < \infty$, and then interpolate with $\dot{L}_1^{p_0}$, where p_0 is chosen such that (G_{p_0}) holds.

With the same arguments as in the proof of Proposition 3.2, one immediately obtains the L^p boundedness of the first term in (3.2). See the proof of [4, Proposition 3.3].

Lemma 3.3. *Assume (UE). Let $p \in (1, \infty)$. Then for every $f \in L^p(M, \mu)$ and every $g \in L^\infty(M, \mu)$, we have*

$$\left\| \int_0^\infty \tilde{Q}_t(tLP_t f \cdot P_t g) \frac{dt}{t} \right\|_p \lesssim \|f\|_p \|g\|_\infty.$$

For the second term, we obviously obtain the symmetric result in f and g . But it is also possible to interchange the roles of f and g .

Lemma 3.4. *Assume (UE). Let $p \in (1, \infty)$. Then for every $g \in L^p(M, \mu)$ and every $f \in L^\infty(M, \mu)$, we have*

$$\left\| \int_0^\infty \tilde{Q}_t(tLP_t f \cdot P_t g) \frac{dt}{t} \right\|_p \lesssim \|f\|_\infty \|g\|_p.$$

A result of this kind was already proven in [8, Theorem 4.2]. For convenience of the reader we give a (different) proof here.

Proof. By Lemma 2.4 applied to $T_t = \tilde{Q}_t$ and [4, Theorem 2.17], we have for every $q \in (p, \infty)$ - with the notation as in [4] -

$$\begin{aligned} \left\| \int_0^\infty \tilde{Q}_t(tLP_t f \cdot P_t g) \frac{dt}{t} \right\|_p &\lesssim \left\| \left(\int_0^\infty |tLP_t f \cdot P_t g|^2 \frac{dt}{t} \right)^{1/2} \right\|_p \\ &\lesssim \|N_*(P_t g)\|_p \|\mathcal{C}_q(tLP_t f)\|_\infty. \end{aligned}$$

We let the reader check that a simple adaptation of [4, Lemma 4.4 (a)] yields $\|N_*(P_t g)\|_p \lesssim \|g\|_p$. Similarly, one can modify the proof of [4, Lemma 4.4 (b)] for the second estimate. To do so, note that by our assumptions,

$$\left\| \left(\int_0^\infty |tLP_t f|^2 \frac{dt}{t} \right)^{1/2} \right\|_q \lesssim \|f\|_q,$$

and that $(tLP_t)_{t>0}$ satisfies L^q off-diagonal estimates of any order. Using this, one obtains $\|\mathcal{C}_q(tLP_t f)\|_\infty \lesssim \|f\|_\infty$. \square

In order to treat the third term in (3.2), we define the operator Π_Γ on \mathcal{S} by

$$\Pi_\Gamma(f, g) := \int_0^\infty \tilde{Q}_t \Gamma(\sqrt{t}P_t f, \sqrt{t}P_t g) \frac{dt}{t}, \quad f, g \in \mathcal{S}.$$

Proposition 3.5. *Assume (UE). Let $p \in (2, \infty)$, and let $g \in L^\infty(M, \mu)$. Then $\Pi_\Gamma(\cdot, g)$ is well-defined on $L^p(M, \mu)$ with for every $f \in L^p(M, \mu)$*

$$\|\Pi_\Gamma(f, g)\|_p \lesssim \|f\|_p \|g\|_\infty.$$

Proof. We can write $\tilde{Q}_t = (tL)^{-1}Q_t^{(D)} = [c_D^{-1}(tL)^{D-1}e^{-t/2L}]e^{-t/2L} =: \tilde{\tilde{Q}}_t P_{t/2}^{(1)}$. By Lemma 2.4 with $T_t = \tilde{\tilde{Q}}_t$ in the first step, Minkowski's inequality in the second, (UE) and (1.1) in the third, and the Cauchy-Schwarz inequality in the last step,

we obtain

$$\begin{aligned}
\|\Pi_\Gamma(f, g)\|_p &\lesssim \left\| \left(\int_0^\infty \left| P_{t/2}^{(1)} \Gamma(\sqrt{t} P_t f, \sqrt{t} P_t g) \right|^2 \frac{dt}{t} \right)^{1/2} \right\|_p \\
&\lesssim \sum_{j=0}^\infty \left\| x \mapsto \left(\int_0^\infty \left| P_{t/2}^{(1)} \mathbf{1}_{S_j(B(x, \sqrt{t}))} \Gamma(\sqrt{t} P_t f, \sqrt{t} P_t g) \right|^2 \frac{dt}{t} \right)^{1/2} \right\|_p \\
&\lesssim \sum_{j=0}^\infty 2^{-2jN} 2^{j\nu} \left\| x \mapsto \left(\int_0^\infty \left(\int_{B(x, 2^j \sqrt{t})} |\sqrt{t} \Gamma P_t f| \cdot |\sqrt{t} \Gamma P_t g| d\mu \right)^2 \frac{dt}{t} \right)^{1/2} \right\|_p \\
(3.3) \quad &\lesssim \sum_{j=0}^\infty 2^{-2jN} 2^{j\nu} \left\| x \mapsto \left(\int_0^\infty \left(\int_{B(x, 2^j \sqrt{t})} |\sqrt{t} \Gamma P_t f|^2 d\mu \right) \left(\int_{B(x, 2^j \sqrt{t})} |\sqrt{t} \Gamma P_t g|^2 d\mu \right) \frac{dt}{t} \right)^{1/2} \right\|_p.
\end{aligned}$$

For all $j \geq 0$ and $x \in M$, L^2 off-diagonal estimates for $(\sqrt{t} \Gamma P_t)_{t>0}$ (see (DG)) yield

$$\begin{aligned}
\left(\int_{B(x, 2^j \sqrt{t})} |\sqrt{t} \Gamma P_t g|^2 d\mu \right)^{1/2} &\leq \sum_{k=0}^\infty \left(\int_{B(x, 2^j \sqrt{t})} |\sqrt{t} \Gamma P_t (\mathbf{1}_{S_k(B(x, 2^{j+k} \sqrt{t}))} g)|^2 d\mu \right)^{1/2} \\
&\lesssim \left(\int_{B(x, 2^j \sqrt{t})} |g|^2 d\mu \right)^{1/2} + \sum_{k=1}^\infty \left(1 + \frac{(2^{j+k} \sqrt{t})^2}{t} \right)^{-N} 2^{k\nu/2} \left(\int_{B(x, 2^{j+k} \sqrt{t})} |g|^2 d\mu \right)^{1/2} \\
(3.4) \quad &\lesssim \|g\|_\infty.
\end{aligned}$$

Using this estimate in (3.3), we get

$$\begin{aligned}
\|\Pi_\Gamma(f, g)\|_p &\lesssim \|g\|_\infty \sum_{j=0}^\infty 2^{-2jN} 2^{j\nu} \left\| x \mapsto \left(\int_0^\infty \int_{B(x, 2^j \sqrt{t})} |\sqrt{t} \Gamma P_t f|^2 d\mu \frac{dt}{t} \right)^{1/2} \right\|_p \\
&= \|g\|_\infty \sum_{j=0}^\infty 2^{-2jN} 2^{j\nu} \|\sqrt{t} \Gamma P_t f\|_{T_{2^j}^{p,2}(M)},
\end{aligned}$$

where $T_{2^j}^{p,2}(M)$ denotes the tent space with angle 2^j and appropriate elliptic scaling. By change of angle in tent spaces [2, Theorem 1.1], $\|\sqrt{t} \Gamma P_t f\|_{T_{2^j}^{p,2}(M)} \lesssim 2^{j\nu/2} \|\sqrt{t} \Gamma P_t f\|_{T^{p,2}(M)}$ for all $p \geq 2$. On the other hand, it is known from e.g. [3, Theorem 3.1] (which extends to our setting) that $\sqrt{t} \Gamma P_t$ satisfies a conical square function estimate for $p \geq 2$. Thus, we finally obtain

$$\|\Pi_\Gamma(f, g)\|_p \lesssim \|g\|_\infty \sum_{j=0}^\infty 2^{-2jN} 2^{j\nu} 2^{j\nu/2} \|\sqrt{t} \Gamma P_t f\|_{T^{p,2}(M)} \lesssim \|f\|_p \|g\|_\infty.$$

□

Putting Lemma 3.3, Lemma 3.4 and Proposition 3.5 together, we obtain

Corollary 3.6. *Assume (UE). Let $p \in (2, \infty)$, and let $g \in L^\infty(M, \mu)$. Then $\Pi(\cdot, g)$ is well-defined on $L^p(M, \mu)$ with for every $f \in L^p(M, \mu)$*

$$\|\Pi(f, g)\|_p \lesssim \|f\|_p \|g\|_\infty.$$

The above result provides us with the required result at one of the endpoints in the interpolation. Let us now have a look at the other endpoint.

One of the terms in (3.2) can be estimated in \dot{L}_α^p without further assumptions. The proof is the same as the one for Proposition 3.2.

Proposition 3.7. *Assume (UE). Let $p \in (1, \infty)$, $\alpha \in (0, 1)$ and $g \in L^\infty(M, \mu)$. Then the integral on the left-hand side is well-defined on \mathcal{S}^p with for every $f \in \mathcal{S}^p$*

$$\left\| \int_0^\infty \tilde{Q}_t(tLP_t f \cdot P_t g) \frac{dt}{t} \right\|_{p, \alpha} \lesssim \|f\|_{p, \alpha} \|g\|_\infty.$$

The result for the resonant term can be obtained similarly to the one in Proposition 3.5, but requires the additional assumption of gradient bounds on the semi-group.

Proposition 3.8. *Assume (UE) and (G_{p_0}) for some $p_0 \in [2, \infty)$. Let $p \in (1, p_0)$, $\alpha \in (0, 1)$ and $g \in L^\infty(M, \mu)$. Then the integral on the left-hand side is well-defined on \mathcal{S}^p with for every $f \in \mathcal{S}^p$*

$$\|\Pi_\Gamma(f, g)\|_{p, \alpha} = \left\| \int_0^\infty \tilde{Q}_t(\sqrt{t}\Gamma P_t f \cdot \sqrt{t}\Gamma P_t g) \frac{dt}{t} \right\|_{p, \alpha} \lesssim \|f\|_{p, \alpha} \|g\|_\infty.$$

Proof. The proof is similar to the one of Proposition 3.5. We first use that by choosing D in the definition of $Q_t = Q_t^{(D)}$ large enough, the operator $(tL)^{\alpha/2} Q_t$ satisfies L^2 off-diagonal estimates of order $N = N(D, \alpha) > \nu$. This allows to follow the steps in (3.3) and (3.4). We obtain

$$\begin{aligned} & \left\| L^{\alpha/2} \int_0^\infty Q_t(\sqrt{t}\Gamma P_t f \cdot \sqrt{t}\Gamma P_t g) \frac{dt}{t} \right\|_p \\ &= \left\| (tL)^{\alpha/2} \int_0^\infty Q_t(t^{-\alpha/2} \sqrt{t}\Gamma P_t f \cdot \sqrt{t}\Gamma P_t g) \frac{dt}{t} \right\|_p \\ &\lesssim \sum_{j=0}^\infty 2^{-2jN} 2^{j\nu} \left\| x \mapsto \left(\int_0^\infty \left(\int_{B(x, 2^j \sqrt{t})} |t^{-\alpha/2} \sqrt{t}\Gamma P_t f|^2 d\mu \right) \left(\int_{B(x, 2^j \sqrt{t})} |\sqrt{t}\Gamma P_t g|^2 d\mu \right) \frac{dt}{t} \right)^{1/2} \right\|_p \\ &\lesssim \|g\|_\infty \sum_{j=0}^\infty 2^{-2jN} 2^{j\nu} \left\| \sqrt{t}\Gamma (tL)^{-\alpha/2} P_t (L^{\alpha/2} f) \right\|_{T_{2^j}^{p, 2}(M)} \\ &\lesssim \|g\|_\infty \left\| \sqrt{t}\Gamma (tL)^{-\alpha/2} P_t (L^{\alpha/2} f) \right\|_{T^{p, 2}(M)}, \end{aligned}$$

where the last line follows from change of angle in tent spaces [2, Theorem 1.1]. If $p \geq 2$, the above conical square function estimate is dominated by its vertical counterpart [3, Proposition 2.1, Remark 2.2]. Invoking Lemma 2.5 for $p \in [2, p_0)$, we therefore have that the above is bounded by

$$\|g\|_\infty \left\| \sqrt{t}\Gamma (tL)^{-\alpha/2} P_t (L^{\alpha/2} f) \right\|_{L^p(M; L^2(\mathbb{R}_+; \frac{dt}{t}))} \lesssim \|g\|_\infty \|L^{\alpha/2} f\|_p.$$

If $p \in (1, 2)$, we use [1, Proposition 6.8] (adapted to our current setting under (UE) and (R_2)), to have the L^p -boundedness of the conical square function and we conclude to the same estimate. \square

Stein's complex interpolation between the estimates in Proposition 3.5 and Proposition 3.8 on the endpoints $(\alpha, p) = (0, \infty)$ and $(\alpha, p) = (1, p_0)$ then yields

Proposition 3.9. *Assume (UE) and (G_{p_0}) for some $p_0 \in [2, \infty)$. Let $p \in (p_0, \infty)$, $\alpha \in (0, \frac{p_0}{p})$ and $g \in L^\infty(M, \mu)$. Then for every $f \in \dot{L}_\alpha^p(M)$, we have*

$$\|\Pi_\Gamma(f, g)\|_{p, \alpha} \lesssim \|f\|_{p, \alpha} \|g\|_\infty.$$

Proof. We apply Stein's complex interpolation [12]. Let $p_1 \in (p_0, \infty)$, and $\beta \in (0, 1)$. Fix $g \in L^\infty(M, \mu)$. Define for $z \in \mathbb{C}$ the operator

$$T_g^z := L^{z/2} \Pi_\Gamma(L^{-z/2} \cdot, g).$$

Recall that under (UE), imaginary powers of L are bounded in L^p for all $p \in (1, \infty)$ (see [4, Proposition 2.1]), with bound

$$\|L^{i\eta}\|_{p \rightarrow p} \lesssim (1 + |\eta|)^s,$$

whenever $s > \frac{\nu}{2}$ and $\eta \in \mathbb{R}$. From Proposition 3.5, we know that $T_g^0 = \Pi_\Gamma(\cdot, g)$ is a bounded operator in L^{p_1} . We thus obtain

$$\sup_{\gamma \in \mathbb{R}} (1 + |\gamma|)^{-s} \|T_g^{i\gamma}\|_{p_1 \rightarrow p_1} \leq C^0,$$

with $s > \frac{\nu}{2}$. On the other hand, Proposition 3.8 yields that $T_g^0 = \Pi_\Gamma(\cdot, g)$ is bounded on \dot{L}_β^p . Hence,

$$\sup_{\gamma \in \mathbb{R}} (1 + |\gamma|)^{-s} \|T_g^{\beta + i\gamma}\|_{p_0 \rightarrow p_0} \leq C_\beta^1.$$

Stein's interpolation [12, Theorem 1] then yields that the operator

$$L^{\alpha/2} \Pi_\Gamma(L^{-\alpha/2} \cdot, g) : L^p \rightarrow L^p$$

is bounded whenever $\alpha = \theta\beta$ and $\frac{1}{p} = \frac{\theta}{p_0} + \frac{1-\theta}{p_1}$. Taking the limit for $\beta \rightarrow 1$ and $p_1 \rightarrow \infty$ yields the result. \square

REFERENCES

- [1] P. Auscher, *On necessary and sufficient conditions for L^p -estimates of Riesz transforms associated to elliptic operators on \mathbb{R}^n and related estimates*, Mem. Amer. Math. Soc., **186** (2007), no. 871.
- [2] P. Auscher, *Change of angle in tent spaces*, C. R. Acad. Sci. Paris, Ser. I, **349** (2011), 297–301.
- [3] P. Auscher, S. Hofmann, and J.-M. Martell, *Vertical versus conical square functions*, Trans. Amer. Math. Soc., **364**, no. 10, (2012), 5469–5489.
- [4] F. Bernicot, T. Coulhon and D. Frey, *Sobolev algebras through heat kernel estimates*, J. Éc. polytech. Math., **3** (2016), 99–161.
- [5] S. Blunck and P.C. Kunstmann, *Generalized Gaussian estimates and the Legendre transform*, J. Operator Theory, **53** (2005), no. 2, 351–365.
- [6] R.R. Coifman and Y. Meyer, *Au-delà des opérateurs pseudo-différentiels*, Astérisque, **57**, Société Math. de France (1978).
- [7] X.-T. Duong and El M. Ouhabaz *Gaussian upper bounds for heat kernels of a class of non-divergence operators*. International Conference on Harmonic Analysis and Related Topics (Sydney, 2002), 35–45, Proc. Centre Math. Appl. Austral. Nat. Univ., 41, Austral. Nat. Univ., Canberra, 2003.

- [8] D. Frey, Paraproducts via H^∞ -functional calculus, *Rev. Matematica Iberoam.*, **29**(2) (2013), 635–663.
- [9] T. Kato and G. Ponce, Commutator estimates and the Euler and Navier-Stokes equations, *Comm. Pure Appl. Math.*, **41** (1988), 891–907.
- [10] A. McIntosh and A. Nahmod, Heat kernel estimates and functional calculi of $-b\Delta$, *Math. Scand.* **87** (2000), no.2, 287–319.
- [11] Y. Meyer, Remarques sur un théorème de J. M. Bony, *Rend. Circ. Mat. Palermo, II. Ser.*, **1** (1981), 1–20.
- [12] E. M. Stein, Interpolation of linear operators, *Trans. Amer. Math. Soc.*, **83** (1956), 482–492.
- [13] R. Strichartz, Multipliers on fractional Sobolev spaces, *J. Math. Mech.*, **16**(9) (1967), 1031–1060.

FRÉDÉRIC BERNICOT, CNRS - UNIVERSITÉ DE NANTES, LABORATOIRE JEAN LERAY, 2
RUE DE LA HOUSSINIÈRE, 44322 NANTES CEDEX 3. FRANCE

E-mail address: frederic.bernicot@univ-nantes.fr

DOROTHEE FREY, DELFT INSTITUTE OF APPLIED MATHEMATICS, DELFT UNIVERSITY OF
TECHNOLOGY, P.O. BOX 5031, 2600 GA DELFT, THE NETHERLANDS

E-mail address: d.frey@tudelft.nl