

Free and projective Banach lattices

de Pagter, Ben; Wickstead, Anthony W.

DOI

[10.1017/S0308210512001709](https://doi.org/10.1017/S0308210512001709)

Publication date

2015

Document Version

Accepted author manuscript

Published in

Royal Society of Edinburgh. Proceedings. Section A(Mathematics)

Citation (APA)

de Pagter, B., & Wickstead, A. W. (2015). Free and projective Banach lattices. *Royal Society of Edinburgh. Proceedings. Section A(Mathematics)*, 145(1), 105-143. <https://doi.org/10.1017/S0308210512001709>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

FREE AND PROJECTIVE BANACH LATTICES.

B. DE PAGTER AND A.W. WICKSTEAD

ABSTRACT. We define and prove the existence of free Banach lattices in the category of Banach lattices and contractive lattice homomorphisms and establish some of their fundamental properties. We give much more detailed results about their structure in the case that there are only a finite number of generators and give several Banach lattice characterizations of the number of generators being, respectively, one, finite or countable. We define a Banach lattice P to be *projective* if whenever X is a Banach lattice, J a closed ideal in X , $Q : X \rightarrow X/J$ the quotient map, $T : P \rightarrow X/J$ a linear lattice homomorphism and $\epsilon > 0$ there is a linear lattice homomorphism $\hat{T} : P \rightarrow X$ such that (i) $T = Q \circ \hat{T}$ and (ii) $\|\hat{T}\| \leq (1 + \epsilon)\|T\|$. We establish the connection between projective Banach lattices and free Banach lattices and describe several families of Banach lattices that are projective as well as proving that some are not.

1. INTRODUCTION.

Free and projective objects have not played anywhere near as important a rôle in analysis as in algebra, nevertheless there has been some work done on these objects, mainly with the results that one would expect. For example, the existence of free and projective Banach spaces is virtually folklore but is uninteresting as both are of the form $\ell_1(I)$ for an arbitrary index set I . The existence of free vector lattices over an arbitrary number of generators is also long established and holds no real surprises, see [1] or [3] for details. In this note we investigate free and projective Banach lattices. Some of our results are rather surprising and although we are able to answer many questions we are forced to leave several unanswered.

It is almost obvious that, if it exists, then the free Banach lattice over \mathfrak{a} generators must be the completion of the free vector lattice over \mathfrak{a} generators for some lattice norm. That the required norm actually exists is easily proved, but describing it in concrete and readily identifiable terms is not so easy. Indeed, except in the case $\mathfrak{a} = 1$, it is

1991 *Mathematics Subject Classification.* 46B42.

Key words and phrases. Banach lattice, vector lattice, free, projective.

not a classical Banach lattice norm at all. In fact it is only in the case that \mathfrak{a} is finite that the free Banach lattice over \mathfrak{a} generators is even isomorphic to an AM-space.

§2 is primarily devoted to establishing notation whilst §3 recapitulates the existing theory of free vector lattices. We then prove the existence of free Banach lattices in §4 and give a representation on a compact Hausdorff space in §5. We establish some of the basic properties of free Banach lattices in §6. The finitely generated free Banach lattices are by far the easiest ones to understand, and we investigate their structure in §7. In §8 we give some characterizations of free Banach lattices over, respectively, one, a finite number or a countable number of generators, amongst all free Banach lattices. Preparatory to looking at projective Banach lattices, in §9 we investigate when disjoint families in quotient Banach lattices X/J can be lifted to disjoint families in X , giving a positive result for countable families and a negative result for larger ones. We prove the connection between free and projective Banach lattices in §10 and in §11 find some classes of Banach lattices that are, or are not, projective. Finally, §12 contains some open problems.

Let us emphasize at this point that this paper is set in the category of Banach lattices and linear lattice homomorphisms. There is a substantial theory of *injective* Banach lattices (and indeed we refer to them later) but this is set in the context of Banach lattices and positive (or regular) operators.¹ Thus there is no reason to expect any kind of duality between the two notions.

¹In fact, although we can find no explicit proof in the literature, there is no non-zero injective in the category of Banach lattices and linear lattice homomorphisms. Indeed, suppose that F were a non-zero injective. Let \mathfrak{a} be strictly greater than the cardinality of F^* and let μ be the product of \mathfrak{a} many copies of the measure which assigns mass $\frac{1}{2}$ to each of 0 and 1 in $\{0, 1\}$. This is a homogenous measure space and each order interval in $\mathcal{L}_1(\mu)$ has the property that the least cardinality of a dense subset is precisely \mathfrak{a} , see [20] §26 for details. In particular every order interval has cardinality at least \mathfrak{a} . As μ is finite, the same is true of $\mathcal{L}_\infty(\mu)$. Pick any non-zero $y \in F_+$. As F is alleged to be injective, there is a linear lattice homomorphism T extending the map that takes the constantly one function in $\mathcal{L}_1(\mu)$, $\mathbf{1}$, to y . The adjoint of this maps F^* into $\mathcal{L}_1(\mu)^* = \mathcal{L}_\infty(\mu)$ and is interval preserving, [14], Theorem 1.4.19. In particular, if $f \in F_+^*$ with $f(y) > 0$ then $T^*f(\mathbf{1}) = f(T\mathbf{1}) = f(y) > 0$, so the image of the order interval $[0, f]$ will be a non-zero order interval in $\mathcal{L}_\infty(\mu)$ which has cardinality at least \mathfrak{a} . This contradicts the fact that $[0, f]$ has cardinality strictly less than \mathfrak{a} .

2. NOTATION.

In this short section we establish the notation that we will use concerning functions and function spaces. If A and X are non-empty sets then, as usual, X^A denotes the set of all maps from A into X . If $\emptyset \neq B \subseteq A$ then we let $r_B : X^A \rightarrow X^B$ denote the restriction map with $r_B \xi = \xi|_B$ for $\xi \in X^A$. Clearly, r_B is surjective. On occasions we will also write ξ_B in place of $r_B(\xi)$.

The space of all real-valued functions on X^A , \mathbb{R}^{X^A} , is a vector lattice under the pointwise operations. Again, we consider the setting where B is a non-empty subset of A and define $j_B : \mathbb{R}^{X^B} \rightarrow \mathbb{R}^{X^A}$ by $(j_B f)(\xi) = f(\xi_B)$ for $\xi \in X^A$ and $f \in \mathbb{R}^{X^B}$. This makes j_B an injective lattice homomorphism. The following description of the image of j_B is easily verified.

Lemma 2.1. *If A, B and X are non-empty sets with $B \subseteq A$ and $f \in \mathbb{R}^{X^A}$ then the following are equivalent:*

- (1) $f \in j_B(\mathbb{R}^{X^B})$.
- (2) If $\xi, \eta \in X^A$ with $\xi_B = \eta_B$ then $f(\xi) = f(\eta)$.

We now specialize somewhat by assuming that $X \subseteq \mathbb{R}$ and that $0 \in X$. This means that if $\xi \in X^A$, $\emptyset \neq B \subseteq A$ and χ_B is the characteristic function of B then the pointwise product $\xi\chi_B \in X^A$.

Lemma 2.2. *If $\emptyset \neq B \subseteq A$ and $0 \in X \subseteq \mathbb{R}$ then the map $P_B : \mathbb{R}^{X^A} \rightarrow j_B(\mathbb{R}^{X^B})$ defined by*

$$(P_B f)(\xi) = f(\xi\chi_B) \quad (\xi \in X^A, f \in \mathbb{R}^{X^A})$$

is a linear lattice homomorphism and a projection onto $j_B(\mathbb{R}^{X^B})$. Furthermore, if $B_1, B_2 \subseteq A$ are non-empty sets with non-empty intersection then $P_{B_1}P_{B_2} = P_{B_2}P_{B_1} = P_{B_1 \cap B_2}$.

Proof. It is clear that P_B is a well-defined vector lattice homomorphism of X^A into itself. If $\xi, \eta \in X^A$ are such that $\xi_B = \eta_B$ then $(P_B f)(\xi) = f(\xi\chi_B) = f(\eta\chi_B) = (P_B f)(\eta)$ so by Lemma 2.1 $P_B f \in j_B(\mathbb{R}^{X^B})$ for all $f \in \mathbb{R}^{X^A}$. If $f \in \mathbb{R}^{X^B}$ then for any $\xi \in X^A$ we have $P_B(j_B f)(\xi) = (j_B f)(\xi\chi_B) = (j_B f)(\xi)$ as ξ and $\xi\chi_B$ coincide on B and using Lemma 2.1 again. Thus P_B is indeed a projection.

Finally, if $f \in \mathbb{R}^{X^A}$ and $\xi \in X^A$ then

$$\begin{aligned} P_{B_1}P_{B_2}f(\xi) &= (P_{B_2})(f\chi_{B_1}) = f(\xi\chi_{B_1}\chi_{B_2}) \\ &= f(\xi\chi_{B_1 \cap B_2}) = (P_{B_1 \cap B_2}f)(\xi), \end{aligned}$$

which shows that $P_{B_1}P_{B_2} = P_{B_1 \cap B_2}$. Similarly $P_{B_2}P_{B_1} = P_{B_2 \cap B_1} = P_{B_1 \cap B_2}$ and the proof is complete. \square

In future, we will identify \mathbb{R}^{X^B} with the vector sublattice $j_B(\mathbb{R}^{X^B})$ of \mathbb{R}^{X^A} .

If L is any vector lattice and D a non-empty subset of L then $\langle D \rangle$ will denote the vector sublattice of L generated by D . All elements of $\langle D \rangle$ can be obtained from those of D by the application of a finite number of multiplications, additions, suprema and infima. The following simple consequence of this observation may also be proved directly:

Lemma 2.3. *If L and M are vector lattices, $T : L \rightarrow M$ is a vector lattice homomorphism and $\emptyset \neq D \subseteq L$ then $\langle T(D) \rangle = T(\langle D \rangle)$.*

We specialize further now to the case that $X = \mathbb{R}$. On the space \mathbb{R}^A we can consider the product topology, which is the topology of pointwise convergence on A . By definition, this is the weakest topology such that all the functions $\delta_a : \xi \mapsto \xi(a)$ are continuous on \mathbb{R}^A for each $a \in A$. As a consequence we certainly have $\{\delta_a : a \in A\} \subset C(\mathbb{R}^A)$. In fact we can do rather better than this. A function $f : \mathbb{R}^A \rightarrow \mathbb{R}$ is *homogeneous* if $f(t\xi) = tf(\xi)$ for $\xi \in \mathbb{R}^A$ and $t \in [0, \infty)$. The space $H(\mathbb{R}^A)$ of continuous homogeneous real-valued functions on \mathbb{R}^A is a vector sublattice of $C(\mathbb{R}^A)$ and clearly $\{\delta_a : a \in A\} \subset H(\mathbb{R}^A)$.

3. FREE VECTOR LATTICES.

In this section we recapitulate much of the theory of free vector lattices, both to make this work as self-contained as possible and in order to establish both our notation (which may not coincide with that used in other papers on free vector lattices) and to point out some properties that we will use later.

Definition 3.1. If A is a non-empty set then a *free vector lattice* over A is a pair (F, ι) where F is a vector lattice and $\iota : A \rightarrow F$ is a map with the property that for any vector lattice E and any map $\phi : A \rightarrow E$ there is a unique vector lattice homomorphism $T : F \rightarrow E$ such that $\phi = T \circ \iota$.

It follows immediately from this definition that the map ι must be injective, as we can certainly choose E and ϕ to make ϕ injective. Many of the results that follow are almost obvious, but we prefer to make them explicit.

Proposition 3.2. *If (F, ι) is a free vector lattice over A then F is generated, as a vector lattice, by $\iota(A)$.*

Proof. Let G be the vector sublattice of F generated by $\iota(A)$. Define $\phi : A \rightarrow G$ by $\phi(a) = \iota(a)$ then it follows from the definition that there

is a unique vector lattice homomorphism $T : F \rightarrow G$ with $T(\iota(a)) = \phi(a) = \iota(a)$ for $a \in A$. If $j : G \rightarrow F$ is the inclusion map, then $j \circ T : F \rightarrow F$ is a vector lattice homomorphism with $(j \circ T)(\iota(a)) = j(\iota(a)) = \iota(a)$ for $a \in A$. The identity on F , I_F , is also a vector lattice homomorphism from F into itself with $I_F(\iota(a)) = \iota(a)$. The uniqueness part of the definition of a free vector lattice applied to the map $a \mapsto \iota(a)$, of A into F , tells us that these two maps are equal so that $j \circ T = I_F$ from which we see that $F \subseteq G$ and therefore $F = G$ as claimed. \square

The definition of a free vector lattice make the following result easy to prove.

Proposition 3.3. *If (F, ι) and (G, κ) are free vector lattices over a non-empty set A then there is a (unique) vector lattice isomorphism $T : F \rightarrow G$ such that $T(\iota(a)) = \kappa(a)$ for $a \in A$.*

In view of this we will just refer to a free vector lattice (F, ι) over a set A as *the free vector lattice over A* (or sometimes as *the free vector lattice generated by A* when we identify A with a subset of that free vector lattice). We will denote it by $FVL(A)$. It will be clear that if A and B are sets of equal cardinality then $FVL(A)$ and $FVL(B)$ are isomorphic vector lattices, so that $FVL(A)$ depends only on the cardinality of the set A . Thus we will also use the notation $FVL(\mathfrak{a})$ for $FVL(A)$ when \mathfrak{a} is the cardinality of A . This is the notation that will be found elsewhere in the literature. We retain both versions so that we can handle proper inclusions of $FVL(B)$ into $FVL(A)$ when $B \subset A$ even when A and B have the same cardinality.

If $\iota : A \rightarrow FVL(A)$ is the embedding of A into $FVL(A)$ specified in the definition then we will often write δ_a for $\iota(a)$ and refer to the set $\{\delta_a : a \in A\}$ as the *free generators* of $FVL(A)$.

A slight rewording of the definition of a free vector lattice is sometimes useful, which trades off uniqueness of the lattice homomorphism for specifying that $\iota(A)$ is a generating set. The proof of this follows immediately from results above.

Proposition 3.4. *If A is a non-empty set then the vector lattice F is the free vector lattice over A if and only if*

- (1) *There is a subset $\{\delta_a : a \in A\} \subset F$, with $\delta_a \neq \delta_b$ if $a \neq b$, which generates F as a vector lattice.*
- (2) *For every vector lattice E and any family $\{x_a : a \in A\} \subset E$ there is a vector lattice homomorphism $T : F \rightarrow E$ such that $T(\delta_a) = x_a$ for $a \in A$.*

We will find the next simple result useful later.

Proposition 3.5. *Let A be a non-empty set and $\{\delta_a : a \in A\}$ be the free generators of $FVL(A)$. Let B and C be non-empty subsets of A with $B \cap C \neq \emptyset$.*

- (1) *The vector sublattice of $FVL(A)$ generated by $\{\delta_b : b \in B\}$ is (isomorphic to) the free vector lattice $FVL(B)$.*
- (2) *There is a lattice homomorphism projection P_B from $FVL(A)$ onto $FVL(B)$.*
- (3) *$P_C P_B = P_B P_C = P_{B \cap C}$.*

Proof. (1) Let F denote the vector sublattice of $FVL(A)$ generated by $\{\delta_b : b \in B\}$. Suppose that E is a vector lattice and $\pi : B \rightarrow E$ is any map. There is a unique vector lattice homomorphism $T : FVL(A) \rightarrow E$ with $T(\delta_b) = \pi(b)$ for $b \in B$ and $T(\delta_a) = 0$ for $a \in A \setminus B$. The restriction S of T to F gives us a vector lattice homomorphism $S : F \rightarrow E$ with $S(\delta_b) = \pi(b)$. It follows from Proposition 3.4 that $F = FVL(B)$.

(2) The free property of $FVL(A)$ gives a (unique) lattice homomorphism $P_B : FVL(A) \rightarrow FVL(A)$ with $P_B(\delta_b) = \delta_b$ if $b \in B$ and $P_B(\delta_a) = 0$ if $a \in A \setminus B$. As P_B maps the generators of $FVL(A)$ into $FVL(B)$, we certainly have $P_B(FVL(A)) \subseteq FVL(B)$. Also, P_B is the identity on the generators of $FVL(B)$ so is the identity linear operator on $FVL(B)$ so that P_B is indeed a projection.

(3) If $a \in B \cap C$ then $P_C P_B \delta_a = P_B P_C \delta_a = P_{B \cap C} \delta_a = \delta_a$ whilst if $a \notin B \cap C$ then $P_C P_B \delta_a = P_B P_C \delta_a = P_{B \cap C} \delta_a = 0$. Thus the three vector lattice homomorphisms $P_B P_C$, $P_C P_B$ and $P_{B \cap C}$ coincide on a set of generators of $FVL(A)$ and are therefore equal. \square

So far all our discussions of free vector lattices have been rather academic as we have not shown that they exist. However it was shown in [1] (see also [3]) that they do exist. In essence we have:

Theorem 3.6. *For any non-empty set A , $FVL(A)$ exists and is the vector sublattice of $\mathbb{R}^{\mathbb{R}^A}$ generated by δ_a ($a \in A$) where $\delta_a(\xi) = \xi(a)$ for $\xi \in \mathbb{R}^A$.*

It is reasonable to ask how this representation of $FVL(A)$ interacts with the properties of free vector lattices noted above. With the notation of §2, if $\emptyset \neq B \subseteq A$ then the map $j_B : \mathbb{R}^{\mathbb{R}^B} \rightarrow \mathbb{R}^{\mathbb{R}^A}$ is a vector lattice embedding of $\mathbb{R}^{\mathbb{R}^B}$ into $\mathbb{R}^{\mathbb{R}^A}$. This corresponds precisely to the embedding of $FVL(B)$ into $FVL(A)$ as indicated in Proposition 3.5. If we use δ_a to denote the map $\xi \mapsto \xi(a)$ on \mathbb{R}^A and η_b for the map

$\xi \mapsto \xi(b)$ on \mathbb{R}^B then we have, for $b \in B$ and $\xi \in \mathbb{R}^A$

$$(j_B \eta_b)(\xi) = \eta_b(\xi_B) = \xi(b) = \delta_b(\xi)$$

so that $j_B \eta_b = \delta_b$. We know from §2 that j_B is a vector lattice homomorphism so that $j_B(FVL(B))$ is the vector sublattice of $FVL(A)$ generated by $\{\delta_b : b \in B\}$ which is precisely what was described in Proposition 3.5.

Also, if $B \subseteq A$ then we may consider $FVL(B) \subseteq FVL(A) \subseteq \mathbb{R}^{\mathbb{R}^A}$. The projection map $P_B : FVL(A) \rightarrow FVL(B)$ defined in Proposition 3.5 (2) is then precisely the restriction to $FVL(A)$ of the projection $P_B : \mathbb{R}^{\mathbb{R}^A} \rightarrow \mathbb{R}^{\mathbb{R}^B}$ described in Lemma 2.2. We will temporarily denote this projection by \tilde{P}_B to distinguish it from the abstract projection. Once we establish equality that distinction will not be required and we will omit the tilde. As P_B and \tilde{P}_B are both vector lattice homomorphisms it suffices to prove this equality for the generators of $FVL(A)$. If $b \in B$ then

$$(\tilde{P}_B \delta_b)(\xi) = \delta_b(\xi \chi_B) = (\xi \chi_B)(b) = \xi(b) = \delta_b(\xi)$$

for $\xi \in \mathbb{R}^A$ so that $\tilde{P}_B \delta_b = \delta_b = P_B \delta_b$. If, on the other hand, $a \in A \setminus B$ then

$$(\tilde{P}_B \delta_a)(\xi) = \delta_a(\xi \chi_B) = 0$$

for $\xi \in \mathbb{R}^A$ so that $\tilde{P}_B \delta_a = 0 = P_B \delta_a$.

A few more observations will be of use later.

Proposition 3.7. *If A is a non-empty set and $\mathcal{F}(A)$ denotes the collection of all non-empty finite subsets of A , then*

$$FVL(A) = \bigcup_{B \in \mathcal{F}(A)} FVL(B).$$

Proof. Any element of $FVL(A)$ is in the vector sublattice of $FVL(A)$ generated by a finite number of generators $\{\delta_{a_1}, \delta_{a_2}, \dots, \delta_{a_n}\}$ so lies in $FVL(\{a_1, a_2, \dots, a_n\})$. \square

Proposition 3.8. *If A is a finite set then $\sum_{a \in A} |\delta_a|$ is a strong order unit for $FVL(A)$.*

Proof. Obvious as $FVL(A)$ is generated by the set $\{\delta_a : a \in A\}$. \square

Lemma 3.9. *The real valued vector lattice homomorphisms on $FVL(A)$ are precisely the evaluations at points of \mathbb{R}^A .*

Proof. It is clear that if $\xi \in \mathbb{R}^A$ then the map $\omega_\xi : f \mapsto f(\xi)$ is a real valued vector lattice homomorphism on $\mathbb{R}^{\mathbb{R}^A}$ and therefore on $FVL(A)$. Note, in particular, that $\omega_\xi(\delta_a) = \delta_a(\xi) = \xi(a)$. Conversely, if ω is a real valued vector lattice homomorphism on $FVL(A)$ then we may

define $\xi \in \mathbb{R}^A$ by $\xi(a) = \omega(\delta_a)$ for $a \in A$. Now we see that for this ξ , ω_ξ is a real valued vector lattice homomorphism on $FVL(A)$ with $\omega_\xi(\delta_a) = \xi(a) = \omega(\delta_a)$. The two maps ω and ω_ξ coincide on a set of generators of $FVL(A)$ so, being vector lattice homomorphisms, are equal. \square

4. FREE BANACH LATTICES.

Definition 4.1. If A is a non-empty set then a *free Banach lattice* over A is a pair (X, ι) where X is a Banach lattice and $\iota : A \rightarrow X$ is a bounded map with the property that for any Banach lattice Y and any bounded map $\kappa : A \rightarrow Y$ there is a unique vector lattice homomorphism $T : X \rightarrow Y$ such that $\kappa = T \circ \iota$ and $\|T\| = \sup\{\|\kappa(a)\| : a \in A\}$.

It is clear that the set $\{\iota(a) : a \in A\}$ generates X as a Banach lattice (cf Proposition 3.2).

Remark 4.2. The definition forces each $\iota(a)$ to have norm precisely one. For if $\kappa(a) = 1 \in \mathbb{R}$ for each $a \in A$ then the map T that is guaranteed to exist has norm 1, so that $1 = \|T(\iota(a))\| \leq \|\iota(a)\|$. On the other hand, if we take $\kappa = \iota$, then T is identity operator, with norm 1, so that $\sup\{\|\iota(a)\| : a \in A\} = 1$.

Proposition 4.3. *If (X, ι) and (Y, κ) are free Banach lattices over a non-empty set A then there is a (unique) isometric order isomorphism $T : X \rightarrow Y$ such that $T(\iota(a)) = \kappa(a)$ for $a \in A$.*

Proof. As (X, ι) is free, there is a vector lattice homomorphism $T : X \rightarrow Y$ with $T(\iota(a)) = \kappa(a)$ for $a \in A$ with $\|T\| = \sup\{\|\kappa(a)\| : a \in A\} = 1$, by the preceding proposition. There is similarly a contractive vector lattice homomorphism $S : Y \rightarrow X$ with $S(\kappa(a)) = \iota(a)$. By uniqueness, the compositions $S \circ T$ and $T \circ S$ must be the identity operators. This suffices to prove our claim. \square

Similarly to the free vector lattice case, we use the notation $FBL(A)$ for the free Banach lattice over A if it exists (which we will shortly show is the case.) Since we know that if A and B have the same cardinality then $FBL(A)$ and $FBL(B)$ are isometrically order isomorphic, we will also use the notation $FBL(\mathfrak{a})$ to denote a free Banach lattice on a set of cardinality \mathfrak{a} . Again, we will also use the notation δ_a for $\iota(a)$ and refer to $\{\delta_a : a \in A\}$ as the *free generators* of $FBL(A)$.

Our first task is to show that free Banach lattices do indeed exist.

Definition 4.4. If A is a non-empty set then we will define a mapping from $FVL(A)^\sim$ into the extended non-negative reals by

$$\|\phi\|^\dagger = \sup\{|\phi|(|\delta_a|) : a \in A\}.$$

We define also

$$FVL(A)^\dagger = \{\phi \in FVL(A)^\sim : \|\phi\|^\dagger < \infty\}$$

which it is clear is a vector lattice ideal in the Dedekind complete vector lattice $FVL(A)^\sim$.

Suppose that a positive functional ϕ vanishes on each $|\delta_a|$. Each element x of $FVL(A)$ lies in the sublattice of $FVL(A)$ generated by a finite set of generators $\{a_k : 1 \leq k \leq n\}$. By Proposition 3.8 $e = \sum_{k=1}^n |\delta_{a_k}|$ is a strong order unit for that sublattice. Thus there is $\lambda \in \mathbb{R}$ with $|x| \leq \lambda e$ so that $|\phi(x)| \leq \phi(|x|) \leq \phi(\lambda e) = \lambda \sum_{k=1}^n \phi(|\delta_{a_k}|) = 0$ and thus $\phi = 0$. It is now clear that $\|\cdot\|^\dagger$ is a lattice norm on $FVL(A)^\dagger$. Given the embedding of $FVL(A)$ in $\mathbb{R}^{\mathbb{R}^A}$ given in Theorem 3.6, if $\xi \in \mathbb{R}^A$ then $\omega_\xi \in FVL(A)^\dagger$ if and only if the map $\xi : A \rightarrow \mathbb{R}$ is bounded and then $\|\omega_\xi\|^\dagger = \sup_{a \in A} |\xi(a)|$. By Lemma 3.9, these maps are lattice homomorphisms. Note that if A is an infinite set then there is an unbounded $\xi \in \mathbb{R}^A$ which induces $\omega_\xi \in FVL(A)^\sim \setminus FVL(A)^\dagger$.

Definition 4.5. For $f \in FVL(A)$, where A is a non-empty set, define

$$\|f\|_F = \sup\{\phi(|f|) : \phi \in FVL(A)_+^\dagger, \|\phi\|^\dagger \leq 1\}.$$

Proposition 4.6. For any non-empty set A , $\|\cdot\|_F$ is a lattice norm on $FVL(A)$.

Proof. Our first step is to show that $\|\cdot\|_F$ is real-valued. By Proposition 3.7, any $f \in FVL(A)$ actually lies in $FVL(B)$ for some finite subset $B \subseteq A$. By Proposition 3.8, $FVL(B)$ has a strong order unit $\sum_{b \in B} |\delta_b|$, so there is λ with $|f| \leq \lambda \sum_{b \in B} |\delta_b|$. If $\phi \in FVL(A)_+^\dagger$ with $\|\phi\|^\dagger \leq 1$ then

$$\phi(|f|) \leq \phi\left(\lambda \sum_{b \in B} |\delta_b|\right) = \lambda \sum_{b \in B} \phi(|\delta_b|) \leq \lambda \sum_{b \in B} 1$$

so that $\|f\|_F$ is certainly finite.

If $\|f\|_F = 0$ then $\phi(|f|) = 0$ for all $\phi \in FVL(A)_+^\dagger$. Using the observation above, $f(\xi) = \omega_\xi(f) = 0$ for any bounded function $\xi : A \rightarrow \mathbb{R}$. But there is a finite set $B \subset A$ such that $f \in FVL(B)$, so that $f(\xi) = f(\xi \chi_B)$ for all $\xi \in \mathbb{R}^A$. As each $\xi \chi_B$ is bounded, $f(\xi) = 0$ for all $\xi \in \mathbb{R}^A$ and therefore $f = 0$.

That $\|\cdot\|_F$ is sublinear and positively homogeneous are obvious, so that $\|\cdot\|_F$ is a norm on $FVL(A)$, which is clearly a lattice norm. \square

Note in particular that we certainly have $\|\delta_a\|_F = 1$ for all $a \in A$. In fact, this construction gives us our desired free Banach lattices.

Theorem 4.7. *For any non-empty set A , the pair consisting of the completion of $FVL(A)$, under the norm $\|\cdot\|_F$, and the map $\iota : a \rightarrow \delta_a$, is the free Banach lattice over A .*

Proof. Suppose that Y is any Banach lattice and $\kappa : A \rightarrow Y_1$, the unit ball of Y . There is a vector lattice homomorphism $T : FVL(A) \rightarrow Y$ with $T(\iota(a)) = \kappa(a)$ for all $a \in A$, as $FVL(A)$ is free. We claim that if $f \in FVL(A)$ with $\|f\|_F \leq 1$ then $\|Tf\| = \||Tf\|| = \||T(|f|)\|| \leq 1$ in Y , where we have used that fact that the norm in Y is a lattice norm and that T is a lattice homomorphism. If this were not the case then we could find $\psi \in Y_{1+}^*$, a positive linear functional on Y with norm at most 1, with $\psi(T(|f|)) > 1$. As $\|T(\iota(a))\| = \|\kappa(a)\| \leq 1$ for all $a \in A$, we have $\||T(\iota(a))\|| = \||T(|\iota(a)|)\|| \leq 1$, using again the fact that T is a lattice homomorphism. Thus $\left| \psi\left(T(|\iota(a)|)\right) \right| \leq 1$ for all $a \in A$. Using the functional $\psi \circ T$ in the definition of $\|f\|_F$, we see that $\|f\|_F \geq \psi(T(|f|)) > 1$, contradicting our assumption that $\|f\|_F \leq 1$.

The completion of $FVL(A)$ is a Banach lattice and T will extend by continuity to it whilst still taking values in Y as Y is complete. \square

We will eventually need to know the relationship between different free Banach lattices, so we record now the following result.

Proposition 4.8. *If B is a non-empty subset of A then $FBL(B)$ is isometrically order isomorphic to the closed sublattice of $FBL(A)$ generated by $\{\delta_b : b \in B\}$. Furthermore there is a contractive lattice homomorphic projection P_B of $FBL(A)$ onto $FBL(B)$.*

Proof. Recall from Proposition 3.5 that $FVL(B)$ is isomorphic to the sublattice of $FVL(A)$ generated by $\{\delta_b : b \in B\}$ that there is a lattice homomorphism projection P_B of $FVL(A)$ onto $FVL(B)$ with $P_B(\delta_b) = \delta_b$ if $b \in B$ and $P_B(\delta_a) = 0$ if $a \in A \setminus B$. As $\|\delta_b\|_F = 1$ in both $FBL(A)$ and $FBL(B)$, there are contractive lattice homomorphisms of $FBL(B)$ into $FBL(A)$ and of $FBL(B)$ onto $FBL(A)$ which act into the same way on the generators so extend these. The conclusion is now clear. \square

There is also a simple relationship between their duals. This is a consequence of the following result which is surely well known but for which we can find no convenient reference, but see [21], IV.12, Problem 6 and [7], Lemma VI.3.3 for similar results.

Proposition 4.9. *If P is a contractive lattice homomorphism projection from a Banach lattice X onto a closed sublattice Y then P^*Y^* is a weak*-closed band in X^* which is isometrically order isomorphic to Y^* .*

Proof. Write $\ker(P)$ for the kernel of P , which is a lattice ideal in X , and $Z = \{\phi \in X^* : \phi|_{\ker(P)} \equiv 0\}$, which is a weak*-closed band in X^* . It is clear that $P^*X^* = Z$.

Define $J : Y^* \rightarrow X^*$ by $J\phi = \phi \circ P$ and note that $J : Y^* \rightarrow Z$ with $\|J\| \leq \|P\|$. If $\phi \in Z$ then $J(\phi|_Y) = \phi$ so that J is actually an isometry of Y^* onto Z . It is clear that both J and J^{-1} are positive. Thus $J : Y^* \rightarrow P^*X^*$ is actually an isometric order isomorphism. \square

Corollary 4.10. *If B is a non-empty subset of A then $FBL(B)^*$ is isometrically order isomorphic to a weak*-closed band in $FBL(A)^*$.*

As in the algebraic case, if B and C are two subsets of A with $B \cap C \neq \emptyset$ of A then $P_B P_C = P_C P_B = P_{B \cap C}$.

In particular, the embedding of the finitely generated free closed sublattices are important.

Proposition 4.11. *Let $\mathcal{F}(A)$ be the collection of all non-empty finite subsets of A , ordered by inclusion. The net of projections $\{P_B : B \in \mathcal{F}(A)\}$ in $FBL(A)$ converges strongly to the identity in $FBL(A)$.*

Proof. If $f \in FVL(A)$ then there is actually $B_0 \in \mathcal{F}(A)$ with $P_{B_0}(f) = f$ whenever $B_0 \subset B$. Recall that each P_B is a contraction. If $\epsilon > 0$ and $f \in FBL(A)$, choose $f' \in FVL(A)$ with $\|f - f'\|_F < \epsilon/2$ and then $B_0 \in \mathcal{F}(A)$ with $P_{B_0}(f') = f'$ for $B_0 \subset B$. Then if $B_0 \subset B$ then

$$\|P_B f - f\|_F \leq \|P_B f - P_B f'\| + \|P_B f' - f'\|_F + \|f' - f\|_F < \epsilon,$$

which completes the proof. \square

Before looking at some properties of $FBL(A)$ in detail, we will ask about its normed dual.

Proposition 4.12. *If A is any non-empty set then the three normed spaces $(FVL(A)^\dagger, \|\cdot\|^\dagger)$, $(FVL(A), \|\cdot\|_F)^*$ and $FBL(A)^*$ are isometrically order isomorphic.*

Proof. If $\phi \in FBL(A)^*$ then the restriction map $\phi \mapsto \phi|_{FVL(A)}$ is an order isomorphism, by continuity, and as $\|\delta_a\| = 1$ we have $|\phi|(\delta_a) \leq \|\phi\|$ so that $\|\phi|_{FVL(A)}\|^\dagger \leq \|\phi\|$. On the other hand, as each $\|\delta_a\| = 1$ we see that

$$\begin{aligned} \|\phi\| &= \|\phi\| = \sup\{|\phi|(f) : \|f\|_F \leq 1\} \\ &\leq \sup\{|\phi|(|\delta_a|) : a \in A\} = \|\phi|_{FVL(A)}\|^\dagger \end{aligned}$$

so the isometric order isomorphism of the first and third spaces is proved. The identification of the second and third follows from the density of $FVL(A)$ in $FBL(A)$. \square

As we noted above, if A is infinite then $FVL(A)^\dagger \neq FVL(A)^\sim$. On the other hand, we have:

Proposition 4.13. *If $n \in \mathbb{N}$ then $FBL(n)^*$ is isometrically order isomorphic to the whole of $FVL(n)^\sim$ under the norm $\|\cdot\|^\dagger$.*

Proof. All that remains to establish is that $\|\phi\|^\dagger$ is finite for all $\phi \in FVL(n)$. Given that $\|\phi\|^\dagger$ is, in this case, a finite supremum of real values $|\phi|(|\delta_a|)$, this is clear. \square

5. A SMALLER REPRESENTATION SPACE

The set $\Delta_A = [-1, 1]^A$ is a compact subset of \mathbb{R}^A . We call a function $f : \Delta_A \rightarrow \mathbb{R}$ *homogeneous* if $f(t\xi) = tf(\xi)$ for $\xi \in \Delta_A$ and $t \in [0, 1]$ (this is consistent with the definition for functions on \mathbb{R}^A). The space of continuous homogeneous real-valued functions on Δ_A is denoted by $H(\Delta_A)$. If we equip $C(\Delta_A)$ with the supremum norm $\|\cdot\|_\infty$, then $H(\Delta_A)$ is a closed vector sublattice of $C(\Delta_A)$ (and hence $H(\Delta_A)$ is itself a Banach lattice with respect to this norm).

Lemma 5.1. *The restriction map $R : H(\mathbb{R}^A) \rightarrow H(\Delta_A)$ is a injective vector lattice homomorphism.*

Proof. The only part of the proof that is not completely trivial is that the map R is injective. Suppose that $f \in H(\mathbb{R}^A)$ and $Rf = 0$. If $\xi \in \mathbb{R}^A$, consider the net $\{\xi\chi_B : B \in \mathcal{F}(A)\}$, where $\mathcal{F}(A)$ is the collection of all non-empty finite subsets of A ordered by inclusion, then we have $\xi\chi_B \rightarrow_{\mathcal{F}(A)} \xi$ in \mathbb{R}^A . For any $B \in \mathcal{F}(A)$, there is $t > 0$ such that $t\xi\chi_B \in [-1, 1]^A$, so that $tf(\xi\chi_B) = f(t\xi\chi_B) = 0$ by homogeneity. Hence, $f(\xi\chi_B) = 0$ and so $f(\xi) = 0$ by the continuity of f , so that $f = 0$. \square

It should be noted that the restriction map is not surjective unless A is a finite set.

Example 5.2. It suffices to prove the non-surjectiveness in the case that $A = \mathbb{N}$. Define $g \in H(\Delta_{\mathbb{N}})$ by $g(\xi) = \sum_{k=1}^{\infty} 2^{-k}\xi(k)$ for $\xi \in \Delta_{\mathbb{N}}$. Suppose that there is $f \in H(\mathbb{R}^{\mathbb{N}})$ with $Rf = g$. Define $\eta \in \mathbb{R}^{\mathbb{N}}$ by $\eta(k) = 2^k$ and let $\eta_n = \eta\chi_{\{1, \dots, n\}}$, for $n \in \mathbb{N}$, so that $\eta_n \rightarrow \eta$ in $\mathbb{R}^{\mathbb{N}}$. But, for each $n \in \mathbb{N}$ we have

$$f(\eta_n) = 2^n f(2^{-n}\eta_n) = 2^n g(2^{-n}\eta_n) = n.$$

As f is supposed to be continuous, this is impossible.

Note that this example also shows that the space $H(\mathbb{R}^A)$, equipped with the sup-norm over Δ_A , is not complete if A is infinite. This is one of the reasons that we shall use the space $H(\Delta_A)$.

In general, $FVL(A)$ may be identified with a vector sublattice of $H(\mathbb{R}^A)$ (see Theorem 3.6), which in turn, courtesy of Lemma 5.1, may be identified with a vector sublattice of $H(\Delta_A)$ via the restriction map R . This identification extends to $FBL(A)$. The proof of this turns out to be slightly more tricky than might have been anticipated.

For sake of convenience, we denote by $J = J_A$ the restriction to $FVL(A)$ of the restriction map $R : H(\mathbb{R}^A) \rightarrow H(\Delta_A)$. Since $\|J\delta_a\|_\infty = 1$ for all $a \in A$, it is clear that $\|J\| = 1$ and so, $\|Jf\|_\infty \leq \|f\|_F$ for all $f \in FVL(A)$. Since $H(\Delta_A)$ is a Banach lattice with respect to $\|\cdot\|_\infty$, J extends by continuity to a lattice homomorphism $J : FBL(A) \rightarrow H(\Delta_A)$ with $\|J\| = 1$. Note that, by the universal property of $FBL(A)$, J is the unique lattice homomorphism from $FBL(A)$ into $H(\Delta_A)$ satisfying $J\delta_a = \delta_{a|\Delta_A}$, $a \in A$. This implies, in particular, that if B is a non-empty subset of A , then J_B is the restriction of J_A to $FBL(B)$ (cf. Proposition 4.8). The problem is to show that this extension J is injective.

First, we consider the situation that A is finite, in which case everything is very nice indeed.

Proposition 5.3. *For any non-empty finite set A , the map $J : FBL(A) \rightarrow H(\Delta_A)$ is a surjective norm and lattice isomorphism.*

Proof. We claim that $\|f\|_F \leq n \|Jf\|_\infty$, $f \in FVL(A)$, where n is the cardinality of A . Indeed, if $f \in FVL(A)$, then

$$|Jf| \leq \|Jf\|_\infty \bigvee_{a \in A} |J(\delta_a)|$$

so that

$$|f| \leq \|Jf\|_\infty \bigvee_{a \in A} |\delta_a|$$

and hence,

$$\|f\|_F \leq \|Jf\|_\infty \left\| \bigvee_{a \in A} |\delta_a| \right\|_F \leq \|Jf\|_\infty \sum_{a \in A} \|\delta_a\|_F = n \|Jf\|_\infty.$$

This proves the claim. Consequently, $\|Jf\|_\infty \leq \|f\|_F \leq n \|Jf\|_\infty$, $f \in FVL(A)$, which implies that $J : FBL(A) \rightarrow H(\Delta_A)$ is a norm and lattice isomorphism. It remains to be shown that J is surjective. For this purpose, denote by S_A the compact subset of Δ_A given by $S_A = \{\xi \in \Delta_A : \|\xi\|_A = 1\}$. Since A is finite, the restriction map $r : H(\Delta_A) \rightarrow C(S_A)$ is a surjective norm and lattice isomorphism. Since the functions $\{\delta_{a|S_A} : a \in A\}$ separate the points of S_A , it follows via

the Stone-Weierstrass theorem that $(r \circ J)(FBL(A)) = C(S_A)$ and hence $J(FBL(A)) = H(\Delta_A)$. The proof is complete. \square

This norm isomorphism is not an isometry unless $n = 1$. In fact, if $a_1, \dots, a_n \in A$ are distinct, then $\left\| \bigvee_{j=1}^n |\delta_{a_j}| \right\|_F = n$ (indeed, consider the lattice homomorphism $T : FBL(A) \rightarrow \ell_1^n$ satisfying $T(\delta_{a_j}) = e_j$, $1 \leq j \leq n$, where e_j denotes the j -th unit vector in ℓ_1^n).

Sometimes it will be convenient to use the following, slightly weaker, description.

Corollary 5.4. *For any non-empty finite set A , $FBL(A)$ is linearly order isomorphic to $H(\mathbb{R}^A)$.*

Proof. We only need to observe that the restriction map $R : H(\mathbb{R}^A) \rightarrow H(\Delta_A)$ is onto whenever A is finite. \square

To show that the lattice homomorphism $J : FBL(A) \rightarrow H(\Delta_A)$ is injective in general, we will make use of real-valued linear lattice homomorphisms on $FBL(A)$ in the course of proving this and it will later allow us to characterize these in general, which must be worth knowing anyway!

Theorem 5.5. *If A is a non-empty set, then $\omega : FBL(A) \rightarrow \mathbb{R}$ is a lattice homomorphism if and only if there exists $\xi \in \Delta_A$ and $0 \leq \lambda \in \mathbb{R}$ such that $\omega(f) = \lambda Jf(\xi)$ for all $f \in FBL(A)$.*

Proof. If ω is a real valued lattice homomorphism on $FBL(A)$, then it follows from Lemma 3.9 that there is $\eta \in \mathbb{R}^A$ such that $\omega(f) = f(\eta)$, $f \in FVL(A)$. As $FBL(A)$ is a Banach lattice, ω is $\|\cdot\|_F$ -bounded and so, $\sup_{a \in A} |\eta(a)| = \sup_{a \in A} |\omega(\delta_a)| = \|\omega\| < \infty$. Hence, there is a $\lambda = \|\omega\| > 0$ such that $\xi = \lambda^{-1}\eta \in \Delta_A$. If $f \in FVL(A)$, then

$$\omega(f) = f(\eta) = \lambda f(\lambda^{-1}\eta) = \lambda Jf(\xi).$$

Given $f \in FBL(A)$, choose a sequence (g_n) in $FVL(A)$ with $\|f - g_n\|_F \rightarrow 0$, so that $\|Jf - Jg_n\|_\infty \rightarrow 0$ and hence $Jg_n(\xi) \rightarrow Jf(\xi)$. Thus,

$$\omega(f) = \lim_{n \rightarrow \infty} \omega(g_n) = \lambda \lim_{n \rightarrow \infty} Jg_n(\xi) = \lambda Jf(\xi).$$

The converse is clear as if $\xi \in \Delta_A$ and $0 \leq \lambda \in \mathbb{R}$, then the formula $\omega(f) = \lambda Jf(\xi)$, $f \in FBL(A)$, defines a lattice homomorphism on $FBL(A)$. \square

It is clear already that, for $f \in FVL(A)$, $f = 0$ if and only if $Jf = 0$. if and only if $\omega(f) = 0$ for every $\|\cdot\|_F$ -bounded real-valued lattice homomorphism on $FVL(A)$. We need this equivalence for $f \in FBL(A)$.

Corollary 5.6. *For any non-empty set A and $f \in FBL(A)$ the following are equivalent:*

- (i) $f = 0$;
- (ii) $\omega(f) = 0$ for all real-valued lattice homomorphisms on $FBL(A)$;
- (iii) $Jf = 0$.

Proof. Clearly, (i) implies (iii) and that (iii) implies (ii) follows directly from Theorem 5.5.

Now assume that (ii) holds. Note firstly that it follows from Proposition 5.3 that for any non-empty finite subset $B \subseteq A$ the restriction of J to $FBL(B)$ is injective. For such a set B , the map $f \mapsto (JP_B f)(\xi)$, $f \in FBL(A)$, is a real-valued lattice homomorphism on $FBL(A)$ for each $\xi \in \Delta_A$, so that $JP_B f = 0$. As J is injective on $FBL(B)$, this shows that $P_B f = 0$. It follows from Proposition 4.11 that $P_B f \rightarrow f$ for $\|\cdot\|_F$, so that $f = 0$. This suffices to complete the proof. \square

Corollary 5.7. *If A is any non-empty set, then the lattice homomorphism $J : FBL(A) \rightarrow H(\Delta_A)$ is injective, so that $FBL(A)$ is linearly order isomorphic to a vector sublattice of $H(\Delta_A)$.*

V. Troitsky has pointed out to the authors that there is not a similar embedding of $FBL(\mathbb{N})$ into $H(\mathbb{R}^{\mathbb{N}})$. Note also that, although we have no need of the fact, the image of $FBL(A)$ is actually a lattice ideal in $H(\Delta_A)$.

In the sequel, we shall identify $FBL(A)$ with the vector sublattice $J(FBL(A))$ of $H(\Delta_A)$.

As we have seen in Proposition 4.8, if B is a non-empty subset of A , then $FBL(B)$ may be identified isometrically with the closed vector sublattice of $FBL(A)$ generated by $\{\delta_b : b \in B\}$ and there is a canonical contractive lattice homomorphic projection P_B in $FBL(A)$ onto $FBL(B)$. It should be observed that we have the following commutative diagram:

$$\begin{array}{ccc} FBL(A) & \xrightarrow{J_A} & H(\Delta_A) \\ k_B \uparrow & & \uparrow j_B \\ FBL(B) & \xrightarrow{J_B} & H(\Delta_B) \end{array}$$

where j_B is the restriction to $H(\Delta_B)$ of the injective lattice homomorphism J_B introduced in Section 2, and k_B is the isometric lattice embedding of $FBL(B)$ into $FBL(A)$ guaranteed by Proposition 4.8. Note that also j_B is an isometry. The commutativity of the diagram follows by considering the action of the maps on the free generators of $FBL(B)$. Consequently, the canonical embedding of $FBL(B)$ into

$FBL(A)$ is compatible with the canonical embedding of $H(\Delta_B)$ into $H(\Delta_A)$. It can similarly be seen that the following diagram also commutes:

$$\begin{array}{ccc} FBL(A) & \xrightarrow{J_A} & H(\Delta_A) \\ P_B \downarrow & & \downarrow (j_B)^{-1} \circ P_B \\ FBL(B) & \xrightarrow{J_B} & H(\Delta_B) \end{array}$$

The next proposition describes this in terms of $FBL(A)$ considered as a vector sublattice of $H(\Delta_A)$. We consider \mathbb{R}^{Δ_B} as a subspace of \mathbb{R}^{Δ_A} as explained in Section 2.

Recall that if B is a non-empty subset of A , then for any $\xi \in \Delta_A$ we denote by ξ_B the restriction of ξ to B , so that $\xi_B \in \Delta_B$.

Proposition 5.8. *Suppose that B is a non-empty subset of A . Considering $FBL(A)$ as a vector sublattice of $H(\Delta_A)$, we have:*

- (i) *the canonical projection P_B of $FBL(A)$ onto $FBL(B)$ is given by $P_B f(\xi) = f(\xi\chi_B)$, $\xi \in \Delta_A$, for all $f \in FBL(A)$;*
- (ii) *if $f \in FBL(A)$, then a necessary and sufficient condition for f to belong to $FBL(B)$ is that $f(\xi) = f(\eta)$ whenever $\xi, \eta \in \Delta_A$ with $\xi_B = \eta_B$.*

Proof. (i). Let P_B be the canonical projection in $FBL(A)$ onto $FBL(B)$ (see Proposition 4.8), so that $P_B \delta_a = \delta_a$ if $a \in B$ and $P_B \delta_a = 0$ if $a \in A \setminus B$. If $f \in FVL(A)$, then it follows from the observations preceding Proposition 3.7 that $P_B f(\xi) = f(\xi\chi_B)$, $\xi \in \Delta_A$. Given $f \in FBL(A)$, let (f_n) be a sequence in $FVL(A)$ such that $\|f - f_n\|_F \rightarrow 0$, which implies that $\|f - f_n\|_\infty \rightarrow 0$ and so, $f_n(\xi) \rightarrow f(\xi)$, $\xi \in \Delta_A$. Furthermore, $\|P_B f - P_B f_n\|_F \rightarrow 0$ and hence $P_B f_n(\xi) \rightarrow P_B f(\xi)$, $\xi \in \Delta_A$. Since $P_B f_n(\xi) = f_n(\xi\chi_B) \rightarrow f(\xi\chi_B)$, we may conclude that $P_B f(\xi) = f(\xi\chi_B)$, $\xi \in \Delta_A$.

(ii). *Necessity.* If $f \in FBL(B)$ and $\xi, \eta \in \Delta_A$ are such that $\xi_B = \eta_B$, then $\xi\chi_B = \eta\chi_B$ and hence it follows from (i) that

$$f(\xi) = P_B f(\xi) = f(\xi\chi_B) = f(\eta\chi_B) = P_B f(\eta) = f(\eta).$$

Sufficiency. If $f \in FBL(A)$ is such that $f(\xi) = f(\eta)$ whenever $\xi, \eta \in \Delta_A$ with $\xi_B = \eta_B$, then $P_B f(\xi) = f(\xi\chi_B) = f(\xi)$, as $(\xi\chi_B)_B = \xi_B$, for all $\xi \in \Delta_A$ and hence $f = P_B f \in FBL(B)$. \square

Recall that a sublattice H of a lattice L is said to be *regularly embedded* if every subset of H with a supremum (resp. infimum) in H has the same supremum (resp. infimum) in L . If we are dealing with vector lattices it suffices to consider only the case of a subset of H that

is downward directed in H to 0 and check that it also has infimum 0 in L .

Proposition 5.9. *If A is any non-empty set and B is a non-empty subset of A , then $FBL(B)$ is regularly embedded in $FBL(A)$.*

Proof. Suppose that $(f_\gamma)_{\gamma \in \Gamma}$ is a downward directed net in $FBL(B)$ such that $f_\gamma \downarrow_\gamma 0$ in $FBL(B)$ and suppose that $g \in FBL(A)$ satisfies $0 < g \leq f_\gamma$ for all $\gamma \in \Gamma$. Let $\xi_0 \in \Delta_A$ be such that $g(\xi_0) > 0$. We claim that we may assume that $\xi_0 \chi_B \neq 0$. If our chosen ξ_0 is such that $\xi_0 \chi_B = 0$, i.e. $\xi_0 = \xi_0 \chi_{A \setminus B}$, then consider $\xi_\epsilon = \xi_0 + \epsilon \chi_B$. Since $\xi_\epsilon \rightarrow \xi_0$ in Δ_A as $\epsilon \downarrow 0$ and g is continuous we may choose $\epsilon \in (0, 1]$ with $g(\xi_\epsilon) > 0$ and then replace ξ_0 by this ξ_ϵ . Given $b \in B$, define $h \in H(\Delta_A)$ by setting

$$h(\xi) = g \left(\xi \chi_B + \frac{|\xi(b)|}{\|\xi_0 \chi_B\|_\infty} \xi_0 \chi_{A \setminus B} \right), \quad \xi \in \Delta_A.$$

We claim that $h \in FBL(A)$. Indeed, define the lattice homomorphism $T : H(\Delta_A) \rightarrow H(\Delta_A)$ by setting

$$Tf(\xi) = f \left(\xi \chi_B + \frac{|\xi(b)|}{\|\xi_0 \chi_B\|_\infty} \xi_0 \chi_{A \setminus B} \right), \quad \xi \in \Delta_A,$$

for all $f \in H(\Delta_A)$. Observing that

$$T\delta_a = \delta_a \chi_B(a) + \frac{|\delta_b|}{\|\xi_0 \chi_B\|_\infty} \delta_a(\xi_0) \chi_{A \setminus B}(a),$$

it follows that $T\delta_a \in FVL(A)$ for all $a \in A$ and that $\sup_{a \in A} \|T\delta_a\|_F < \infty$. Consequently, there exists a unique lattice homomorphism $S : FBL(A) \rightarrow FBL(A)$ such that $S\delta_a = T\delta_a$ for all $a \in A$. Evidently, $Tf = Sf$ for all $f \in FVL(A)$. Given $f \in FBL(A)$, we may approximate f with a sequence (f_n) with respect to $\|\cdot\|_F$. Using that convergence with respect to $\|\cdot\|_F$ implies pointwise convergence on Δ_A , it follows that $Sf = Tf$ (cf. the proof of Proposition 5.8). This implies, in particular, that $h = Tg = Sg \in FBL(A)$, by which our claim is proved.

If $\xi, \eta \in \Delta_A$ are such that $\xi_B = \eta_B$, then $h(\xi) = h(\eta)$ and so, by Proposition 5.8 and Lemma 2.1, it follows that $h \in FBL(B)$. If $\xi \in \Delta_A$, then

$$\xi_B = \left(\xi \chi_B + \frac{|\xi(b)|}{\|\xi_0 \chi_B\|_\infty} \xi_0 \chi_{A \setminus B} \right)_B$$

(recall that the subscript B indicates taking the restriction to the subset B) and hence

$$\begin{aligned} f_\gamma(\xi) &= f_\gamma\left(\xi\chi_B + \frac{|\xi(b)|}{\|\xi_0\chi_B\|_\infty}\xi_0\chi_{A\setminus B}\right) \\ &\geq g\left(\xi\chi_B + \frac{|\xi(b)|}{\|\xi_0\chi_B\|_\infty}\xi_0\chi_{A\setminus B}\right) = h(\xi), \quad \xi \in \Delta_A, \end{aligned}$$

that is, $f_\gamma \geq h \geq 0$ for all $\gamma \in \Gamma$. We may conclude that $h = 0$.

It follows, in particular, that

$$g\left(\xi_0\chi_B + \frac{|\xi_0(b)|}{\|\xi_0\chi_B\|_\infty}\xi_0\chi_{A\setminus B}\right) = 0, \quad b \in B.$$

Applying this to $b = b_n$, where (b_n) is a sequence in B satisfying $|\xi_0(b_n)| \rightarrow \|\xi_0\chi_B\|_\infty$, the continuity of g implies that

$$g(\xi_0) = g(\xi_0\chi_B + \xi_0\chi_{A\setminus B}) = 0,$$

which is a contradiction. The proof is complete. \square

6. SOME PROPERTIES OF FREE BANACH LATTICES.

If X is a non-empty set and $f : X \rightarrow \mathbb{R}$ then we let $O_f = \{x \in X : f(x) \neq 0\}$ and if W is a non-empty subset of \mathbb{R}^X then we define $O_W = \bigcup\{O_f : f \in W\}$. Although probably well known we know of no convenient reference for the following result.

Proposition 6.1. *If X is a Hausdorff topological space, L a vector sublattice of $C(X)$ and the open set O_L is connected then the only projection bands in L are $\{0\}$ and L .*

Proof. Suppose that B is a projection band in L , so that $L = B \oplus B^d$. If $f \in B$ and $g \in B^d$ then $f \perp g$ and hence $O_f \cap O_g = \emptyset$ and therefore $O_B \cap O_{B^d} = \emptyset$. Given $x \in O_L$ there is $0 \neq f \in L_+$ with $f(x) > 0$. We may write $f = f_1 \oplus f_2$ with $0 \leq f_1 \in B$ and $0 \leq f_2 \in B^d$. Clearly, either $f_1(x) > 0$ or $f_2(x) > 0$. I.e. $x \in O_{f_1} \cup O_{f_2} \subset O_B \cup O_{B^d}$. Hence $O_L \subset O_B \cup O_{B^d}$ and therefore $O_L = O_B \cup O_{B^d}$. The sets O_B and O_{B^d} are both open and disjoint and O_L is, by hypothesis, connected. This is only possible if either O_B or O_{B^d} is empty which says that either $L = B^d$ or $L = B$. \square

Corollary 6.2. *If $|A| \geq 2$ then the only projection bands in $FBL(A)$ are $\{0\}$ and $FBL(A)$.*

Proof. By Corollary 5.7 we may identify $FBL(A)$ with a vector sublattice of $H(\Delta_A) \subset C(\Delta_A)$. Observe that

$$O_{FBL(A)} \supset \bigcup_{a \in A} O_{\delta_a} = \bigcup_{a \in A} \{\xi \in \Delta_A : \xi(a) \neq 0\} = \Delta_A \setminus \{0\}.$$

Clearly, $O_{FBL(A)} \subset \Delta_A \setminus \{0\}$ so that $O_{FBL(A)} = \Delta_A \setminus \{0\}$ which, provided $|A| \geq 2$, is (pathwise) connected. \square

Corollary 6.3. *If $|A| \geq 2$ then $FBL(A)$ is not Dedekind σ -complete.*

Corollary 6.4. *If $|A| \geq 2$ then $FBL(A)$ has no atoms.*

Proof. The linear span of an atom is always a projection band. \square

Corollary 6.5. *If $a \in A$ then $|\delta_a|$ is a weak order unit for $FBL(A)$.*

Proof. If $f \in FBL(A)$ and $f \perp |\delta_a|$ then $O_f \subset \{\xi \in \Delta_A : \xi(a) = 0\}$, and the latter set has an empty interior so that $O_f = \emptyset$ and hence $f = 0$. \square

Corollary 6.6. *Every disjoint system in $FBL(A)$ is at most countable.*

Proof. If $\{u_i : i \in I\}$ is a disjoint family of strictly positive elements of $FBL(A)$ then the corresponding sets O_{u_i} are non-empty disjoint open subsets of Δ_A . As $\Delta_A = [-1, 1]^A$ is a product of separable spaces, Theorem 2 of [19] tells us that Δ_A can contain only countably many disjoint non-empty open sets so that the families of all O_{u_i} and of all u_i are indeed countable. \square

The same result is true for $FVL(A)$, being first proved by Weinberg in [25]. It can also be found, with essentially the current proof, in [1].

Recall that an Archimedean vector lattice is *order separable* if every subset $D \subset L$ contains an at most countable subset with the same upper bounds in L as D has. This is equivalent to every order bounded disjoint family of non-zero elements being at most countable, [13], Theorem 29.3. Corollary 6.6 thus actually tells us that the universal completion of $FBL(A)$, [13], Definition 50.4, is always order separable.

Every Banach lattice is a quotient of a free Banach lattice. We can actually make this statement quite precise. The following lemma is well known dating back, in the case that $\mathfrak{a} = \aleph_0$, to a result of Banach and Mazur [2]. A more accessible proof, again in the case that $\mathfrak{a} = \aleph_0$ (although the modifications needed for the general case are minor), are given as part of the proof of Theorem 5 of Chapter VII of [6].

Lemma 6.7. *Let X be a Banach space and D a dense subset of the unit ball of X . If $x \in X$ and $\|x\| < 1$ then there are sequences (x_n) in D and (α_n) in \mathbb{R} such that $\sum_{n=1}^{\infty} |\alpha_n| < 1$ and $x = \sum_{n=1}^{\infty} \alpha_n x_n$.*

Proposition 6.8. *Let X be a Banach lattice. If D is a dense subset of the unit ball of X of cardinality \mathfrak{a} , then there is a closed ideal J in $FBL(\mathfrak{a})$ such that X is isometrically order isomorphic to $FBL(\mathfrak{a})/J$.*

Proof. Let $D = \{x_a : a \in \mathfrak{a}\}$. By the definition of a free Banach lattice there is a unique contractive lattice homomorphism $T : FBL(\mathfrak{a}) \rightarrow X$ with $T(\delta_a) = x_a$ for each $a \in \mathfrak{a}$. If $x \in X$ with $\|x\| < 1$ then Lemma 6.7 gives us sequences (x_{a_n}) in D and (α_n) in \mathbb{R} with $\sum_{n=1}^{\infty} |\alpha_n| < 1$ and $x = \sum_{n=1}^{\infty} \alpha_n x_{a_n}$. If we define $f \in FBL(\mathfrak{a})$ by $f = \sum_{n=1}^{\infty} \alpha_n \delta_{a_n}$, noting that this series converges absolutely, then $\|f\|_F < 1$ and $Tf = x$. This shows that T maps the open unit ball in $FBL(\mathfrak{a})$ onto the open unit ball in X . In particular, T is surjective.

Take J to be the kernel of T and let $Q : FBL(\mathfrak{a}) \rightarrow FBL(\mathfrak{a})/J$ be the quotient map. Let $U : FBL(\mathfrak{a})/J \rightarrow X$ be defined by $U(Qf) = Tf$ for $f \in FBL(\mathfrak{a})$, which is clearly well-defined. It is also clear that U is a contractive lattice isomorphism. As T maps the open unit ball of $FBL(\mathfrak{a})$ onto the open unit ball of X and Q maps the open unit ball of $FBL(\mathfrak{a})$ onto the open unit ball of $FBL(\mathfrak{a})/J$, it follows that U maps the open unit ball of $FBL(\mathfrak{a})/J$ onto the open unit ball of X so that U is an isometry. \square

Corollary 6.9. *Let X be a Banach lattice. If D is a dense subset of the unit ball of X of cardinality \mathfrak{a} , then $FBL(\mathfrak{a})^*$ contains a weak*-closed band which is isometrically order isomorphic to X^* .*

Proof. If $T : FBL(\mathfrak{a}) \rightarrow X$ is the quotient map from Proposition 6.8 then $T^* : X^* \rightarrow FBL(\mathfrak{a})^*$ is an isometry and its range, which is $\ker(T)^\perp$, is a weak*-closed band. As T is a surjective lattice homomorphism, T^* is actually a lattice isomorphism. \square

In particular note:

Corollary 6.10. *If \mathfrak{a} is any cardinal then there is a weak*-closed band in $FBL(\mathfrak{a})^*$ which is isometrically order isomorphic to $\ell_\infty(\mathfrak{a})$.*

Proof. If \mathfrak{a} is infinite then we need merely note that the unit ball of $\ell_1(\mathfrak{a})$ has a dense subset of cardinality \mathfrak{a} and that $\ell_\infty(\mathfrak{a})$ may be identified with $\ell_1(\mathfrak{a})^*$.

Suppose that $\text{card}(A) = \mathfrak{a}$ is finite. For $a \in A$ we will write ξ_a for that element of $\Delta_A = [-1, 1]^A$ with $\xi_a(a) = 1$ and $\xi_a(b) = 0$ if $a \neq b$. If $b \in A$ then $|\delta_b|(\xi_a) = |\delta_b(\xi_a)| = 1$ if $a = b$ and is zero if $a \neq b$. It follows from the Theorem 5.5 that the functional $f \mapsto f(\xi_a)$ is a lattice homomorphism on $FBL(A)$, and therefore an atom of $FBL(A)^*$, of norm one. Finite sums of such maps also have norm one. This embeds

a copy of $\ell_\infty(A)$ isometrically onto an order ideal in $FBL(A)^*$ which, as it is finite dimensional, is certainly a weak*-closed band. \square

Corollary 6.11. *If X is a separable Banach lattice then X is isometrically order isomorphic to a Banach lattice quotient of $FBL(\aleph_0)$ and X^* is isometrically order isomorphic to a weak*-closed band in $FBL(\aleph_0)^*$.*

This illustrates quite effectively what a rich structure free Banach lattices and their duals have. For example if X and Y are separable Banach lattices such that no two non-zero bands in X^* and Y^* are isometrically isomorphic then the isometrically order isomorphic bands in $FBL(\aleph_0)^*$ must be disjoint in the lattice theoretical sense. So, for example, we have:

Corollary 6.12. *In $FBL(\aleph_0)^*$ there are mutually disjoint weak*-closed bands A and B_p ($p \in (1, \infty]$) with B_p isometrically order isomorphic to $L_p([0, 1])$ and A to ℓ_∞ .*

This gives continuum many disjoint non-zero elements in $FBL(\aleph_0)^*$, which should be contrasted with Corollary 6.6.

7. THE STRUCTURE OF FINITELY GENERATED FREE BANACH LATTICES.

We will see shortly that $FBL(n)$ is not an AM-space unless $n = 1$, but it does have a lot of AM-structure provided that n is finite.

If we have only a finite number of generators, n , say then we may identify $FBL(n)$ with $H(\Delta_n)$, where Δ_n is now a product of n copies of $[-1, 1]$. In this setting, it might be more useful to consider the restriction of these homogeneous functions to the union of all the proper faces of Δ_n , which we will denote by F_n . An alternative description of this set is that it is the points in \mathbb{R}^n with supremum norm equal to 1. Each of the generators δ_k ($1 \leq k \leq n$) takes the value +1 on one maximal proper face of F_n of dimension $n - 1$ and the value -1 on the complementary face. These faces exhaust the maximal proper faces of Δ_n . The restriction map from $H(\Delta_n)$ to $C(F_n)$ is a surjective vector lattice isomorphism and an isometry from the supremum norm over Δ_n to the supremum norm over F_n . We know also that these norms are equivalent to the free norm. Thus when we identify $FBL(n)$ with $C(F_n)$, even though the norms are not the same, the closed ideals, band, quotients etc remain the same so that we can read many structural results off from those for $C(K)$ spaces. Whenever we refer to the free norm on $C(F_n)$ we refer to the free norm generated using the generators which take value ± 1 on the maximal proper faces.

In particular, we may identify the dual of $FBL(n)$ with the space of regular Borel measures on F_n , $\mathcal{M}(F_n)$. We will see in Theorem 8.1 that unless $n = 1$ the dual of the free norm, $\|\cdot\|^\dagger$ is definitely not the usual norm, $\|\cdot\|_1$, under which $\mathcal{M}(F_n)$ is an AL-space. However, there remains a lot of AL-structure in this dual.

Proposition 7.1. *If $\mu \in \mathcal{M}(F_n)$ is supported by a maximal proper face of Δ_n then $\|\mu\|^\dagger = \|\mu\|_1$.*

Proof. Suppose first that $\mu \geq 0$. Let the free generators be denoted by $\delta_1, \delta_2, \dots, \delta_n$. If G is the maximal proper face in question, we may suppose that $G \subset \delta_1^{-1}(1)$. As $|\delta_k| \leq 1$ on F_n , for $1 \leq k \leq n$, we have

$$\int |\delta_k| d\mu \leq \int \mathbf{1} d\mu = \|\mu\|_1,$$

and on taking the maximum we have $\|\mu\|^\dagger \leq \|\mu\|_1$. On the other hand, $|\delta_1| \equiv 1$ on G so that

$$\|\mu\|^\dagger \geq \int |\delta_1| d\mu = \|\mu\|_1,$$

so we have equality. Both $\|\cdot\|_1$ and $\|\cdot\|^\dagger$ are lattice norms, so in the general case we have

$$\|\mu\|^\dagger = \|\|\mu\|^\dagger\|_1 = \|\mu\|_1$$

and the proof is complete. \square

Corollary 7.2. *If $f \in C(F_n)$ and there is a maximal proper face G such that f vanishes off G then $\|f\|_F = \|f\|_\infty$.*

Proof. If $\mu \in \mathcal{M}(F_n)$ then we may write $\mu = \mu_G + \mu_{F_n \setminus G}$, where $\mu_A(X) = \mu(A \cap X)$, and note that $\int f d\mu = \int f d\mu_G$. If $\|\mu\|^\dagger \leq 1$ then $\|\mu_G\|^\dagger = \|\mu_G\|_1 \leq 1$ as $|\mu_G| \leq |\mu|$. Thus

$$\begin{aligned} \|f\|_F &= \sup\left\{ \int |f| d|\mu| : \|\mu\|^\dagger \leq 1 \right\} \\ &\leq \sup\left\{ \int |f| d|\mu| : \|\mu\|_1 \leq 1 \right\} = \|f\|_\infty \end{aligned}$$

and the embedding of $FBL(n)$ into $H(\Delta_n)$ is a contraction so that $\|f\|_F \geq \|f\|_\infty$. \square

This means that certain closed ideals in $FBL(n)$ are actually AM-spaces, namely those that may be identified with functions on F_n which vanish on a closed set A whose complement is contained in a single proper face of F_n . Rather more interesting is an analogous result for quotients.

In general, if J is a closed ideal in a Banach lattice X then $(X/J)^*$ may be identified, both in terms of order and norm, with the ideal $J^\circ = \{f \in X^* : f|_J \equiv 0\}$. We know that if A is a closed subset of a compact Hausdorff space K and J^A denotes the closed ideal $J^A = \{f \in C(K) : f|_A \equiv 0\}$ then when $C(K)$ is given the supremum norm the normed quotient $C(K)/J^A$ is isometrically order isomorphic to $C(A)$ under its supremum norm and its dual is isometrically order isomorphic to the space of measures on K which are supported by A . In the particular case that $K = F_n$ we may still identify quotients algebraically in the same way, but the description of the quotient norm has to be modified slightly. That means that the quotient norm may be described in a similar manner to our original description of the free norm:

Proposition 7.3. *If A is a closed subset of F_n and $C(F_n)$ is normed by its canonical free norm then $C(F_n)/J^A$ is isometrically order isomorphic to $C(A)$, where $C(A)$ is normed by*

$$\|f\|_A = \sup\left\{ \int |f| d|\mu_A| : \|\mu\|^\dagger \leq 1 \right\}.$$

In this supremum we may restrict to measures μ supported by A .

In particular we have, using Proposition 7.1:

Corollary 7.4. *If A is a closed subset of a proper face of F_n and $C(F_n)$ is normed by its canonical free norm then $C(F_n)/J^A$ is isometrically order isomorphic to $C(A)$ under its supremum norm.*

The free vector lattices over a finite number of generators exhibit a lot of symmetry. For example it is not difficult to see that $FVL(n)$ is invariant under rotations. In studying symmetry of $FBL(n)$ it makes things clearer to identify $FBL(n)$ with the space $C(S^{n-1})$ rather than $C(F_n)$, where S^{n-1} is the Euclidean unit sphere in \mathbb{R}^n , even though the description of the free norm is made slightly more difficult. In the case $n = 2$, we are looking at continuous functions on the unit circle and the dual free norm is given by

$$\|\mu\|^\dagger = \int_{S^1} |\sin(t)| d|\mu|(t) \vee \int_{S^1} |\cos(t)| d|\mu|(t).$$

In particular, if η_x denotes the unit measure concentrated at x then

$$\|\eta_x\|^\dagger = |\sin(x)| \vee |\cos(x)|$$

which is certainly not rotation invariant. Note also that

$$\|\eta_x + \eta_{x+\pi/2}\|^\dagger = (|\sin(x)| + |\sin(x+\pi/2)|) \vee (|\cos(x)| + |\cos(x+\pi/2)|).$$

In fact only rotations through multiples of $\pi/2$ are isometries on $C(S^1)$ for the free norm. Of course, all rotations of $FBL(n)$ will be isomorphisms.

There is an obvious procedure for obtaining a rotation invariant norm from the free norm, namely to take the average, with respect to Haar measure on the group of rotations, of the free norms of rotations of a given element. Although this will certainly not be the free norm, given that it is derived in a canonical manner from the free norm we might expect that either it is a familiar norm or else is of some independent interest. It turns out not to be familiar. This is again easiest to see in the dual.

If we denote this symmetric free norm by $\|\cdot\|_S$ and its dual norm by $\|\cdot\|^S$ then we have

$$\|\eta_x\|^S = \|\eta_{x+\pi/2}\|^S = \frac{1}{2\pi} \int_0^{2\pi} |\sin(t)| \vee |\cos(t)| dt = \frac{2\sqrt{2}}{\pi}$$

and

$$\begin{aligned} \|\eta_x + \eta_{x+\pi/2}\|^S &= \frac{1}{2\pi} \int_0^{2\pi} (|\sin(x)| + |\sin(x + \pi/2)|) \vee (|\cos(x)| + |\cos(x+\pi/2)|) dt \\ &= \frac{4}{\pi} \end{aligned}$$

so that the symmetric free norm is not an AL-norm, which is the natural symmetric norm on $C(S^1)^*$, nor an AM-norm. In fact $\|\eta_x + \eta_{x+t}\|^S$ can take any value between $\frac{4}{\pi}$ and $\frac{4\sqrt{2}}{\pi}$ so the symmetric free norm cannot be any L^p norm either, implausible though that would be anyway.

8. CHARACTERIZING THE NUMBER OF GENERATORS.

Apart from wanting to understand how the number of generators affects the Banach lattice structure of $FBL(A)$, we would like to know when $FBL(A)$ is a classical Banach lattice or has various properties generally considered desirable. The answer to this is “not very often”! It turns out that such properties can be used to characterize the number of generators, at least in a rather coarse manner.

In fact several properties that are normally considered “good” are only possessed by a free Banach lattice if it has only one generator. We gather several of these into our first result. We know that in the finitely generated case, both $FBL(n)$ and its dual have a certain amount of AM-structure. There is another area of Banach lattice theory where the same is true, namely in injective Banach lattices in the category

of Banach lattices and contractive positive operators, see [10]. As injective Banach lattices are certainly Dedekind complete we cannot have $FBL(n)$ being injective if $n > 1$. It might be thought possible that $FBL(A)^*$ was injective, but that also turns out to be false unless $|A| = 1$.

Theorem 8.1. *If A is a non-empty set then the following are equivalent:*

- (1) $|A| = 1$.
- (2) $FBL(A)$ is isometrically an AM-space.
- (3) $FBL(A)$ is isomorphic to an AL-space.
- (4) Every bounded linear functional on $FBL(A)$ is order continuous.
- (5) There is a non-zero order continuous linear functional on $FBL(A)$.
- (6) $FBL(A)^*$ is an injective Banach lattice.

Proof. If A is a singleton then $\Delta_A = [-1, 1]$ and $FBL(A)$ may be identified with $H(\Delta_A)$ which in turn may be identified with \mathbb{R}^2 . The generator is the pair $g = (-1, 1)$. The positive linear functionals ϕ such that $\phi(|g|) \leq 1$ are those described by pairs of reals (ϕ_1, ϕ_2) with $|\phi_1| + |\phi_2| \leq 1$. The free norm that they induce on \mathbb{R}^2 is precisely the supremum norm.

If $|A| > 1$ then by Corollary 6.10 $FBL(A)^*$ contains an order isometric copy of $\ell_\infty(A)$ so is not an AL-space and therefore $FBL(A)$ is not an AM-space. This establishes that (1) \Leftrightarrow (2).

It is clear that (1) \Rightarrow (3) although even in this case it is clear that $FBL(1)$ is not isometrically an AL-space. $FBL(2)$, on the other hand is isomorphic to continuous functions on a square so is certainly not isomorphic to an AL-space. In view of Proposition 4.8 and the fact that every closed sublattice of an AL-space is itself an AL-space we see that (3) \Rightarrow (1).

It is clear that (1) \Rightarrow (4) \Rightarrow (5). To show that (5) \Rightarrow (1), suppose that $|A| > 1$ and that ϕ is a non-zero order continuous linear functional on $FBL(A)$. By continuity of ϕ and density of $FVL(A)$ in $FBL(A)$, $\phi|_{FVL(A)} \neq 0$. Similarly, as $FVL(A) = \bigcup\{FVL(F) : F \subseteq A, |F| < \infty\}$ we may choose a finite subset $F \subseteq A$ with $\phi|_{FVL(F)} \neq 0$ so certainly $\phi|_{FBL(F)} \neq 0$. Without loss of generality, as long as $|A| > 1$ we may assume that $|F| > 1$. As $FBL(F)$ is regularly embedded in $FBL(A)$, by Proposition 5.9, $\phi|_{FBL(F)}$ is order continuous. As a vector lattice, we may identify $FBL(F)$ with $C(S_F)$, where S_F is the ℓ_∞ unit sphere in Δ_F . Certainly S_F is a dense in itself, metrizable (and hence separable) compact Hausdorff space so it follows from Proposition 19.9.4 of [20] that $\phi|_{FBL(F)} = 0$, contradicting our original claim.

Certainly $FBL(1)^*$, being an AL-space, is injective, [12] Proposition 3.2. We know from Corollary 4.10 that if $|A| > 1$ then $FBL(2)^*$ is isometrically order isomorphic to a (projection) band in $FBL(A)^*$. If $FBL(A)^*$ were injective then certainly $FBL(2)^*$ would also be injective. Recall that Proposition 3G of [10] tells us that an injective Banach lattice either contains a sublattice isometric to ℓ_∞ , or else is isometrically isomorphic to a finite AM-direct sum of AL-spaces. We know that $FBL(2)$ is order and norm isomorphic to continuous functions on the square F_2 so that $FBL(2)^*$ is norm and order isomorphic to the space of measures on F_2 so certainly has an order continuous norm. Thus it does not contain even an isomorphic copy of ℓ_∞ by Corollary 2.4.3 of [14], so it certainly suffices to show that $FBL(2)^*$ cannot be decomposed into a non-trivial finite AM-direct sum of bands of any nature.

The dual of $FBL(2)$ can be identified, as a vector lattice, with the regular Borel measures on F_2 . The dual free norm amounts to $\|\mu\| = \max\{\int |\delta_1| d|\mu|, \int |\delta_2| d|\mu|\}$, where δ_i is the projection onto the i 'th coordinate. It is clear that $\int |\delta_1| d|\mu| = 0$ if and only if μ is supported by $S_1 = \{\langle 0, -1 \rangle, \langle 0, 1 \rangle\}$ whilst $\int |\delta_2| d|\mu| = 0$ if and only if μ is supported by $S_2 = \{\langle -1, 0 \rangle, \langle 1, 0 \rangle\}$. If any non-trivial AM-decomposition of $FBL(2)^*$ were possible, into $J \oplus K$ (say), then we can pick $0 \neq \mu \in J_+$ and $0 \neq \nu \in K_+$. We may assume that $\|\mu\| = \|\nu\| = 1$ and therefore $\|\mu + \nu\| = 1$. The fact that $\|\mu\| = \|\nu\| = 1$ means that

$$\int |\delta_1| d\mu \vee \int |\delta_2| d\mu = \int |\delta_1| d\nu \vee \int |\delta_2| d\nu = 1.$$

Suppose that $\int |\delta_1| d\mu = \int |\delta_1| d\nu = 1$, then we have $1 = \|\mu + \nu\| \geq \int |\delta_1| d(\mu + \nu) = \int |\delta_1| d\mu + \int |\delta_1| d\nu = 2$, which is impossible. Similarly, we cannot have $\int |\delta_2| d\mu = \int |\delta_2| d\nu = 1$. If $\int |\delta_1| d\mu = \int |\delta_2| d\nu = 1$ then the fact that $1 = \|\mu + \nu\| \leq \int |\delta_1| d(\mu + \nu)$ tells us that $\int |\delta_1| d\nu = 0$ so that ν is supported by S_1 . Similarly we see that $\int |\delta_2| d\mu = 0$ so that μ is supported by S_2 . This implies that $FBL(2)^*$ is supported by $S_1 \cup S_2$ which is impossible. A similar contradiction arises if $\int |\delta_2| d\mu = \int |\delta_1| d\nu = 1$. □

It is already clear that free Banach lattices on more than one generator are not going to be amongst the classical Banach lattices. Isomorphism with AM-spaces is still possible and turns out to determine whether or not the number of generators is finite.

Theorem 8.2. *If A is any non-empty set then the following are equivalent:*

- (1) A is finite.
- (2) $FBL(A)$ is isomorphic to $H(\Delta_A)$ under the supremum norm.
- (3) $FBL(A)$ has a strong order unit.
- (4) $FBL(A)$ is isomorphic to an AM-space.
- (5) $FBL(A)^*$ has an order continuous norm.

Proof. We have already seen that (1) \Rightarrow (2) \Rightarrow (3). It is well known and simple to prove that (3) \Rightarrow (4). That (4) \Rightarrow (5) is because the dual of an AM-space is an AL-space which has an order continuous norm and the fact that order continuity of the norm is preserved under (not necessarily isometric) isomorphisms. In order to complete the proof we need only prove that (5) \Rightarrow (1).

If A is infinite then $FBL(A)^*$ contains a weak*-closed band that is isometrically order isomorphic to ℓ_∞ , by Corollary 6.10. By Theorem 2.4.14 of [14] this is equivalent to $FBL(\mathfrak{a})^*$ not having an order continuous norm (and to many other conditions as well.) \square

In a similar vein, we can characterize, amongst free Banach lattices, those with a countable number of generators. Before doing so, though, we note that once there are infinitely many generators then there is an immediate connection between the number of generators and the cardinality of dense subsets. Perhaps not entirely unexpectedly, given Corollary 6.6, the same result holds for order intervals. Recall that the *density character* of a topological space is the least cardinal of a dense subset.

Theorem 8.3. *If \mathfrak{a} is an infinite cardinal then the following conditions on a set A are equivalent:*

- (1) $\text{card}(A) = \mathfrak{a}$.
- (2) $FBL(A)$ has density character \mathfrak{a} .
- (3) The smallest cardinal \mathfrak{b} such that every order interval in $FBL(A)$ has density character at most \mathfrak{b} is \mathfrak{a} .

Proof. Let $\mathfrak{a} = \text{card}(A)$, \mathfrak{b} be the density character of $FBL(A)$ and \mathfrak{c} the smallest cardinal which is at least as large as the density character of every order interval in $FBL(A)$. We need to show that $\mathfrak{a} = \mathfrak{b} = \mathfrak{c}$.

The free vector lattice over \mathbb{Q} with \mathfrak{a} many generators has cardinality precisely \mathfrak{a} , given that \mathfrak{a} is infinite. That is dense in $FVL(A)$ and hence in $FBL(A)$ for the free norm, so $\mathfrak{b} \leq \mathfrak{a}$. Clearly $\mathfrak{c} \leq \mathfrak{b}$. let K be a compact Hausdorff space such that the smallest cardinality of a dense subset of $C(K)$, and hence of the unit ball in $C(K)$, is \mathfrak{a} . For example we could take $K = [0, 1]^\mathfrak{a}$. There is a bounded lattice homomorphism $T : FBL(A) \rightarrow C(K)$ which maps the generators of A onto a dense subset of the unit ball of $C(K)$. The proof of Proposition 6.8 shows that

T is onto. Let $\mathbf{1}_K$ denote the constantly one function on K . The order interval $[-T^{-1}\mathbf{1}_K, T^{-1}\mathbf{1}_K]$ has a dense subset of cardinality at most \mathfrak{c} . As T is a surjective lattice homomorphism, $T([-T^{-1}\mathbf{1}_K, T^{-1}\mathbf{1}_K]) = [-\mathbf{1}_K, \mathbf{1}_K]$, and this will have a dense subset of cardinality at most \mathfrak{c} . Hence $\mathfrak{a} \leq \mathfrak{c}$. This establishes that $\mathfrak{a} = \mathfrak{b} = \mathfrak{c}$. \square

For the statement of the next result which characterizes a free Banach lattice having countably many generators, we need to recall some definitions. A *topological order unit* e of a Banach lattice E is an element of the positive cone such that the closed order ideal generated by e is the whole of E . These are also referred to as *quasi-interior points*. Separable Banach lattices always possess topological order units. The *centre* of E , $Z(E)$, is the space of all linear operators on E lying between two real multiples of the identity. The centre is termed *topologically full* if whenever $x, y \in E$ with $0 \leq x \leq y$ here is a sequence (T_n) in $Z(E)$ with $T_n y \rightarrow x$ in norm. If E has a topological order unit then its centre is topologically full. At the other extreme there are AM-spaces in which the centre is *trivial*, i.e. it consists only of multiples of the identity.

Theorem 8.4. *If A is a non-empty set, then the following are equivalent:*

- (1) *A is finite or countably infinite.*
- (2) *$FBL(A)$ is separable.*
- (3) *Every order interval in $FBL(A)$ is separable.*
- (4) *$FBL(A)$ has a topological order unit.*
- (5) *$Z(FBL(A))$ is topologically full.*
- (6) *$Z(FBL(A))$ is non-trivial.*

Proof. If A is finite then it follows from the isomorphism seen in Theorem 8.2 that $FBL(A)$, and hence its order intervals, is separable. Combining this observation with the preceding theorem shows that (1), (2) and (3) are equivalent.

We noted earlier that separable Banach lattices always have a topological order unit. The fact that Banach lattices with a topological order unit have a topologically full centre is also widely known, but finding a complete proof in the literature is not easy. The earliest is in Example 1 of [16], but that proof is more complicated than it need be. A simpler version is in Proposition 1.1 of [28] and see also Lemma 1 of [17].

Even if $\mathfrak{a} = 1$, $FBL(\mathfrak{a})$ is not one-dimensional so that if the centre is topologically full then it cannot be trivial.

We know from Proposition 5.7 that we may identify $FBL(A)$ with a sublattice of $H(\Delta_A)$. It is clear, as it contains the coordinate projections, that it separates points of Δ_A . If $|A|$ is uncountable then $\{0\}$ is not a G_δ subset of Δ_A . It follows from Theorem 3.1 of [26] that the centre of this sublattice, and therefore of $FBL(\mathfrak{a})$, is then trivial. \square

Corollary 8.5. *If A is an uncountable set then $FBL(A)$ has trivial centre.*

Note that this would seem to be the first “natural” example of a Banach lattice with a trivial centre. If $\mathfrak{a} > 1$ then $FVL(A)$ always has trivial centre. The details are left to the interested reader.

9. LIFTING DISJOINT FAMILIES IN QUOTIENT BANACH LATTICES.

In [24], Weinberg asked what were the projective objects in the category of abelian ℓ -groups, pointing out, for example, that a summand of a free ℓ -group was projective. Topping studied projective vector lattices in [23] but the reader should be warned that Theorem 8, claiming that countable positive disjoint families in quotients L/J of vector lattices L lift to positive disjoint families in L , is false. In fact that is only possible for an Archimedean Riesz space if the space is a direct sum of copies of the reals, see [5] and [15].

Later in this paper we will study projective Banach lattices, which are intimately connected with quotient spaces. We will need to know when disjoint families in a quotient Banach lattice X/J can be lifted to disjoint families in X . As this is a question of considerable interest in its own right and also because the results that we need do not seem to be in the literature already, we present them in a separate section here.

It is well-known, although we know of no explicit reference, that any finite disjoint family $(y_k)_{k=1}^n$ in a quotient Riesz space X/J can be lifted to a disjoint family $(x_k)_{k=1}^n$ in X with $Qx_k = y_k$, where $Q : X \rightarrow X/J$ is the quotient map.

Proposition 9.1. *If X is a vector lattice, J a vector lattice ideal in X , $Q : X \rightarrow X/J$ the quotient map and $(y_k)_{k=1}^n$ is a disjoint family in X/J then there is a disjoint family $(x_k)_{k=1}^n$ in X with $Qx_k = y_k$ for $1 \leq k \leq n$.*

Proof. It suffices to consider the case that each $y_k \geq 0$. The proof will be by induction, the case $n = 1$ being trivial. Assume the result is true for $n = m$ and we verify it for $n = m + 1$. If $(y_k)_{k=1}^{m+1}$ is a disjoint non-negative family in X/J we may find $(\tilde{x}_k)_{k=1}^{m+1}$ in X with $Q\tilde{x}_k = y_k$ ($1 \leq k \leq m + 1$) and $\tilde{x}_j \perp \tilde{x}_k$ for $j \neq k$ and $1 \leq j, k \leq m$. Let

$x_k = \tilde{x}_k = \tilde{x}_k \wedge \tilde{x}_{m+1}$, for $1 \leq k \leq m$, and $x_{m+1} = \tilde{x}_{m+1}$. Then for $i \leq k \leq m$ we have

$$Qx_k = Q\tilde{x}_k - Q\tilde{x}_k \wedge Q(\tilde{x}_{m+1} = y_k - y_k \wedge y_{m+1} = y_k).$$

Clearly if $1 \leq k \leq m$ then $x_k \perp x_{m+1}$ whilst if $j \neq k$ and $1 \leq j, k \leq m$ then $0 \leq x_j \wedge x_k \leq \tilde{x}_j \tilde{x}_k = 0$. This establishes the result for $n = m + 1$. \square

If we restrict attention to norm closed ideals in Banach lattices then, unlike the vector lattice case, we can handle countably infinite disjoint liftings, but not larger ones. This does not contradict the vector lattice result cited above as there are many non-closed ideals in a Banach lattice.

Theorem 9.2. *If X is a Banach lattice, J a closed ideal in X , $Q : X \rightarrow X/J$ the quotient map and $(y_k)_{k=1}^\infty$ is a disjoint sequence in X/J then there is a disjoint sequence (x_k) in X with $Qx_k = y_k$ for all $k \in \mathbb{N}$.*

Proof. It suffices to consider the case that each $y_n \geq 0$ and $\|\sum_{k=1}^\infty y_k\| < \infty$. Define $z_n = \sum_{k=n+1}^\infty y_k \in X/J$ and note that z_n is disjoint from y_1, \dots, y_n . The sequence (x_n) will be constructed inductively.

For $n = 1$ we start by choosing $x_1, u_1 \in X$ with $x_1 \perp u_1$, $Qx_1 = y_1$ and $Qu_1 = z_1$ using the Proposition 9.1.

Now suppose that we have constructed a disjoint system $\{x_1, \dots, x_n, u_n\}$ with $Qx_j = y_j$ ($1 \leq j \leq n$) and $Qu_n = z_n$. As $0 \leq y_{n+1} \perp z_{n+1}$, there are disjoint $\tilde{x}_{n+1}, \tilde{u}_{n+1} \in X_+$ with $Q\tilde{x}_{n+1} = y_{n+1}$ and $Q\tilde{u}_{n+1} = z_{n+1}$. Let $x_{n+1} = \tilde{x}_{n+1} \wedge u_n$ and $u_{n+1} = \tilde{u}_{n+1} \wedge u_n$ so that, for example, $Qx_{n+1} = Q\tilde{x}_{n+1} \wedge Qu_n = y_{n+1} \wedge z_n = y_{n+1}$. Obviously, $x_{n+1} \perp u_{n+1}$ whilst if $1 \leq k \leq n$ we have, for example, $0 \leq x_k \wedge u_{n+1} \leq x_k \wedge u_n = 0$. \square

Even in Banach lattices, Theorem 9.2 is as far as we can go.

Example 9.3. Given any uncountable disjoint family in a Banach lattice X , we know from Proposition 6.8 that there is a free Banach lattice $FBL(\mathfrak{a})$ and a closed ideal J in $FBL(\mathfrak{a})$ such that X is isometrically order isomorphic to $FBL(\mathfrak{a})/J$. As a disjoint family in a free Banach lattice has to be countable, Corollary 6.6, the disjoint family cannot possibly be lifted to $FBL(\mathfrak{a})$.

A slightly more concrete example may be found using Problem 6S of [9] where it is shown that $\beta\mathbb{N} \setminus \mathbb{N}$ contains continuum many disjoint non-empty open and closed subsets. I.e. ℓ_∞/c_0 contains continuum many non-zero disjoint positive elements. As ℓ_∞ contains only countably many disjoint elements, we cannot possibly lift each of this continuum of disjoint elements in ℓ_∞/c_0 to disjoint elements in ℓ_∞ . The same will be true of any uncountable subset of these disjoint positive

elements of ℓ_∞/c_0 , so this shows that lifting of disjoint positive families of cardinality \aleph_1 is not possible.

An apparently simpler problem is to start with two subsets A and B in X/J with $A \perp B$ and seek subsets A', B' of X with $A' \perp B'$, $Q(A') = A$ and $Q(B') = B$. Again countability is vital to the success of this attempt, in fact it allows us to do much more.

Proposition 9.4. *If X is a Banach lattice, J a closed ideal in X , $Q : X \rightarrow X/J$ the quotient map and $(A_n)_{n=1}^\infty$ be a sequence of countable subsets of X/J with $A_m \perp A_n$ if $m \neq n$ then there are subsets (B_n) of X , with $B_m \perp B_n$ if $m \neq n$ and $Q(B_n) = A_n$ for each $n \in \mathbb{N}$.*

Proof. As above, there is no loss of generality in assuming that each $A_n \subset (X/J)_+$. Enumerate each set as $A_n = \{a_k^n : k \in \mathbb{N}\}$ (there is no difference, apart from notation, if one or both set is finite). Let $v_n = \sum_{k=1}^\infty a_k^n / (2^k \|a_k\|)$ so that $v_m \perp v_n$ if $m \neq n$ and $0 \leq a_k^n \leq 2^k \|a_k\| v_n$ for $k, n \in \mathbb{N}$. We know from Theorem 9.2 that there is a disjoint sequence (u_n) in X_+ with $Q(u_n) = v_n$. For any $a_k^n \in A_n$ we can find $c_k^n \in X_+$ with $Q(c_k^n) = a_k^n$. Now set $b_k^n = c_k^n \wedge (2^k \|a_k\| u_n)$ so that we still have

$$Q(b_k^n) = Q(c_k^n) \wedge (2^k \|a_k\| Q(u_n)) = a_k^n \wedge (2^k \|a_k\| v_n) = a_k^n.$$

Also each $b_k^n \in u_n^\perp$ so that if $m \neq n$ then for any choice of j and k we see that $b_j^m \perp b_k^n$ as $u_m \perp u_n$. Now defining $B_n = \{b_k^n : k \in \mathbb{N}\}$ gives the required sets. \square

Considering the case of singleton sets, the example above shows that we cannot allow an uncountable number of disjoint families. Nor can we allow even one of the families to be uncountable.

In the case that $X = C(K)$, for K a compact Hausdorff space, a closed ideal J is of the form $F = \{f \in C(K) : f|_A \equiv 0\}$ for some closed subset $A \subset K$ and the quotient X/J may be identified with $C(A)$ in the obvious manner. For two elements $f, g \in C(K)$, $f \perp g$ if and only if the two sets $f^{-1}(\mathbb{R} \setminus \{0\})$ and $g^{-1}(\mathbb{R} \setminus \{0\})$ are disjoint.

Example 9.5. The *Tychonoff plank* K is the topological space $[0, \omega] \times [0, \omega_1] \setminus \{(\omega, \omega_1)\}$ where ω is the first infinite ordinal and ω_1 the first uncountable ordinal. This is renowned as an example of a non-normal Hausdorff space. The sets $U = [0, \omega) \times \{\omega_1\}$ and $V = \{\omega\} \times [0, \omega_1)$ are disjoint closed subsets which cannot be separated by disjoint open sets. See for example §8.20 of [9]. If we add back in the removed corner point, and define $A = U \cup V \cup \{(\omega, \omega_1)\}$ then U and V become open subsets of A .

Each point of U is isolated so their characteristic functions lie in $C(A)$ giving a (countable) family F with $U = \bigcup \{f^{-1}(\mathbb{R} \setminus \{0\}) : f \in F\}$. Let

G be a family of functions in $C(A)$ such that $V = \bigcup\{g^{-1}(\mathbb{R} \setminus \{0\}) : g \in G\}$, which is certainly possible using Urysohn's lemma. If these could be lifted to disjoint families L and M in $C(K)$ then $\bigcup\{f^{-1}(\mathbb{R} \setminus \{0\}) : f \in L\}$ and $\bigcup\{f^{-1}(\mathbb{R} \setminus \{0\}) : f \in M\}$ would be disjoint open subsets of K which intersected A in the disjoint open sets U and V .

But any disjoint open subsets of the whole product space which intersected A in U and V respectively would, with the corner point removed if necessary, separate the closed sets U and V in the plank. This contradiction shows that the lifting is not possible.

10. PROJECTIVE BANACH LATTICES.

Definition 10.1. A Banach lattice P is *projective* if whenever X is a Banach lattice, J a closed ideal in X and $Q : X \rightarrow X/J$ the quotient map then for every linear lattice homomorphism $T : P \rightarrow X/J$ and $\epsilon > 0$ there is a linear lattice homomorphism $\hat{T} : P \rightarrow X$ such that

- (1) $T = Q \circ \hat{T}$,
- (2) $\|\hat{T}\| \leq (1 + \epsilon)\|T\|$.

Even if we take $P = \mathbb{R}$, which is easily seen to be projective given this definition, it is clear that we cannot replace $1 + \epsilon$ by 1 as the quotient norm is an infimum which need not be attained. There are projective Banach lattices, because:

Proposition 10.2. *A free Banach lattice is projective.*

Proof. Let $(\delta_a)_{a \in \mathfrak{a}}$ be the generators of the free Banach lattice F . Suppose that X is a Banach lattice, J a closed ideal in X , $Q : X \rightarrow X/J$ the quotient map, $T : F \rightarrow X/J$ a lattice homomorphism and $\epsilon > 0$. For each $a \in \mathfrak{a}$, there is $x_a \in X$ with $Qx_a = T\delta_a$ and $\|x_a\| \leq (1 + \epsilon)\|T\delta_a\| \leq (1 + \epsilon)\|T\|$, using the definition of the quotient norm. As F is free there is a linear lattice homomorphism $\hat{T} : F \rightarrow X$ with $\hat{T}\delta_a = x_a$ for all $a \in \mathfrak{a}$ and $\|\hat{T}\| \leq \sup\{\|x_a\| : a \in \mathfrak{a}\} \leq (1 + \epsilon)\|T\|$. As $(Q \circ \hat{T})\delta_a = T\delta_a$ for all $a \in \mathfrak{a}$ and both $Q \circ \hat{T}$ and T are linear lattice homomorphisms they must coincide on the vector lattice generated by the δ_a and, by continuity, on F . \square

We can characterize projective Banach lattices in a reasonably familiar manner.

Theorem 10.3. *The following conditions on a Banach lattice P are equivalent.*

- (1) P is projective.
- (2) For all $\epsilon > 0$ there are:

- (a) a free Banach lattice F ,
 - (b) a closed sublattice H of F and a lattice isomorphism $\mathcal{J} : H \rightarrow P$ with $\|\mathcal{J}\|, \|\mathcal{J}^{-1}\| \leq 1 + \epsilon$, and
 - (c) a lattice homomorphism projection $R : F \rightarrow H$ with $\|R\| \leq 1 + \epsilon$.
- (3) For all $\epsilon > 0$ there are:
- (a) a projective Banach lattice F ,
 - (b) a closed sublattice H of F and a lattice isomorphism $\mathcal{J} : H \rightarrow P$ with $\|\mathcal{J}\|, \|\mathcal{J}^{-1}\| \leq 1 + \epsilon$, and
 - (c) a lattice homomorphism projection $R : F \rightarrow H$ with $\|R\| \leq 1 + \epsilon$.

Proof. To see that (1) \Rightarrow (2), suppose that P is projective, let F be a free Banach lattice and J a closed ideal in F such that P is isometrically order isomorphic to the quotient F/J via the linear lattice isomorphism $\mathcal{J} : P \rightarrow F/J$, which is always possible using Proposition 6.8. Let $Q : F \rightarrow F/J$ be the quotient map. As P is projective, for any $\epsilon > 0$ there is a linear lattice homomorphism $\hat{\mathcal{J}} : P \rightarrow F$ with $Q \circ \hat{\mathcal{J}} = \mathcal{J}$ and $\|\hat{\mathcal{J}}\| \leq (1 + \epsilon)\|\mathcal{J}\| = 1 + \epsilon$. As $Q \circ \hat{\mathcal{J}}$ is injective, $\hat{\mathcal{J}}$ is also injective and $\hat{\mathcal{J}}P$ is a closed sublattice of F as $\|\hat{\mathcal{J}}p\| \geq \|Q(\hat{\mathcal{J}}p)\| = \|\mathcal{J}p\| = \|p\|$. The map $\hat{\mathcal{J}} \circ \mathcal{J}^{-1} \circ Q$ is a lattice homomorphism which projects F onto $\hat{\mathcal{J}}(P)$ and $\|\hat{\mathcal{J}} \circ \mathcal{J}^{-1} \circ Q\| \leq \|\hat{\mathcal{J}}\| \leq 1 + \epsilon$, so (2)(b) holds. We know that $\|\hat{\mathcal{J}}\| \leq 1 + \epsilon$ and $\hat{\mathcal{J}}^{-1} = \mathcal{J}^{-1} \circ Q$ so that $\|\hat{\mathcal{J}}^{-1}\| = 1$ and (2)(c) holds.

In view of Proposition 10.2, clearly (2) \Rightarrow (3).

Suppose that (3) holds, and in particular that (a), (b) and (c) hold for the real number η . Suppose that X is any Banach lattice, J a closed ideal in X , $Q : X \rightarrow X/J$ the quotient map, $\eta > 0$ and that $T : P \rightarrow X/J$ is a linear lattice homomorphism. The map $T \circ \mathcal{J} \circ R : F \rightarrow X/J$ is also a linear lattice homomorphism with $\|T \circ \mathcal{J} \circ R\| \leq \|T\|\|\mathcal{J}\|\|R\| \leq (1 + \eta)^2\|T\|$. As F is projective there is a linear lattice homomorphism $S : F \rightarrow X$ with $Q \circ S = T \circ \mathcal{J} \circ R$ and $\|S\| \leq (1 + \eta)\|T \circ \mathcal{J} \circ R\| \leq (1 + \eta)^3\|T\|$. Now let $\hat{T} = S \circ \mathcal{J}^{-1} : P \rightarrow X$, which is also a linear lattice homomorphism, so that $\|\hat{T}\| \leq \|S\|\|\mathcal{J}^{-1}\| \leq (1 + \eta)^4\|T\|$ and

$$Q \circ \hat{T} = Q \circ (S \circ \mathcal{J}^{-1}) = (Q \circ S) \circ \mathcal{J}^{-1} = (T \circ \mathcal{J} \circ R) \circ \mathcal{J}^{-1} = T.$$

By choosing η small enough we can ensure that $(1 + \eta)^4 \leq 1 + \epsilon$ and we have shown that P is projective. \square

In particular, in light of Corollary 6.11, all the separable projective Banach lattices that we produce later will (almost) embed in $FBL(\aleph_0)$ reinforcing the richness of its structure.

Combining Theorem 10.3 with Corollary 5.6 we have:

Corollary 10.4. *The real-valued lattice homomorphisms on a projective Banach lattice separate points.*

In particular this tells us that, for finite p , the Banach lattice $L_p([0, 1])$ is *not* projective.

Similarly, from Corollary 6.6 and Theorem 10.3, using the lattice homomorphism projection from a free Banach lattice onto a projective, we see:

Corollary 10.5. *Every disjoint system in a projective Banach lattice is at most countable.*

Although, in a sense, Theorem 10.3 gives a complete description of projective Banach lattices, given that we know little about free Banach lattices it actually tells us very little. One immediate consequence, given that $FBL(1)$ may be identified with $\ell_\infty(2)$, is:

Corollary 10.6. *The one dimensional Banach lattice \mathbb{R} is projective.*

Of course, this is easy to verify directly, but it does show that there are projective Banach lattices which are not free.

Let us note also one rather simple consequence of the characterization of projectives in Theorem 10.3.

Corollary 10.7. *If X is a projective Banach lattice, H a closed sublattice of X for which there is a contractive lattice homomorphism projecting X onto H , then H is a projective Banach lattice.*

11. WHICH BANACH LATTICES ARE PROJECTIVE?

We will now approach matters from the other end. We try to find out as much as we can about projective Banach lattices and deduce information about the structure of free Banach lattices. We will start by identifying some “small” Banach lattices, apart from free ones, which are projective. After that we will show that certain AL-sums of projectives are again projective.

Our first positive result may be slightly surprising, given that when dealing with Banach spaces the free and projective objects are precisely the spaces $\ell_1(I)$, [20], Theorem 27.4.2.

Theorem 11.1. *Every finite dimensional Banach lattice is projective.*

Proof. Let P be a finite dimensional Banach lattice, X an arbitrary Banach lattice, J a closed ideal in X , $Q : X \rightarrow X/J$ the quotient map, $T : P \rightarrow X/J$ a lattice homomorphism and $1 \geq \epsilon > 0$. We identify P with \mathbb{R}^n with the pointwise order and normed by some lattice norm $\|\cdot\|_P$. Without loss of generality we may assume that the standard

basic vectors in \mathbb{R}^n , e_k , all have $\|e_k\|_P = 1$. Let $\{p_k : 1 \leq k \leq m\}$ be an ϵ -net for the compact set $\{p \in \mathbb{R}_+^n : \|p\|_P = 1\}$. We write $p_k = (p_k^1, p_k^2, \dots, p_k^n)$.

As T is a lattice homomorphism, the family $(Te_k)_{k=1}^n$ is a disjoint family in $(X/J)_+$ so by Proposition 9.1 there is a disjoint family $(s_k)_{k=1}^n$ in X_+ with $Qs_k = Te_k$ for $1 \leq k \leq n$. By the definition of the quotient norm, for each k there is $t_k \in X$ with $Qt_k = Te_k$ and $\|t_k\| \leq \|Te_k\| + \epsilon \leq \|T\| + \epsilon$. Now, let $x_k = s_k \wedge t_k^+$, so that the family (x_k) remains disjoint. As Q is a lattice homomorphism, $Qx_k = Qs_k \wedge Qt_k^+ = (Te_k) \wedge (Te_k)^+ = Te_k$. Also, we now have $\|x_k\| \leq \|t_k^+\| \leq \|t_k\| \leq \|T\| + \epsilon$.

Also, for each $i \in \{1, 2, \dots, m\}$ there is $q_i \in X_+$ with $Qq_i = Tp_i$ and $\|q_i\| \leq \|Tp_i\| + \epsilon \leq \|T\| + \epsilon$.

Define $z_k = x_k \wedge \bigwedge_{i=1}^m (p_i^k)^{-1} q_i$ where the $'$ indicates that terms where $p_i^k = 0$ are omitted. As the family (x_k) is disjoint, the same is true for the family (z_k) . If $p_i^k > 0$ then $(p_i^k)^{-1} p_i \geq e_k$ so that $(p_i^k)^{-1} Qq_i = (p_i^k)^{-1} Tp_i \geq Te_k$ so that $Qz_k = Qx_k = Te_k$.

Define $Se_k = z_k$ and extend S linearly to a lattice homomorphism (because the (z_k) are disjoint) of $\mathbb{R}^n \rightarrow X$. Clearly $Q \circ S_k = T$. As \mathbb{R}^n is finite dimensional, there is a constant $K \in \mathbb{R}_+$ such that $\|x\|_1 \leq K\|x\|_P$ for all $x \in \mathbb{R}^n$. It follows that

$$\begin{aligned} \left\| S \left(\sum_{k=1}^n \lambda_k e_k \right) \right\| &\leq \sum_{k=1}^n |\lambda_k| \|Se_k\| \\ &= \sum_{k=1}^n |\lambda_k| \|z_k\| \\ &\leq \sum_{k=1}^n |\lambda_k| \|x_k\| \\ &\leq \left\| \sum_{k=1}^n \lambda_k e_k \right\| (\|T\| + 1) \\ &\leq K(\|T\| + 1) \left\| \sum_{k=1}^n \lambda_k e_k \right\|_P \end{aligned}$$

so that $\|S\| \leq K(\|T\| + 1)$. Note that this estimate is independent of the choice of ϵ .

In order better to estimate the norm of S , we write $p_i = \sum_{k=1}^n p_i^k e_k$ and see that

$$\begin{aligned} Sp_i &= \sum_{k=1}^n S(p_i^k e_k) \\ &= \sum_{k=1}^n p_i^k S e_k \\ &= \sum_{k=1}^n p_i^k z_k. \end{aligned}$$

Also, if $p_i^k = 0$ then certainly $p_i^k z_k \leq q_i$, whilst if $p_i^k > 0$ then $p_i^k z_k \leq p_i^k (p_i^k)^{-1} q_i = q_i$. As $p_i^j z_j \perp p_i^k z_k$ if $j \neq k$ we see that $\sum_{k=1}^n p_i^k z_k \leq q_i$ so that $Sp_i \leq q_i$ and $\|Sp_i\| \leq \|q_i\| \leq \|T\| + \epsilon$. Now if we take an arbitrary $p \in \{P_+ : \|p\| = 1\}$ then we can choose i with $\|p - p_i\|_P < \epsilon$, so that

$$\begin{aligned} \|Sp\| &\leq \|Sp_i\| + \|S\| \|p - p_i\|_P \\ &\leq \|T\| + \epsilon + K(\|T\| + 1)\epsilon \end{aligned}$$

which can be made as close to $\|T\|$ as we desire. \square

The spaces $C(K)$, for K a compact Hausdorff space, play a distinguished rôle in the general theory of Banach lattices so it is worth knowing which $C(K)$ spaces are projective. We give here a partial answer, which is already of substantial interest. We refer the reader to [4] for basic concepts about retracts, but include the basic definitions here for the convenience of the reader.

Definition 11.2. If X is a topological space and K a subset of X then

- (1) K is a *retract* of X if there is a continuous function $\pi : X \rightarrow K$ with $\pi(k) = k$ for all $k \in K$.
- (2) K is a *neighbourhood retract* of X if there is a neighbourhood U of K in X and a continuous function $\sigma : U \rightarrow K$ with $\sigma(k) = k$ for all $k \in K$.

Definition 11.3. In a category \mathcal{C} of topological spaces,

- (1) A space K is an *absolute retract* if K is a retract of X whenever $K \subseteq X \in \mathcal{C}$.
- (2) A space K is an *absolute neighbourhood retract* if K is a neighbourhood retract of X whenever $K \subseteq X \in \mathcal{C}$.

Theorem 11.4. *If K is a compact subset of \mathbb{R}^n for some $n \in \mathbb{N}$ then the following are equivalent:*

- (1) $C(K)$ is a projective Banach lattice under some norm.

- (2) $C(K)$ is projective under the supremum norm.
- (3) K is a neighbourhood retract of \mathbb{R}^n .

Proof. Without loss of generality we may suppose that K is a subset of the unit ball in \mathbb{R}^n for the supremum norm. We write p_k for the restriction to K of the k 'th coordinate projection in \mathbb{R}^n and p_0 for the constantly one function on K . The vector sublattice generated by the $\{p_k : 0 \leq k \leq n\}$ is certainly dense in $C(K)$ by the Stone-Weierstrass theorem. As $FBL(n+1)$ is free there is a bounded vector lattice homomorphism $T : FBL(n+1) \rightarrow C(K)$ with $T(\delta_k) = p_{n-1}$. We know that, algebraically, we may identify $FBL(n+1)$ with $C(F_{n+1})$ and that the constantly one function on F_{n+1} is precisely $\bigvee_{k=1}^{n+1} |\delta_k|$. As $\bigvee_{k=0}^n |p_k| = p_0$, here is where we use the boundedness assumption on K , we may regard T as a unital lattice homomorphism from $C(F_{n+1})$ to $C(K)$. Such maps are of the form $f \mapsto f \circ \phi$ where $\phi : K \rightarrow F_{n+1}$ is continuous. The image of $C(F_{n+1})$ is dense in $C(K)$ and it is well known that the image of such composition maps is closed so that T is onto. This is equivalent to ϕ being injective. I.e. we have a topological embedding of K into F_{n+1} and we may regard T as simply being the restriction map from $C(F_{n+1})$ to $C(K)$. So far we have not used the assumption that $C(K)$ is projective.

If J is the kernel of T then $C(F_{n+1})/J$ is isomorphic to $C(K)$. If $C(K)$ is projective (even in a purely algebraic sense) then there is a vector lattice homomorphism $U : C(K) \rightarrow C(F_{n+1})$ with $Uf|_K = f$ for all $f \in C(K)$. But U is of the form

$$Uf(p) = \begin{cases} w(p)f(\pi p) & (p \in U) \\ 0 & (p \notin U) \end{cases}$$

where w is a non-negative continuous real-valued function on F_{n+1} and $\pi : F_{n+1} \setminus w^{-1}(0) \rightarrow K$, so we must have $w(p) = 1$ and $\pi p = p$ for $p \in K$. Thus $F_{n+1} \setminus w^{-1}(0)$ is open and contains K so that π is a neighbourhood retract of F_{n+1} onto K . If we remove any single point from F_{n+1} that is not in K then what remains is homeomorphic to \mathbb{R}^n so we have a neighbourhood retraction from \mathbb{R}^n onto K . This only fails to be possible if $K = F_{n+1}$, and that is not homeomorphic to a subset of \mathbb{R}^n by the Borsuk-Ulam Theorem, see for example Theorem 5.8.9 of [22]. Thus (1) implies (3).

Clearly (2) implies (1), so we need only prove that (3) implies (2). The blanket assumption on K tells us that it is homeomorphic to a subset of one face G of F_{n+1} . By scaling it if necessary, we may assume that it is a neighbourhood retract of G and therefore of the whole of F_{n+1} . That allows us to construct a continuous $w : F_{n+1} \rightarrow [0, 1]$ with

$U = \{p \in F_{n+1} : w(p) > 0\} \subset G$, $K \subset w^{-1}(1)$ and a continuous retract $\pi : U \rightarrow K$. The vector lattice homomorphism $U : C(F_{n+1}) \rightarrow C(F_{n+1})$ defined by

$$Uf(p) = \begin{cases} w(p)f(\pi p) & (p \in U) \\ 0 & (p \notin U) \end{cases}$$

is certainly a projection. For any $p \in F_{n+1}$ we have, writing $J^K = \{f \in C(F_{n+1}) : f|_K \equiv 0\}$,

$$\begin{aligned} \|Uf\|_F &= \|Uf\|_\infty && \text{(Corollary 7.2)} \\ &= \sup\{|w(p)f(\pi p)| : p \in U\} \\ &\leq \sup\{|f(\pi p)| : p \in U\} \\ &= \sup\{|f(k)| : k \in K\} = \|f|_K\|_\infty \\ &= \|f + J^K\| && \text{(Corollary 7.4)} \\ &\leq \|f\|_F \end{aligned}$$

so that U is a contraction.

We claim also that the image $UC(F_{n+1})$ is isometrically order isomorphic to $C(K)$ under its supremum norm. To prove this, it suffices to prove that $Uf \mapsto f|_K$ is an isometry for the free norm on Uf , which is equal to its supremum norm, and the supremum norm on $f|_K$. The calculation above shows that $\|Uf\|_\infty \leq \|f|_K\|_\infty$. We also have, for $p \in U$, $|Uf(p)| = |w(p)||f(\pi p)| \leq \|f|_K\|_\infty$ as $|w(p)| \leq 1$ and $\pi p \in K$. Thus $\|Uf\|_\infty \leq \|f|_K\|_\infty$ and we have our desired isometry.

In view of Theorem 10.3, this shows that $C(K)$ is projective. \square

The reader will notice that the first implication would actually work for an isomorphic version of projectivity. We allude further to this in §12.

Corollary 11.5. *$C([0, 1])$ under the usual supremum norm is a projective Banach lattice.*

Notice that some $C(K)$ -spaces can be projective for different (necessarily equivalent) Banach lattice norms. E.g. $C(F_n)$ will be projective both under the free and supremum norms.

Recall that, as closed bounded convex subsets of \mathbb{R}^n are absolute retracts in the category of compact Hausdorff spaces, any compact neighborhood retract of \mathbb{R}^n will necessarily be an absolute neighbourhood retract in the category of compact Hausdorff spaces and therefore certainly in the category of compact metric spaces.

Descriptions of absolute neighbourhood retracts in the category of compact metric spaces may be found in Chapter V of [4]. We note two

particular properties that they have. Firstly, absolute neighbourhood retracts have only finitely many components ([4], V.2.7) and if K is an absolute neighbourhood retract subset of \mathbb{R}^n then $\mathbb{R}^n \setminus K$ has only finitely many components ([4], V.2.20).

In particular, we have

Corollary 11.6. *The sequence space c is not projective.*

Proof. We can identify c with $C(K_0)$ where $K_0 = \{\frac{1}{n} : n \in \mathbb{N}\} \cup \{0\}$. As $K_0 \subset \mathbb{R}$ and K_0 has infinitely many components it is not an absolute neighbourhood retract. \square

There seems little hope of removing the assumption of finite dimensionality from K in Theorem 11.4. We can rescue one implication.

Proposition 11.7. *If $C(K)$ is a projective Banach lattice under the supremum, or an equivalent, norm then K is an absolute neighbourhood retract in the category of compact Hausdorff spaces.*

Proof. Suppose that K is a closed subset of a compact Hausdorff space X . We need to show that there is a continuous retraction π of U onto K , where U is an open subset of X with $K \subset U$.

The restriction map $R : C(X) \rightarrow C(K)$ may be identified with the canonical quotient map of $C(X)$ onto $C(X)/J$ where J is the closed ideal $\{f \in C(X) : f|_K \equiv 0\}$. If $C(K)$ is projective then the identity on $C(K)$ lifts to a lattice homomorphism $T : C(K) \rightarrow C(X)$ with $R \circ T = I_{C(K)}$. There is a continuous function w from X into \mathbb{R}_+ and a continuous map $\pi : U = \{x \in X : w(x) > 0\} \rightarrow K$ such that

$$Tf(x) = \begin{cases} w(x)f(\pi x) & [w(x) > 0] \\ 0 & [w(x) = 0]. \end{cases}$$

If $k \in K$ then $Tf(k) = f(k)$ so that $\pi k = k$ and $w(k) = 1$ showing that $K \subset U$ and that π is a retraction of the open set U onto K . \square

Without knowledge of the properties of absolute neighbourhood retracts in the category of compact Hausdorff spaces, this does not tell us a lot. There seems to be very little material in the literature on absolute neighbourhood retracts in this setting, so we make our own modest contribution here.

Lemma 11.8. *If C is a compact convex subset of a locally convex space, K a closed subset of C and U an open subset of C with $K \subseteq U \subset C$ then there is an open set V with $K \subseteq V \subseteq U \subseteq C$ such that V has finitely many components.*

Proof. As U is open, if $k \in K$ there is a convex (and therefore connected) open set W_k with $k \in W_k \subseteq U$, using local convexity. The open sets W_k , for $k \in K$, cover the compact set K so there is a finite subcover, W_1, W_2, \dots, W_n . Take $V = \bigcup_{k=1}^n W_k$. \square

Proposition 11.9. *If K is an absolute neighbourhood retract in the category of compact Hausdorff spaces then K has only finitely many components.*

Proof. Let $C = P(K)$, the space of probability measures on K , with the weak* topology induced by $C(K)$, which is a locally convex topology under which C is compact as well as certainly being convex. The mapping which takes k to the point mass at k is a homeomorphism of K onto the set of extreme points of C . If K is an absolute neighbourhood retract then there is a retraction $\pi : U \rightarrow K$ where U is an open subset of C with $K \subseteq U$. By the preceding lemma, there is an open set V , with finitely many components, such that $K \subseteq V \subseteq U$. The image of each component of V under π is connected and their union is K , so that K has only finitely many components. \square

Thus if $C(K)$ is a projective Banach lattice under any norm then K has only finitely many components. In particular:

Corollary 11.10. *The sequence space ℓ_∞ is not a projective Banach lattice.*

In [1] Baker characterized projective vector lattices with n generators as being quotients of $FVL(n)$ by a principal ideal. If we embed K_0 into one of the faces of F_2 then we know that c is isometrically order isomorphic to $FBL(2)/J^{K_0}$. It is clear that J^{K_0} is a principal closed ideal of $FBL(2)$ and that c has two generators as a Banach lattice, so the natural analogue of Baker's result fails in the Banach lattice setting.

The obvious candidate for a projective Banach lattice, as in the Banach space case, is $\ell_1(I)$ for an arbitrary index set I , however Corollary 10.5 tells us that if I is an uncountable index set then $\ell_1(I)$ is definitely *not* a projective Banach lattice. Similarly $\ell_p(I)$ ($1 \leq p < \infty$) and $c_0(I)$ are not projective if I is uncountable.

Given that we can lift disjoint sequences it is not difficult to show that ℓ_1 is projective. In fact we can show much more.

Theorem 11.11. *If, for each $n \in \mathbb{N}$, P_n is a projective Banach lattice with a topological order unit then the countable sum $\ell_1(P_n)$, under the coordinate-wise order and normed by $\|(p_n)\|_1 = \sum_{n=1}^{\infty} \|p_n\|$, is a projective Banach lattice.*

Proof. Let e_n be a topological order unit for P_n . We will identify P_n with the subspace of $\ell_1(P_n)$ in which all entries apart from the n 'th are zero and e_n with the corresponding member of that subspace so that the e_n are all disjoint. If X is a Banach lattice, J a closed ideal in X , $Q : X \rightarrow X/J$ the quotient map, $T : \oplus_1(P_n) \rightarrow X/J$ a lattice homomorphism and $\epsilon > 0$ then we start by noting that the Te_n are disjoint, so by Theorem 9.2 we can find disjoint u_n in X_+ with $Qu_n = Te_n$. If we write X_n for the closed ideal in X generated by u_n then the family (X_n) is disjoint in X .

Note that the natural embedding of $X_n/(J \cap X_n)$ into X/J is an isometry onto an ideal and that $T(P_n) \subset X_n/(J \cap X_n)$ as e_n is a topological order unit for P_n and T is a lattice homomorphism. The projectivity of P_n allows us to lift T_n to a lattice homomorphism $\hat{T}_n : P_n \rightarrow X_n$ with $\|\hat{T}_n\| \leq \|T_n\| + \epsilon \leq \|T\| + \epsilon$ with $Q \circ \hat{T}_n = T_n$. Piecing together this sequence of operators in the obvious way will give us the desired lifting of T . \square

Recall that if \mathfrak{a} is finite or countably infinite then $FBL(\mathfrak{a})$ has a topological order unit as do finite dimensional Banach lattices and $C(K)$ -spaces. This gives us a source of building blocks to create other projectives.

We already have some examples of Banach lattices which are not projective. It is interesting to note that the free Banach lattices on uncountably many generators seem to be, in some sense at least, maximal projectives.

Example 11.12. *If \mathfrak{a} is uncountable then there is no non-zero Banach lattice X for which $X \oplus FBL(\mathfrak{a})$ is projective under any norm.*

Proof. Suppose that, under some norm, $FBL(\mathfrak{a}) \oplus X$ is projective, where X is a Banach lattice and \mathfrak{a} is uncountable.

Consider $C(K)$, where $K = [0, \omega] \times [0, \omega_1]$, and (with the notation of Example 9.5) $J = \{f \in C(K) : f|_A \equiv 0\}$ so that $C(K)/J$ is isometrically order isomorphic to $C(A)$.

For each $v \in V$ there is $f_v \in C(A)$ with $0 \leq f_v(a) \leq 1$ for all $a \in V$, $f_v(v) = 1$ and f_v identically zero on $A \setminus V$. As V has cardinality \aleph_1 there will be a map of the set of generators $\{\delta_a : a \in \mathfrak{a}\}$ of $FBL(\mathfrak{a})$ onto $\{f_v : v \in V\}$, which extends to a lattice homomorphism of $FBL(\mathfrak{a})$ into $C(A)$. The image of every generator vanishes on U , hence the same is true for elements of $T(FVL(\mathfrak{a}))$ and, by continuity, for elements of $T(FBL(\mathfrak{a}))$. Note that $\bigcup_{f \in FBL(\mathfrak{a})} \{a \in A : f(a) \neq 0\} = V$.

As U is an F_σ there is $g \in C(A)$ with $g(u) > 0$ for all $u \in U$ and with g identically zero on $A \setminus U$. If $X \oplus FBL(\mathfrak{a})$ were projective and

$x_0 \in X_+ \setminus \{0\}$ there would be a real-valued lattice homomorphism on $X \oplus FBL(\mathfrak{a})$ with $\phi(x_0) > 0$ (and necessarily $\phi|_{FBL(\mathfrak{a})} \equiv 0$.) Define $Sx = \phi(x)g$ for $x \in X$ so that S is a lattice homomorphism of X into $C(A)$. The disjointness of the images of $S(X)$ and $T(FBL(\mathfrak{a}))$ shows that the direct sum operator $S \oplus T : X \oplus FBL(\mathfrak{a}) \rightarrow C(A) = C(K)/J$ is also a lattice homomorphism. If $X \oplus FBL(\mathfrak{a})$ were projective we could find a lattice homomorphism $\hat{S} \oplus \hat{T} : X \oplus FBL(\mathfrak{a}) \rightarrow C(K)$ with $Q \circ (\hat{S} \oplus \hat{T}) = S \oplus T$. The images of $X \oplus \{0\}$ and $\{0\} \oplus FBL(\mathfrak{a})$ will be disjoint in $C(K)$ and their open supports will give disjoint open sets with traces on A equal to U and V respectively, which we know is impossible. \square

The family of projective Banach lattices seems to possess very few stability properties beyond those that we have already noted. In particular, closed sublattices of projectives need not be projective as the non-projective c may be isometrically embedded as a closed sublattice of the projective Banach lattice $C([0, 1])$, by mapping the sequence (a_n) to the function that is linear on each interval $[1/(n+1), 1/n]$ and takes the value a_n at $1/n$. Similarly, we may realize c as the quotient of $C([0, 1])$ by the closed ideal $\{f \in C([0, 1]) : f(1/n) = 0 \forall n \in \mathbb{N}\}$, showing that the class of projective Banach lattices is not closed under quotients.

12. SOME OPEN PROBLEMS.

We start with a few questions on free Banach lattices.

Question 12.1. Must the norm on a free Banach lattice be Fatou, or even Nakano? See [27] for the definition of a Nakano norm. We are not sure of the answer even when there are only finitely many generators.

The following question is rather a long shot as we have very little evidence for it beyond the case of a finite number of generators (see below).

Question 12.2. If the free Banach lattice $FBL(\mathfrak{a})$ is embedded as a closed ideal in a Banach lattice must it be a projection band?

The reason that this holds in the case of a finite number of generators is because this (isomorphic) property of Banach lattices is possessed by Banach lattices with a strong order unit. The following is undoubtedly well-known but we know of no convenient reference for it.

Proposition 12.3. *Let Y be a Banach lattice with the property that every upward directed norm bounded subset of Y_+ is bounded above. If*

Y is embedded as a closed ideal in a Banach lattice *X* then it must be a projection band.

Proof. It suffices to prove that if $x \in X_+$ then the set $B = \{y \in Y : 0 \leq y \leq x\}$ has a supremum in *Y*. As *B* is upward directed and norm bounded, it has an upper bound $u \in Y_+$. As $u \wedge x \in Y_+$, since *Y* is an ideal, $u \wedge x$ is an upper bound for *B* in *Y*. As we also have $0 \leq u \wedge x \leq x$, $u \wedge x \in B$ so it is actually the maximum element of *B*. \square

We have seen that, unless $|A| = 1$, $FBL(A)^*$ is not an injective Banach lattice. However, in the case of finite *A*, $FBL(A)^*$ is isomorphic to an AL-space and therefore to an injective Banach lattice. We suspect that the following question might lead to another characterization of finitely generated free Banach lattices.

Question 12.4. When is $FBL(\mathfrak{a})^*$ isomorphic to an injective Banach lattice?

In Theorem 8.3 we showed that the density character of $FBL(A)$ was equal to the cardinality of *A* and related this to the density character or order intervals in $FBL(A)$. This is something of importance in the study of regular operators between Banach lattices, so an answer to the following question would have implications in that field.

Question 12.5. Does every order interval in $FBL(A)$ have the same density character?

In the light of Theorem 8.3 that density character would have to be the cardinality of *A*.

Question 12.6. Investigate the structure of the symmetric free norm on $FBL(n)$.

Question 12.7. Can the construction of a free Banach lattice be generalized to give a free Banach lattice over a metric space? Here a metric space *S* embeds in a “free” Banach lattice in some sense and any isometry of the generators into a Banach lattice extends to a lattice homomorphism with some restriction on the norm. See [18] for the Banach space case.

We have seen in Corollary 6.10 that $FBL(\mathfrak{a})^*$ contains a disjoint family of cardinality \mathfrak{a} which contrasts strongly with the fact that disjoint families in $FBL(A)$ itself can only be at most countably infinite.

Question 12.8. How large can disjoint families of non-zero elements in $FBL(\mathfrak{a})^*$ be?

At present we have no feel at all for what kinds of Banach lattice are likely to be projective. Clearly, there are a lot of “small” ones, where small means either separable or having a topological order unit. A major and obvious question to pose is:

Question 12.9. Determine the structure of the class of projective Banach lattices.

In particular,

Question 12.10. Are separable atomic Banach lattices with an order continuous norm projective?

In particular,

Question 12.11. Is c_0 , under the supremum norm, a projective Banach lattice?

Question 12.12. For what compact Hausdorff spaces K is $C(K)$ projective under the supremum norm?

We know the answer to the preceding question for compact subsets of \mathbb{R}^n by Theorem 11.4.

The following two questions were posed by G. Buskes. An apparently simple question to answer is:

Question 12.13. If P_k ($1 \leq k \leq n$) are projective Banach lattices with topological order units then is their ℓ_∞ sum also projective?

It is not difficult to lift a lattice homomorphism $T : \bigoplus_{k=1}^n nP_k \rightarrow Y/J$ to a lattice homomorphism $\hat{T} : \bigoplus_{k=1}^n P_k \rightarrow Y$ by lifting the images of the topological order units first. The problem seems to be the norm condition on \hat{T} .

It is clear that the Fremlin tensor product, see [8], of two projective Banach lattices need not be projective in general. Example 11.12 shows that this cannot be true for the product of ℓ_1 and $FBL(\mathfrak{a})$ when \mathfrak{a} is uncountable. There seems no good structural reason to expect a positive result to the next question, but a counterexample has eluded us so far.

Question 12.14. If X and Y are projective Banach lattices with topological order units, is their Fremlin tensor product projective?

The building blocks that we can use in Theorem 11.11 to build new projectives include finite dimensional spaces, $FBL(\mathfrak{a})$ for \mathfrak{a} either finite or countably infinite and certain $C(K)$ -spaces. Any of these, and the space that is produced by that theorem, will be separable and hence will have a topological order unit. Some (possibly rather rash) conjectures that we might make are:

Conjecture 12.15. If a projective Banach lattice has a topological order unit then it is separable.

Conjecture 12.16. A projective Banach lattice which does not have a topological order unit must be free.

Even if this conjecture were to fail, we can ask for an improvement of Example 11.12 by asking:

Conjecture 12.17. If \mathfrak{a} is uncountable and a projective Banach lattice X contains a closed ideal isomorphic to $FBL(\mathfrak{a})$, do we actually have $X = FBL(\mathfrak{a})$?

Question 12.18. The ℓ_1 sum of a sequence of finite dimensional Banach lattices is a Dedekind complete projective. Are these the only Dedekind (σ -)complete projectives?

Conjecture 12.19. All order continuous functionals on a projective Banach lattice determined by its atoms.

Question 12.20. Assuming a positive answer to Question 12.10, we can further ask if there is a result similar to Theorem 11.11 for ℓ_p sums ($1 < p < \infty$) or for c_0 sums.

The whole of this paper has been written in an isometric setting. All of our results may be reproved in an isomorphic setting, where we replace an (almost) isometric condition on operators with mere norm boundedness. It is not difficult to see that there will automatically be uniform bounds to the norms of operators and that isometrically free (resp. projective) Banach lattices will be isomorphically free (resp. projective). Isomorphically free Banach lattices will certainly be isomorphic to isometrically free Banach lattices. At present it does not seem worth recording such a theory, unless there is negative answer to the following question.

Question 12.21. Is every isomorphically projective Banach lattice isomorphic to an isometrically projective Banach lattice?

REFERENCES

- [1] Kirby A. Baker, *Free vector lattices*, *Canad. J. Math.* **20** (1968), 58–66. MR0224524 (37 #123)
- [2] S. Banach and S. Mazur, *Zuhr Theorie der linearen Dimension*, *Studia. Math.* **4** (1933), 100–112.
- [3] Roger D. Bleier, *Free vector lattices*, *Trans. Amer. Math. Soc.* **176** (1973), 73–87. MR0311541 (47 #103)
- [4] Karol Borsuk, *Theory of retracts*, *Monografie Matematyczne, Tom 44*, Państwowe Wydawnictwo Naukowe, Warsaw, 1967. MR0216473 (35 #7306)

- [5] Paul F. Conrad, *Lifting disjoint sets in vector lattices*, *Canad. J. Math.* **20** (1968), 1362–1364. MR0232720 (38 #1043)
- [6] Joseph Diestel, *Sequences and series in Banach spaces*, Graduate Texts in Mathematics, vol. 92, Springer-Verlag, New York, 1984. MR737004 (85i:46020)
- [7] Nelson Dunford and Jacob T. Schwartz, *Linear operators. Part I*, Wiley Classics Library, John Wiley & Sons Inc., New York, 1988. General theory; With the assistance of William G. Bade and Robert G. Bartle; Reprint of the 1958 original; A Wiley-Interscience Publication. MR1009162 (90g:47001a)
- [8] D. H. Fremlin, *Tensor products of Banach lattices*, *Math. Ann.* **211** (1974), 87–106. MR0367620 (51 #3862)
- [9] Leonard Gillman and Meyer Jerison, *Rings of continuous functions*, The University Series in Higher Mathematics, D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto-London-New York, 1960. MR0116199 (22 #6994)
- [10] Richard Haydon, *Injective Banach lattices*, *Math. Z.* **156** (1977), no. 1, 19–47. MR0473776 (57 #13438)
- [11] Joram Lindenstrauss and Lior Tzafriri, *Classical Banach spaces. II*, *Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas]*, vol. 97, Springer-Verlag, Berlin, 1979. Function spaces. MR540367 (81c:46001)
- [12] Heinrich P. Lotz, *Extensions and liftings of positive linear mappings on Banach lattices*, *Trans. Amer. Math. Soc.* **211** (1975), 85–100. MR0383141 (52 #4022)
- [13] W. A. J. Luxemburg and A. C. Zaanen, *Riesz spaces. Vol. I*, North-Holland Publishing Co., Amsterdam, 1971. North-Holland Mathematical Library. MR0511676 (58 #23483)
- [14] Peter Meyer-Nieberg, *Banach lattices*, Universitext, Springer-Verlag, Berlin, 1991. MR1128093 (93f:46025)
- [15] L. C. Moore Jr., *The lifting property in Archimedean Riesz spaces*, *Nederl. Akad. Wetensch. Proc. Ser. A* **73**=*Indag. Math.* **32** (1970), 141–150. MR0258707 (41 #3353)
- [16] Mehmet Orhon, *The ideal center of the dual of a Banach lattice*, Technische Universität Darmstadt preprint **1182** (1988).
- [17] Ben de Pagter, *Irreducible compact operators*, *Math. Z.* **192** (1986), no. 1, 149–153, DOI 10.1007/BF01162028. MR835399 (87d:47052)
- [18] V. G. Pestov, *Free Banach spaces and representations of topological groups*, *Funktsional. Anal. i Prilozhen.* **20** (1986), no. 1, 81–82 (Russian). MR831059 (87h:22004)
- [19] K. A. Ross and A. H. Stone, *Products of separable spaces*, *Amer. Math. Monthly* **71** (1964), 398–403. MR0164314 (29 #1611)
- [20] Zbigniew Semadeni, *Banach spaces of continuous functions. Vol. I*, PWN—Polish Scientific Publishers, Warsaw, 1971. Monografie Matematyczne, Tom 55. MR0296671 (45 #5730)
- [21] Angus E. Taylor and David C. Lay, *Introduction to functional analysis*, 2nd ed., Robert E. Krieger Publishing Co. Inc., Melbourne, FL, 1986. MR862116 (87k:46001)
- [22] Edwin H. Spanier, *Algebraic topology*, Springer-Verlag, New York, 1981. Corrected reprint. MR666554 (83i:55001)
- [23] David M. Topping, *Some homological pathology in vector lattices*, *Canad. J. Math.* **17** (1965), 411–428. MR0174499 (30 #4700)

- [24] Elliot Carl Weinberg, *Free lattice-ordered abelian groups*, Math. Ann. **151** (1963), 187–199. MR0153759 (27 #3720)
- [25] ———, *Free lattice-ordered abelian groups. II*, Math. Ann. **159** (1965), 217–222. MR0181668 (31 #5895)
- [26] A. W. Wickstead, *Banach lattices with trivial centre*, Proc. Roy. Irish Acad. Sect. A **88** (1988), no. 1, 71–83. MR974286 (89m:46030)
- [27] ———, *An isomorphic version of Nakano's characterisation of $C_0(\Sigma)$* , Positivity **11** (2007), no. 4, 609–615, DOI 10.1007/s11117-007-2078-6. MR2346446 (2008g:46032)
- [28] ———, *Banach lattices with topologically full centre*, Vladikavkaz. Mat. Zh. **11** (2009), no. 2, 50–60 (English, with English and Russian summaries). MR2529410

DELFT INSTITUTE OF APPLIED MATHEMATICS, FACULTY EEMCS, DELFT
UNIVERSITY OF TECHNOLOGY, PO BOX 5031, 2600 GA DELFT, THE NETHER-
LANDS

E-mail address: B.dePagter@tudelft.nl

PURE MATHEMATICS RESEARCH CENTRE, QUEEN'S UNIVERSITY BELFAST,
BELFAST BT7 1NN, NORTHERN IRELAND.

E-mail address: A.Wickstead@qub.ac.uk