

Impetiginized dyshidrotic eczema

Tchernev, Georgi; Zanardelli, Matteo; Voicu, Cristiana; Bakardzhiev, Ilko; Lotti, Torello; Lotti, Jacopo; França, Katlein; Batashki, Atanas; Wollina, Uwe

DOI

[10.3889/oamjms.2017.081](https://doi.org/10.3889/oamjms.2017.081)

Publication date

2017

Document Version

Final published version

Published in

Open Access Macedonian Journal of Medical Sciences

Citation (APA)

Tchernev, G., Zanardelli, M., Voicu, C., Bakardzhiev, I., Lotti, T., Lotti, J., França, K., Batashki, A., & Wollina, U. (2017). Impetiginized dyshidrotic eczema. *Open Access Macedonian Journal of Medical Sciences*, 5(4 Special Issue GlobalDermatology), 539-540. <https://doi.org/10.3889/oamjms.2017.081>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Impetiginized Dyshidrotic Eczema

Georgi Tchernev^{1*}, Matteo Zanardelli², Cristiana Voicu³, Ilko Bakardzhiev⁴, Torello Lotti⁵, Jacopo Lotti⁶, Katlein França⁷, Atanas Batashki⁸, Uwe Wollina⁹

¹Department of Dermatology, Venereology and Dermatologic Surgery, Medical Institute of Ministry of Interior, and Onkodermatologic Polyclinic for Dermatology and Dermatologic Surgery, Sofia, Bulgaria; ²Pharmacology, Toxicology and Innovative Treatments, Rome, Italy; ³Dermatology Department, Polisano Clinic, 26Z Timisoara Blvd, Bucharest, Romania; ⁴Medical College, Medical University of Varna, Varna 9000, Bulgaria; ⁵Department of Dermatology, University of Rome "G. Marconi"; Rome, Italy; ⁶Department of Biotechnology, Delft University of Technology, 2628 BC, Delft, The Netherlands; ⁷Department of Nuclear, Subnuclear and Radiation Physics, University of Rome "G. Marconi", Rome, Italy; ⁸Institute for Bioethics & Health Policy; Department of Dermatology & Cutaneous Surgery; Department of Psychiatry & Behavioral Sciences, University of Miami Miller School of Medicine - Miami, FL, USA; ⁹Abdominal and Thoracic Surgery, Department of Special Surgery, Medical University of Plovdiv, bul. "Peshtersko shose" Nr 66, 4000, Plovdiv, Bulgaria; ⁹Department of Dermatology and Allergology, Academic Teaching Hospital Dresden-Friedrichstadt, Friedrichstrasse 41, 01067, Dresden, Germany

Abstract

Citation: Tchernev G, Zanardelli M, Voicu C, Bakardzhiev I, Lotti T, Lotti J, França K, Batashki A, Wollina U. Impetiginized Dyshidrotic Eczema. Open Access Maced J Med Sci. 2017 Jul 25; 5(4):539-540. <https://doi.org/10.3889/oamjms.2017.081>

Keywords: hand dermatitis; superinfections; St. aureus; clobetasol; skin barrier.

***Correspondence:** Professor Dr Georgi Tchernev PhD, Chief of 1) Department of Dermatology, venereology and Dermatologic Surgery, Medical Institute of Ministry of Interior (MVR), General Skobelev 79, 1806 Sofia; 2) Onkodermatologic Polyclinic for Dermatology and Dermatologic Surgery, General Skobelev 26, Sofia, Bulgaria. GSM: 00359885588424. E-mail: georgi_tchernev@yahoo.de

Received: 30-Mar-2017; **Revised:** 04-Apr-2017; **Accepted:** 05-Apr-2017; **Online first:** 19-Jul-2017

Copyright: © 2017 Georgi Tchernev, Matteo Zanardelli, Cristiana Voicu, Ilko Bakardzhiev, Torello Lotti, Jacopo Lotti, Katlein França, Atanas Batashki, Uwe Wollina. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

Funding: This research did not receive any financial support.

Competing Interests: The authors have declared that no competing interests exist.

A 16 years old female patient, affected by atopic dermatitis and rhinoconjunctivitis allergica since childhood, requested a dermatologic consultation for lesions which had appeared after 3 months of local treatment with clobetasole propionate. The histological analysis confirmed the diagnosis of dyshidrotic eczema and the microbiological smears demonstrated a significant infection with *Staphylococcus aureus*. The risk of developing corticosteroids' side-effects depends on the potency of the product, extended period of use and the volume of product applied. Clobetasol propionate is a group I- highly potent corticosteroid, which should be used for a maximum period of 2 weeks. Several authors have found that this agent has cumulative depot effect, persisting in the epidermis for 4 days after only one application. Taking together these observations, sustained by the clinical case presented above, we can conclude that the infectious risks associated with topical corticosteroid treatment must not be neglected, particularly since treated patients are fragile, and frequently have multiple well-known risk factors.

A 16 years old female patient, affected by atopic dermatitis and rhinoconjunctivitis allergica since childhood, requested a dermatologic consultation for the lesions presented in Fig. 1a, 1b, which had appeared after 3 months of local treatment with clobetasole propionate for dyshidrotic eczema. The histological analysis confirmed the diagnosis of dyshidrotic eczema and the microbiological smears demonstrated a significant infection with *Staphylococcus aureus*. The pictures show the corpus inflammation and maceration of the skin (Fig. 1a, 1b). After a 10 days treatment regimen with systemic

antibiotics (Clarithromycin Retard 500 mg once daily), antihistamine tablets (Bilastine 20 mg) and topical antibacterial agents (silver sulfadiazine), the symptomatology improved significantly.

This complex case reveals a significant immune system dysregulation since the patient had been affected by atopic dermatitis and rhinoconjunctivitis allergica for a long period. As demonstrated by Ambach A et al., these pathologies can be linked to T-cell dysfunction [1].

The risk of cutaneous infections due to topical

corticosteroids is known but has never been thoroughly studied [2]. Z. Boughrara et al. noted cutaneous super-infection in nine of 30 patients receiving topical corticosteroids for bullous pemphigoid, among which there were three cases of fatal necrotizing fasciitis due to *Streptococcus A* [2].

The risk of developing corticosteroids' side-effects depends on the potency of the product, extended period of use and the volume of product applied [6]. Clobetasol propionate is a group I- highly potent corticosteroid, which should be used for a maximum period of 2 weeks [6]. Research has found that this agent has cumulative depot effect, persisting in the epidermis for 4 days after only one application [3]. Recently, del Rosso highlighted the fact that topical corticosteroids may disrupt the lipid synthesis in stratum corneum, interfering with epidermal barrier recovery [4].



Figure 1: Clinical pictures of a female patient with macerated dyshidrotic eczema and massive superinfection with *St. aureus*. Yellow-brown colour of the lesions and massive edema

The biomolecular events that facilitate the infection suggest a systemic dysfunction, not limited to the skin district. Indeed, Boudhir H et al. underlined that nodular lesions of Kaposi's disease appeared during treatment of bullous pemphigoid with topical corticosteroids [5].

Taking together these observations, evoked after the analysis of the clinical case, indicate that the infectious risks associated with topical corticosteroid treatment must not be neglected, particularly since treated patients are fragile, and frequently have multiple well-known risk factors.

Taking together these observations, sustained by the clinical case presented above, we can conclude that the infectious risks associated with topical corticosteroid treatment must not be neglected, particularly since treated patients are fragile, and frequently have multiple well-known risk factors.

References

1. Ambach A, Bonnekoh B, Gollnick H. Perforin hyperreleasability and depletion in cytotoxic T cells from patients with exacerbated atopic dermatitis and asymptomatic rhinoconjunctivitis allergica. *J Allergy Clin Immunol*. 2001;107(5):878-86. <https://doi.org/10.1067/mai.2001.114240> PMID:11344356
2. Boughrara Z, Ingen-Housz-Oro S, Legrand P, Duong TA, Roujeau JC. Cutaneous infections in bullous pemphigoid patients treated with topical corticosteroids. *Ann Dermatol Venerol*. 2010;137(5):345-51. <https://doi.org/10.1016/j.annder.2010.03.015> PMID:20470914
3. Abidi A, Ahmad F, Singh SK, Kumar A. Study of reservoir effect of clobetasol propionate cream in an experimental animal study using histamine-induced wheal suppression test. *Indian J. Dermatol*. 2010; 55:329-33. <https://doi.org/10.4103/0019-5154.74531> PMID:21430883 PMCID:PMC3051290
4. Del Rosso JQ. Topical corticosteroid application and the structural and functional integrity of the epidermal barrier. *J Clin Aesthet Dermatol*. 2013; 6(11): 20-27.
5. Boudhir H, Mael-Ainin M, Senouci K, Hassam B, Benzekri L. Kaposi's disease: an unusual side-effect of topical corticosteroids. *Ann Dermatol Venerol*. 2013;140(6-7):459-61. <https://doi.org/10.1016/j.annder.2013.04.078> PMID:23773746
6. Tadicherla S, Ross K, Shenefelt PD, Fenshe NA. Topical corticosteroids in dermatology. *J Drugs Dermatol*. 2009; 8:1093-105. PMID:20027937