

**LpLq Off-diagonal estimates for the Ornstein-Uhlenbeck semigroup
Some positive and negative results**

Amenta, Alex; Teuwen, Jonas

DOI

[10.1017/S0004972716001386](https://doi.org/10.1017/S0004972716001386)

Publication date

2017

Document Version

Final published version

Published in

Bulletin of the Australian Mathematical Society

Citation (APA)

Amenta, A., & Teuwen, J. (2017). LpLq Off-diagonal estimates for the Ornstein-Uhlenbeck semigroup: Some positive and negative results. *Bulletin of the Australian Mathematical Society*, 96(1), 154-161. <https://doi.org/10.1017/S0004972716001386>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

L^p – L^q OFF-DIAGONAL ESTIMATES FOR THE ORNSTEIN–UHLENBECK SEMIGROUP: SOME POSITIVE AND NEGATIVE RESULTS

ALEX AMENTA  and **JONAS TEUWEN**

(Received 17 October 2016; accepted 7 November 2016; first published online 6 February 2017)

Abstract

We investigate $L^p(\gamma)$ – $L^q(\gamma)$ off-diagonal estimates for the Ornstein–Uhlenbeck semigroup $(e^{tL})_{t>0}$. For sufficiently large t (quantified in terms of p and q), these estimates hold in an unrestricted sense, while, for sufficiently small t , they fail when restricted to maximal admissible balls and sufficiently small annuli. Our counterexample uses Mehler kernel estimates.

2010 *Mathematics subject classification*: primary 47D06; secondary 43A99.

Keywords and phrases: Ornstein–Uhlenbeck semigroup, off-diagonal estimates, Mehler kernel.

1. Introduction

Consider the Gaussian measure

$$d\gamma(x) := \pi^{-n/2} e^{-|x|^2} dx$$

on the Euclidean space \mathbb{R}^n , where $n \geq 1$. Naturally associated with this measure space is the Ornstein–Uhlenbeck operator

$$L := \frac{1}{2}\Delta - \langle x, \nabla \rangle = -\frac{1}{2}\nabla^* \nabla,$$

where ∇^* is the adjoint of the gradient operator ∇ with respect to the Gaussian measure. This operator generates a heat semigroup $(e^{tL})_{t>0}$ on $L^2(\gamma) = L^2(\mathbb{R}^n, \gamma)$, called the Ornstein–Uhlenbeck semigroup, with an explicit kernel: for all $u \in L^2(\gamma)$ and all $x \in \mathbb{R}^n$,

$$e^{tL}u(x) = \int_{\mathbb{R}^n} M_t(x, y)u(y) d\gamma(y),$$

The first author acknowledges financial support from the Australian Research Council Discovery Grant DP120103692 and the ANR project ‘Harmonic analysis at its boundaries’ ANR-12-BS01-0013. The second author acknowledges partial financial support from the Netherlands Organisation for Scientific Research (NWO) by the NWO-VICI grant 639.033.604.

© 2017 Australian Mathematical Publishing Association Inc. 0004-9727/2017 \$16.00

where

$$M_t(x, y) = \frac{1}{(1 - e^{-2t})^{n/2}} \exp\left(-e^{-t} \frac{|x - y|^2}{1 - e^{-2t}}\right) \exp\left(2e^{-t} \frac{\langle x, y \rangle}{1 + e^{-t}}\right) \quad (1.1)$$

is the Mehler kernel. If we equip \mathbb{R}^n with the Euclidean distance and the Gaussian measure, and if we consider operators associated with the Ornstein–Uhlenbeck operator, we find ourselves within the realm of *Gaussian harmonic analysis*, where the Ornstein–Uhlenbeck operator takes the place of the Laplace operator Δ . The multiplicative factor $1/2$, which is not present in the usual definition of the Laplacian, arises naturally from the probabilistic interpretation of the Ornstein–Uhlenbeck operator. For a deeper introduction to Gaussian harmonic analysis, see the review of Sjögren [10] and the introduction of [11].

In this paper, we investigate whether the Ornstein–Uhlenbeck semigroup satisfies $L^p(\gamma)$ – $L^q(\gamma)$ off-diagonal estimates: that is, estimates of (or similar to) the form

$$\left(\int_F |e^{tL} \mathbf{1}_E f|^q d\gamma\right)^{1/q} \lesssim t^{-\theta} \exp\left(-c \frac{\text{dist}(E, F)^2}{t}\right) \left(\int_E |f|^p d\gamma\right)^{1/p}, \quad (1.2)$$

for some parameters $c > 0$ and $\theta \geq 0$, where $1 \leq p < q \leq \infty$, $f \in L^p(\gamma)$ and for some class of *testing sets* $E, F \subset X$. Often such estimates hold whenever E and F are Borel, but, in applications, we generally only need E to be a ball and F to be an annulus associated with E . Such estimates serve as a replacement for pointwise kernel estimates in the harmonic analysis of operators whose heat semigroups have rough kernels, or no kernels at all, most notably in the solution to the Kato square root problem [2] (see also [4]). Even though the Ornstein–Uhlenbeck semigroup has a smooth kernel, it would be useful to show that it satisfies some form of off-diagonal estimates, as this would suggest potential generalisation to perturbations of the Ornstein–Uhlenbeck operator, whose heat semigroups need not have nice kernels.

Various notions of off-diagonal estimates, including (1.2), have been considered by Auscher and Martell [3]. However, they only consider doubling metric measure spaces, ruling out the nondoubling Gaussian measure. Mauceri and Meda [7] observed that γ is doubling when restricted to *admissible balls* in the sense that $\gamma(B(x, 2r)) \lesssim \gamma(B(x, r))$ when $r \leq \min(1, |x|^{-1})$. Therefore it is reasonable to expect that the Ornstein–Uhlenbeck semigroup may satisfy some form of $L^p(\gamma)$ – $L^q(\gamma)$ off-diagonal estimates if we restrict the testing sets E, F to admissible balls and sufficiently small annuli.

Here we demonstrate both the success and failure of off-diagonal estimates of the form of (1.2), as a first step in the search for the ‘right’ off-diagonal estimates. First, we give a simple positive result (Theorem 2.3): for $p \in (1, 2)$, and for t sufficiently large (depending on p), (1.2) is satisfied for all Borel $E, F \subset \mathbb{R}^n$. This is proved by interpolating between $L^2(\gamma)$ – $L^2(\gamma)$ Davies–Gaffney-type estimates and Nelson’s $L^p(\gamma)$ – $L^2(\gamma)$ hypercontractivity. We follow with a negative result (Theorem 3.1): for $1 \leq p < q < \infty$ and for t sufficiently small (again depending on p and q), (1.2) fails when E is a ‘maximal’ admissible ball $B(c_B, |c_B|^{-1})$ and when F is a sufficiently small annulus $C_k(B)$, in the sense that the implicit constant in (1.2) must blow up exponentially in $|c_B|$. This is shown by direct estimates of the Mehler kernel.

Notation. Throughout the paper, we will work in finite dimension $n \geq 1$. We will write $L^p(\gamma) = L^p(\mathbb{R}^n, \gamma)$. Every ball $B \subset \mathbb{R}^n$ is of the form

$$B = B(c_B, r_B) = \{x \in \mathbb{R}^n : |x - c_B| < r_B\}$$

for some unique centre $c_B \in \mathbb{R}^n$ and radius $r_B > 0$. For each ball B and each scalar $\lambda > 0$, we define the expansion $\lambda B = \lambda B(c_B, r_B) := B(c_B, \lambda r_B)$, and we define annuli $(C_k(B))_{k \in \mathbb{N}}$ by

$$C_k(B) := \begin{cases} 2B & \text{if } k = 0, \\ 2^{k+1}B \setminus 2^k B & \text{if } k \geq 1. \end{cases}$$

For two sets $E, F \subset \mathbb{R}^n$ we write

$$\text{dist}(E, F) := \inf\{|x - y| : x \in E, y \in F\}.$$

For two nonnegative numbers A and B , we write $A \lesssim_{a_1, a_2, \dots} B$ to mean that $A \leq CB$, where C is a positive constant depending on the quantities a_1, a_2, \dots . This constant will generally change from line to line.

2. A positive result

The Ornstein–Uhlenbeck semigroup satisfies the following ‘Davies–Gaffney-type’ $L^2(\gamma)$ – $L^2(\gamma)$ off-diagonal estimates. These appear in [13, Example 6.1], where they are attributed to Alan McIntosh.

THEOREM 2.1 (McIntosh). *There exists a constant $C > 0$ such that, for all Borel subsets E, F of \mathbb{R}^n and all $u \in L^2(\gamma)$,*

$$\|\mathbf{1}_F e^{tL}(\mathbf{1}_E u)\|_{L^2(\gamma)} \leq C \frac{t}{\text{dist}(E, F)} \exp\left(-\frac{\text{dist}(E, F)^2}{2t}\right) \|\mathbf{1}_E u\|_{L^2(\gamma)}.$$

Furthermore, Nelson [8] established the following hypercontractive behaviour of the semigroup. This is done only for $n = 1$ in [8]. A full proof for general n is given in Nelson’s seminal 1973 paper [9]. These papers won him the 1995 Steele prize.

THEOREM 2.2 (Nelson). *Let $t > 0$ and $p \in (1 + e^{-2t}, 2]$. Then e^{tL} is a contraction from $L^p(\gamma)$ to $L^2(\gamma)$.*

Note that $p > 1 + e^{-2t}$ if and only if $t > \frac{1}{2} \log(1/(p - 1))$. Thus the hypercontractive behaviour of the Ornstein–Uhlenbeck semigroup is much more delicate than that of the usual heat semigroup $e^{t\Delta}$ on \mathbb{R}^n , which is a contraction from $L^p(\mathbb{R}^n)$ into $L^q(\mathbb{R}^n)$ for all $1 \leq p \leq q \leq \infty$ and all $t > 0$.

As indicated in the proof of [1, Proposition 3.2], one can interpolate between Theorems 2.1 and 2.2 to deduce certain $L^p(\gamma)$ – $L^2(\gamma)$ off-diagonal estimates for the Ornstein–Uhlenbeck semigroup.

THEOREM 2.3. *Suppose that E, F are Borel subsets of \mathbb{R}^n . Let $t > 0$ and $p \in (1 + e^{-2t}, 2]$. Then, for all $u \in L^p(\gamma)$,*

$$\|\mathbf{1}_F e^{tL}(\mathbf{1}_E u)\|_{L^2(\gamma)} \leq \left(\frac{Ct}{\text{dist}(E, F)} \exp\left(-\frac{\text{dist}(E, F)^2}{2t}\right) \right)^{1-\delta(p,t)} \|\mathbf{1}_E u\|_{L^p(\gamma)},$$

where C is the constant from Theorem 2.1 and where

$$\delta(p, t) := \left(\frac{1}{2} - \frac{1}{p}\right) \left/ \left(\frac{1}{2} - \frac{1}{1 + e^{-2t}}\right) \right. \in [0, 1).$$

PROOF. Write

$$C_M := \frac{Ct}{\text{dist}(E, F)} \exp\left(\frac{\text{dist}(E, F)^2}{2t}\right).$$

Theorem 2.1 says that

$$\|e^{tL}\|_{L^2(\gamma, E) \rightarrow L^2(\gamma, F)} \leq C_M.$$

For all $p_0 \in (1 + e^{-2t}, p)$,

$$\|e^{tL}\|_{L^{p_0}(\gamma, E) \rightarrow L^2(\gamma, F)} \leq \|e^{tL}\|_{L^{p_0}(\gamma) \rightarrow L^2(\gamma)} \leq 1,$$

by Theorem 2.2. Therefore, by the Riesz–Thorin theorem,

$$\|e^{tL}\|_{L^p(\gamma, E) \rightarrow L^p(\gamma, F)} \leq C_M^{\theta(p_0)},$$

where $p^{-1} = (1 - \theta(p_0))/p_0 + \theta(p_0)/2$ or, equivalently,

$$\theta(p_0) = \left(\frac{1}{p} - \frac{1}{p_0}\right) \left/ \left(\frac{1}{2} - \frac{1}{p_0}\right) \right. = 1 - \left(\frac{1}{2} - \frac{1}{p}\right) \left/ \left(\frac{1}{2} - \frac{1}{p_0}\right) \right.$$

Taking the limit as $p_0 \rightarrow 1 + e^{-2t}$ gives

$$\|e^{tL}\|_{L^p(\gamma, E) \rightarrow L^p(\gamma, F)} \leq C_M^{1-\delta(p,t)}$$

and completes the proof. □

REMARK 2.4. For $1 < p < q < \infty$, a $L^p(\gamma)$ – $L^q(\gamma)$ version of Theorem 2.3 could be proved by first establishing $L^q(\gamma)$ – $L^q(\gamma)$ off-diagonal estimates—which may be obtained by interpolating between boundedness on $L^q(\gamma)$ and the Davies–Gaffney type estimates—and then arguing by the $L^p(\gamma)$ – $L^q(\gamma)$ version of Nelson’s theorem.

This positive result does not rule out the possibility of some *restricted* $L^p(\gamma)$ – $L^2(\gamma)$ off-diagonal estimates for $p \leq 1 + e^{-2t}$. In the next section, we show one way in which these can fail.

3. Lower bounds and negative results

In this section, we show that the $L^p(\gamma)$ – $L^q(\gamma)$ off-diagonal estimates of (1.2) are not satisfied for admissible balls and small annuli when t is sufficiently small (depending

on p and q). More precisely, we show that (1.2) fails when E is a maximal admissible ball B (that is, a ball for which $r_B = \min(1, |c_B|^{-1})$) and F is an annulus $C_k(B)$ with k sufficiently small. These sets typically appear in applications of off-diagonal estimates.

THEOREM 3.1. *Suppose that $1 \leq p < q < \infty$ and that*

$$\frac{2}{e^t + 1} > 1 - \left(\frac{1}{p} - \frac{1}{q}\right) \tag{3.1}$$

or, equivalently, that

$$t < \log\left(\left(1 + \left(\frac{1}{p} - \frac{1}{q}\right)\right) / \left(1 - \left(\frac{1}{p} - \frac{1}{q}\right)\right)\right).$$

Then the off-diagonal estimates (1.2) do not hold for the class of testing sets

$$\{(E, F) : E = B(c_B, |c_B|^{-1}), F = C_k(B), 2^k \leq |c_B|\}.$$

Note that $1/p - 1/q \in (0, 1)$, so we always obtain some range of t for which the off-diagonal estimates (1.2) fail.

Let us compare Theorems 3.1 and 2.3. Having fixed $p \in (1, 2)$, we get failure of $L^p(\gamma)$ – $L^2(\gamma)$ off-diagonal estimates for maximal admissible balls and small annuli for e^{tL} when

$$t < \log\left(\left(1 + \left(\frac{1}{p} - \frac{1}{2}\right)\right) / \left(1 - \left(\frac{1}{p} - \frac{1}{2}\right)\right)\right),$$

and, when $t > \frac{1}{2} \log(1/(p - 1))$, the off-diagonal estimates hold for all Borel sets. We do not know what happens for the remaining values of t .

To prove Theorem 3.1, we rely on the following lower bound.

LEMMA 3.2. *Suppose that $k \geq 1$ is a natural number and that $1 < q < \infty$, and let B be a maximal admissible ball with $|c_B| \geq 2^k$. Then*

$$\left(\int_{C_k(B)} |(e^{tL} \mathbf{1}_B)(y)|^q \, d\gamma(y)\right)^{1/q} \gtrsim_{k,n,t} |c_B|^{-n(1+1/q)} \exp\left(|c_B|^2 \left(\frac{2}{e^t + 1} - 1 - \frac{1}{q}\right)\right).$$

PROOF OF LEMMA 3.2. Suppose that $x \in B$ and $y \in C_j(B)$. We argue by computing a lower bound for the Mehler kernel $M_t(x, y)$, as given in (1.1).

Write $x = (x_1, x_2, \dots, x_n)$ and $y = (y_1, y_2, \dots, y_n)$. First, we focus on the factor involving the inner product $\langle x, y \rangle$. By symmetry, we may assume that $c_B = |c_B|e_1$. Using $r_B = |c_B|^{-1}$,

$$x_1 y_1 \geq (|c_B| - r_B)(|c_B| - 2^{k+1} r_B) \geq |c_B|^2 + O(1),$$

where we use the big- O notation $O(1)$ to mean that $x_1 y_1 - |c_B|^2$ is bounded as $|c_B| \rightarrow \infty$. If $n \geq 2$, then, by using $x_i y_i = O(1)$ for $i \geq 2$, we deduce that

$$\langle x, y \rangle \geq |c_B|^2 + O(1).$$

Evidently, this estimate remains true when $n = 1$.

Using the Mehler kernel representation of e^{tL} , for all $y \in C_k(B)$,

$$e^{tL}\mathbf{1}_B(y) \gtrsim_{n,t} \int_B \exp\left(-e^{-t} \frac{|x-y|^2}{1-e^{-2t}}\right) \exp\left(\frac{2|c_B|^2}{e^t+1}\right) d\gamma(x).$$

Since $|x-y| < 2^{k+1}r_B \leq 2$, using $r_B = |c_B|^{-1} \leq 2^{-k}$,

$$\begin{aligned} e^{tL}\mathbf{1}_B(y) &\gtrsim_{n,t} \exp\left(\frac{2|c_B|^2}{e^t+1}\right) \gamma(B) \\ &\gtrsim_n |c_B|^{-n} \exp\left(\frac{2|c_B|^2}{e^t+1} - (|c_B| + |c_B|^{-1})^2\right) \\ &\simeq |c_B|^{-n} \exp\left(|c_B|^2\left(\frac{2}{e^t+1} - 1\right)\right), \end{aligned} \tag{3.2}$$

using a straightforward estimate on $\gamma(B)$. Next, we estimate

$$\begin{aligned} \gamma(C_k(B)) &\gtrsim_n |C_k(B)| e^{-(|c_B|+2^{k+1}r_B)^2} \\ &\simeq_n 2^{kn} r_B^n \exp(-(|c_B|^2 + 2^{k+2}|c_B|r_B + 2^{k+2}r_B^2)) \\ &\simeq_{k,n} |c_B|^{-n} e^{-|c_B|^2}. \end{aligned}$$

Combining this with (3.2) gives

$$\begin{aligned} \left(\int_{C_k(B)} |(e^{tL}\mathbf{1}_B(y))|^q d\gamma(y)\right)^{1/q} &\gtrsim_{n,t} |c_B|^{-n} \exp\left(|c_B|^2\left(\frac{2}{e^t+1} - 1\right)\right) \gamma(C_k(B))^{1/q} \\ &\gtrsim_{k,n} |c_B|^{-n(1+1/q)} \exp\left(|c_B|^2\left(\frac{2}{e^t+1} - 1 - \frac{1}{q}\right)\right), \end{aligned}$$

as claimed. □

PROOF OF THEOREM 3.1. We argue by contradiction. Suppose that e^{tL} satisfies the $L^p(\gamma)$ – $L^q(\gamma)$ off-diagonal estimates (1.2) for some $\theta \geq 0$ and for (E, F) , as stated. Fix a natural number $k \geq 1$ and let B be a maximal admissible ball with $|c_B| > 2^k$. Lemma 3.2 and the off-diagonal estimates for $E = B, F = C_k(B)$ and $f = \mathbf{1}_B$ imply that

$$\begin{aligned} |c_B|^{-n(1+1/q)} \exp\left(|c_B|^2\left(\frac{2}{e^t+1} - 1 - \frac{1}{q}\right)\right) &\lesssim_{k,n,t,\theta} \exp\left(-c \frac{(2^{k+1}-1)^2 r_B^2}{t}\right) \gamma(B)^{1/p} \\ &\simeq \gamma(B)^{1/p} \end{aligned}$$

for some $c > 0$. Since

$$\gamma(B)^{1/p} \lesssim_n |B|^{1/p} e^{-(|c_B|-r_B)^2/p} \simeq_n |c_B|^{-n/p} \exp\left(-\frac{|c_B|^2}{p}\right),$$

this implies that

$$\exp\left(|c_B|^2\left(\frac{2}{e^t+1} - 1 + \frac{1}{p} - \frac{1}{q}\right)\right) \lesssim_{k,n,t,\theta} |c_B|^{n(1-(1/p-1/q))}.$$

The left-hand side grows exponentially in $|c_B|$ when (3.1) is satisfied. However, the right-hand side only grows polynomially in $|c_B|$. Thus we have a contradiction. □

REMARK 3.3. By the same argument, we can prove failure of $L^p(\gamma)$ – $L^q(\gamma)$ off-diagonal estimates for the derivatives $(L^m e^{tL})_{m \in \mathbb{N}}$ of the Ornstein–Uhlenbeck semigroup, with the same conditions on (p, q, t) and the same class of testing sets (E, F) . This relies on an identification of the kernel of $L^m e^{tL}$, which has been done by the second author in [12].

In this paper we only considered off-diagonal estimates with respect to the Gaussian measure γ . In future work, it would be very interesting to consider appropriate weighted measures, following, in particular, [5] and [6], in which (among many other things) it is shown that estimates of the form $\|e^{tL} f\|_{L^2(\gamma)} \lesssim \|f V_t\|_{L^1(\gamma)}$ hold, where V_t is a certain weight depending on t . Thus the Ornstein–Uhlenbeck semigroup does satisfy a form of ‘ultracontractivity’, but with the caveat that one must keep track of t -dependent weights. It seems that this has not yet been explored in the context of Gaussian harmonic analysis.

Acknowledgements

The authors thank Mikko Kempainen, Jan van Neerven and Pierre Portal for valuable discussions and encouragement on this topic. We also thank an anonymous referee for the suggested simplification of the proof of Lemma 3.2.

References

- [1] P. Auscher, ‘On necessary and sufficient conditions for L^p -estimates of Riesz transforms associated to elliptic operators on \mathbb{R}^n and related estimates’, *Mem. Amer. Math. Soc.* **186**(871) (2007), xviii+75.
- [2] P. Auscher, S. Hofmann, M. Lacey, A. McIntosh and P. Tchamitchian, ‘The solution of the Kato square root problem for second order elliptic operators on \mathbb{R}^n ’, *Ann. of Math. (2)* **156** (2002), 633–654.
- [3] P. Auscher and J. M. Martell, ‘Weighted norm inequalities, off-diagonal estimates and elliptic operators part ii: off-diagonal estimates on spaces of homogeneous type’, *J. Evol. Equ.* **7**(2) (2007), 265–316.
- [4] A. Axelsson, S. Keith and A. McIntosh, ‘Quadratic estimates and functional calculi of perturbed Dirac operators’, *Invent. Math.* **163** (2006), 455–497.
- [5] D. Bakry, F. Bolley and I. Gentil, ‘Dimension dependent hypercontractivity for Gaussian kernels’, *Probab. Theory Related Fields* **154**(3–4) (2012), 845–874.
- [6] D. Bakry, F. Bolley, I. Gentil and P. Maheux, ‘Weighted Nash inequalities’, *Rev. Mat. Iberoam.* **28**(3) (2012), 879–906.
- [7] G. Mauceri and S. Meda, ‘BMO and H^1 for the Ornstein–Uhlenbeck operator’, *J. Funct. Anal.* **252**(1) (2007), 278–313.
- [8] E. Nelson, ‘A quartic interaction in two dimensions’, in: *Mathematical Theory of Elementary Particles (Proc. Conf., Dedham, MA, 1965)* (MIT Press, Cambridge, MA, 1966), 69–73.
- [9] E. Nelson, ‘Construction of quantum fields from Markoff fields’, *J. Funct. Anal.* **12** (1973), 97–112.
- [10] P. Sjögren, ‘Operators associated with the Hermite semigroup—a survey’, in: *Proceedings of the conference dedicated to Professor Miguel de Guzmán (El Escorial, 1996)*, *J. Fourier Anal. Appl.* **3** (1997), Supplement 1, 813–823.
- [11] J. Teuwen, ‘A note on Gaussian maximal functions’, *Indag. Math. (N.S.)* **26** (2015), 106–112.

- [12] J. Teuwen, ‘On the integral kernels of derivatives of the Ornstein–Uhlenbeck semigroup’, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **19** (2016), Article ID 1650030, 13 pages.
- [13] J. van Neerven and P. Portal, ‘Finite speed of propagation and off-diagonal bounds for Ornstein–Uhlenbeck operators in infinite dimensions’, *Ann. Mat. Pura Appl. (4)* **195**(6) (2016), 1889–1915.

ALEX AMENTA, Delft Institute of Applied Mathematics,
Delft University of Technology, PO Box 5031, 2628 CD Delft,
The Netherlands
e-mail: amenta@fastmail.fm

JONAS TEUWEN, Division of Radiation Oncology,
Netherlands Cancer Institute/Antoni van Leeuwenhoek,
Plesmanlaan 121, 1066 CX Amsterdam, The Netherlands
and
Department of Imaging Physics, Optics Research Group,
Delft University of Technology, PO Box 5031, 2628 CD Delft,
The Netherlands
e-mail: jonasteuwen@gmail.com