

Microbiome, resistome and mobilome of chlorine-free drinking water treatment systems

Calderon Franco, D.; Corbera Rubio, F.; Cuesta Sanz, M.; Pieterse, Brent; de Ridder, David; van Loosdrecht, Mark C.M.; van Halem, D.; Laureni, M.; Weissbrodt, D.G.

DOI

[10.1016/j.watres.2023.119905](https://doi.org/10.1016/j.watres.2023.119905)

Publication date

2023

Document Version

Final published version

Published in

Water Research

Citation (APA)

Calderon Franco, D., Corbera Rubio, F., Cuesta Sanz, M., Pieterse, B., de Ridder, D., van Loosdrecht, M. C. M., van Halem, D., Laureni, M., & Weissbrodt, D. G. (2023). Microbiome, resistome and mobilome of chlorine-free drinking water treatment systems. *Water Research*, 235. <https://doi.org/10.1016/j.watres.2023.119905>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.



Microbiome, resistome and mobilome of chlorine-free drinking water treatment systems

David Calderón-Franco^{a,1}, Francesc Corbera-Rubio^{a,1}, Marcos Cuesta-Sanz^a, Brent Pieterse^b, David de Ridder^c, Mark C.M. van Loosdrecht^{a,*}, Doris van Halem^a, Michele Laurenzi^{a,2}, David G. Weissbrodt^{a,d,2}

^a Delft University of Technology, Delft, the Netherlands

^b Dunea, Utility for drinking water and nature conservancy, Plein van de Verenigde Naties 11-15, 2719 EG Zoetermeer, the Netherlands

^c Evides Water Company N.V., Schaarwijk 150, 3063 NH, Rotterdam, the Netherlands

^d Department of Biotechnology and Food Science, Division of Analysis and Control of Microbial Systems, Norwegian University of Science and Technology, Trondheim, Norway

ARTICLE INFO

Keywords:

Drinking water treatment plants
Chlorine-free
Sand filtration
Microbiome
Resistome
Mobilome

ABSTRACT

Drinking water treatment plants (DWTPs) are designed to remove physical, chemical, and biological contaminants. However, until recently, the role of DWTPs in minimizing the cycling of antibiotic resistance determinants has got limited attention. In particular, the risk of selecting antibiotic-resistant bacteria (ARB) is largely overlooked in chlorine-free DWTPs where biological processes are applied. Here, we combined high-throughput quantitative PCR and metagenomics to analyze the abundance and dynamics of microbial communities, antibiotic resistance genes (ARGs), and mobile genetic elements (MGEs) across the treatment trains of two chlorine-free DWTPs involving dune-based and reservoir-based systems. The microbial diversity of the water increased after all biological unit operations, namely rapid and slow sand filtration (SSF), and granular activated carbon filtration. Both DWTPs reduced the concentration of ARGs and MGEs in the water by circa 2.5 log gene copies mL⁻¹, despite their relative increase in the disinfection sub-units (SSF in dune-based and UV treatment in reservoir-based DWTPs). The total microbial concentration was also reduced (2.5 log units), and none of the DWTPs enriched for bacteria containing genes linked to antibiotic resistance. Our findings highlight the effectiveness of chlorine-free DWTPs in supplying safe drinking water while reducing the concentration of antibiotic resistance determinants. To the best of our knowledge, this is the first study that monitors the presence and dynamics of antibiotic resistance determinants in chlorine-free DWTPs.

1. Introduction

Access to safe water and sanitation is a key Sustainable Development Goal (United Nations, 2015) and a central objective of the Water Action Decade (United Nations, 2018) of the United Nations. Drinking water treatment plants (DWTPs) are used to remove water contaminants and deliver safe water for consumption. The sources and types of contaminants in drinking water can vary based on multiple factors such as the water source (Yu et al., 2018), geographical location (UNEP - United Nations Environment Programme, 2016), season (Kumpel et al., 2017), and human activity in the water basin (Khatri and Tyagi, 2015).

Contaminants can be classified into two categories: physical-chemical agents such as suspended particles, iron, and ammonia, and biological agents such as pathogens and antimicrobial resistances (AMR) (United States Environmental Protection Agency, 2021).

The process configuration of DWTPs is mainly dictated by the water source, either groundwater or surface water. While confined groundwater is generally microbiologically safe, surface water may contain pathogenic organisms that must be eliminated (Smeets et al., 2009). Chemical disinfectants, such as chlorine, are usually applied to disinfect drinking water (i.e., inactivate pathogenic microorganisms) and/or to prevent microbial (re)growth in the distribution network (Sedlak and

* Corresponding author.

E-mail address: M.C.M.vanLoosdrecht@tudelft.nl (M.C.M. van Loosdrecht).

¹ Shared first co-authorship

² Shared co-last authorship

<https://doi.org/10.1016/j.watres.2023.119905>

Received 29 November 2022; Received in revised form 16 March 2023; Accepted 20 March 2023

Available online 27 March 2023

0043-1354/© 2023 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

von Gunten, 2011). However, the use of disinfectants can generate by-products with mutagenic and carcinogenic effects (Rook, 1976) and selects for antibiotic-resistant bacteria (ARB) (Shi et al., 2013). A few countries (e.g., The Netherlands, Denmark, or Switzerland) ceased disinfectants use and rely on strict source-to-consumer production standards and engineering solutions for drinking water supply (Smeets et al., 2009). For the chlorine-free drinking water production from surface water, two main DWTP configurations (dune-based and reservoir-based) are employed in the Netherlands. In both cases, a large fraction of the treatment consists of biological biofilm-based unit operations such as dune infiltration, rapid sand filtration (RSF), slow sand filtration (SSF), or granular activated carbon (GAC) filtration, which combine biological and physical-chemical processes. In these systems, chemical and biological water contaminants are converted by microbial communities (Mouchet, 1992; Tekerlekopoulou et al., 2013), which shape the microbiome of the drinking water that reaches consumers (Pinto et al., 2012). Therefore, the biological safety of the microbial communities harbored in DWTPs is of utmost importance for public health.

In contrast to wastewater environments (Calderón-Franco et al., 2022; Miobedzka et al., 2022; Pallares-Vega et al., 2019), few studies focus on the fate and removal of ARGs and ARB in DWTPs. Biofilms are known reservoirs of ARB and antibiotic resistance genes (ARGs) (Balcázar et al., 2015). There is limited knowledge on the impact of biofilm-based DWTPs on the generation and/or persistence of ARB in drinking water. While antibiotic concentrations are very low or non-existent (Stackelberg et al., 2004), the generation of ARB in biofilms by horizontal gene transfer (HGT) is a well-known phenomenon (Parkas et al., 2013). HGT is the transfer of genetic material from one organisms to another that is not its offspring including mechanisms such as bacterial conjugation, transduction, or transformation, allowing for the spread of traits such as antibiotic resistance among microorganisms (Soucy et al., 2015). Therefore, the effect of biofilms present in DWTPs operational-units on ARB development needs to be uncovered. To date, molecular studies of microbial communities, ARB, and ARGs in DWTPs have been limited by the low biomass concentration present in these systems for DNA extraction, sample collection logistics, and sampling standardization (Ma et al., 2017). The results often rely on either lab-scale experiments (Stange et al., 2019; Wan et al., 2019) or specific treatment processes such as biological activated carbon filters (Wan et al., 2021) or tertiary treatments such as chlorine, UV, or a combination of them (Destiani and Templeton, 2019; Shi et al., 2013). Therefore, information about how biological treatments affect the fate of ARGs and MGEs in full-scale DWTPs from an integral consideration of the treatment train and different geographical areas is missing.

Most integrative studies have been carried out in China (Hu et al., 2019; Jia et al., 2020, 2015; Su et al., 2018; Xu et al., 2016; Zhang et al., 2019; S. 2016), i.e., one of the largest antibiotic-producing and consuming countries world-wide (Huang et al., 2019). The studies have used qPCR (Hu et al., 2019; Su et al., 2018; S. Zhang et al., 2016), high-throughput qPCR (Xu et al., 2016) or sequencing methods like amplicon sequencing and metagenomics (Jia et al., 2020, 2015; Zhang et al., 2019) to investigate the concentration and richness of ARGs in full-scale DWTPs. Sevillano et al. (2020) have compared the effect of disinfection systems on antimicrobial resistance determinants on tap water samples in DWTPs from The Netherlands, UK, and USA. Yet, no detailed information about individual process units was provided. Moreover, none of these studies have combined qualitative (metagenomics) and quantitative (HT-qPCR) approaches for their analysis to get information about the total amount and diversity of AMR determinants in chlorine-free DWTPs.

In this work, we qualitatively and quantitatively resolve the role of chlorine-free DWTPs in the control of ARB and ARGs throughout the entire treatment train of two full-scale DWTPs in a low-antibiotic-consuming country. Specifically, we aim at deciphering how the biofilms in biological unit operations shape the resistome and mobilome of the drinking water. To do so, we compared the contribution of different

methods for water storage, physical-chemical contaminant removal, and disinfection in one dune-based DWTP and one reservoir-based DWTP.

2. Material and methods

2.1. Sampling of two full-scale DWTPs

Water samples were collected from two different chlorine-free DWTPs supplying drinking water to the South Holland and Zeeland provinces in the Netherlands (Fig. 1). The dune-based DWTP in this study infiltrates pre-treated river water into the sand dunes for storage and water quality improvement (e.g., disinfection). Subsequently, dune water is abstracted with wells and treated with pellet softening, powdered activated carbon (PAC), and rapid sand filtration (RSF) to remove hardness, organic (micro)contaminants, iron, and ammonium. Slow sand filtration (SSF) is deployed as a final disinfection step, as well as to ensure the biological stability of the water (i.e., remove trace nutrients). The reservoir-based DWTP in this study stores water in open reservoirs and uses a treatment train of coagulation-flocculation and RSF to eliminate physical-chemical contaminants, UV treatment for disinfection, and granular activated carbon (GAC) to remove organic contaminants (e.g., color, odor, pesticides).

The first, dune-based DWTP (N 52° 7' 1.9992; E 4° 18' 23.9184) treats surface water from the Meuse River. The second, reservoir-based DWTP (N 51° 48' 44.4132; E 4° 20' 0.2112) processes surface water from the Meuse River as well (The Netherlands), after storage in a reservoir. Multiple wastewater treatment plants and an industrial complex discharge effluent water upstream of the intake point. Five sampling process stages per DWTP were targeted for water collection across their process stages (Fig. 2, Table 1).

The dune-based DWTP samples consisted of (D1) influent from the Meuse River water (N 51° 55' 41.7288; E 4° 46' 15.7404), (D2) outlet of the first rapid sand filtration, (D3) dune outlet (2 months hydraulic residence time), (D4) outlet of the second rapid sand filtration, and (D5) outlet of the slow sand filtration. The reservoir-based DWTP samples consisted of: (R1) influent of the reservoir from the Meuse River water (N 51° 45' 39.3228; E 4° 46' 8.6664), (R2) a sample of water after being stored in the reservoir for 2 months, (R3) rapid sand filtration treatment outlet, (R4) UV treatment outlet, and (R5) GAC outlet. Water quality parameters were provided by the DWTPs (Figure S2). The volume of each water sample depended on the expected biomass concentration at each stage, based on the author's experience and knowledge of DWTP personnel.

2.2. DNA extraction

Volumes indicated in Table 1 above 10 L were filtered on the DWTP site through a pressure filter holder containing sterile 0.22 µm polyethersulfone membrane filters, 293 mm diameter (PALL, USA) with a vacuum pump. The membrane containing the biological retentate was folded and introduced into the DNA extraction tubes. The biomass collected in the membranes were stored at 4 °C until DNA extraction. Total DNA was extracted using the DNeasy PowerWater DNA extraction kit (Qiagen, The Netherlands) following the manufacturers' instructions. DNA qualities of the extracts were measured as absorbance ratio at 260 and 280 nm using a NanoDrop spectrophotometer. DNA concentrations were measured with a Qubit4 fluorometer (Thermo Fisher Scientific, USA). The DNA quality and concentration obtained for each sample is given in Table 1.

2.3. Library preparation, sequencing, quality control, and assembly

2.3.1. Preparation of metagenome libraries

The DNA analytes were sent to Novogene (Cambridge, United Kingdom) for metagenome library preparation and sequencing. A total amount of 1 µg DNA per sample was used as input material to prepare




Fig. 1. Geographic map of sampling sites. Numbering in the figure: (1) represents the river or reservoir from which the influent sample to the DWTP was taken, and (2) represents the location of the DWTP.




Fig. 2. Schemes of the dune-based and reservoir-based drinking water treatment processes (DWTPs). The dunes and the reservoir are the storage water steps (underlined). The dune-based DWTP consists of a first rapid sand filtration (RSF1) of the Meuse River water followed by infiltration and storage in dunes (HRT = 2 months). Subsequently, the dune water is processed by pellet softening to regulate hardness, and iron (Fe) and powdered activated carbon (PAC) is dosed to improve color, odor and the performance of the second rapid sand filtration (RSF2). Finally, the water is disinfected via slow sand filtration (SSF). In the reservoir-based DWTP, the Meuse River water is stored in a reservoir (HRT = 2 months) followed by lime dosage to regulate hardness, flocculation, precipitation, and rapid sand filtration to reduce turbidity and disinfection with UV and taste and odor correction with granular activated carbon (GAC). The overall hydraulic residence times of the waters across the treatment trains amount to *circa* 2 months in both processes. Sampling points are represented by numbers D1-D5 for the dune-based DWTP (top) and R1-R5 for the reservoir-based DWTP (bottom). Numbers next to the sampling point represent the biomass concentration in

the water phase in 16S rRNA gene copies per mL. Figure adapted from (van Halem and Rietveld, 2014).

Table 1

Water samples were collected from the different stages of the dune-based and reservoir-based DWTPs processing surface water from the Meuse River and Meuse River, respectively, in the Netherlands. The metadata for the molecular biology analyses are given. Sample collection dates were selected according to the residence of the dune infiltration and reservoir steps.

DWTP system	Sample collection date	Sample description	Water volume extracted (L)	DNA extract quality (A260/A280, ± 0.1)	DNA concentration in extract (ng/ μ L)	Metagenomic analyte identifier	BioSample IDs
Dune-based	2020–10–26	Meuse River influent	5.8	1.8	71	INF-DB-DWTP	SAMN31143551
	2020–10–26	RSF1 outlet	9	1.8	27	RSF1-DB-DWTP	SAMN31143552
	2020–11–30	After dune infiltration	1200	1.8	30	DUNE-DB-DWTP	SAMN31143553
	2020–12–14	RSF2 outlet	1200	1.8	15.5	RSF2-DB-DWTP	4SAMN31143555
	2020–12–16	SSF outlet	1200	1.8	98	SSF-DB-DWTP	SAMN31143555
Reservoir-based	2020–12–19	Meuse river influent	1.5	1.8	34	INF-RB-DWTP	SAMN31143556
	2021–02–24	Reservoir outlet	1.5	1.8	10	RES-RB-DWTP	SAMN31143557
	2021–03–02	Before UV	100	1.8	11	BefUV-RB-DWTP	SAMN31143558
	2021–03–02	After UV	100	1.8	22	AftUV-RB-DWTP	SAMN31143559
	2021–03–05	GAC outlet	210	1.8	11	GAC-RB-DWTP	SAMN31143560

libraries that were generated using the NEBNext® Ultra™ DNA Library Prep Kit for Illumina (NEB, USA) following manufacturer's instructions; index codes were added to attribute sequences to each sample. In short, the DNA sample was fragmented by sonication into fragment sizes of 350 bp; the DNA fragments were end-polished, A-tailed, and ligated with the full-length adaptor for Illumina sequencing with further PCR amplification to add the sequence adapters; the PCR products were purified on AMPure XP magnetic beads (Beckman Coulter, USA).

2.3.2. Sequencing of libraries

The library preparations were sequenced with an Illumina HiSeq PE150 system. Ten raw sequencing files with 150 bp paired-reads were obtained, with an average of 5.6 Gb per sample (41 million reads). More details are given in Table S1.

2.3.3. Quality control of sequenced reads

The quality of the sequenced raw reads was assessed by FastQC (version 0.11.7) with default parameters (Andrews, 2010) and visualized with MultiQC (version 1.0) (Figure S1). Low-quality paired-end reads were trimmed and filtered by Trimmomatic version 0.39 on the paired-end mode (Bolger et al., 2014).

2.3.4. Assembly of sequence reads

Clean reads were assembled into contigs using MetaSPAdes (version 3.13.0) with default parameters (Nurk et al., 2017). An average of 37,744.2 total contigs of 4761 bp were obtained (Table S1).

2.3.5. Microbiome profiling obtained from dune and reservoir-based DWTPs

Taxonomic classification of raw reads was performed to profile the microbiome from each sample using the standard Kraken2 (version 2) database (uses all complete bacterial, archaeal, and viral genomes in NCBI Refseq database) with default parameters (Wood et al., 2019). Raw reads, divided into k-mers (substrings of length k contained within a biological sequence, determined by Kraken2), were matched with the NCBI database (Agarwala et al., 2018). The absolute abundance of each taxonomic group was indicated as the number of k-mers aligned to a specific taxonomic group. The relative abundance is the normalization of the total number of k-mers aligned in each sample.

The estimated average coverage of the samples was calculated using the online version of Nonpareil (<http://enve-omics.ce.gatech.edu/nonpareil/submit>, February 2023) with k -mer kernel and k -mer length 24. Species richness (S) was measured as the number of different species detected in the raw datasets. The Shannon (H') diversity index was calculated with the following equation:

$$H' = - \sum_{i=1}^S p_i \cdot \ln p_i$$

where p_i represents the relative abundance of species i with respect to the total amount of species (S). Microbial community distance estimation was calculated using MinHash in Mash v2.3 (Ondov et al., 2016) with “-k” 18, the minimum value required for distance estimation.

2.3.6. Resistome and mobilome profiling of DNA analytes obtained from both DWTPs

ARGs were annotated by aligning the assembled contigs > 500 bp to the ResFinder 4.0 resistance gene database using the BLASTn (version 2.6.0) nucleotide alignment tool with a cut-off E-value < 10^{-5} and sequence identity above 90% (Bortolaia et al., 2020). The richness of ARGs was defined as the number of different detected ARGs. Coverage (%) was manually checked on the identified hits, being most of them above the standard 70% required for correct identification (Garber et al., 2020).

The mobilome was analyzed on the same set of contigs >500 bp using BLASTn (version 2.6.0) with the following specific databases of MGEs with sequence identity >95% and an e-value < 10^{-20} . The presence of plasmids was studied with the PLSDb database (Galata et al., 2019). Integrons were detected with the INTEGRALL database (Moura et al., 2009). The ISfinder database was used to identify bacterial insertion sequences (Siguier et al., 2006). The ICEberg database (version May 2, 2018) detected bacterial integrative and conjugative elements (Liu et al., 2019). For all queries, the ARG or MGE identified with the best score was selected to annotate the query.

Co-occurrence (or co-localization) of MGEs and ARGs within the same contig was identified. It was checked with the BLASTn outputs if a contig contained both ARGs and MGEs. Contigs >500 bp that simultaneously included hits from the ResFinder 4.0 database and at least one of the different MGE databases were considered to have co-localized. Afterward, a specific Kraken2 taxonomic analysis was performed with these contigs to identify the potential microbial host that might carry the co-localized ARG and MGE.

2.3.7. High-throughput quantitative PCR analysis

Aliquots of the DNA extracts were sent in parallel to Resistomap (Helsinki, Finland) for high-throughput quantitative PCR (HT-qPCR) to detect and quantify the presence and abundance of 295 genes (listed in Table S2). These genes belonged to ARGs (238 genes), MGEs (51), and pathogens (6). A concentration of 2 ng DNA μ L⁻¹ in a reaction volume of 0.05 μ L was used to obtain the number of gene copies of the different biomarkers. HT-qPCR results were corrected (detailed explanation in supplementary material) to get the number of gene copies existing per

volume of filtered water from the DWTPs sampling points.

Gene detection and quantification were performed using the SmartChip™ real-Time PCR system (TakaraBio, CA, USA) by Resisto-map Oy (Helsinki, Finland). qPCR cycling conditions and processing of raw data were previously described (Lai et al., 2021; Muurinen et al., 2017; W.I. Muziasari et al., 2016; Wang et al., 2014). Briefly, the qPCR conditions included initial enzyme activation at 95 °C for 10 min, 40 cycles of denaturation at 95 °C for 30 s and then annealing at 60 °C for 30 s for amplification. Melting curve analysis was performed for each primer set of all the samples. Amplicons with unspecific melting curves and multiple peaks based on the slope of melting profiles were false positive data and therefore discarded from the analysis. Melting curve analysis was processed using the SmartChip™ qPCR software. The threshold cycle (C_T) of 27 was set as the detection limit (Muurinen et al., 2017; Muziasari et al., 2017; W.I. 2016). Standard curves and no-template controls were run alongside samples. Each DNA sample was analysed in three qPCR reactions (*i.e.* technical replicates). When a gene was detected in at least two technical replicates, mean C_T of three technical replicates in each qPCR reaction was calculated.

The gene abundance results were expressed in different ways. The absolute abundance of ARGs and MGEs was calculated as a number of gene copies per mL of filtered water as done in Xu et al. (2016). The absolute abundances of ARGs and MGEs sorted by antibiotic class and MGE type were averaged over all genetic components belonging to each group. The relative abundance of ARGs and MGEs was calculated based on the ARG or MGE copies per number of 16S rRNA gene copies. Abundance values were logarithmically transformed for comprehensive data calculation and visualization.

2.3.8. Statistics and data visualization

Graphs were made with RStudio (version 1.3.1093). Microbiome absolute and relative abundances were calculated by R package Pavian v.1.1.2.0. (Breitwieser and Salzberg, 2020). Linear correlations between absolute abundances of ARGs and MGEs were analyzed using Pearson

correlation coefficient ("ggpubr" R package) at value < 0.05. Pearson correlations between ARGs and each specific type of MGE (plasmid, insertion sequence, integron, and transposon) were also calculated. This gives the first-hint proxy for examining the co-localization of ARGs and MGEs.

Due to the logistical problems inherent to full-scale studies, we could not take a larger number of samples, which compromises the statistical significance of the results. Consequently, the analysis and discussion of results relies on trends and not on comparison between two stand-alone values.

3. Results

3.1. Microbial community composition

3.1.1. Richness and alpha diversity of the water metagenomes

The metagenomes of the microbial communities present at the different sampling points across the dune-based and reservoir-based DWTPs were sequenced to obtain first their taxonomic profiles. All sequenced samples had high-quality rates (quality rate per sequence base > 30; $Q = -10 \times \log_{10}(P)$, where P is the probability that a base call is erroneous) (Figure S1). In analogy to other engineered ecosystems such as wastewater treatment plants, and likely owing to current data-bases incompleteness, an average of $24.6 \pm 6.5\%$ of the raw reads were taxonomically classified.

The alpha diversity of the water microbiomes was assessed using the richness and the Shannon index (Fig. 3A). The former measures the number of different populations (at the genus level) in the community, and the latter accounts for the number, relative abundance, and evenness of species (Hill et al., 2003). Richness was stable throughout both DWTPs, ranging between 7027 and 7959 different classified species detected from the water metagenomes (Fig. 3). The richness of the influent of the reservoir-based DWTP was 7% higher than the dune-based DWTP influent water. Rapid sand filtration (RSF1 and




Fig. 3. A) Richness and Shannon diversity indices (x-axes) of the taxonomically classified metagenomics datasets of the waters sampled at the different locations (y-axes) within the dune-based and reservoir-based DWTP trains. B) Microbial community distance estimation with MinHash between each sampling point and the influent (p-value < 0.05). RSF: rapid sand filtration; SSF: slow sand filtration; GAC: granular activated carbon filtration.

3.2. Pathogenic bacteria decreased across both chlorine-free drinking water treatment plants

HT-qPCR was used to detect the presence of *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and Enterobacteriaceae, the three most critical antibiotic-resistant pathogenic bacteria and family containing ARBs as designated by the World Health Organization (Tacconelli and Magrini, 2017). Overall, the absolute abundance of the pathogenic bacteria detected by qPCR was low ($< 10^6$ gene copies mL⁻¹) and further reduced along the two DWTPs. *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and *Enterococci* were detected ($> 10^2$ genes copies mL⁻¹), while *Klebsiella pneumoniae*, *Campylobacter*, and *Staphylococci* were not. *A. baumannii* was detected across both plants. *P. aeruginosa* was present across the treatment train of the dune-based DWTP but was not detected after UV disinfection in the reservoir-based DWTP. Interestingly, *Enterococci* was found only after RSF1 in the dune-based DWTP but this could be an artifact due to low sample size.

3.3. Gram-negative bacteria as potential carriers of ARGs in DWTPs

The resistance determinants from the two DWTPs exhibited a large diversity of ARGs, highlighted by both qualitative (metagenomics) and quantitative (HT-qPCR) analyses.

The resistome richness ranged from 3 to 20 different ARGs detected per sample. In total, 34 different ARGs were detected in the water of the dune-based DWTP and 58 in the water of the reservoir-based DWTP (Fig. 6). The most abundant ARGs related to resistance against macrolides (MLS_B, 57 different ARGs), followed by beta-lactams (13), aminoglycosides (10), quinolones (7), sulfonamide (2), tetracycline (2), and trimethoprim (1). Some ARGs were present across the DWTPs: notably, the *msr(D)_2_AF27302* gene conferring macrolide resistance remained in the treated water of the reservoir-based DWTP. In the dune-based DWTP, the dune infiltration step was most prominently increasing the diversity of ARGs, likely due to the pronounced shift in microbial community (Fig. 4). In the reservoir-based DWTP, the rapid sand filtration (before the UV step) and the GAC filtration introduced the highest variability in the resistome profile.

We linked ARG contigs to potential microbial origins by assigning taxonomies to contigs carrying ARGs. The results of this analysis at genus level are given in Figure S3. Generally, contigs containing ARGs mainly affiliated with *Limnohabitans* in the influent water samples of both DWTPs. Other genera included *Paracoccus* and “*Ca. Fonsibacter*” in reservoir-based DWTP and *Polynucleobacter*, *Acidovorax*, *Hydrogenophaga*, and “*Ca. Fonsibacter*” in dunes-based DWTP. Most of these populations but “*Ca. Fonsibacter*” decreased across the treatment train in reservoir-based DWTP, while “*Ca. Fonsibacter*” and *Limnohabitans* persisted within the dune-based DWTP.

In the reservoir-based DWTP, the last GAC filtration step mostly increased the number of potential hosts carrying ARGs. This released bacteria potentially carrying ARGs, such as *Pseudomonas*, *Kaistella*, *Microbacterium*, *Cellulosimicrobium*, *Caulobacter*, *Methylobacterium*, *Rhodoplanes*, *Messorzhibium*, and *Rhodoferax*, among others. In the dune-based DWTP, the dune infiltration step introduced the potential hosts carrying ARGs in the drinking water treatment train. *Acinetobacter*, *Rhodoferax*, and *Pseudomonas* were the microbial genera that persisted throughout the process after infiltration in the dune.

3.4. Chlorine-free DWTPs achieve 2–3 logs removal of ARGs and MGEs

HT-qPCR was used to assess the ARG and MGE removal efficiencies from both DWTPs by quantifying the number of gene copies per volume of filtered water in each sampling point. The absolute concentration of ARGs decreased along the treatment trains down to 2.2 log gene copies mL⁻¹ (dune-based DWTP) and 2.6 log gene copies mL⁻¹ (reservoir-based DWTP) (Fig. 7a). MGEs decreased by 2.7 log gene copies mL⁻¹ (dune-based DWTP) and 2.6 log gene copies mL⁻¹ (reservoir-based

DWTP) (Fig. 7b). Similarly, the bacterial proxy *16S rRNA* gene decreased by 2.5 (dune-based DWTP) and 2.6 (reservoir-based DWTP) log gene copies mL⁻¹.

The influent water samples from two water treatment plants had similar amounts of ARGs: 6.4 ± 0.9 log ARG copies mL⁻¹ in the dune-based plant and 6.8 ± 0.9 log ARG copies mL⁻¹ in the reservoir-based plant. In the dune-based plant, the concentration of ARGs decreased from 6.0 ± 0.9 log ARG copies mL⁻¹ after the first rapid sand filtration to 4.2 ± 0.9 log ARG copies mL⁻¹ after the last slow sand filtration. In the reservoir-based plant, the ARG concentration decreased from 5.7 ± 0.7 log ARG copies mL⁻¹ after 2 months in the reservoir to 4.2 ± 0.9 log ARG copies mL⁻¹ after GAC filtration.

The concentration of MGEs in the dune-based plant was 7.1 ± 1.2 log MGE copies mL⁻¹ in the influent and decreased to 4.5 ± 1.1 log MGE copies mL⁻¹ after the last slow sand filtration. In the reservoir-based plant, the MGE concentration decreased from 6.0 ± 0.9 log MGE copies mL⁻¹ after 2 months in the reservoir to 4.4 ± 1.2 log MGE copies/mL after GAC filtration (Fig. 7b).

Some process stages increased the concentration of ARGs and MGEs in water, such as the slow sand filtration in the dune-based DWTP (22% in ARGs and 20% in MGEs) and the UV treatment in the reservoir-based DWTP (7% in ARGs and 2% in MGEs). However, the decrease in the concentration of ARGs and MGEs was progressive across both DWTPs. Detailed information on the ARGs and MGEs reduction throughout the processes per sampling point is given in Table S2.

Several ARGs persisted across both DWTPs. The *aadA7* (aminoglycoside resistance; 6.0 ± 1.2 log gene copies mL⁻¹), *mexF* (multi-drug resistance; 5.6 ± 1.2) and *fox5* (beta-lactam resistance; and 5.5 ± 1.3 log gene copies mL⁻¹) genes were the 3 most abundant ARGs in both DWTPs (Figure S4). Other ARGs were not present in the influent but appeared across the DWTPs such as *blaTEM*, *blaPAO* and *vanWG*. From the 238 ARGs tested, 72 (i.e., 30%) were not detected in any sampling point.

Regarding MGEs, the integron genes were the most abundant in both DWTPs (5.8 ± 1.7 log gene copies mL⁻¹): the *intI1_1* (integron), *repA* (plasmid), *intI3* (integron), and *Tn5403* (transposon) genes were the most abundant (Figure S5). The conjugative plasmid sequences such as *IncP_oriT* and *trbC* and promiscuous plasmid *IncQ_oriT* gene sequences were also abundant.

Despite the reduction in ARGs and MGEs concentration in water, the ratios of ARGs and MGEs to the *16S rRNA* gene remained stable throughout both DWTPs (Fig. 7c). This indicates that the DWTP process did not enrich for bacterial populations carrying ARGs or MGEs.

3.5. ARGs and MGEs co-localized on contigs of DWTP metagenomes

When ARGs and MGEs co-localize on the same genetic fragment, there is an increased chance that the fragment can be transferred between bacterial cells. Since facilitating the transfer, conjugation, integration, and transposition of genes in genomes, MGE can pose a risk for the dissemination of ARGs. Sets of 7 (dune-based DWTP) and 12 (reservoir-based DWTP) events of co-localization of ARGs and MGEs were detected on contigs retrieved from the sequenced metagenomes. A detailed description of the co-localization events is provided in Table S3. Co-localizations were detected in all sampling points from both DWTPs, except for the influent of the reservoir-based DWTP. The ARGs involved coded for mainly aminoglycosides and beta-lactam resistance.

For instance, the *ant(3'')-Ia* gene is an aminoglycoside resistant gene broadly described in *Klebsiella pneumoniae*. This persistent ARG was annotated from the metagenome of the dune-based DWTP influent and in the outlets of the RSF1, reservoir, UV, and GAC units. This ARG was embedded in plasmids, integrons, and bacterial integrative conjugative elements (ICEs), affiliating with *Polynucleobacter* and *Pseudomonas* genera. The *blaVIM* ARG against last-resource beta-lactamases (carbapenem antibiotics) was embedded in a plasmid, between insertion sequences, and in conjugative elements in different stages of the process:

in the reservoir-based DWTP, it taxonomically affiliated with *Sphingobium* and *Sphingomonas* before UV, and with *Pseudomonas* after UV and GAC.

Other ARGs only appeared once. The *sul2* and *blaOXA-287* genes appeared after RSF1 and dune infiltration, respectively, both affiliating with *Acinetobacter*. The *sul2* gene was carried by plasmid, integrons, and bacterial integrative and conjugative elements. A plasmid carried the *blaOXA-287* gene. The *sul1* gene appeared before UV, linked to *Sphingobium*. The *blaTEM-181* gene after UV linked to *Bacillus*. Both *sul1* and *blaTEM-181* genes were potentially carried by plasmids, integrons, and bacterial integrative and conjugative elements. The *srm(B)* gene after GAC was already assigned in *Rhodoferrax*. The *mef(C)* and *mph(G)* genes appeared after GAC as well, with *Kaistella* as the potential host. These 2 ARGs were detected on the same contig (NODE_16870), which implied the possible existence of a plasmid co-containing multiple resistance genes.

4. Discussion

4.1. Operational units shape the structure of the drinking water microbiome

In order to study the impact of each unit operation on the dynamics of the water microbiome, we applied shotgun metagenomics on water samples collected along the treatment train of two chlorine-free DWTPs, namely a dune-based and a reservoir-based plant. The alpha diversities of both effluents, calculated as H' Shannon diversity indices, were varied from 4.9 to 7.7 (average 6.3 ± 0.8). These values are comparable to other chlorine-free DWTPs ($4.37 \pm 0.1 - 6.02 \pm 0.4$; (Palomo et al., 2016), and significantly higher than the ones in plants with chemical disinfection (ca. 2 – 4) (Tiwari et al., 2021; Waak et al., 2019) (Dai et al., 2019). Noticeably, the alpha diversity increased after every biological filter (RSF1, RSF2, SSF, and GAC filtration), a likely consequence of direct seeding from biofilm detachment (Lautenschlager et al., 2014; Velten et al., 2011).

Proteobacteria, *Actinobacteria*, and *Bacteroidetes* were the most abundant phyla in both DWTPs. This matched with previous observations (Lin et al., 2014; Oh et al., 2017; Su et al., 2018), and with the presence of *Actinobacteria* and *Bacteroidetes* in freshwater ecosystems (Neuenschwander et al., 2018; Warnecke et al., 2004). Interestingly, the microbial community after the first operational unit of both DWTPs, namely RSF in dune-based DWTP and reservoir in reservoir-based DWTP, was similar to the influent. However, the similarity decreased in the downstream stages of the DWTPs (Fig. 3B). This aligned with the observations of Pinto et al. (2012), who highlighted that even though the source water seeds the drinking water microbiome, the unit operations shape the structure of the effluent microbial community.

The conditions within biological sand filters have different impacts

on microorganisms fitness (Hu et al., 2020), yet Webster & Fierer (2019) postulated that changes in community composition before and after lab-scale biological sand filters are largely predictable. In this line, we found higher abundances of *Actinobacteria* the RSF and SSF effluents as compared to their influents in the dune-based DWTP (Figure S2), in analogy to the high *Actinobacteria* abundance in bench-scale sand filters (Xu et al., al.(2020)). In contrast, the relative abundance of *Bacteroidetes* decreased after every biological sand filtration unit, similar to earlier reports (Mukherjee et al., 2016; Pfannes et al., 2015). Another example are the common freshwater bacteria *Limnohabitans*, "*Ca. Planktophila*", "*Ca. Nanopelagicus*" and *Rhodoluna* (Hahn, 2016; Kasalický et al., 2013; Neuenschwander et al., 2018), abundant in the water influent but almost absent in the DWTPs effluents. Overall, our findings showcase common patterns in the effect of biological operational units on the water microbiome dynamics of full-scale DWTP, paving the way to predict and modulate the microbial community in the drinking water effluent.

4.2. Chlorine-free DWTPs remove antibiotic resistance determinants

To the best of our knowledge, this is the first study in which the fate of ARG and ARB is monitored throughout the treatment train of chlorine-free DWTPs. Both DWTPs effectively reduced the concentration of ARGs by ca. 2.5 log gene copies mL^{-1} . These removals are comparable to the highest reported in chlorine-amended DWTPs, between <0.1 log ARG copies mL^{-1} (Su et al., 2018; S. Zhang et al., 2016) and 2.4 log ARG copies mL^{-1} (Hu et al., 2019). Moreover, the total ARG concentration in the water effluent of both chlorine-free DWTPs was ca. 4 log ARG copies mL^{-1} , similar to what Hu et al. (2019) found in a chlorine-amended DWTPs. Overall, both chlorine-free DWTP proved at least as effective as chlorine-amended DWTP at reducing ARGs. Additionally, the decrease in ARGs and MGEs concentration was linearly correlated with that of 16S rRNA (Fig. 7c, Figure S6), proving that none of the biological unit operations in chlorine-free DWTPs selected for ARB, i.e. the ARB/16S ratio did not increase.

The water storage steps yielded the highest ARG and ARB removal in both plants (2.1 and 1.6 ARG copies mL^{-1} in dune- and reservoir-based DWTPs, respectively), likely due to biomass decay and plasmid degradation due to the high hydraulic retention times and low nutrient availability in these systems (Amarasiri et al., 2020; Griffiths et al., 1990; Zhang et al., 2021). Likewise, RSFs reduced the ARG and ARB concentration by decreasing biomass concentration, as previously reported (Hu et al., 2019; Su et al., 2018; S. Zhang et al., 2016). Unexpectedly, GAC filtration also decreased the concentration of ARGs (Fig. 7a), in contrast to previous studies describing GAC filtration as the critical step where resistance determinants increase (Su et al., 2018; Wan et al., 2021; Zhang et al., 2019). However, the decrease in ARGs concentration in this study contrasted with the increase in ARG richness




Fig. 5. Heat map of absolute gene abundances (number of gene copies mL^{-1}) of 6 pathogenic microorganisms in dune-based and reservoir-based DWTPs, displayed in logarithmic scale. Y-axis represents the abundance of the pathogens in the different sampling points of both DWTPs. In X-axis, the 16S rRNA absolute abundance is provided. RSF: rapid sand filtration; SSF: slow sand filtration; GAC: granular activated carbon filtration.




Fig. 6. Resistome profile of dune-based and reservoir-based DWTP microbiome sorted by antibiotic class. The number of the different ARGs sorted per antibiotic class is represented in the different sampling points. The gray, horizontal line discriminates the results coming from each of the analyzed DWTPs.




Fig. 7. (a) Absolute abundance of difference ARGs mL^{-1} (sorted by antibiotic class) from both DWTPs. Values are represented on a logarithmic scale. Black dots indicate the average ARGs abundance per sampling point (b) Absolute abundance of difference MGEs mL^{-1} (sorted by antibiotic class) from both DWTPs. Values are represented on a logarithmic scale. Black dots indicate the average ARGs abundance per sampling point. (c) The ratio of ARGs or MGEs / 16S rRNA in both DWTPs. Each boxplot represents (from top to bottom) maximum, upper quartile, median, lower quartile, and minimum values. Note: RSF: rapid sand filtration; SSF: slow sand filtration; GAC: granular activated carbon.

(Fig. 6), which suggests that the microbiome in the GAC effluent is seeded by the GAC biofilm.

The final treatment step before discharge to the environment is disinfection, which is intended to suppress or inactivate harmful microorganisms and prevent the regrowth of opportunistic bacteria (National Research Council Safe Drinking Water Committee, 1980). However, the SSF ($0.75 \log \text{ gene copies mL}^{-1}$) and UV treatments (0.32

$\log \text{ gene copies mL}^{-1}$) in this study increased the concentration ARGs (Fig. 7a-b). The fate of ARG in SSF has not been directly studied before. However, Xu et al. (2020) showed that SSF hardly decreases the concentration of antibiotics in water, and Ciric (2022) reported that while their SSF removed most of the microorganisms, those in the effluent were more prone to resistance to antibiotics. In the case of UV treatment, previous studies proved its efficacy for cell reduction (plate counting)

but not for ARGs removal (Chen and Zhang, 2013; Stange et al., 2019). Gram-negative bacteria, and specifically *Pseudomonas*, tolerate UV by efficient repair mechanisms, high growth rates, or the use of low-molecular-weight organic carbon (generated by UV illumination) as an energy source (Chen et al., 2020). This can explain the rise in relative abundance of *Pseudomonas*, a common multi-drug resistant bacteria in drinking water systems (Su et al., 2018), and the quantitative increase in the 16S rRNA gene marker after UV disinfection (Fig. 7a). Nevertheless, despite the intermediate increase of ARGs, MGEs, and pathogenic bacteria after disinfection, DWTPs successfully reduced their effluent concentration (Fig. 5, Fig. 7, Figure S7).

4.3. Clinical implications of gene transfer in chlorine-free DWTPs

In concert with wastewater treatment plants (WWTPs), DWTPs are the ultimate barriers preventing the spread of waterborne diseases, and the release of antibiotics, ARB, and ARGs into water systems (Collignon and McEwen, 2019; Miłobedzka et al., 2022). One crucial aspect is the presence of ARGs against last-resort antibiotics such as carbapenems or colistin. Carbapenem resistance genes like *blaIMP* or *blaVIM* (class B beta-lactamases resistant genes) have been described in pathogenic bacteria such as *Pseudomonas*, *Acinetobacter*, or *Enterobacteriaceae* (Nordmann et al., 2012; Shanthi Amudhan et al., 2012). Carbapenem is a beta-lactam antibiotic with a broad antimicrobial spectrum and administered as a last resort for treating drug-resistant bacterial infections. However, the number of carbapenem-resistant bacteria has steadily increased (WHO, 2017), and represents a primary concern in drinking water. *blaIMP* genes were rarely detected along both DWTPs. However, *blaVIM* was detected along the dune-based DWTP and in the effluent (after GAC treatment) of the reservoir-based DWTP (Figure S4). The taxonomic annotation of the contigs containing *blaVIM* genes revealed their potential co-localization with multiple plasmids affiliating with *Sphingobium* in the dune-based DWTP, and with *Sphingobium* (plant pathogen) and *Pseudomonas* in the reservoir-based DWTP after GAC filtration. The carbapenem-resistant *Pseudomonas* is accounted by WHO within the list of critical priority pathogens for which new antibiotics are required (Tacconelli and Magrini, 2017). Colistin resistance genes, such as *mcr1* variants, were also highly abundant in both the dune-based and the reservoir-based DWTPs. However, this is not unique to chlorine-free DWTPs as multiple last-resort ARGs have also been identified in conventional DWTP (with chlorine use) as well as in tap water (Dias et al., 2020). Importantly, the *mcr1* gene load decreased significantly along the water treatment trains and neither co-localized with any MGE nor affiliated with any pathogenic bacteria. Further research should underpin the regrowth capacity of such pathogens in chlorine-free DWTP effluents.

5. Conclusions

In this work, we characterized for the very first time the abundance and dynamics of microbial communities, antibiotic resistance genes (ARGs), and mobile genetic elements (MGEs) across the treatment trains of two chlorine-free drinking water treatment plants. The in-depth analysis of the metagenomes and resistomes led to the following main conclusions:

- 1 Chlorine-free DWTPs do not select for antibiotic resistant bacteria, as supported by the linear correlated between ARGs and MGEs, and the 16S rRNA gene concentrations.
- 2 The measured reduction in ARGs concentration by ca. 2.5 log gene copies mL⁻¹ in both chlorine-free DWTPs is comparable to the highest removals reported so far for chlorine-amended DWTPs.
- 3 Water storage systems alone reduced the abundance of the 16S rRNA gene, ARGs, and MGEs by ca. 1.6 log gene copies mL⁻¹, and dune infiltration achieved the highest removal.

- 4 Despite a ca. 2.5 log 16S rRNA gene copies mL⁻¹ reduction, the effluent microbial diversity increased likely due to the seeding from the biofilms actively growing in the rapid and slow sand filters, and the granular activated carbon.
- 5 Despite the overall ARG decrease in the DWTP, disinfection (slow sand filtration and UV radiation) internally increased the concentration of ARGs, MGEs, and 16S rRNA genes by ca. 0.5 log gene copies mL⁻¹, yet with no impact on overall reduction.

Overall, our findings confirm the effectiveness of chlorine-free DWTPs in providing safe drinking water and reducing the load of antibiotic resistance determinants, offering the Water Authorities the possibility to establish centralized risk management around these specific treatment steps.

Data availability

Metagenome sequencing data were deposited in the NCBI database with the BioProject ID: PRJNA886863.

Authors' contributions

DCF and FCR designed the study with inputs of MvL, DvH, ML, and DGW. Fieldwork was supported by BP and DdR. DCF, FCR, and MCS performed the experimental investigations. DCF and FCR wrote the manuscript with direct contribution, edits, and critical feedback by all authors. DCF and FCR are first co-authors of this manuscript.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. Francesc Corbera-Rubio reports financial support was provided by Dunea Duin & Water. Brent Pieterse reports a relationship with Dunea Duin & Water that includes: employment. David de Ridder reports a relationship with Evides NV that includes: employment.

Data availability

Available in NCBI BioProject PRJNA886863

Acknowledgements

This work is part of the research project "Transmission of Antimicrobial Resistance Genes and Engineered DNA from Transgenic Biosystems in Nature" (TARGETBIO) funded by the program Biotechnology & Safety of the Ministry of Infrastructure and Water Management (grant no. 15812) of the Applied and Engineering Sciences (TTW) Division of the Dutch Research Council (NWO) (PhD thesis of David Calderón Franco). The work was further supported by the partnership program Dunea-Vitens: Sand Filtration partnership (17830) of the Dutch Research Council (NWO) and the drinking water companies Vitens and Dunea (PhD thesis of Francesc Corbera Rubio). Michele Laurenzi was supported by a VENI grant from the Dutch Research Council (NWO) (project number VI.Veni.192.252). Marcos Cuesta Sanz benefitted from an Erasmus+ grant.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.watres.2023.119905.

References

- Amarasiri, M., Sano, D., Suzuki, S., 2020. Understanding human health risks caused by antibiotic resistant bacteria (ARB) and antibiotic resistance genes (ARG) in water environments: current knowledge and questions to be answered. *Crit. Rev. Environ. Sci. Technol.* 50, 2016–2059. <https://doi.org/10.1080/10643389.2019.1692611>.
- Balcázar, J.L., Subirats, J., Borrego, C.M., 2015. The role of biofilms as environmental reservoirs of antibiotic resistance. *Front. Microbiol.* 6, 1216. <https://doi.org/10.3389/FMICB.2015.01216/BIBTEX>.
- Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30, 2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bortolai, V., Kaas, R.S., Ruppe, E., Roberts, M.C., Schwarz, S., Cattoir, V., Philippon, A., Allesoe, R.L., Rebelo, A.R., Florensa, A.F., Fagelhauer, L., Chakraborty, T., Neumann, B., Werner, G., Bender, J.K., Stingsl, K., Nguyen, M., Coppens, J., Xavier, B. B., Malhotra-Kumar, S., Westh, H., Pinholt, M., Anjum, M.F., Duggett, N.A., Kempf, I., Nykäsenoja, S., Olkkola, S., Wiczorek, K., Amaro, A., Clemente, L., Mossong, J., Losch, S., Ragimbeau, C., Lund, O., Aarestrup, F.M., 2020. ResFinder 4.0 for predictions of phenotypes from genotypes. *J. Antimicrob. Chemother.* 75. <https://doi.org/10.1093/jac/dkaa345>.
- Breitwieser, F.P., Salzberg, S.L., 2020. Pavian: interactive analysis of metagenomics data for microbiome studies and pathogen identification. *Bioinformatics* 36, 1303–1304. <https://doi.org/10.1093/bioinformatics/btz715>.
- Calderón-Franco, D., Sarelse, R., Christou, S., Pronk, M., van Loosdrecht, M.C.M., Abeel, T., Weissbrodt, D.G., 2022. Metagenomic profiling and transfer dynamics of antibiotic resistance determinants in a full-scale granular sludge wastewater treatment plant. *Water Res.* <https://doi.org/10.1016/j.watres.2022.118571>.
- Chen, H., Zhang, M., 2013. Effects of advanced treatment systems on the removal of antibiotic resistance genes in wastewater treatment plants from Hangzhou, China. *Environ. Sci. Technol.* 47, 8157–8163. <https://doi.org/10.1021/es401091y>.
- Chen, P.F., Zhang, R.J., Huang, S.B., Shao, J.H., Cui, B., Du, Z.L., Xue, L., Zhou, N., Hou, B., Lin, C., 2020. UV dose effects on the revival characteristics of microorganisms in darkness after UV disinfection: evidence from a pilot study. *Sci. Total Environ.* 713, 136582. <https://doi.org/10.1016/j.scitotenv.2020.136582>.
- Ciric, L., 2022. Are victorian water treatment technologies fit for the amr era? [WWW Document]. *Microbiol. Soc. URL* <https://microbiology.society.org/our-work/75th-anniversary-a-sustainable-future/antimicrobial-resistance-amr/antimicrobial-resistance-amr-case-studies/victorian-water-treatment-technologies-amr.html>.
- Collignon, P.J., McEwen, S.A., 2019. One health-its importance in helping to better control antimicrobial resistance. *Trop. Med. Infect. Dis.* 4. <https://doi.org/10.3390/tropicalmed4010022>.
- Dai, Z., Sevilano-Rivera, M.C., Calus, S.T., Melina Bautista-de los Santos, Q., Murat Eren, A., van der Wielen, P.W.J.J., Ijaz, U.Z., Pinto, A.J., 2019. Disinfection exhibits systematic impacts on the drinking water microbiome. *bioRxiv* 1–19. [10.1101/828970](https://doi.org/10.1101/828970).
- Destiani, R., Templeton, M.R., 2019. Chlorination and ultraviolet disinfection of antibiotic-resistant bacteria and antibiotic resistance genes in drinking water. *AIMS Environ. Sci.* 6, 222–241. <https://doi.org/10.3934/envirosci.2019.3.222>.
- Dias, M.F., da Rocha Fernandes, G., Cristina de Paiva, M., Christina de Matos Salim, A., Santos, A.B., Amaral Nascimento, A.M., 2020. Exploring the resistome, virulome and microbiome of drinking water in environmental and clinical settings. *Water Res* 174, 115630. <https://doi.org/10.1016/j.watres.2020.115630>.
- Farkas, A., Butiuc-Keul, A., Ciatară, D., Neamțu, C., Crăciunaș, C., Podar, D., Drăgan-Bularda, M., 2013. Microbiological contamination and resistance genes in biofilms occurring during the drinking water treatment process. *Sci. Total Environ.* 443, 932–938. <https://doi.org/10.1016/j.scitotenv.2012.11.068>.
- Galata, V., Fehlmann, T., Backes, C., Keller, A., 2019. PLSDB: a resource of complete bacterial plasmids. *Nucleic Acids Res* 47, D195–D202. <https://doi.org/10.1093/nar/gky1050>.
- Garber, A.I., Nealon, K.H., Okamoto, A., McAllister, S.M., Chan, C.S., Barco, R.A., Merino, N., 2020. FeGenie: a comprehensive tool for the identification of iron genes and iron gene neighborhoods in genome and metagenome assemblies. *Front. Microbiol.* 11, 1–23. <https://doi.org/10.3389/fmicb.2020.00037>.
- Griffiths, R.P., Moyer, C.L., Caldwell, B.A., Ye, C., Morita, R.Y., 1990. Long-term starvation-induced loss of antibiotic resistance in bacteria. *Microb. Ecol.* 19, 251–257. <https://doi.org/10.1007/BF02017169>.
- Hill, T.C.J., Walsh, K.A., Harris, J.A., Moffett, B.F., 2003. Using ecological diversity measures with bacterial communities. *FEMS Microbiol. Ecol.* 43, 1–11. [https://doi.org/10.1016/S0168-6496\(02\)00449-X](https://doi.org/10.1016/S0168-6496(02)00449-X).
- Hu, Y., Zhang, T., Jiang, L., Luo, Y., Yao, S., Zhang, D., Lin, K., Cui, C., 2019. Occurrence and reduction of antibiotic resistance genes in conventional and advanced drinking water treatment processes. *Sci. Total Environ.* 669, 777–784. <https://doi.org/10.1016/j.scitotenv.2019.03.143>.
- Huang, Z., Zhao, W., Xu, T., Zheng, B., Yin, D., 2019. Occurrence and distribution of antibiotic resistance genes in the water and sediments of Qingcaosha Reservoir, Shanghai, China. *Environ. Sci. Eur.* 31. <https://doi.org/10.1186/s12302-019-0265-2>.
- Jia, S., Bian, K., Shi, P., Ye, L., Liu, C.H., 2020. Metagenomic profiling of antibiotic resistance genes and their associations with bacterial community during multiple disinfection regimes in a full-scale drinking water treatment plant. *Water Res* 176, 115721. <https://doi.org/10.1016/j.watres.2020.115721>.
- Jia, S., Shi, P., Hu, Q., Li, B., Zhang, T., Zhang, X.X., 2015. Bacterial community shift drives antibiotic resistance promotion during drinking water chlorination. *Environ. Sci. Technol.* 49, 12271–12279. <https://doi.org/10.1021/acs.est.5b03521>.
- Khatiri, N., Tyagi, S., 2015. Influences of natural and anthropogenic factors on surface and groundwater quality in rural and urban areas. *Front. Life Sci.* 8, 23–39. <https://doi.org/10.1080/21553769.2014.933716>.
- Kumpel, E., Cock-Esteb, A., Duret, M., Waal, O.De, Khush, R., 2017. Seasonal variation in drinking and domestic water sources and quality in port harcourt, Nigeria. *Am. J. Trop. Med. Hyg.* 96, 437–445. <https://doi.org/10.4269/ajtmh.16-0175>.
- Lai, F.Y., Muziasari, W., Virta, M., Wiberg, K., Ahrens, L., 2021. Profiles of environmental antibiotic resistomes in the urban aquatic recipients of Sweden using high-throughput quantitative PCR analysis. *Environ. Pollut.* 287, 117651. <https://doi.org/10.1016/j.envpol.2021.117651>.
- Lautenschlager, K., Hwang, C., Ling, F., Liu, W.T., Boon, N., Köster, O., Egli, T., Hammes, F., 2014. Abundance and composition of indigenous bacterial communities in a multi-step biofiltration-based drinking water treatment plant. *Water Res* 62, 40–52. <https://doi.org/10.1016/j.watres.2014.05.035>.
- Lin, W., Yu, Z., Zhang, H., Thompson, I.P., 2014. Diversity and dynamics of microbial communities at each step of treatment plant for potable water generation. *Water Res* 52, 218–230. <https://doi.org/10.1016/j.watres.2013.10.071>.
- Liu, M., Li, X., Xie, Y., Bi, D., Sun, J., Li, J., Tai, C., Deng, Z., Ou, H.Y., 2019. ICEberg 2.0: an updated database of bacterial integrative and conjugative elements. *Nucleic Acids Res* 47, D660–D665. <https://doi.org/10.1093/nar/gky1123>.
- Ma, L., Li, B., Jiang, X.T., Wang, Y.L., Xia, Y., Li, A.D., Zhang, T., 2017. Catalogue of antibiotic resistance and host-tracking in drinking water deciphered by a large scale survey. *Microbiome* 5, 154. <https://doi.org/10.1186/s40168-017-0369-0>.
- Miłobedzka, A., Ferreira, C., Vaz-Moreira, I., Calderón-Franco, D., Gorecki, A., Purkrtova, S., Bartacek, Jan, Dziewit, L., Singleton, C.M., Nielsen, P.H., Weissbrodt, D.G., Manaia, C.M., 2022. Monitoring antibiotic resistance genes in wastewater environments: the challenges of filling a gap in the One-Health cycle. *J. Hazard. Mater.* 424. <https://doi.org/10.1016/j.jhazmat.2021.127407>.
- Mouchet, P., 1992. From conventional to biological removal of iron and manganese in France. *J. Am. Water Work. Assoc.* 84, 158–167. <https://doi.org/10.1002/j.1551-8833.1992.tb07342.x>.
- Moura, A., Soares, M., Pereira, C., Leitão, N., Henriques, I., Correia, A., 2009. INTEGRAL: a database and search engine for integrons, integrases and gene cassettes. *Bioinformatics* 25, 1096–1098. <https://doi.org/10.1093/bioinformatics/btp105>.
- Mukherjee, N., Bartelli, D., Patra, C., Chauhan, B.V., Dowd, S.E., Banerjee, P., 2016. Microbial diversity of source and point-of-use water in rural Haiti - a pyrosequencing-based metagenomic survey. *PLoS ONE* 11, e0167353. <https://doi.org/10.1371/journal.pone.0167353>.
- Muurinen, J., Stedtfeld, R., Karkman, A., Pärnänen, K., Tiedje, J., Virta, M., 2017. Influence of manure application on the environmental resistome under finnish agricultural practice with restricted antibiotic use. *Environ. Sci. Technol.* 51, 5989–5999. <https://doi.org/10.1021/acs.est.7b00551>.
- Muziasari, W.I., Pärnänen, K., Johnson, T.A., Lyra, C., Karkman, A., Stedtfeld, R.D., Tamminen, M., Tiedje, J.M., Virta, M., 2016. Aquaculture changes the profile of antibiotic resistance and mobile genetic element associated genes in Baltic Sea sediments. *FEMS Microbiol. Ecol.* 92, 1–7. <https://doi.org/10.1093/femsec/fiw052>.
- Muziasari, W.I., Pitkänen, L.K., Sørum, H., Stedtfeld, R.D., Tiedje, J.M., Virta, M., 2017. The resistome of farmed fish feces contributes to the enrichment of antibiotic resistance genes in sediments below baltic sea fish farms. *Front. Microbiol.* 7, 1–10. <https://doi.org/10.3389/fmicb.2016.02137>.
- National Research Council (US) Safe Drinking Water Committee, 1980. *Drinking Water and Health*. In: *Drinking Water and Health*, 2. National Academies Press, US, pp. 955–956. <https://doi.org/10.1111/j.1752-1688.1984.tb04810.x>.
- Neuenschwander, S.M., Ghai, R., Pernthaler, J., Salcher, M.M., 2018. Microdiversification in genome-streamlined ubiquitous freshwater Actinobacteria. *ISME J* 12, 185–198. <https://doi.org/10.1038/ismej.2017.156>.
- Nordmann, P., Dortet, L., Poirel, L., 2012. Carbapenem resistance in Enterobacteriaceae: here is the storm! *Trends Mol. Med.* 18, 263–272. <https://doi.org/10.1016/j.molmed.2012.03.003>.
- Oh, S., Hammes, F., Liu, W.T., 2017. Metagenomic characterization of biofilter microbial communities in a full-scale drinking water treatment plant. *Water Res* 128, 278–285. <https://doi.org/10.1016/j.watres.2017.10.054>.
- Ondov, B.D., Treangen, T.J., Melsted, P., Mallonee, A.B., Bergman, N.H., Koren, S., Phillippy, A.M., 2016. Mash: fast genome and metagenome distance estimation using MinHash. *Genome Biol* 17, 1–14. <https://doi.org/10.1186/S13059-016-0997-X/FIGURES/5>.
- Pallares-Vega, R., Blaak, H., van der Plaats, R., de Roda Husman, A.M., Hernandez Leal, L., van Loosdrecht, M.C.M., Weissbrodt, D.G., Schmitt, H., 2019. Determinants of presence and removal of antibiotic resistance genes during WWTP treatment: a cross-sectional study. *Water Res* 161, 319–328. <https://doi.org/10.1016/j.watres.2019.05.100>.
- Palomo, A., Jane Fowler, S., Gülay, A., Rasmussen, S., Sicheritz-Ponten, T., Smets, B.F., 2016. Metagenomic analysis of rapid gravity sand filter microbial communities suggests novel physiology of Nitrospira spp. *ISME J* 10, 2569–2581. <https://doi.org/10.1038/ismej.2016.63>.
- Pfannes, K.R., Langenbach, K.M.W., Pilloni, G., Stührmann, T., Euringer, K., Lueders, T., Neu, T.R., Müller, J.A., Kästner, M., Meckenstock, R.U., 2015. Selective elimination of bacterial faecal indicators in the Schmutzdecke of slow sand filtration columns. *Appl. Microbiol. Biotechnol.* 99, 10323–10332. <https://doi.org/10.1007/s00253-015-6882-9>.
- Pinto, A.J., Xi, C., Raskin, L., 2012. Bacterial community structure in the drinking water microbiome is governed by filtration processes. *Environ. Sci. Technol.* 46, 8851–8859. <https://doi.org/10.1021/es302042t>.
- Rodríguez-R, L.M., Konstantinidis, K.T., 2014. Estimating coverage in metagenomic data sets and why it matters. *ISME J* 2349–2351. <https://doi.org/10.1038/ISMEJ.2014.76>, 2014 811 8.
- Rook, J.J., 1976. Haloforms in drinking water. *J. Am. Water Works Assoc.* 68, 168–172. <https://doi.org/10.1002/J.1551-8833.1976.TB02376.X>.

- Sedlak, D.L., Von Gunten, U., 2011. The chlorine dilemma. *Science* (80-). <https://doi.org/10.1126/science.1196397>.
- Sevillano, M., Dai, Z., Calus, S., Bautista-de los Santos, Q.M., Eren, A.M., van der Wielen, P.W.J.J., Ijaz, U.Z., Pinto, A.J., 2020. Differential prevalence and host-association of antimicrobial resistance traits in disinfected and non-disinfected drinking water systems. *Sci. Total Environ.* 749, 141451 <https://doi.org/10.1016/j.scitotenv.2020.141451>.
- Shanthi Amudhan, M., Sekar, U., Kamalanathan, A., Balaraman, S., 2012. blaIMP and blaVIM mediated carbapenem resistance in pseudomonas and acinetobacter species in India. *J. Infect. Dev. Ctries.* 6, 757–762. <https://doi.org/10.3855/jidc.2268>.
- Shi, P., Jia, S., Zhang, X.X., Zhang, T., Cheng, S., Li, A., 2013. Metagenomic insights into chlorination effects on microbial antibiotic resistance in drinking water. *Water Res* 47, 111–120. <https://doi.org/10.1016/j.watres.2012.09.046>.
- Siguié, P., Perochon, J., Lestrade, L., Mahillon, J., Chandler*, M., 2006. ISfinder: the reference centre for bacterial insertion sequences. *Nucleic Acids Res* 34, D32–D36. <https://doi.org/10.1093/nar/gkj014>.
- Smeets, P.W.M.H., Medema, G.J., Van Dijk, J.C., 2009. The Dutch secret: how to provide safe drinking water without chlorine in the Netherlands. *Drink. Water Eng. Sci.* 2, 1–14. <https://doi.org/10.5194/dwes-2-1-2009>.
- Soucy, S.M., Huang, J., Gogarten, J.P., 2015. Horizontal gene transfer: building the web of life. *Nat. Rev. Genet.* 16, 472–482. <https://doi.org/10.1038/nrg3962>.
- Stackelberg, P.E., Furlong, E.T., Meyer, M.T., Zaugg, S.D., Henderson, A.K., Reissman, D. B., 2004. Persistence of pharmaceutical compounds and other organic wastewater contaminants in a conventional drinking-water-treatment plant. *Sci. Total Environ.* 329, 99–113. <https://doi.org/10.1016/j.scitotenv.2004.03.015>.
- Stange, C., Sidhu, J.P.S., Toze, S., Tiehm, A., 2019. Comparative removal of antibiotic resistance genes during chlorination, ozonation, and UV treatment. *Int. J. Hyg. Environ. Health* 222, 541–548. <https://doi.org/10.1016/j.ijheh.2019.02.002>.
- Su, H.C., Liu, Y.S., Pan, C.G., Chen, J., He, L.Y., Ying, G.G., 2018. Persistence of antibiotic resistance genes and bacterial community changes in drinking water treatment system: from drinking water source to tap water. *Sci. Total Environ.* 616–617, 453–461. <https://doi.org/10.1016/j.scitotenv.2017.10.318>.
- Tacconelli, E., Magrini, N., 2017. *Global Priority List of Antibiotic-Resistant Bacteria to Guide research, discovery, and Development of New Antibiotics*. World Health Organization.
- Tekerlekopoulou, A.G., Pavlou, S., Vayenas, D.V., 2013. Removal of ammonium, iron and manganese from potable water in biofiltration units: a review. *J. Chem. Technol. Biotechnol.* 88, 751–773. <https://doi.org/10.1002/jctb.4031>.
- Tiwari, A., Hokajärvi, A.M., Domingo, J.S., Elk, M., Jayaprakash, B., Ryu, H., Siponen, S., Vepsäläinen, A., Kauppinen, A., Puurunen, O., Artimo, A., Perkola, N., Huttula, T., Miettinen, I.T., Pitkänen, T., 2021. Bacterial diversity and predicted enzymatic function in a multipurpose surface water system – from wastewater effluent discharges to drinking water production. *Environ. Microbiomes* 16, 1–17. <https://doi.org/10.1186/s40793-021-00379-w>.
- UNEP - United Nations Environment Programme, 2016. *A Snapshot of the World's Water Quality : Towards a global Assessment*. United Nations Environment Programme.
- United Nations, 2018. *Water Action Decade* [WWW Document]. URL <https://wateractiondecade.org/>(accessed 1.26.22).
- United Nations, 2015. *Sustainable development goals* [WWW Document]. URL <https://www.un.org/sustainabledevelopment/water-and-sanitation/>(accessed 1.26.22).
- United States Environmental Protection Agency, 2021. *Types of drinking water contaminants* [WWW Document]. URL <https://www.epa.gov/ccl/types-drinking-water-contaminants> (accessed 1.26.22).
- van Halem, D., Rietveld, L.C., 2014. *Introduction to drinking water treatment* [WWW Document]. URL <https://online-learning.tudelft.nl/courses/introduction-to-drinking-water-treatment/>(accessed 11.4.20).
- Velten, S., Boller, M., Köster, O., Helbing, J., Weilenmann, H.U., Hammes, F., 2011. Development of biomass in a drinking water granular active carbon (GAC) filter. *Water Res* 45, 6347–6354. <https://doi.org/10.1016/j.watres.2011.09.017>.
- Waak, M.B., Hozalski, R.M., Hallé, C., Lapara, T.M., 2019. Comparison of the microbiomes of two drinking water distribution systems - With and without residual chloramine disinfection. *Microbiome* 7. <https://doi.org/10.1186/s40168-019-0707-5>.
- Wan, K., Guo, L., Ye, C., Zhu, J., Zhang, M., Yu, X., 2021. Accumulation of antibiotic resistance genes in full-scale drinking water biological activated carbon (BAC) filters during backwash cycles. *Water Res* 190, 116744. <https://doi.org/10.1016/j.watres.2020.116744>.
- Wan, K., Zhang, M., Ye, C., Lin, W., Guo, L., Chen, S., Yu, X., 2019. Organic carbon: an overlooked factor that determines the antibiotic resistome in drinking water sand filter biofilm. *Environ. Int.* 125, 117–124. <https://doi.org/10.1016/j.envint.2019.01.054>.
- Wang, F.H., Qiao, M., Su, J.Q., Chen, Z., Zhou, X., Zhu, Y.G., 2014. High throughput profiling of antibiotic resistance genes in urban park soils with reclaimed water irrigation. *Environ. Sci. Technol.* 48, 9079–9085. <https://doi.org/10.1021/es502615e>.
- Warnecke, F., Amann, R., Pernthaler, J., 2004. Actinobacterial 16S rRNA genes from freshwater habitats cluster in four distinct lineages. *Environ. Microbiol.* 6, 242–253. <https://doi.org/10.1111/j.1462-2920.2004.00561.x>.
- WHO, 2017. *Guidelines for the prevention and control of carbapenem-resistant Enterobacteriaceae, Acinetobacter baumannii and Pseudomonas aeruginosa in health care facilities*.
- Xu, L., Campos, L.C., Canales, M., Ciric, L., 2020. Drinking water biofiltration: behaviour of antibiotic resistance genes and the association with bacterial community. *Water Res* 182, 115954. <https://doi.org/10.1016/j.watres.2020.115954>.
- Xu, L., Ouyang, W., Qian, Y., Su, C., Su, J., Chen, H., 2016. High-throughput profiling of antibiotic resistance genes in drinking water treatment plants and distribution systems. *Environ. Pollut.* 213, 119–126. <https://doi.org/10.1016/j.envpol.2016.02.013>.
- Yu, L., Rozemeijer, J., Van Breukelen, B.M., Ouboter, M., Van Der Vlugt, C., Broers, H.P., 2018. Groundwater impacts on surface water quality and nutrient loads in lowland polder catchments: monitoring the greater Amsterdam area. *Hydrol. Earth Syst. Sci.* 22, 487–508. <https://doi.org/10.5194/hess-22-487-2018>.
- Zhang, H., Chang, F., Shi, P., Ye, L., Zhou, Q., Pan, Y., Li, A., 2019. Antibiotic resistome alteration by different disinfection strategies in a full-scale drinking water treatment plant deciphered by metagenomic assembly. *Environ. Sci. Technol.* 53, 2141–2150. <https://doi.org/10.1021/acs.est.8b05907>.
- Zhang, S., Lin, W., Yu, X., 2016. Effects of full-scale advanced water treatment on antibiotic resistance genes in the Yangtze Delta area in China. *FEMS Microbiol. Ecol.* 92, 1–9. <https://doi.org/10.1093/femsec/fiw065>.
- Zhang, T., Lv, K., Lu, Q., Wang, L., Liu, X., 2021. Removal of antibiotic-resistant genes during drinking water treatment: a review. *J. Environ. Sci. (China)* 104, 415–429. <https://doi.org/10.1016/j.jes.2020.12.023>.