

Scatterer identification and analysis using combined InSAR and laser data

Hanssen, Ramon; van Natijne, Adriaan; Lindenberg, Roderik; Dheenathayalan, Prabu; Yang, Mengshi; Chang, Ling; van Leijen, Freek; Lopez Dekker, Paco; van der Maaden, Jippe; van Oosterom, P.J.M.

Publication date

2018

Document Version

Final published version

Published in

Geophysical Research Abstracts (online)

Citation (APA)

Hanssen, R., van Natijne, A., Lindenberg, R., Dheenathayalan, P., Yang, M., Chang, L., van Leijen, F., Lopez Dekker, P., van der Maaden, J., van Oosterom, P. J. M., Xiong, H., Hu, P., Zhan, Z., & Yang, B. (2018). Scatterer identification and analysis using combined InSAR and laser data. *Geophysical Research Abstracts (online)*, 20, Article EGU2018-17008.

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.



Scatterer identification and analysis using combined InSAR and laser data

Ramon Hanssen (1), Adriaan van Natiijne (1), Roderik Lindenbergh (1), Prabu Dheenathayalan (1), Mengshi Yang (1,2), Ling Chang (1), Freek van Leijen (1), Paco Lopez-Dekker (1), Jippe van der Maaden (1), Peter van Oosterom (1), Hanjiang Xiong (2), PingBo Hu (2), Zhang Zhan (2), and Bisheng Yang (2)

(1) Delft University of Technology, Netherlands (r.f.hanssen@tudelft.nl), (2) Wuhan University, China (bshyang@whu.edu.cn)

The geolocation of coherent radar scatterers, used for InSAR deformation analysis, is often not accurate enough to associate them to physical geo-objects. The imaging geometry of satellite InSAR results in (i) biases in the entire point field, and (ii) quite elongated and skewed confidence ellipsoids in the range, azimuth and cross-range direction. The metric defined by the covariance matrix of the InSAR results defines the optimal way to associate scatterers with geo-objects.

Laser scanning point clouds, stemming from aerial or terrestrial laser surveys, yield very dense geometry of geo-objects and topography.

Here we combine InSAR and laser point clouds, taking the covariance metrics of the InSAR data into account. This enables us to correct the positions of InSAR data, to provide a geometric match with geo-objects. We demonstrate how this allows for adding contextual information as attributes to individual scatterers, which improves the interpretation of the InSAR results.