



Delft University of Technology

Document Version

Accepted author manuscript

Citation (APA)

Hartstra, I., Almagro Vidal, C., & Wapenaar, CPA. (2016). Full-field MDD for body-wave reflections from passive transient-sources under severely limited and irregular illumination conditions. In *Proceedings of the AGU fall meeting 2016 AGU*.

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

In case the licence states "Dutch Copyright Act (Article 25fa)", this publication was made available Green Open Access via the TU Delft Institutional Repository pursuant to Dutch Copyright Act (Article 25fa, the Taverne amendment). This provision does not affect copyright ownership.
Unless copyright is transferred by contract or statute, it remains with the copyright holder.

Sharing and reuse

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

This work is downloaded from Delft University of Technology.

Full-field MDD for body-wave reflections from passive transient-sources under severely limited and irregular illumination conditions

Authors: Iris Hartstra, Carlos Almagro Vidal and Kees Wapenaar

Seismic interferometry (SI) presents a set of inexpensive and noninvasive methods that can be applied to any array at the surface to retrieve virtual body-wave reflection responses from earthquake recordings. Conventional SI by cross-correlation requires recordings of wavefields in lossless media generated by a smooth continuous distribution of passive sources with isotropic source radiation patterns and similar power spectra. These conditions are unlikely to be met in the lithosphere: earthquakes are distributed sparsely and generated by complex mechanisms. The resulting anisotropy in the illumination of the receiver array causes the retrieved virtual-source radiation patterns to be irregular, leading to artifacts which can obscure the desired body-wave reflections.

SI by multidimensional deconvolution (MDD) can inherently correct for anisotropic illumination of the array and does not rely on the medium being lossless. We propose an alternative formulation of MDD for two-way wavefields: full-field MDD. Different from previous MDD methods for passive two-way wavefield recordings, full-field MDD uses multiples in the passive data to construct the reflection response without free-surface interaction. Therefore, this MDD method profits from additional wavenumbers provided by scattering to compensate for sparse earthquake distributions. Besides, this method does not require wavefield decomposition, which is sensitive to velocity variations at the receiver level.

We compare the reflection retrieval by full-field MDD and cross-correlation for a limited passive source distribution in a lithospheric model with a discontinuous Moho at a depth of 50 km. We simulate earthquakes generated by dipole sources along a listric fault-system with power spectra varying within bandwidth 0.2-2.6 Hz. The reflection response retrieved by full-field MDD shows a continuous high-resolution Moho reflection, while cross-correlation yields a very low resolution response obscured by artifacts.