

A continuous-time distributed generalized Nash equilibrium seeking algorithm over networks for double-integrator agents

Bianchi, Mattia; Grammatico, Sergio

DOI

[10.23919/ECC51009.2020.9143714](https://doi.org/10.23919/ECC51009.2020.9143714)

Publication date

2020

Document Version

Final published version

Published in

Proceedings of the European Control Conference 2020, ECC 2020

Citation (APA)

Bianchi, M., & Grammatico, S. (2020). A continuous-time distributed generalized Nash equilibrium seeking algorithm over networks for double-integrator agents. In *Proceedings of the European Control Conference 2020, ECC 2020* (pp. 1474-1479). IEEE. <https://doi.org/10.23919/ECC51009.2020.9143714>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

Green Open Access added to TU Delft Institutional Repository

'You share, we take care!' - Taverne project

<https://www.openaccess.nl/en/you-share-we-take-care>

Otherwise as indicated in the copyright section: the publisher is the copyright holder of this work and the author uses the Dutch legislation to make this work public.

A continuous-time distributed generalized Nash equilibrium seeking algorithm over networks for double-integrator agents

Mattia Bianchi and Sergio Grammatico

Abstract—We consider a system of single- or double-integrator agents playing a generalized Nash game over a network, in a partial-information scenario. We address the generalized Nash equilibrium seeking problem by designing a fully-distributed dynamic controller, based on continuous-time consensus and primal-dual gradient dynamics. Our main technical contribution is to show convergence of the closed-loop system to a variational equilibrium, under strong monotonicity and Lipschitz continuity of the game mapping, by leveraging monotonicity properties and stability theory for projected dynamical systems.

I. INTRODUCTION

Generalized Nash equilibrium (GNE) problems arise in several network systems, where multiple selfish decision-makers, or agents, aim at optimizing their individual, yet inter-dependent, objective functions, subject to shared constraints. Engineering applications include demand-side management in the smart grid [1], charging/discharging of electric vehicles [2], formation control [3] and communication networks [4]. From a game-theoretic perspective, the aim is to design distributed GNE seeking algorithms, using the local information available to each agent. Moreover, in the cyber-physical systems framework, games are often played by agents with their own dynamics [5], [3], and controllers have to be conceived to steer the physical process to a Nash equilibrium, while ensuring closed-loop stability. This stimulates the development of continuous-time schemes [6], [7], for which control-theoretic properties are more easily unraveled.

Literature review: A variety of different methods have been proposed to seek GNE in a distributed way [8], [9], [10]. These works refer to a full-information setting, where each agent can access the decision of all other agents, for example if a coordinator broadcasts the data to the network. Nevertheless, there are applications where the existence of a central node must be excluded and each agent only relies on the information exchanged over a network, via peer-to-peer communication. To deal with this partial-information scenario, payoff-based algorithms for Nash equilibrium (NE) seeking have been studied, [11], [3]. In this paper, we are instead interested in a different, model-based, approach. We assume that the agents agree on sharing their strategies with their neighbors; each agent keeps an estimate of other agents'

The authors are with the Delft Center for Systems and Control (DCSC), TU Delft, The Netherlands E-mail addresses: {m.bianchi, s.grammatico}@tudelft.nl. This work was partially supported by NWO under research project OMEGA (grant n. 613.001.702) and by the ERC under research project COSMOS (802348).

actions and asymptotically reconstructs the true values, exploiting the information exchanged over the network. This solution has been examined extensively for games without coupling constraints, both in discrete time [12], [13], and continuous-time [6], [14]. However, fewer works deal with generalized games. Remarkably, Pavel in [15] derived a single-timescale, fixed step sizes GNE learning algorithm, by leveraging an elegant operator splitting approach. The authors in [16] proposed a continuous-time design for aggregative games with equality constraints. All the results mentioned above consider single-integrator agents only. Distributively driving a network of more complex physical systems to game theoretic solutions is still a relatively unexplored problem. With regard to aggregative games, a proportional integral feedback algorithm was developed in [5] to seek a NE in networks of passive nonlinear second-order systems. In [17], continuous-time gradient-based controllers were introduced, for some classes of nonlinear systems with uncertainties. The authors of [3] considered generally coupled costs games played by linear time-invariant agents, via a discrete-time extremum seeking approach. NE problems arising in systems of multi-integrator agents, in the presence of deterministic disturbances, were addressed in [18]. In all the references cited, the assumption is made of unconstrained action sets and absence of coupling constraints.

Contribution: Motivated by the above, in this paper we investigate continuous-time GNE seeking for networks of single- or double-integrator agents. We consider games with affine coupling constraints, played under partial-decision information. Specifically:

- We introduce a primal-dual projected-gradient controller for single-integrator agents, which is a continuous-time version of the one proposed in [15]. We show convergence of both primal and dual variables, under strong monotonicity and Lipschitz continuity of the game mapping. We are not aware of other continuous-time GNE seeking algorithms for games with generally coupled costs, whose convergence is guaranteed under such mild assumptions. With respect to the setup (for aggregative game only) in [16], we can also handle inequality constraints.
- We show how our controller can be adapted to learning GNE in games with shared constraints, played by double-integrator agents. To the best of our knowledge, this is the first equilibrium-seeking algorithm for *generalized* games where the agents have second-order dynamics.

Basic notation: \mathbb{R} ($\mathbb{R}_{\geq 0}$) denotes the set of (nonnegative) real numbers. $\mathbf{0}$ ($\mathbf{1}$) denotes a matrix/vector with all elements

equal to 0 (1); we may add the dimension as subscript for clarity. $\|\cdot\|$ denotes the Euclidean norm. $I_n \in \mathbb{R}^{n \times n}$ denotes the identity matrix of dimension n . For a matrix $A \in \mathbb{R}^{n \times m}$, its transpose is A^\top , $[A]_{i,j}$ represents the element on the row i and column j . $A \otimes B$ denotes the Kronecker product of the matrices A and B . $A \succ 0$ stands for symmetric positive definite matrix. If A is symmetric, $\lambda_{\min}(A) := \lambda_1(A) \leq \dots \leq \lambda_n(A) := \lambda_{\max}(A)$ denote its eigenvalues. Given N vectors x_1, \dots, x_N , $x := \text{col}(x_1, \dots, x_N) = [x_1^\top \dots x_N^\top]^\top$, and for each $i = 1, \dots, N$, $x_{-i} := \text{col}(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_N)$. For a differentiable function $g : \mathbb{R}^n \rightarrow \mathbb{R}$, $\nabla_x g(x)$ denotes its gradient.

Operator-theoretic definitions: \overline{H} denotes the closure of a set $H \subseteq \mathbb{R}^n$. A set-valued mapping $\mathcal{F} : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is (μ -strongly) monotone if $(u - v)^\top(x - y) \geq 0$ ($\geq \mu\|x - y\|^2$) for all $x \neq y \in \mathbb{R}^n$, $u \in \mathcal{F}(x)$, $v \in \mathcal{F}(y)$. Given a closed, convex set $S \subseteq \mathbb{R}^n$, the mapping $\text{proj}_S : \mathbb{R}^n \rightarrow S$ denotes the projection onto S , i.e., $\text{proj}_S(v) := \text{argmin}_{y \in S} \|y - v\|$. The set-valued mapping $N_S : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ denotes the normal cone operator for the set S , i.e., $N_S(x) = \emptyset$ if $x \notin S$, $\{v \in \mathbb{R}^n \mid \sup_{z \in S} v^\top(z - x) \leq 0\}$ otherwise. The tangent cone of S at a point $x \in S$ is defined as $T_S(x) = \bigcup_{\delta > 0} \frac{1}{\delta}(S - x)$. $\Pi_S(x, v) := \text{proj}_{T_S(x)}(v)$ denotes the projection on the tangent cone of S at x of a vector $v \in \mathbb{R}^n$. By Moreau's Decomposition Theorem [19, Th. 6.30], it holds that $v = \text{proj}_{T_S(x)}(v) + \text{proj}_{N_S(x)}(v)$ and $\text{proj}_{T_S(x)}(v)^\top \text{proj}_{N_S(x)}(v) = 0$.

Lemma 1: For any closed convex set $S \subseteq \mathbb{R}^q$, any $y, y' \in S$ and any $\xi \in \mathbb{R}^q$, it holds that

$$(y - y')^\top \Pi_S(y, \xi) \leq (y - y')^\top \xi.$$

In particular, if $\Pi_S(y, \xi) = 0$, then $(y - y')^\top \xi \geq 0$. \square

Proof: By Moreau's theorem, $(\xi - \Pi_C(y, \xi)) \in N_S(y)$, hence for any $y, y' \in C$, $(y' - y)^\top(\xi - \Pi_C(y, \xi)) \leq 0$. \blacksquare

II. MATHEMATICAL SETUP

We consider a set of noncooperative agents, $\mathcal{I} := \{1, \dots, N\}$, where each agent $i \in \mathcal{I}$ shall choose its decision variable (i.e., strategy) x_i from its local decision set $\Omega_i \subseteq \mathbb{R}^{n_i}$. Let $x = \text{col}((x_i)_{i \in \mathcal{I}}) \in \Omega$ denote the stacked vector of all the agents' decisions, $\Omega = \times_{i \in \mathcal{I}} \Omega_i \subseteq \mathbb{R}^n$ the overall action space and $n := \sum_{i=1}^N n_i$. Moreover, let $x_{-i} = \text{col}((x_j)_{j \in \mathcal{I} \setminus \{i\}})$ denote the collective strategy of the all the agents, except that of agent i . The goal of each agent $i \in \mathcal{I}$ is to minimize its objective function $J_i(x_i, x_{-i})$ which depends on both the local variable x_i and on the decision variables of the other agents x_{-i} .

Furthermore, we consider *generalized* games, where the agents' strategies are also coupled via some shared affine constraints. Thus the overall feasible set is

$$\mathcal{X} := \Omega \cap \{x \in \mathbb{R}^n \mid Ax \leq b\}, \quad (1)$$

where $A := [A_1, \dots, A_N]$ and $b := \sum_{i=1}^N b_i$, with $A_i \in \mathbb{R}^{m \times n_i}$ and $b_i \in \mathbb{R}^m$ being local data. The game then is

represented by the inter-dependent optimization problems:

$$\forall i \in \mathcal{I} : \quad \underset{y_i \in \mathbb{R}^{n_i}}{\text{argmin}} J_i(y_i, x_{-i}) \quad \text{s.t.} \quad (y_i, x_{-i}) \in \mathcal{X}. \quad (2)$$

The technical problem we consider in this paper is the computation of a GNE, i.e., a set of strategies from which no agent has an incentive to unilaterally deviate.

Definition 1: A collective strategy $x^* = \text{col}((x_i^*)_{i \in \mathcal{I}})$ is a generalized Nash equilibrium if, for all $i \in \mathcal{I}$,

$$x_i^* \in \underset{y_i}{\text{argmin}} J_i(y_i, x_{-i}^*) \quad \text{s.t.} \quad (y_i, x_{-i}^*) \in \mathcal{X}. \quad \square$$

Next, we postulate standard regularity assumptions for the constraint sets and cost functions, [15, Ass. 1], [14, Ass. 1].

Standing Assumption 1: For each $i \in \mathcal{I}$, the set Ω_i is non-empty, closed and convex; \mathcal{X} is non-empty and satisfies Slater's constraint qualification; J_i is continuously differentiable and $J_i(\cdot, x_{-i})$ is convex for every x_{-i} . \square

Among all the possible GNEs, we focus on the subclass of variational GNE (v-GNE) [20, Def. 3.11]. Under the previous assumption, x^* is a v-GNE of the game in (2) if and only if there exist a dual variable $\lambda^* \in \mathbb{R}^m$ such that the following Karush-Kuhn-Tucker (KKT) conditions are satisfied [20, Th. 4.8]:

$$\begin{aligned} \mathbf{0}_n &\in F(x^*) + A^\top \lambda^* + N_\Omega(x^*) \\ \mathbf{0}_m &\in -(Ax^* - b) + N_{\mathbb{R}_{\geq 0}^m}(\lambda^*), \end{aligned} \quad (3)$$

where F is the *pseudo-gradient* mapping of the game:

$$F(x) := \text{col}((\nabla_{x_i} J_i(x_i, x_{-i}))_{i \in \mathcal{I}}). \quad (4)$$

We will assume strong monotonicity of the pseudo-gradient [6, Ass. 2], [10, Ass. 3], [14, Ass. 4], which is a sufficient condition for the existence of a unique v-GNE for the game in (2) [21, Th. 2.3.3]. This condition requires strong convexity of the functions $J_i(\cdot, x_{-i})$, for every x_{-i} , but not necessarily convexity of J_i in its full argument.

Standing Assumption 2: The pseudo-gradient mapping in (4) is μ -strongly monotone and θ_0 -Lipschitz continuous, for some $\mu, \theta > 0$: for any pair $x, y \in \mathbb{R}^n$, $(x - y)^\top(F(x) - F(y)) \geq \mu\|x - y\|^2$ and $\|F(x) - F(y)\| \leq \theta_0\|x - y\|$. \square

III. DISTRIBUTED GENERALIZED EQUILIBRIUM SEEKING

In this section, we consider the game in (2), where each agent is associated with a dynamical system:

$$\forall i \in \mathcal{I} : \quad \dot{x}_i = \Pi_{\Omega_i}(x_i, u_i), \quad x_i(0) \in \Omega_i. \quad (5)$$

Our aim is to design the inputs u_i to seek a v-GNE in a fully distributed way. Specifically, agent i does not have full knowledge of x_{-i} , and only relies on the information exchanged locally with neighbors over a communication network $\mathcal{G}(\mathcal{I}, \mathcal{E})$, with weighted symmetric Laplacian $L \in \mathbb{R}^{N \times N}$. The unordered pair (i, j) belongs to the set of edges, \mathcal{E} , if and only if agent j and i can exchange information.

Standing Assumption 3: The communication graph $\mathcal{G}(\mathcal{I}, \mathcal{E})$ is undirected and connected. \square

To cope with partial-information, each agent keeps an estimate of all other agents' actions. We denote $x^i =$

Algorithm 1 Distributed GNE seeking

$$\begin{aligned}
 \dot{x}_i &= \Pi_{\Omega_i}(x_i, u_i) \\
 u_i &= -(\nabla_{x_i} J_i(x_i, \mathbf{x}_{-i}^i) + A_i^\top \lambda_i + c \sum_{j \in \mathcal{N}_i} w_{ij}(x_i - \mathbf{x}_j^i)) \\
 \dot{\mathbf{x}}_{-i}^i &= -c \sum_{j \in \mathcal{N}_i} w_{ij}(\mathbf{x}_{-i}^i - \mathbf{x}_{-i}^j) \\
 \dot{z}_i &= \sum_{j \in \mathcal{N}_i} w_{ij}(\lambda_i - \lambda_j) \\
 \dot{\lambda}_i &= \Pi_{\mathbb{R}_{\geq 0}^m}(\lambda_i, A_i x_i - b_i - \sum_{j \in \mathcal{N}_i} w_{ij}(z_i - z_j + \lambda_i - \lambda_j))
 \end{aligned}$$

$\text{col}((\mathbf{x}_j^i)_{j \in \mathcal{I}}) \in \mathbb{R}^{Nn}$, where $\mathbf{x}_i^i := x_i$ and \mathbf{x}_j^i is i 's estimate of agent j 's action, for all $j \neq i$; $\mathbf{x}_{-i}^j = \text{col}((\mathbf{x}_\ell^j)_{\ell \in \mathcal{I} \setminus \{i\}})$. Also, each agent keeps an estimate $\lambda_i \in \mathbb{R}_{\geq 0}^m$ of the Lagrangian multiplier and an auxiliary variable $z_i \in \mathbb{R}^m$ to allow for distributed consensus of the multiplier estimates. Our closed-loop dynamics are shown in Algorithm 1, where $c > 0$ is a global constant parameter, $W = [w_{ij}]_{i,j \in \{1, \dots, N\}} \in \mathbb{R}^{N \times N}$ is the weighted adjacency matrix of the graph \mathcal{G} , \mathcal{N}_i is the set of neighbors of agent i and $\mathbf{x}_{-i}^i(0) \in \mathbb{R}^{n-1}$, $\lambda_i(0) \in \mathbb{R}_{\geq 0}^m$, $z_i(0) \in \mathbb{R}^m$ can be chosen arbitrarily.

The agents exchange $\{x^i, z_i, \lambda_i\}$ with their neighbors only, therefore the controller can be implemented distributedly. In steady state, the agents should agree on their estimates, i.e., $x^i = x^j$, $\lambda_i = \lambda_j$, for all $i, j \in \mathcal{I}$. This motivates the presence of consensual terms for both the primal and dual variables. We denote $\mathbf{E}_q := \{\mathbf{y} \in \mathbb{R}^{Nq} : \mathbf{y} = \mathbf{1}_N \otimes y, y \in \mathbb{R}^q\}$ the consensual subspace of dimension q , for some $q > 0$, and \mathbf{E}_q^\perp its orthogonal complement. Specifically, \mathbf{E}_n is the estimate consensus subspace and \mathbf{E}_m is the multiplier consensus subspace. To write the dynamics in compact form, let us define $\mathbf{x} = \text{col}((x^i)_{i \in \mathcal{I}})$, and, as in [15, Eq. 13-14], for all $i \in \mathcal{I}$,

$$\mathcal{R}_i := \begin{bmatrix} 0_{n_i \times n_{<i}} & I_{n_i} & \mathbf{0}_{n_i \times n_{>i}} \end{bmatrix}, \quad (6a)$$

$$\mathcal{S}_i := \begin{bmatrix} I_{n_{<i}} & \mathbf{0}_{n_{<i} \times n_i} & \mathbf{0}_{n_{<i} \times n_{>i}} \\ \mathbf{0}_{n_{>i} \times n_{<i}} & \mathbf{0}_{n_{>i} \times n_i} & I_{n_{>i}} \end{bmatrix}, \quad (6b)$$

where $n_{<i} := \sum_{j < i, j \in \mathcal{I}} n_j$, $n_{>i} := \sum_{j > i, j \in \mathcal{I}} n_j$. In simple terms, \mathcal{R}_i selects the i -th n_i dimensional component from an n -dimensional vector, while \mathcal{S}_i removes it. Thus, $\mathcal{R}_i \mathbf{x}^i = x_i$ and $\mathcal{S}_i \mathbf{x}^i = \mathbf{x}_{-i}^i$. We define $\mathcal{R} := \text{diag}((\mathcal{R}_i)_{i \in \mathcal{I}})$, $\mathcal{S} := \text{diag}((\mathcal{S}_i)_{i \in \mathcal{I}})$. It follows that $x = \mathcal{R} \mathbf{x}$, $\text{col}((\mathbf{x}_{-i}^i)_{i \in \mathcal{I}}) = \mathcal{S} \mathbf{x} \in \mathbb{R}^{(N-1)n}$, and $\mathbf{x} = \mathcal{R}^\top x + \mathcal{S}^\top \mathcal{S} \mathbf{x}$. Let $\boldsymbol{\lambda} := \text{col}((\lambda_i)_{i \in \mathcal{I}})$, $\boldsymbol{\Lambda} := \text{diag}((A_i)_{i \in \mathcal{I}})$, $\mathbf{b} := \text{col}((b_i)_{i \in \mathcal{I}})$, $\mathbf{L}_x := L \otimes I_n$, $\mathbf{L}_\lambda := L \otimes I_m$, $\mathbf{z} := \text{col}((z_i)_{i \in \mathcal{I}})$. Further, we define the *extended pseudo-gradient* mapping \mathbf{F} as:

$$\mathbf{F}(\mathbf{x}) := \text{col}((\nabla_{x_i} J_i(x_i, \mathbf{x}_{-i}^i))_{i \in \mathcal{I}}). \quad (7)$$

The overall closed-loop system, in compact form, reads as:

$$\dot{\mathbf{x}} = \mathcal{R}^\top \Pi_{\Omega}(\mathcal{R} \mathbf{x}, -(\mathbf{F}(\mathbf{x}) + \boldsymbol{\Lambda}^\top \boldsymbol{\lambda} + c \mathcal{R} \mathbf{L}_x \mathbf{x})) + \mathcal{S}^\top (-c \mathcal{S} \mathbf{L}_x \mathbf{x}) \quad (8a)$$

$$\dot{\mathbf{z}} = \mathbf{L}_\lambda \boldsymbol{\lambda} \quad (8b)$$

$$\dot{\boldsymbol{\lambda}} = \Pi_{\mathbb{R}_{\geq 0}^m}(\boldsymbol{\lambda}, (\boldsymbol{\Lambda} \mathcal{R} \mathbf{x} - \mathbf{b} - \mathbf{L}_\lambda \boldsymbol{\lambda} - \mathbf{L}_\lambda \mathbf{z})). \quad (8c)$$

The following Lemma relates the equilibria of the system in (8) to the v-GNE of the game in (2).

The proof is analogous to [15, Th. 1], hence it is omitted.

Lemma 2: The following statements hold:

- i) Any equilibrium point $\bar{\omega} = \text{col}(\bar{x}, \bar{z}, \bar{\lambda})$ of the dynamics in (8) is such that $\bar{x} = \mathbf{1}_N \otimes x^*$, $\bar{\lambda} = \mathbf{1}_N \otimes \lambda^*$, where the pair (x^*, λ^*) satisfies the KKT conditions in (3), hence x^* is the v-GNE of the game in (2);
- ii) The set of equilibrium points of (8) is nonempty. \square

Remark 1: When considering Algorithm 1 in absence of coupling constraints, we retrieve the controller in [6, Eq. 47]. In Algorithm 1, each agent evaluates the gradient of its cost function in its local estimate, not on the actual collective strategy. In fact, only when the estimates belong to the consensus subspace, i.e., $\mathbf{x} = \mathbf{1} \otimes x$ (i.e., the estimate \mathbf{x}^i of each agent coincide with the real actions x , for example in the case of full-information), we have that $\mathbf{F}(\mathbf{x}) = F(x)$. It follows that the operator $\mathcal{R}^\top \mathbf{F}$ is not necessarily monotone, not even if the pseudo gradient F in (4) is strongly monotone, as in Standing Assumption 2. This is the main technical difficulty that arises in (G)NE problems under partial-information. \square

Lemma 3: The extended pseudo-gradient mapping \mathbf{F} in (7) is θ -Lipschitz continuous, for some $\mu \leq \theta \leq \theta_0$: for any $\mathbf{x}, \mathbf{y} \in \mathbb{R}^{Nn}$, $\|\mathbf{F}(\mathbf{x}) - \mathbf{F}(\mathbf{y})\| \leq \theta \|\mathbf{x} - \mathbf{y}\|$. \square

Proof: See Appendix VI-A. \blacksquare

Under Lipschitz continuity of \mathbf{F} , the work [15] showed the following *restricted strong monotonicity* property, which is crucial to prove convergence of the dynamics in (8).

Lemma 4 ([15, Lem. 3]): Let

$$\mathcal{M} := \begin{bmatrix} \frac{\mu}{N} & -\frac{\theta_0 + \theta}{2\sqrt{N}} \\ -\frac{\theta_0 + \theta}{2\sqrt{N}} & c\lambda_2(L) - \theta \end{bmatrix}, \quad \underline{c} := \frac{(\theta_0 + \theta)^2 + 4\mu\theta}{4\mu\lambda_2(L)}. \quad (9)$$

For any $c > \underline{c}$, for any \mathbf{x} and any $\mathbf{x}' \in \mathbf{E}_n$, it holds that $\mathcal{M} \succ 0$ and that

$$\begin{aligned}
 (\mathbf{x} - \mathbf{x}')^\top (\mathcal{R}^\top (\mathbf{F}(\mathbf{x}) - \mathbf{F}(\mathbf{x}')) + c \mathbf{L}_x (\mathbf{x} - \mathbf{x}')) \\
 \geq \lambda_{\min}(\mathcal{M}) \|\mathbf{x} - \mathbf{x}'\|^2. \quad \square
 \end{aligned}$$

By leveraging Lemma 4, we can now prove the main result of this section, i.e., the convergence of the dynamics in (8) to a v-GNE.

Theorem 1: Let $c > \underline{c}$, with \underline{c} as in (9), and let $\Omega := \{\mathbf{x} \in \mathbb{R}^{Nn} \mid \mathcal{R} \mathbf{x} \in \Omega\}$. For any initial condition in $\Xi = \Omega \times \mathbb{R}^{mN} \times \mathbb{R}_{\geq 0}^m$, the dynamics in (8) have a unique Carathéodory solution, which belongs to Ξ for all $t \geq 0$. The solution converges to an equilibrium $\text{col}(\bar{x}, \bar{z}, \bar{\lambda})$, with $\bar{x} = \mathbf{1}_N \otimes x^*$, $\bar{\lambda} = \mathbf{1}_N \otimes \lambda^*$, where the pair (x^*, λ^*) satisfies the KKT conditions in (3), hence x^* is the v-GNE of the game in (2). \square

Proof: See Appendix VI-B. \blacksquare

Remark 2: In Algorithm 1, each agent keeps and exchanges an estimate of the strategies of all other agents. Thus, the computation and communication costs increase with the number of agents. An open research direction is to design dynamics that allow each agent to estimate the strategies of only some of its competitors, when the inference graph is sparse (i.e., when the cost of each agent only depends on the action of a limited subset of other agents). \square

IV. DOUBLE-INTEGRATOR AGENTS

In this section, we make the following additional assumption.

Assumption 1 ([18, Ass. 1]): $\Omega = \mathbb{R}^n$. \square

Moreover, we model each agent as a double integrator:

$$\forall i \in \mathcal{I} : \begin{cases} \dot{x}_i = v_i & (10a) \\ \dot{v}_i = u_i, & (10b) \end{cases}$$

where (x_i, v_i) is the state of agent i , and $u_i \in \mathbb{R}^{n_i}$ its control input. Our objective is to drive the agents' actions, i.e., the x_i coordinates of their state, to a v-GNE of the game in (2). We emphasize that in (10) we cannot directly control the agent's action. Moreover, at steady state, the velocities v_i of all the agents must be zero. This scenario has been considered recently in [18], for games *without* coupling constraints.

In (10), we consider the input $u_i = \frac{1}{h_i}(\tilde{u}_i - v_i)$, where $h_i > 0$ is a positive scalar and \tilde{u}_i has to be chosen appropriately, for all $i \in \mathcal{I}$; moreover, as in [18], let us define the coordinates transformation

$$\zeta_i := x_i + h_i v_i. \quad (11)$$

The quantity ζ_i can be interpreted as a prediction of the position of agent i , given a forward step h_i . The closed-loop system in the new coordinates then reads as

$$\forall i \in \mathcal{I} : \begin{cases} \dot{v}_i = \frac{1}{h_i}(\tilde{u}_i - v_i) & (12a) \\ \dot{\zeta}_i = \tilde{u}_i. & (12b) \end{cases}$$

We note that the dynamics of the variable ζ_i in (12b), under Assumption 1, are identical to the single-integrator in (5), with translated input \tilde{u}_i . As such, we are able to design the input \tilde{u}_i , according to Algorithm 1, to drive $\zeta := \text{col}((\zeta_i)_{i \in \mathcal{I}})$ to an equilibrium $\bar{\zeta} = x^*$, where x^* is the v-GNE for the game in (2). Moreover, the velocity dynamics (12a) are Input-to-state-stable (ISS) with respect to the input u_i [22, Lemma 4.6]. Finally, we remark that, at any equilibrium of (10), $v_i = \mathbf{0}_{n_i}$, hence $\zeta_i = x_i$, for all $i \in \mathcal{I}$. Building on this considerations, we propose Algorithm 2 to drive the double-integrator agents (10) towards a v-GNE.

Differently from Algorithm 1, the agents are not keeping an estimate of other agents' actions, but of other agents' *predictions*. Here, $\zeta^i = \text{col}((\zeta_j^i)_{j \in \mathcal{I}})$, and ζ_j^i represents agent i 's estimation of the quantity $\zeta_j = x_j + h_j v_j$ for $j \neq i$, while $\zeta_i^i := x_i + h_i v_i = \zeta_i$. Let us denote $\zeta = \text{col}((\zeta^i)_{i \in \mathcal{I}})$, $\zeta_{-i}^j = \text{col}((\zeta_\ell^j)_{\ell \in \mathcal{I} \setminus \{i\}})$, $H = \text{diag}((h_i I_{n_i})_{i \in \mathcal{I}})$.

Algorithm 2 Distributed GNE seeking (double-integrators)

$$\begin{aligned} \dot{x}_i &= v_i, & \dot{v}_i &= u_i \\ u_i &= -\frac{1}{h_i}(\nabla_{x_i} J_i(\zeta^i, \zeta_{-i}^i) + A_i^\top \lambda_i \\ &\quad + c \sum_{j \in \mathcal{N}_i} w_{ij}(\zeta^i - \zeta^j)) - \frac{1}{h_i} v_i \\ \dot{\zeta}_{-i}^i &= -\sum_{j \in \mathcal{N}_i} w_{ij}(\zeta_{-i}^i - \zeta_{-i}^j), & \zeta^i &= x_i + h_i v_i \\ \dot{z}_i &= \sum_{j \in \mathcal{N}_i} w_{ij}(\lambda_i - \lambda_j) \\ \dot{\lambda}_i &= \Pi_{\mathbb{R}_{\geq 0}^m}(\lambda_i, A_i \zeta^i - b_i - \sum_{j \in \mathcal{N}_i} w_{ij}(z_i - z_j + \lambda_i - \lambda_j)) \end{aligned}$$

In compact form, the closed-loop system reads as

$$\dot{x} = v \quad (13a)$$

$$H \dot{v} = -(\mathbf{F}(\zeta) + \Lambda^\top \lambda + c \mathcal{R} L_x \zeta) - v \quad (13b)$$

$$S \dot{\zeta} = -c S L_x \zeta, \quad \mathcal{R} \zeta = x + H v \quad (13c)$$

$$\dot{z} = L_\lambda \lambda \quad (13d)$$

$$\dot{\lambda} = \Pi_{\mathbb{R}_{\geq 0}^{Nm}}(\lambda, (\Lambda \mathcal{R} \zeta - b - L_\lambda \lambda - L_\lambda z)). \quad (13e)$$

Theorem 2: Let Assumption 1 hold. For any initial condition with $\lambda(0) \in \mathbb{R}_{\geq 0}^{Nm}$, the equations in (13) have a unique Carathéodory solution, such that $\lambda(t) \in \mathbb{R}_{\geq 0}^{Nm}$, for all $t \geq 0$. The solution converges to an equilibrium $\text{col}(\bar{x}, \bar{v}, \bar{\zeta}, \bar{z}, \bar{\lambda})$, with $\bar{x} = x^*$, $\bar{v} = \mathbf{0}_n$, $\bar{\zeta} = \mathbf{1}_N \otimes x^*$, $\bar{\lambda} = \mathbf{1}_N \otimes \lambda^*$, where the pair (x^*, λ^*) satisfies the KKT conditions in (3), so x^* is a v-GNE for the game in (2). \square

Proof: See Appendix VI-C. \blacksquare

Algorithm 2 is derived by choosing \tilde{u}_i in (12) according to Algorithm 1. The proof of Theorem 2 is not based on the specific structure of Algorithm 1, but only on its convergence properties, hence the result still holds if another controller with similar features is selected in place of Algorithm 1. In [18], the authors addressed NE problems and chose the inputs \tilde{u}_i according to the algorithm presented in [6, Eq. 47]. The controller in [6] achieves exponential convergence to a NE, hence ISS with respect to possible additive disturbances [22, Lemma 4.6]. Therefore, in [18], the authors were able to tackle the presence of deterministic disturbances, via an asymptotic observer and by leveraging ISS arguments. We have not guaranteed this robustness, i.e., exponential convergence, for the primal-dual dynamics in (8). However, the controller in [18] is designed for games without any constraints (local or shared). On the contrary, the controller in Algorithm 2 drives the system in (10) to a v-GNE of a generalized game, and ensures for the coupling constraints to be satisfied asymptotically. Also, like in [18], we assumed the absence of local constraints (Assumption 1). Nevertheless, if some are present, they can be included in the coupling constraints, hence dualized and satisfied asymptotically.

V. NUMERICAL EXAMPLE: MOBILE SENSOR NETWORK

We consider a numerical example, inspired by connectivity control problems for sensor networks [3], [18]. Each of five agent is represented by a robot/vehicle, moving in a plane, designed to optimize some private primary objective related to its position, provided that overall connectivity is preserved over the network. For each agent $i \in \mathcal{I} = \{1, \dots, 5\}$, its cost function is $J_i(p_i, p_{-i}) := p_i^\top p_i + p_i^\top r_i + \sum_{j \in \mathcal{I}} \|p_i - p_j\|^2$, with $p_i = \text{col}(x_i, y_i)$ its cartesian coordinates, $r_i \in \mathbb{R}^2$ a local parameter. We assume the local constraints $0.1 \leq y_i \leq 0.5$, $\forall i \in \mathcal{I}$. In order for all the agents to maintain communication with their neighbors, we impose the Chebyshev distance between any two neighboring robots to be smaller than $0.2m$. Hence the (affine) coupling constraints are represented by $\max\{|x_i - x_j|, |y_i - y_j|\} \leq 0.2, \forall (i, j) \in \mathcal{E}$. As common for autonomous vehicles, we model the agents as single- or double-integrators.




Fig. 1. Results of Alg. 1 for single- and double-integrator agents.




Fig. 2. Velocity-actuated and force-actuated robots trajectories.

Velocity-actuated robots: each agent is modeled as in (5) and we apply Algorithm 1.

Force-actuated robots : Each agent has a dynamic as in (10), under Algorithm 2. The local constraints are dualized and will be satisfied asymptotically (see Section IV).

The initial conditions are chosen randomly and we fix $c = 30$ to satisfy the condition in Theorem 1. Figure 1 illustrates the results for the two cases and shows convergence to the GNE of the game and asymptotic satisfaction of the coupling constraints. Finally, in Figure 2, we compare the trajectories of the five robots in the velocity-actuated and force-actuated scenario. In the two cases, the agents are converging to the same, unique v-GNE. However, the local constraints are satisfied along the whole trajectory for single integrator agents, only asymptotically for the double integrator agents.

VI. CONCLUSION AND OUTLOOK

Generalized games played by double-integrator agents can be solved via a fully distributed primal-dual projected-pseudogradient dynamic controller, if the game mapping is strongly monotone and Lipschitz continuous. Seeking an equilibrium in games with compact action sets or constrained dynamics is currently an unexplored problem. The extension of our results to networks of heterogeneous dynamical systems is left as future research.

APPENDIX

A. Proof of Lemma 3

Let us define $\mathbf{x} = \text{col}((\mathbf{x}^i)_{i \in \mathcal{I}})$, $\mathbf{y} = \text{col}((\mathbf{y}^i)_{i \in \mathcal{I}})$. By Standing Assumption 2, we have, for all $i \in \mathcal{I}$, $\|\nabla_{\mathbf{x}_i} J_i(\mathbf{x}^i) - \nabla_{\mathbf{x}_i} J_i(\mathbf{y}^i)\| \leq \|F(\mathbf{x}^i) - F(\mathbf{y}^i)\| \leq \theta_0 \|\mathbf{x}^i - \mathbf{y}^i\|$. Therefore it holds that

$$\begin{aligned} \|\mathbf{F}(\mathbf{x}) - \mathbf{F}(\mathbf{y})\|^2 &= \sum_{i \in \mathcal{I}} \|\nabla_{\mathbf{x}_i} J_i(\mathbf{x}^i) - \nabla_{\mathbf{x}_i} J_i(\mathbf{y}^i)\|^2 \\ &\leq \theta_0^2 \sum_{i \in \mathcal{I}} \|\mathbf{x}^i - \mathbf{y}^i\|^2 = \theta_0^2 \|\mathbf{x} - \mathbf{y}\|^2. \end{aligned}$$

That $\theta \geq \mu$ follows by choosing $\mathcal{S}\mathbf{x} = \mathcal{S}\mathbf{y}$, $\mathbf{x} \neq \mathbf{y}$. \blacksquare

B. Proof of Theorem 1

Under Standing Assumption 3, we have, for any $q > 0$,

$$\text{Null}(L \otimes I_q) = \text{Range}(\mathbf{1}_N \otimes I_q) = \mathbf{E}_q. \quad (14)$$

We first rewrite the dynamics as

$$\dot{\boldsymbol{\omega}} = \Pi_{\Xi}(\boldsymbol{\omega}, -\mathcal{B}(\boldsymbol{\omega}) - \Phi\boldsymbol{\omega}), \quad (15)$$

where $\boldsymbol{\omega} = \text{col}(\mathbf{x}, \mathbf{z}, \boldsymbol{\lambda})$,

$$\Phi = \begin{bmatrix} 0 & 0 & \mathcal{R}^\top \Lambda^\top \\ 0 & 0 & -L_\lambda \\ -\Lambda \mathcal{R} & L_\lambda & 0 \end{bmatrix}, \quad \mathcal{B}(\boldsymbol{\omega}) = \begin{bmatrix} \mathcal{R}^\top \mathbf{F}(\mathbf{x}) + cL_x \mathbf{x} \\ \mathbf{0}_{N \times m} \\ L_\lambda \boldsymbol{\lambda} + \mathbf{b} \end{bmatrix}.$$

\mathcal{B} is Lipschitz continuous by Standing Assumption 2 and Ξ is closed and convex by Standing Assumption 1. We conclude that there exists a unique Carathéodory solution to (15), that belongs to Ξ for every $t \geq 0$. Consider the quadratic Lyapunov function

$$V(\boldsymbol{\omega}) = \frac{1}{2} \|\boldsymbol{\omega} - \bar{\boldsymbol{\omega}}\|^2,$$

where $\bar{\boldsymbol{\omega}} := \text{col}(\bar{\mathbf{x}}, \bar{\mathbf{z}}, \bar{\boldsymbol{\lambda}})$ is any equilibrium of (8). We remark that, by Lemma 2, an equilibrium exists, and $\bar{\mathbf{x}} = \mathbf{1}_N \otimes x^*$, $\bar{\boldsymbol{\lambda}} = \mathbf{1}_N \otimes \boldsymbol{\lambda}^*$, with $(x^*, \boldsymbol{\lambda}^*)$ satisfying the KKT conditions in (3). We can apply Lemma 1 to obtain

$$\begin{aligned} \dot{V}(\boldsymbol{\omega}) &:= \nabla V(\boldsymbol{\omega}) \dot{\boldsymbol{\omega}} = (\boldsymbol{\omega} - \bar{\boldsymbol{\omega}})^\top \Pi_{\Xi}(\boldsymbol{\omega}, -\mathcal{B}(\boldsymbol{\omega}) - \Phi\boldsymbol{\omega}) \\ &\leq (\boldsymbol{\omega} - \bar{\boldsymbol{\omega}})^\top (-\mathcal{B}(\boldsymbol{\omega}) - \Phi\boldsymbol{\omega}). \end{aligned} \quad (16)$$

By Lemma 1, it also holds that $(\boldsymbol{\omega} - \bar{\boldsymbol{\omega}})^\top (-\mathcal{B}(\bar{\boldsymbol{\omega}}) - \Phi\bar{\boldsymbol{\omega}}) \leq 0$. By subtracting this term from (16), we obtain

$$\begin{aligned} \dot{V} &\leq -(\boldsymbol{\omega} - \bar{\boldsymbol{\omega}})^\top (\mathcal{B}(\boldsymbol{\omega}) - \mathcal{B}(\bar{\boldsymbol{\omega}}) + \Phi(\boldsymbol{\omega} - \bar{\boldsymbol{\omega}})) \\ &= -(\mathbf{x} - \bar{\mathbf{x}})^\top \mathcal{R}^\top (\mathbf{F}(\mathbf{x}) - \mathbf{F}(\bar{\mathbf{x}})) \\ &\quad - (\mathbf{x} - \bar{\mathbf{x}})^\top cL_x (\mathbf{x} - \bar{\mathbf{x}}) - (\boldsymbol{\lambda} - \bar{\boldsymbol{\lambda}})^\top L_\lambda (\boldsymbol{\lambda} - \bar{\boldsymbol{\lambda}}), \end{aligned} \quad (17)$$

where, in the last equality, we used that, $\Phi^\top = -\Phi$. By (14) and [19, Cor. 18.16], we have that $(\boldsymbol{\lambda} - \bar{\boldsymbol{\lambda}})^\top L_\lambda (\boldsymbol{\lambda} - \bar{\boldsymbol{\lambda}}) \geq \frac{1}{2\lambda_{\max}(L)} \|\mathbf{L}_\lambda \boldsymbol{\lambda}\|^2$. Finally, by Lemma 4, we obtain

$$\dot{V} \leq -\lambda_{\min}(\mathcal{M}) \|\mathbf{x} - \bar{\mathbf{x}}\|^2 - \frac{1}{2\lambda_{\max}(L)} \|\mathbf{L}_\lambda \boldsymbol{\lambda}\|^2 \leq 0, \quad (18)$$

with $\mathcal{M} \succ 0$ as in (9). By noticing that V is radially unbounded, we conclude that the solution to (8) is bounded. Besides, by [14, Th. 2], the solution converges to the largest invariant set \mathcal{O} contained in $\mathcal{Z} := \{\boldsymbol{\omega} \text{ s.t. } \dot{V}(\boldsymbol{\omega}) = 0\}$.

We first characterize any point $\text{col}(\hat{\mathbf{x}}, \hat{\mathbf{z}}, \hat{\boldsymbol{\lambda}}) \in \mathcal{Z}$, for which the quantities in (16)-(18) must be zero. By (18), $\hat{\mathbf{x}} = \bar{\mathbf{x}} = \mathbf{1}_N \otimes x^*$, and $\hat{\boldsymbol{\lambda}} \in \mathbf{E}_m$, i.e. $\hat{\boldsymbol{\lambda}} = \mathbf{1}_N \otimes \hat{\boldsymbol{\lambda}}$, for

some $\hat{\lambda} \in \mathbb{R}_{\geq 0}^m$. Also, by expanding (16), and by $\hat{x} = \bar{x}$, $L_\lambda \hat{\lambda} = 0$, we have

$$\begin{aligned} 0 &= (\hat{\lambda} - \bar{\lambda})^\top (\Lambda \mathcal{R} \bar{x} - \mathbf{b} - L_\lambda \hat{z}) = (\hat{\lambda} - \lambda^*)^\top (Ax^* - b) \\ &= \hat{\lambda}^\top (Ax^* - b) = \hat{\lambda}^\top (\Lambda \mathcal{R} \bar{x} - \mathbf{b} - L_\lambda \hat{\lambda} - L_\lambda \hat{z}), \end{aligned} \quad (19)$$

where in the second equality we have used (14) and the fourth equality follows from the KKT conditions in (3). This concludes the characterization of the set \mathcal{Z} .

By invariance, any trajectory $\omega(t) = \text{col}(\mathbf{x}(t), \mathbf{z}(t), \boldsymbol{\lambda}(t))$ starting at any $\text{col}(\mathbf{x}, \mathbf{z}, \boldsymbol{\lambda}) \in \mathcal{O}$ must lie in $\mathcal{Z} \supset \mathcal{O}$ for all $t \geq 0$. Therefore, $\mathbf{x}(t) \equiv \bar{x}$ and $\boldsymbol{\lambda}(t) \in \mathbf{E}_m$ for all t . Moreover, $\dot{\mathbf{z}}(t) = 0$, for all t , by (8b), or $\mathbf{z}(t) \equiv \bar{z}$. Hence the quantity $v := (\Lambda \mathcal{R} \mathbf{x}(t) - \mathbf{b} - L_\lambda \boldsymbol{\lambda}(t) - L_\lambda \mathbf{z}(t))$ is constant along the trajectory ω . Suppose by contradiction that $v_k > 0$, where v_k denotes the k -th component of v . Then, by (8c), $\dot{\boldsymbol{\lambda}}(t)_k = v_k$ for all t , and $\boldsymbol{\lambda}(t)$ grows indefinitely. Since all the solutions of (8) are bounded, this is a contradiction. Therefore, $v \leq 0$, and $\boldsymbol{\lambda}(t)^\top v = 0$ by (19). Equivalently, $v \in \mathbb{N}_{\mathbb{R}_{\geq 0}^m}(\boldsymbol{\lambda}(t))$, hence $\boldsymbol{\lambda}(t) = 0$, for all t . We conclude that all the points in the set \mathcal{O} are equilibria.

The set $\Lambda(\omega_0)$ of ω -limit points¹ of the solution to (8) starting from any $\omega_0 \in \Xi$ is nonempty (by Bolzano-Weierstrass theorem, since all the trajectories of (8) are bounded) and invariant (as in proof of [14, Lemma 5]). By $\dot{V} \leq 0$ it follows that V must be constant on $\Lambda(\omega_0)$, hence $\Lambda(\omega_0) \subseteq \mathcal{Z}$ (see proof of [14, Th.2]). Also $\Lambda(\omega_0)$ is invariant, so $\Lambda(\omega_0) \subseteq \mathcal{O}$. Since the distance to any equilibrium point along any trajectory of (8) is non-increasing by (18), it follows that if a solution of (8) has an ω -limit point at an equilibrium, then the solution converges to that equilibrium. ■

C. Proof of Theorem 2

By applying the coordinate transformation $x \mapsto \mathcal{R}\zeta = x + Hv$ to the system in (13), we obtain:

$$\dot{v} = -H^{-1}(\mathbf{F}(\zeta) + \Lambda^\top \boldsymbol{\lambda} + c\mathcal{R}L_x \zeta) - H^{-1}v \quad (20a)$$

$$\dot{\zeta} = -\mathcal{R}^\top(\mathbf{F}(\zeta) + \Lambda^\top \boldsymbol{\lambda} + c\mathcal{R}L_x \zeta) - c\mathcal{S}^\top \mathcal{S}L_x \zeta \quad (20b)$$

$$\dot{z} = L_\lambda \boldsymbol{\lambda} \quad (20c)$$

$$\dot{\boldsymbol{\lambda}} = \Pi_{\mathbb{R}_{\geq 0}^m}(\boldsymbol{\lambda}, (\Lambda \mathcal{R} \zeta - \mathbf{b} - L_\lambda \boldsymbol{\lambda} - L_\lambda z)). \quad (20d)$$

The system (20) is in cascade form for (20a) with respect to (20b)-(20d). Notice also that, under Assumption 1, the subsystem (20b)-(20d) is exactly (8). Hence, there exists a unique solution to (20b)-(20d), that is bounded and converges to an equilibrium point $\text{col}(\mathbf{1}_N \otimes x^*, \bar{z}, \mathbf{1}_N \otimes \lambda^*)$, where the pair (x^*, λ^*) satisfies the KKT conditions in (3), by Theorem 1. On the other hand, the dynamic (20a) is ISS with respect to the input $\tilde{u} := -H^{-1}(\mathbf{F}(\zeta) + \Lambda^\top \boldsymbol{\lambda} + c\mathcal{R}L_x \zeta)$ [22, Lemma 4.6], and this input is bounded, by boundedness of the trajectory $(\zeta, \mathbf{k}, z, \boldsymbol{\lambda})$ and Lemma 3. Moreover, since $\bar{\zeta} = \mathbf{1}_N \otimes x^*$, $\bar{\boldsymbol{\lambda}} = \mathbf{1}_N \otimes \lambda^*$, by the KKT conditions in (3) and by continuity, we have $\tilde{u} \rightarrow \mathbf{0}_n$ for $t \rightarrow \infty$. Therefore, $v(t) \rightarrow \mathbf{0}_n$ for $t \rightarrow \infty$ [22, Ex. 4.58]. By definition of $\zeta_i = \mathcal{R}_i \zeta^i$ in (11), we can also conclude that $x \rightarrow x^*$. ■

¹ $z : [0, \infty) \rightarrow \mathbb{R}^n$ has an ω -limit point at \bar{z} if there exists a nonnegative diverging sequence $\{t_k\}_{k \in \mathbb{N}}$ such that $z(t_k) \rightarrow \bar{z}$.

REFERENCES

- [1] W. Saad, Z. Han, H. V. Poor, and T. Basar, "Game-theoretic methods for the smart grid: An overview of microgrid systems, demand-side management, and smart grid communications," *IEEE Signal Processing Magazine*, vol. 29, no. 5, pp. 86–105, 2012.
- [2] S. Grammatico, "Dynamic control of agents playing aggregative games with coupling constraints," *IEEE Transactions on Automatic Control*, vol. 62, no. 9, pp. 4537–4548, 2017.
- [3] M. S. Stankovic, K. H. Johansson, and D. M. Stipanovic, "Distributed seeking of Nash equilibria with applications to mobile sensor networks," *IEEE Transactions on Automatic Control*, vol. 57, no. 4, pp. 904–919, 2012.
- [4] F. Facchinei and J. Pang, "Nash equilibria: the variational approach," in *Convex Optimization in Signal Processing and Communications*, D. P. Palomar and Y. C. Eldar, Eds. Cambridge University Press, 2009, p. 443–493.
- [5] C. De Persis and N. Monshizadeh, "A feedback control algorithm to steer networks to a Cournot–Nash equilibrium," *IEEE Transactions on Control of Network Systems*, vol. 6, no. 4, pp. 1486–1497, 2019.
- [6] D. Gadjov and L. Pavel, "A passivity-based approach to Nash equilibrium seeking over networks," *IEEE Transactions on Automatic Control*, vol. 64, no. 3, pp. 1077–1092, 2019.
- [7] C. De Persis and S. Grammatico, "Continuous-time integral dynamics for a class of aggregative games with coupling constraints," *IEEE Transactions on Automatic Control*, DOI: 10.1109/TAC.2019.2939639, 2019.
- [8] P. Yi and L. Pavel, "An operator splitting approach for distributed generalized Nash equilibria computation," *Automatica*, vol. 102, pp. 111 – 121, 2019.
- [9] C. Yu, M. van der Schaar, and A. H. Sayed, "Distributed learning for stochastic generalized Nash equilibrium problems," *IEEE Transactions on Signal Processing*, vol. 65, no. 15, pp. 3893–3908, 2017.
- [10] G. Belgioioso and S. Grammatico, "Projected-gradient algorithms for generalized equilibrium seeking in aggregative games are preconditioned forward-backward methods," *2018 European Control Conference (ECC)*, pp. 2188–2193, 2018.
- [11] P. Frihauf, M. Krstic, and T. Basar, "Nash equilibrium seeking in noncooperative games," *IEEE Transactions on Automatic Control*, vol. 57, no. 5, pp. 1192–1207, 2012.
- [12] F. Salehisadaghiani, W. Shi, and L. Pavel, "Distributed Nash equilibrium seeking under partial-decision information via the alternating direction method of multipliers," *Automatica*, vol. 103, pp. 27 – 35, 2019.
- [13] J. Koshal, A. Nedic, and U. V. Shanbhag, "Distributed algorithms for aggregative games on graphs," *Operations Research*, vol. 64, pp. 680–704, 2016.
- [14] C. De Persis and S. Grammatico, "Distributed averaging integral Nash equilibrium seeking on networks," *Automatica*, DOI: 10.1016/j.automatica.2019.108548, 2019.
- [15] L. Pavel, "Distributed GNE seeking under partial-decision information over networks via a doubly-augmented operator splitting approach," *IEEE Transactions on Automatic Control*, DOI: 10.1109/TAC.2019.2922953, 2019.
- [16] Z. Deng and X. Nian, "Distributed generalized Nash equilibrium seeking algorithm design for aggregative games over weight-balanced digraphs," *IEEE Transactions on Neural Networks and Learning Systems*, vol. 30, no. 3, pp. 695–706, 2019.
- [17] Y. Zhang, S. Liang, X. Wang, and H. Ji, "Distributed Nash equilibrium seeking for aggregative games with nonlinear dynamics under external disturbances," *IEEE Transactions on Cybernetics*, pp. 1–10, 2019.
- [18] A. R. Romano and L. Pavel, "Dynamic Nash equilibrium seeking for higher-order integrators in networks," in *2019 18th European Control Conference (ECC)*, 2019, pp. 1029–1035.
- [19] H. H. Bauschke and P. L. Combettes, *Convex analysis and monotone operator theory in Hilbert spaces*. Springer, 2017, vol. 2011.
- [20] F. Facchinei and C. Kanzow, "Generalized Nash equilibrium problems," *Annals of Operations Research*, vol. 175, pp. 177–211, 2010.
- [21] F. Facchinei and J. Pang, *Finite-dimensional variational inequalities and complementarity problems*. Springer Science & Business Media, 2007.
- [22] H. K. Khalil, *Nonlinear Systems*, ser. Pearson Education. Prentice Hall, 2002.