

Maintaining Tropical Beaches with Seagrass and Algae A Promising Alternative to Engineering Solutions

James, Rebecca K.; Pietrzak, Julie; Candy, Adam; Katsman, Caroline; van der Boog, C.G.; Riva, Riccardo; Slobbe, Cornelis; Klees, Roland; More Authors

DOI

[10.1093/biosci/biy154](https://doi.org/10.1093/biosci/biy154)

Publication date

2019

Document Version

Accepted author manuscript

Published in

BioScience

Citation (APA)

James, R. K., Pietrzak, J., Candy, A., Katsman, C., van der Boog, C. G., Riva, R., Slobbe, C., Klees, R., & More Authors (2019). Maintaining Tropical Beaches with Seagrass and Algae: A Promising Alternative to Engineering Solutions. *BioScience*, 69(2), 136-142. Article 154. <https://doi.org/10.1093/biosci/biy154>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Maintaining Tropical Beaches with Seagrass and Algae: A Promising Alternative to Engineering Solutions

Running head: Maintaining beaches with seagrass and algae

Authors:

Rebecca K. James (MSc) Department of Estuarine and Delta Systems, NIOZ Royal Netherlands Institute for Sea Research and Utrecht University, Korringaweg 7, 4401 NT Yerseke, The Netherlands. Rebecca.james@nioz.nl

Rodolfo Silva Casarín (Dr.), Instituto de Ingeniería, Universidad Nacional Autónoma de México, Circuito Escolar s/n, Edificio 5, Mexico City, 04510, Mexico.

rsilvac@iingen.unam.mx

Brigitta I. van Tussenbroek (Dr.), Unidad Académica de Sistemas Arrecifales-Puerto Morelos, Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, Prolongación Avenida Niños Héroes S/N, Puerto Morelos, Quintana Roo 77580, Mexico.

vantuss@cmarl.unam.mx

Mireille Escudero-Castillo (Dr.), Instituto de Ingeniería, Universidad Nacional Autónoma de México, Circuito Escolar s/n, Edificio 5, Mexico City, 04510, Mexico.

mescuderoc@iingen.unam.mx

Ismael Mariño-Tapia (Dr.), Departamento de Recursos del Mar, Cinvestav, km 6 Carretera Antigua a Progreso, Cordemex, 97310, A.S.P. 73, Mérida, Yuc., Mexico.

imarinotapia@gmail.com

23 Henk A. Dijkstra (Prof. Dr. ir.), Institute for Marine and Atmospheric research Utrecht,
24 Department of Physics, Utrecht University, Princetonplein 5, 3584 CC Utrecht, The
25 Netherlands. h.a.dijkstra@uu.nl

26 René M. van Westen (MSc.), Institute for Marine and Atmospheric research Utrecht,
27 Department of Physics, Utrecht University, Princetonplein 5, 3584 CC Utrecht, The
28 Netherlands. r.m.vanwesten@uu.nl

29 Julie D. Pietrzak (Prof. Dr.), Department of Hydraulic Engineering, Delft University of
30 Technology, Stevinweg 1, 2628 CN Delft, The Netherlands. J.D.Pietrzak@tudelft.nl

31 Adam S. Candy (Dr.), Department of Hydraulic Engineering, Delft University of Technology,
32 Stevinweg 1, 2628 CN Delft, The Netherlands. A.S.Candy@tudelft.nl

33 Caroline A. Katsman (Assoc. Prof. Dr.), Department of Hydraulic Engineering, Delft
34 University of Technology, Stevinweg 1, 2628 CN Delft, The Netherlands.
35 c.a.katsman@tudelft.nl

36 Carine G. van der Boog (MSc.), Department of Hydraulic Engineering, Delft University of
37 Technology, Stevinweg 1, 2628 CN Delft, The Netherlands.
38 c.g.vanderboog@tudelft.nl

39 Riccardo E. M. Riva (Dr.), Department of Geoscience and Remote Sensing, Delft University of
40 Technology, Stevinweg 1, 2628 CN Delft, The Netherlands. R.E.M.Riva@tudelft.nl

41 Cornelis Slobbe (Dr. ir.), Department of Geoscience and Remote Sensing, Delft University of
42 Technology, Stevinweg 1, 2628 CN Delft, The Netherlands. D.C.Slobbe@tudelft.nl

43 Roland Klees (Prof. Dr), Department of Geoscience and Remote Sensing, Delft University of
44 Technology, Stevinweg 1, 2628 CN Delft, The Netherlands. r.klees@tudelft.nl

Johan Stapel (Dr.), Caribbean Netherlands Science Institute, L.E. Saddlerweg, St Eustatius,
Caribbean Netherlands. johan.stapel@cnsi.nl

Tjisse van der Heide (Dr.), Department of Coastal Systems, NIOZ Royal Netherlands Institute
for Sea Research and Utrecht University, P.O. Box 59, 1790 AB Den Burg, Texel, The
Netherlands. tjisse.van.der.heide@nioz.nl

Marieke M. van Katwijk (Dr.), Department of Environmental Science, Institute for Water and
Wetland Research, Faculty of Science, Radboud University Nijmegen, Heyendaalseweg 135,
6525 Nijmegen, The Netherlands. M.vanKatwijk@science.ru.nl

Peter M. J. Herman (Prof. Dr.), Department of Hydraulic Engineering, Delft University of
Technology, Stevinweg 1, 2628 CN Delft, The Netherlands. peter.herman@deltares.nl

Tjeerd J. Bouma (Prof. Dr.), Department of Estuarine and Delta Systems, NIOZ Royal
Netherlands Institute for Sea Research and Utrecht University, Korringaweg 7, 4401 NT
Yerseke, The Netherlands. tjeerd.bouma@nioz.nl

Corresponding author: Rebecca K. James, rebecca.james@nioz.nl, 0031 113 577 491

Keywords: coastal erosion, nature-based engineering, ecosystem services, coastal protection,
coastal management

Paper type: Forum article

Abstract

Tropical beaches provide coastal flood protection, income from tourism and habitat for ‘flag-ship’ species. They urgently need protection from erosion, which is being exacerbated by changing climate and coastal development. Traditional coastal engineering solutions are expensive, provide unstable temporary solutions and often disrupt natural sediment transport. Instead, natural foreshore stabilization and nourishment may provide a sustainable and resilient, long-term solution. Field flume and ecosystem process measurements along with data from the literature, show that sediment stabilization by seagrass in combination with sediment-producing calcifying algae in the foreshore, form an effective mechanism for maintaining tropical beaches worldwide. The long-term efficacy of this type of nature-based beach management is shown at a large scale by comparing vegetated and unvegetated coastal profiles. We argue that preserving and restoring vegetated beach foreshore ecosystems offers a viable, self-sustaining alternative to traditional engineering solutions, increasing the resilience of coastal areas to climate change.

Introduction:

Beaches are key ecosystems in coastal zones, making up 31% of the world’s shoreline in ice-free regions of the world (Luijendijk et al. 2018). They have a vital role in flood defence, provide a source of income as a tourist attraction, and are essential habitats for various tropical “flag-ship” species, such as sea turtles and sea birds (Defeo et al. 2009). Beach erosion, however, has become a major global problem, with a recent analysis showing that 24% of the world’s sandy beaches experience chronic erosion (Luijendijk et al. 2018). The development of human infrastructure along the coast and waterways (Fig. 1a-c) has led to the rapid loss of natural systems that accumulate and stabilize sediment - such as coastal dunes, seagrass

meadows and mangroves - disrupting the regular pathways of sediment transport (Feagin et al. 2015; Luijendijk et al. 2018). Moreover, the combination of sea level rise with increasing storm occurrence and intensity will exacerbate beach erosion in the future (Defeo et al. 2009; Nicholls & Cazenave 2010). This is of great concern for many tropical areas, which typically have a high dependency on beaches for flood safety, and also economically for local tourism (red shading in Fig. 1d). For example, Caribbean islands together received over 23 million tourist visitors in 2015, creating a revenue of 26.5 billion USD (UNWTO 2016). On average, 23% of the gross domestic product (GDP) of countries within the Caribbean is obtained from tourism (Fig. 1d), with most tourists being attracted by the sandy beaches. Cost effective solutions to prevent or mitigate beach erosion are thus urgently needed for the long-term economic sustainability in these countries (Secretary-General 2016; Morris et al. 2018).

Many tropical countries lack the infrastructure and finances to undertake engineering solutions for beach protection. Hence, beaches continue to disappear into the sea, increasing the vulnerability of coastal areas to flooding, and threatening coastal structures and beach tourism (Fig. 1b). Where there are sufficient resources, two schemes of coastal engineering strategies are used to counter beach erosion: hard and soft (Finkl & Walker 2005; Castelle et al. 2009; Stive et al. 2013; Silva et al. 2016), both incurring a high capital cost. Hard coastal defence schemes are employed to mitigate wave attack and reduce local erosion (Fig. 1a; Ranasinghe and Turner 2006; Ruiz-Martínez et al. 2015; Walker, Dong and Anastasiou 1991). Such physical barriers typically inhibit the natural sand transport pathways, thereby depleting sand from neighbouring areas (Ranasinghe & Turner 2006; Ruiz-Martínez et al. 2015; Luijendijk et al. 2018). Soft defence schemes, such as beach or foreshore nourishments, have recently become more popular (Fig. 1c; Bishop et al. 2006; Castelle et al. 2009; Ruiz-Martínez et al. 2015; Stive et al. 2013). Although effective, soft engineering requires continuous maintenance, resulting in repeated smothering and disturbance of the natural beach communities (Bishop et

al. 2006; Defeo et al. 2009) and neighbouring ecosystems (e.g. coral reefs). In the long-term, nourishments can alter beach grain characteristics (Hanson et al. 2002), which can potentially cause permanent changes to the benthic community (Bishop et al. 2006).

By combining experimental field measurements with data from the literature, we demonstrate that the combination of foreshore stabilization by seagrass and natural foreshore nourishment by calcifying macroalgae can provide long-term maintenance of tropical beaches. In general, foreshore nourishment (both natural or engineered) is effective in beach protection, as a shallow foreshore reduces wave attack on the beach (Hanson et al. 2002; Christianen et al. 2013). Because a natural foreshore stabilization-and-nourishment regime requires no maintenance and operates gradually over long timescales with locally-produced sediment, it offers a cost-effective and sustainable alternative to human-engineered solutions. Comparing unique long-term beach profiles of vegetated, transitioning and unvegetated coasts illustrate the effectiveness of this approach.

Natural foreshore nourishment by vegetation: sediment stabilization and production

Shallow inter- and sub-tidal foreshores of natural tropical sandy beaches are predominately composed of locally produced calcium carbonate (CaCO_3) sediments. These carbonate sediments are biogenically produced and need to be continually captured and retained within the foreshore for a beach to resist erosion and remain stable, something that seagrass is extremely effectively at achieving.

With a newly developed portable flume, designed to be used in the field, the ability of different vegetation types (bare, vegetated with only calcifying macroalgae, sparse seagrass: 50% cover of *T. testudinum*, and dense seagrass: 100% cover of *T. testudinum*) to stabilize sediment was

measured directly within Galion Bay, St Martin (Caribbean). Regulating the speed of two motor-driven propellers allowed the flow velocity within the flume tunnel to be modified (see photo in Fig. 2a, and further methods in Suppl. 1). The point at which the surface sediment began to move was recorded as the threshold shear velocity. We found that in bare areas and areas with only calcifying macroalgae, the coarse carbonate sediments (median grain size: 337 μm , SE = 33) that are present in these areas start eroding already at flow speeds caused by moderate breezes (i.e. a wind of 10 m s^{-1} can cause flow speeds of 0.2 m s^{-1} within shallow areas (Hughes 1956)). However, where a sparse cover of seagrass is present, the sediment is finer (median grain size: 297 μm , SE = 17) as the protected seagrass canopy promotes fine grains to settle (De Boer 2007), but the flow required to erode the carbonate sediment doubles. And when *T. testudinum* seagrass cover is dense, the sediment is finer again (median grain size: 129 μm , SE = 7), but remains stable at flows stronger than 1.0 m s^{-1} (Fig. 2a); the maximum flow velocity of the flume. These flume results were confirmed by the seven times longer retention time of stained sediment that was placed in dense seagrass beds as compared to bare areas, in a high uni-directional flow environment within Galion Bay, and the four times higher retention time in a wave-exposed area (Fig. 2b).

Although relatively few studies have directly measured the sediment stabilizing effect of seagrass (Scoffin 1970; Widdows et al. 2008), the available literature widely supports our findings. For example, Christianen et al. (2013) found that even low density, heavily grazed seagrass meadows significantly reduce sediment erosion in Indonesia. A global review by Potouroglou et al., (2017) shows an average accretion rate of 5.33 mm year^{-1} occurring within seagrass meadows compared to adjacent unvegetated areas that experience an average erosion rate of 21.3 mm year^{-1} . Seagrasses reduce erosion and cause sediment accretion by stabilizing the sediment with their root-rhizome mat (Potouroglou et al. 2017), and by attenuating water

flow and waves. Hansen & Reidenbach (2012) reported that dense seagrass canopies of *Zostera marina* can attenuate flow velocity by 70-90%, whereas Fonseca & Cahalan (1992) showed a wave energy reduction of 34-44% for four varying species of seagrass, including *T. testudinum*. Flow and wave attenuation cause sediment particles to settle and reduces their resuspension, while additionally, seagrass leaves can bend over the sediment surface, further stabilizing the sediments. For a beach to remain stable over the long-term, however, a continuous supply of sediment is required to offset any erosion that occurs during storm events or from seaward currents that may transport unprotected sediment out of the beach system.

The breakdown and erosion of nearby coral reefs can provide a large contribution of sediment when the reefs are present (Chave et al. 1972; Hallock 1981). Another sediment contributor is calcifying macroalgae from the Halimedaceae family, which are composed of 70-90% CaCO_3 (van Tussenbroek & Van Dijk 2007). Because they grow directly within and adjacent to seagrass meadows on tropical beach foreshores, the sediment they produce is deposited where it is most valuable for providing a natural foreshore nourishment. This sediment production does vary significantly depending on the season, species and their abundance, however, the fast growth and rapid turn-over rates mean that the average sediment production reported for *Halimeda* spp. growing within seagrass meadows in the Pacific region is $337 \text{ g}_{\text{dwt}} \text{ CaCO}_3 \text{ m}^{-2} \text{ year}^{-1}$ (SE = 70, n = 10) (Suppl. 2; Garrigue 1991; Merten 1971; Payri 1988), and in Caribbean region, $166 \text{ g}_{\text{dwt}} \text{ CaCO}_3 \text{ m}^{-2} \text{ year}^{-1}$ (SE = 93, n = 8) (Suppl. 2; Armstrong and Miller 1988; Freile 2004; Multer 1988; Neumann and Land 1975; van Tussenbroek and Van Dijk 2007; Wefer 1980). Although this average rate contributes less than 0.28 (Pacific) and 0.15 mm (Caribbean) of sediment to the bed level per year (assuming a dry bulk density of 1.08 g per cm^3), the deposition of this CaCO_3 occurs directly within the foreshore where seagrass is present. The algae-produced sediment is therefore immediately captured and retained within

the beach foreshore ecosystem by the seagrass, thereby supplying a continuous and natural nourishment.

Engineering and natural nourishment as contrasting management regimes

We postulate engineering solutions and natural foreshore nourishment as contrasting management regimes, each having its own positive feedback (Fig. 3a). The engineered regime, where there is an unvegetated disturbed foreshore ecosystem with little or no biogenic sand production and highly mobile sediments. Such a regime results in a beach vulnerable to erosion, and therefore, requires regular engineering nourishments of the beach foreshore system to maintain its form. The alternative regime, a natural self-sustaining foreshore ecosystem with seagrass and calcifying macroalgae fronting a stable beach, which forms a self-stabilizing and self-nourishing system.

The combined sediment-stabilization by seagrass and sediment-production by calcifying algae yields a biologically-driven landscape with self-maintaining feedbacks. Specifically, by attenuating waves, preventing excessive erosion, and replenishing lost sediments, seagrass meadows and calcifying algae together create a self-reinforcing loop (Maxwell et al. 2017). Stable sediment has been shown to be a main requirement for the long-term persistence of seagrass meadows (Reise & Kohlus 2008; Christianen et al. 2014; Suykerbuyk et al. 2016), and in areas with fine sediment, can lead to a higher water transparency needed to sustain growth (van der Heide et al. 2007; Adams et al. 2018). This means that disruption of these self-reinforcing feedbacks may result in rapid losses of the seagrass-algae community (Maxwell et al. 2017). That is, in beach foreshore systems without seagrasses and algae, the sediment surface is freely agitated by currents and waves, yielding highly mobile sediments (Widdows et al. 2008; Marbà et al. 2015). Such unstable sediment conditions make it very difficult for

seagrasses and algae to (re-)establish (Williams 1990; Infantes et al. 2011; Balke et al. 2014; Suykerbuyk et al. 2016), and can increase turbidity levels if smaller sediment particles become suspended in the water column (van der Heide et al. 2007; Adams et al. 2018).

Human engineering through frequent beach nourishments can increase the sand supply to such disturbed beach foreshore systems (Finkl & Walker 2005; Castelle et al. 2009; Stive et al. 2013). However, these repeated nourishments smother establishing seagrasses and algae, and create an unstable sediment surface which is more likely to erode (Fig. 3a). Thus, although engineered nourishments may save the beach in the short term, it paradoxically may generate the necessity for recurrent beach nourishments in the long run (Trembanis & Pilkey 1998), creating an expensive and unsustainable management cycle in developing tropical regions (Silva et al. 2014).

Examples of the two alternative management regimes and one in transition, are found along the coast of Mexico (see Suppl. 1). In coastal areas where seagrass and calcifying macroalgae dominate the system, beach shore profiles conducted from 2008 to 2012 (methods detailed in Suppl. 1) are stable (Fig. 3b). In contrast, areas devoid of these species are typified by continuous erosion, which persists after engineered nourishments (Fig. 3d). A transition between these contrasting management regimes is observed in a third area. Here, extensive seagrass meadows of *T. testudinum* disappeared from the first 60 meters of the foreshore in 2015 due to a large brown tide of drifting *Sargassum* spp. (van Tussenbroek et al. 2017). As a result of these losses, beach profiles taken in 2007 and 2017 show the beach foreshore experienced strong vertical erosion, up to 0.4 m in some areas (Fig. 3c). However, a small area of the beach foreshore where seagrass was not lost, experienced only minor erosion and remained relatively stable (Fig. 3c). Overall these examples impressively illustrate the effectiveness of vegetated foreshore ecosystems for maintaining stable beaches and shorelines.

Implications & challenges for future management of tropical beaches

To create stable long-term management solutions for tropical beaches, beach management would benefit from shifting away from frequent engineered nourishments and hard structures, towards maintenance by natural ecosystems. With current insights, anthropogenic use of beaches could be designed to halt and reverse current decline of natural foreshore ecosystems. Tropical seagrass and *Halimeda* spp. usually co-occur and can be found in tropical sandy regions all around the world (Fig. 1d; Green and Short 2003; UNEP-WCMC and Short 2005), so there is widespread potential to restore these systems (Orth et al. 2006) to create a natural, self-sustaining beach management regime.

Conservation of areas where natural foreshore vegetation still persists will improve the condition of foreshore ecosystems, maximising their ability to protect beaches against erosion. Where foreshore vegetation has become degraded, an effort to protect what remains and to restore the ecosystem to a healthy self-reinforcing state may be necessary to implement effective natural beach management regimes. Preserving and restoring foreshore vegetation that still exists is especially important as climate-driven disturbance events - such as extreme wave action, cyclones (Saunders & Lea 2008), and the occurrence of brown tides from *Sargassum* spp. drifts (van Tussenbroek et al. 2017) - become more frequent with rising global temperatures. As climate-driven factors are hard to manage at a local scale, management should primarily aim at reducing local human-induced impacts (Scheffer et al. 2001). Local impacts, like greater turbidity (Orth et al. 2006), nutrient enrichment and pollution (Kemp et al. 2005), physical damage to seagrass meadows from trampling and boat anchoring (Eckrich & Holmquist 2000), and modification of natural sediment transport and increased wave reflection caused by the construction of hard structures (Defeo et al. 2009; Ruiz-Martínez et al. 2015; Luijendijk et al. 2018), are all intensifying as coastlines develop further. The installation of sewage treatment plants and limiting construction of hard structures along the coast are the most

obvious steps to help protect and restore natural foreshore vegetation. Another is to limit accessibility of people to vulnerable areas, and provide boat anchoring facilities outside regions of vegetation. Ensuring coral reefs remain in abundance and their sediment input to tropical beaches persists, would also improve the prospects of tropical beaches to keep up with sea level rise.

Given that the engineering management regime of a disturbed beach is self-reinforced by a feedback that maintains sediment instability (Fig. 3a), it will be difficult to induce a transition to the natural beach systems in areas where engineering management regimes already take place and/or vegetation has been completely lost. Developing ways to stimulate natural vegetation development may be necessary, such as utilising temporary structures that protect establishing seagrass and calcifying macroalgae, until they grow to a point that they can self-stabilize the sediment (Suykerbuyk et al. 2016; van Katwijk et al. 2016). Engineered nourishments will need to either cease, or be modified to ensure that any added sediment encourages the growth of the natural ecosystem rather than smothers it (Cheong et al. 2013). This may be achieved by using methods that give a gradual sediment flux, like the sand engine in The Netherlands (Stive et al. 2013), or by using smaller doses of sediment.

It is imperative that we recognize the benefits of a vegetated foreshore ecosystem in preventing beach erosion, and thus increase the resistance of coastal areas to storm surges and flooding. Switching disturbed beach systems to natural self-sustaining ecosystems for coastal defence will require financial investments (e.g. from the World Bank, in the context of climate adaptation (Secretary-General 2016; World Bank 2017)), development of effective restoration methods, as well as altered governance. Only a collaborative approach of many stakeholders will ensure both economic and ecological benefits. This will require interdisciplinary collaboration between economists focusing on tourism, ecologists focusing on ecosystem functioning and natural values, engineers focusing on physical processes and design measures,

and sociologists focusing on governance processes and public support. With this paper, we aim to provide an alternative beach management regime to traditional engineering solutions, by highlighting the viable and self-sustaining capacity of vegetated beach foreshore ecosystem in preventing erosion. Utilising an effective natural solution to coastal erosion will help to increase the resilience of tropical coastal areas to climate change in a sustainable way.

Acknowledgments

This work was primarily funded by the NWO ‘Caribbean Research: a Multidisciplinary Approach’ grant, which was awarded to the SCENES project (Grant number 858.14.063). Permits for the work in St Martin were obtained from the Reserve Naturelle Saint Martin, and we thank them for their advice and allowing us to conduct our research there. We would also like to thank Carlos Gonzales Godoy for taking the 2007 beach profiles of Mirador Nizuc, and Edgar Escalante-Mancera & Miguel-Angel Gomez from SAMMO, ICML for the 2017 Mirador Nizuc profiles.

References

- Adams MP, Ghisalberti M, Lowe RJ, Callaghan DP, Baird ME, Infantes E, O'Brien KR. 2018. Water residence time controls the feedback between seagrass, sediment and light: Implications for restoration. *Advances in Water Resources* 117: 14–26.
- Armstrong ME, Miller AI. 1988. Modern carbonate sediment production and its relation to bottom variability Grahams Harbor, San Salvador, Bahamas. Pages 23–32 in Mylroie JE, Gerace DT, ed. *Proceedings of the fourth symposium on the geology of the Bahamas*. Bahamian Field Station.
- Balke T, Herman PMJ, Bouma TJ. 2014. Critical transitions in disturbance-driven ecosystems: Identifying Windows of Opportunity for recovery. *Journal of Ecology* 102: 700–708.
- Bishop MJ, Peterson CH, Summerson HC, Lenihan HS, Grabowski JH. 2006. Deposition and long-shore transport of dredge spoils to nourish beaches: Impacts on benthic infauna of an ebb-tidal delta. *Journal of Coastal Research* 223: 530–546.
- De Boer WF. 2007. Seagrass-sediment interactions, positive feedbacks and critical thresholds for occurrence: A review. *Hydrobiologia* 591: 5–24.
- Castelle B, Turner IL, Bertin X, Tomlinson R. 2009. Beach nourishments at Coolangatta Bay over the period 1987-2005: Impacts and lessons. *Coastal Engineering* 56: 940–950.
- Chave KE, Smith S V., Roy KJ. 1972. Carbonate production by coral reefs. *Marine Geology* 12: 123–140.
- Cheong SM, Silliman B, Wong PP, Van Wesenbeeck B, Kim CK, Guannel G. 2013. Coastal adaptation with ecological engineering. *Nature Climate Change* 3: 787–791.
- Christianen MJA, van Belzen J, Herman PMJ, van Katwijk MM, Lamers LPM, van Leent

327 PJM, Bouma TJ. 2013. Low-canopy seagrass beds still provide important coastal
328 protection services. PLoS ONE 8.

329 Christianen MJA, Herman PMJ, Bouma TJ, Lamers LPM, van Katwijk MM, van der Heide T,
330 Mumby PJ, Silliman BR, Engelhard SL, van de Kerk M, Kiswara W, van de Koppel J.
331 2014. Habitat collapse due to overgrazing threatens turtle conservation in marine
332 protected areas. Proceedings of the Royal Society B: Biological Sciences 281:
333 20132890–20132890.

334 Defeo O, McLachlan A, Schoeman DS, Schlacher TA, Dugan J, Jones A, Lastra M, Scapini
335 F. 2009. Threats to sandy beach ecosystems: A review. Estuarine, Coastal and Shelf
336 Science 81: 1–12.

337 Eckrich CE, Holmquist JG. 2000. Trampling in a seagrass assemblage: Direct effects,
338 response of associated fauna, and the role of substrate characteristics. Marine Ecology
339 Progress Series 201: 199–209.

340 Feagin RA, Figlus J, Zinnert JC, Sigren J, Martínez ML, Silva R, Smith WK, Cox D, Young
341 DR, Carter G. 2015. Going with the flow or against the grain? The promise of vegetation
342 for protecting beaches, dunes, and barrier islands from erosion. Frontiers in Ecology and
343 the Environment 13: 203–210.

344 Finkl CW, Walker HJ. 2005. Beach Nourishment. Pages 147–161 in Schwartz ML ed.
345 Encyclopedia of Coastal Sciences. Springer-Verlag, Dordrecht, The Netherlands.

346 Fonseca MS, Cahalan JA. 1992. A preliminary evaluation of wave attenuation by four species
347 of seagrass. Estuarine, Coastal and Shelf Science 35: 565–576.

348 Freile D. 2004. Carbonate productivity rates of Halimeda in two different locations, San
349 Salvador Island, Bahamas. Pages 95–106 in Lewis RD, Panuska BC ed. Proceedings of

350 the 11th symposium on the geology of the Bahamas and other carbonate regions. Gerace
351 Research Centre, Auburn University, Auburn, AL.

352 Garrigue C. 1991. Biomass and production of two *Halimeda* species in the southwest new
353 caledonian lagoon. *Oceanologica Acta* 14: 581–588.

354 Green EP, Short FT. 2003. World atlas of seagrasses. Berkley, USA

355 Hallock P. 1981. Production of Carbonate Sediments by Selected Large Benthic Foraminifera
356 on Two Pacific Coral Reefs. *Journal of Sedimentary Research* Vol. 51: 467–474.

357 Hansen JCR, Reidenbach MA. 2012. Wave and tidally driven flows in eelgrass beds and their
358 effect on sediment suspension. *Marine Ecology Progress Series* 448: 271–287.

359 Hanson H, Brampton A, Capobianco M, Dette HH, Hamm L, Laustrup C, Lechuga A,
360 Spanhoff R. 2002. Beach nourishment projects, practices, and objectives - A European
361 overview. *Coastal Engineering* 47: 81–111.

362 van der Heide T, Van Nes EH, Geerling GW, Smolders AJP, Bouma TJ, Van Katwijk MM.
363 2007. Positive feedbacks in seagrass ecosystems: Implications for success in
364 conservation and restoration. *Ecosystems* 10: 1311–1322.

365 Hughes P. 1956. A determination of the relation between wind and sea-surface drift.
366 *Quarterly Journal of the Royal Meteorological Society* 82: 494–502.

367 Infantes E, Orfila A, Bouma TJ, Simarro G, Terrados J. 2011. *Posidonia oceanica* and
368 *Cymodocea nodosa* seedling tolerance to wave exposure. *Limnology and Oceanography*
369 56: 2223–2232.

370 van Katwijk MM, et al. 2016. Global analysis of seagrass restoration: The importance of
371 large-scale planting. *Journal of Applied Ecology* 53: 567–578.

372 Kemp WM, et al. 2005. Eutrophication of Chesapeake Bay: Historical trends and ecological
373 interactions. *Marine Ecology Progress Series* 303: 1–29.

374 Luijendijk A, Hagenaars G, Ranasinghe R, Baart F, Donchyts G, Aarninkhof S. 2018. The
375 State of the World's Beaches. *Scientific Reports*: 1–11.

376 Marbà N, Arias-Ortiz A, Masqué P, Kendrick GA, Mazarrasa I, Bastyan GR, Garcia-Orellana
377 J, Duarte CM. 2015. Impact of seagrass loss and subsequent revegetation on carbon
378 sequestration and stocks. *Journal of Ecology* 103: 296–302.

379 Maxwell PS, et al. 2017. The fundamental role of ecological feedback mechanisms for the
380 adaptive management of seagrass ecosystems – a review. *Biological Reviews* 92: 1521–
381 1538.

382 Merten M. 1971. Ecological observations of *Halimeda macroloba* Decaisne (Chlorophyta) on
383 Guam. *Micronesica* 7: 27–44.

384 Morris RL, Konlechner TM, Ghisalberti M, Swearer SE. 2018. From grey to green: Efficacy
385 of eco-engineering solutions for nature-based coastal defence. *Global Change Biology*
386 24: 1827–1842.

387 Multer HG. 1988. Growth rate, ultrastructure and sediment contribution of *Halimeda*
388 *incrassata* and *Halimeda monile*, Nonsuch and Falmouth Bays, Antigua, W.I. *Coral*
389 *Reefs* 6: 179–186.

390 Neumann ACC, Land LS. 1975. Lime mud deposition and calcareous algae in the Bight of
391 Abaco, Bahamas: A budget. *Journal of Sedimentary Research* 45: 763–786.

392 Nicholls RJ, Cazenave A. 2010. Sea Level Rise and Its Impact on Coastal Zones. *Science*
393 328: 1517–1520.

394 Orth RJ, et al. 2006. A Global Crisis for Seagrass Ecosystems. *Bioscience* 56: 987–996.

395 Payri CE. 1988. Halimeda contribution to organic and inorganic production in a Tahitian reef
396 system. *Coral Reefs* 6: 251–262.

397 Potouroglou M, Bull JC, Krauss KW, Kennedy HA, Fusi M, Daffonchio D, Mangora MM,
398 Githaiga MN, Diele K, Huxham M. 2017. Measuring the role of seagrasses in regulating
399 sediment surface elevation. *Scientific Reports*: 1–11.

400 Ranasinghe R, Turner IL. 2006. Shoreline response to submerged structures: A review.
401 *Coastal Engineering* 53: 65–79.

402 Reise K, Kohlus J. 2008. Seagrass recovery in the Northern Wadden Sea? *Helgoland Marine*
403 *Research* 62: 77–84.

404 Ruiz-Martínez G, Mariño-Tapia I, Mendoza Baldwin EG, Silva Casarín R, Enríquez Ortiz
405 CE. 2015. Identifying Coastal Defence Schemes through Morphodynamic Numerical
406 Simulations along the Northern Coast of Yucatan, Mexico. *Journal of Coastal Research*:
407 651–670.

408 Saunders MA, Lea AS. 2008. Large contribution of sea surface warming to recent increase in
409 Atlantic hurricane activity. *Nature* 451: 557–560.

410 Saxby T. Saxby. Integration and Application Network, University of Maryland Center for
411 Environmental Science. ian.umces.edu/imagelibrary/.

412 Scheffer M, Carpenter S, Foley JA, Folke C, Walker B. 2001. Catastrophic shifts in
413 ecosystems. *Nature* 413: 591–596.

414 Scoffin TP. 1970. The trapping and binding of subtidal carbonate sediments by marine
415 vegetation in Bimini Lagoon, Bahamas. *Journal of Sedimentary Petrology* 40: 249–273.

416 Secretary-General UN. 2016. United Nations Economic and Social Council Progress towards
417 the Sustainable Development Goals.

418 Silva R, et al. 2014. Present and future challenges of coastal erosion in Latin America. Journal
419 of Coastal Research 71: 1–16.

420 Silva R, Mendoza E, Mariño-Tapia I, Martínez ML, Escalante E. 2016. An artificial reef
421 improves coastal protection and provides a base for coral recovery. Journal of Coastal
422 Research 75: 467–471.

423 Stive MJF, et al. 2013. A new alternative to saving our beaches from sea-level rise: the sand
424 engine. Journal of Coastal Research 29: 1001–1008.

425 Suykerbuyk W, Bouma TJ, Govers LL, Giesen K, de Jong DJ, Herman P, Hendriks J, van
426 Katwijk MM. 2016. Surviving in Changing Seascapes: Sediment Dynamics as
427 Bottleneck for Long-Term Seagrass Presence. Ecosystems 19: 296–310.

428 Trembanis AC, Pilkey OH. 1998. Summary of Beach Nourishment along the U.S. Gulf of
429 Mexico Shoreline. Journal of Coastal Research 14: 407–417.

430 van Tussenbroek BI, Van Dijk JK. 2007. Spatial and temporal variability in biomass and
431 production of psammophytic *Halimeda incrassata* (Bryopsidales, Chlorophyta) in a
432 Caribbean reef lagoon. Journal of Phycology 43: 69–77.

433 van Tussenbroek BI, Hernández Arana HA, Rodríguez-Martínez RE, Espinoza-Avalos J,
434 Canizales-Flores HM, González-Godoy CE, Barba-Santos MG, Vega-Zepeda A,
435 Collado-Vides L. 2017. Severe impacts of brown tides caused by *Sargassum* spp. on
436 near-shore Caribbean seagrass communities. Marine Pollution Bulletin.

437 UNEP-WCMC, Short FT. 2005. Global distribution of seagrasses (version 3.0). Third update
438 to the data layer used in Green and Short (2003). url:
439 <http://data.unepwcmc.org/datasets/7>.

440 UNWTO. 2016. UNWTO Tourism Highlights 2016 Edition. Madrid, Spain

Walker DJ, Dong P, Anastasiou K. 1991. Sediment Transport Near Groynes in the Nearshore Zone. *Journal of Coastal Research* 7: 1003–1011.

Wefer G. 1980. Carbonate production by algae *Halimeda*, *Pencillus* and *Padina*. *Nature* 285: 323–324.

Widdows J, Pope ND, Brinsley MD, Asmus H, Asmus RM. 2008. Effects of seagrass beds (*Zostera noltii* and *Z. marina*) on near-bed hydrodynamics and sediment resuspension. *Marine Ecology Progress Series* 358: 125–136.

Williams SL. 1990. Experimental studies of Caribbean seagrass bed development. *Ecological Monographs* 60: 449–469.

World Bank. 2017. Atlas of Sustainable Development Goals 2017 : From World Development Indicators. License: C. World Bank ed. ©World Bank, Washington, DC.

Figures




Figure 1. The building of hard structures to prevent coastal erosion, such as seawalls (a), the

over-development of coastlines (b), and beach nourishments (c) only serve to exacerbate coastal erosion. The global map (d) shows the proportion of GDP obtained from tourism in 2015 (data sourced from World Bank and World Tourism Organization), with the darker red shading indicating a higher proportion of the gross domestic product (GDP) is obtained from tourism for that country. The effective sediment-stabilizing seagrass *Thalassia* spp. is globally distributed (green circles, sourced from UNEP-WCMC & Short (2005)), and can be found alongside the sediment-producing calcifying macroalgae *Halimeda* spp. (blue squares, sightings reported in peer reviewed literature).




Figure 2. Carbonate sediment is stabilized by seagrass, as indicated by measuring the critical threshold for bed-load transport with a field flume in contrasting vegetation types: bare, calcifying algae only, sparse *Thalassia* (50% cover of *T. testudinum*), dense *Thalassia* (100% cover of *T. testudinum*) (a). This was corroborated by measuring the retention time of stained sediments for contrasting vegetation types in the different physical environments (b): wave sheltered (mean wave height = 0.15 m, SE = 0.004, n = 370), uni-directional (mean flow rate = 0.15 m s⁻¹, SE = 0.025, n = 18), and wave exposed (mean wave height = 0.22 m, SE = 0.005,

n = 429). Bars represent means $\pm$ SE ($n_{\text{sed.stab}} = 3$, $n_{\text{sed.ret}} = 5$) and black points indicate individual data points. Different letters above bars denote significant difference ($p < 0.05$), tested with Tukey HSD pair-wise comparisons.




Figure 3. Self-reinforcing feedbacks drive the contrasting beach management regimes as schematised in (a). The natural beach is driven by seagrass stabilizing the sediment, which encourages further ecosystem development. Whereas the system devoid of vegetation has increasingly mobile sediment, discouraging the growth of vegetation and leading to an unstable beach system, requiring engineering which further contributes to sediment mobility and erosion. These types of beach regimes can be seen in examples from the coastline of Mexico (map in S1). Regular beach profiles taken from two transects at the natural beach of Puerto Morelos from June 2008 (dashed lines) to May 2012 (solid line) show that this relatively undisturbed beach with extensive seagrass-calcifying algae meadows has remained stable over many years (b). While beach profiles at Mirador Nizuc in 2007 (dashed line) and June 2017

(solid line) show that the beach had significant erosion after a Sargassum brown tide that persisted from July 2015 to May 2016 resulted in the loss of seagrass (c, upper graph), however in an area of the same beach where seagrass persisted, very little erosion occurred (c, lower graph). While Cancun has no natural reef or seagrass meadows and development along the sand dunes has led to constant beach erosion, a sand nourishment in 2010 helped to restore the beach, but this continues to erode (d). Elevations are relative to mean sea level. (Thalassia illustration sourced from IAN image library (Saxby)).