

## Recent changes in extreme floods across multiple continents

Berghuijs, Wouter; Woods, Ross A.; Aalbers, Emma; Larsen, Joshua R.; Trancoso, Ralph

**Publication date**

2017

**Document Version**

Final published version

**Published in**

Geophysical Research Abstracts (online)

**Citation (APA)**

Berghuijs, W., Woods, R. A., Aalbers, E., Larsen, J. R., & Trancoso, R. (2017). Recent changes in extreme floods across multiple continents. *Geophysical Research Abstracts (online)*, 19, Article EGU2017-10597-1.

**Important note**

To cite this publication, please use the final published version (if applicable).  
Please check the document version above.

**Copyright**

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

**Takedown policy**

Please contact us and provide details if you believe this document breaches copyrights.  
We will remove access to the work immediately and investigate your claim.



## **Recent changes in extreme floods across multiple continents**

Wouter Berghuijs (1), Ross Woods (1), Emma Aalbers (2), Josh Larsen (3,4), Ralph Trancoso (3,5)

(1) Department of Civil Engineering, University of Bristol, University Walk, Bristol BS8 1TR, United Kingdom, (2) Royal Netherlands Meteorological Institute (KNMI), De Bilt, Netherlands, (3) School of Earth and Environmental Science, University of Queensland, Brisbane 4072, Australia, (4) Institute of Earth Surface Dynamics, University of Lausanne, Lausanne, Switzerland, (5) Department of Environment and Heritage Protection, Queensland Government, Australia.

Analyses of trends in observed floods often focus on relatively frequent events, whereas changes in rare floods are only studied for a small number of locations that have exceptionally long observational records. Understanding changes in rare floods is especially relevant as these events are often most damaging and influence the design of major structures. Here we provide an assessment of changes in the largest flood events ( $\sim 0.033$  annual exceedance probability) observed during the period 1980-2009 for 1744 catchments located in Australia, Brazil, Europe and the United States. The occurrence of rare floods in spatial aggregate shows strong decadal variability and peaked around 1995. During the 30-year period there are overall increases in both the frequency and magnitude of extreme floods. These currently unattributed increases are strongest in Europe and the United States, and weakest in Brazil and Australia.