

Achieving at-scale seascape restoration by optimising cross-habitat facilitative processes

Vozzo, Maria L.; Buelow, Christina A.; Sievers, Michael; Adame, M. Fernanda; Branson, Paul; Brown, Maddison; Crosswell, Joseph R.; ter Hofstede, Remment; van Koningsveld, Mark; More Authors

DOI

[10.1038/s44183-024-00095-1](https://doi.org/10.1038/s44183-024-00095-1)

Publication date

2024

Document Version

Final published version

Published in

NPJ Ocean Sustainability

Citation (APA)

Vozzo, M. L., Buelow, C. A., Sievers, M., Adame, M. F., Branson, P., Brown, M., Crosswell, J. R., ter Hofstede, R., van Koningsveld, M., & More Authors (2024). Achieving at-scale seascape restoration by optimising cross-habitat facilitative processes. *NPJ Ocean Sustainability*, 3(1), Article 57. <https://doi.org/10.1038/s44183-024-00095-1>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright


Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

<https://doi.org/10.1038/s44183-024-00095-1>

Achieving at-scale seascape restoration by optimising cross-habitat facilitative processes



Maria L. Vozzo¹ ✉, Christina A. Buelow², Michael Sievers², M. Fernanda Adame³, Paul Branson⁴, Maddison Brown^{1,5}, Joseph R. Crosswell⁶, Christopher Doropoulos¹, Ben L. Gilby⁷, Francisco Martinez-Baena⁸, Simon Reeves⁹, Vera Rullens⁹, Andy Steven¹, Ziyu Xiao⁶, Kirk Dahle¹⁰, Brian Silliman¹¹, Remment ter Hofstede^{12,13}, Mark van Koningsveld^{12,13} & Megan I. Saunders^{14,15}

Cross-habitat facilitative processes can enhance seascape restoration outcomes but there is uncertainty around the spatial dependencies of these processes across habitats. We synthesised the influence of environmental parameters on six processes underpinning cross-habitat facilitation and identified the linear distances over which they operate between habitats. All six process types occur at distances commonly used in seascape restoration demonstrating how harnessing facilitation can scale-up restoration to meet national and international goals.

Unless new ways to scale and accelerate restoration of seascapes (comprised of multiple coastal marine ecosystems) are developed and implemented, it is unlikely that global and international targets for restoration (e.g., UN Decade on Ecological Restoration and EU Nature Restoration Law), and the reinstatement of ecosystem services derived from these habitats, will be met. Restoration, defined as the assisted recovery of habitat types that have been damaged, degraded or destroyed¹, has occurred successfully for many decades on single habitats². Moving from isolated, single-habitat restoration approaches to those that restore multiple habitats simultaneously or sequentially, or that aim to restore connections within the broader seascape (collectively and hereafter ‘seascape restoration’; Box 1), should help scale-up restoration efforts and maximise benefits³. However, most published coastal marine restoration research has focused on restoration of a single habitat type rather than multiple habitat types, simultaneously⁴ and only 12% of restoration projects are strategically designed with seascape connectivity in mind⁵.

To optimise seascape restoration, we need to better understand and be cognisant of the facilitative processes that exist between habitats when designing and implementing restoration⁶. Facilitative processes are positive interactions, such as wave attenuation, sediment stabilisation or nutrient

cycling, that occur between habitats and help support the characteristic distributions of habitats across seascapes⁷. Whilst several facilitative processes that can enhance seascape connectivity, ecological functioning, and resilience are well established (e.g., coral reefs attenuating wave energy, enabling seagrass establishment⁸; mangroves retaining nutrients and sediment, improving water quality in the surrounding areas for seagrass and coral reefs⁹), we lack a clear understanding of the spatio-temporal dependencies of facilitative processes across coastal habitats. To achieve seascape restoration at-scale, practitioners, scientists, managers, engineers, policy-makers and stakeholders need to move from siloed approaches focusing on a single habitat in isolation, to those that incorporate cross-disciplinary expertise on how processes occur across different spatial scales and habitat types and include governance and permitting arrangements that enable multi-habitat restoration.

An understanding of how facilitative processes operate over time and space is needed so that they can be accounted for and harnessed in restoration practice. The strength of a facilitative process generated in one habitat and its impact on the maintenance or recovery of another habitat will depend on the distance between, size and spatial arrangement of those habitats^{7,10}, and a suite of abiotic drivers¹¹. For example, the extent of

¹CSIRO Environment, St Lucia, QLD, Australia. ²Coastal and Marine Research Centre, Australian Rivers Institute, School of Environment and Science, Griffith University, Gold Coast, QLD, Australia. ³Australian Rivers Institute, Centre for Marine and Coastal Research, School of Environment and Science, Griffith University, Brisbane, QLD, Australia. ⁴CSIRO Environment, Indian Ocean Marine Research Centre, Crawley, WA, Australia. ⁵School of the Environment, Steele Building (Building 3), The University of Queensland, St Lucia, QLD, Australia. ⁶CSIRO Environment, Dutton Park, QLD, Australia. ⁷School of Science, Technology and Engineering, University of the Sunshine Coast, Petrie, QLD, Australia. ⁸Sydney Institute of Marine Science, Mosman, NSW, Australia. ⁹The Nature Conservancy Australia, Carlton, VIC, Australia. ¹⁰The Nature Conservancy Australia, Pyrmont, NSW, Australia. ¹¹Nicholas School of the Environment, Duke University, Beaufort, NC, USA. ¹²Delft University of Technology, Civil Engineering and Geosciences, Delft, the Netherlands. ¹³Van Oord DMC, Rotterdam, the Netherlands. ¹⁴CSIRO Environment, Hobart, TAS, Australia. ¹⁵Centre for Biodiversity Conservation Science, The University of Queensland, St Lucia, QLD, Australia.

✉ e-mail: maria.vozzo@csiro.au

Box 1 | Seascape restoration typologies

Multi-habitat restoration is an emerging field within the discipline of restoration ecology and holds promise to help scale up restoration in seascapes. However, there are multiple potential applications and interpretations of multi-habitat restoration in seascapes, resulting in a need for refinement and nuance in definitions. We outline three restoration typologies, including two where restoration actions explicitly target

more than one habitat type (*simultaneous multi-habitat* and *sequential multi-habitat*), and one where actions explicitly target one habitat type whilst being explicitly cognisant of the spatial configuration of habitats in the seascape and the cross-habitat facilitative processes amongst those habitats (*single-habitat acknowledging seascape*).

Box 1 Table 1 Three coastal restoration typologies for seascape restoration approaches

Restoration typology	Description	Examples
Simultaneous multi-habitat	Restoring multiple habitat types simultaneously. This includes situations where multiple habitat types are intended to be restored at the same time but in practice, may be slightly asynchronous due to permitting or logistics (e.g., site access, season, contractor availability) rather than habitat suitability issues.	Simultaneous restoration of mangroves (planting) and oysters (reef construction) (Florida, USA) ¹²⁴ . Natural regeneration of mangroves, saltmarsh and seagrass following removal of bund wall to restore hydrology and tidal inundation of an area (Blue Heart Restoration Project, Queensland, Australia) ¹²⁵ .
Sequential multi-habitat	Intentionally restoring multiple habitat types in succession due to ecological requirements such as habitat suitability issues (i.e., where habitat X <i>needs</i> to be restored <i>before</i> habitat Y can be restored).	Restoration of boulder reef to stabilise sediment for transplanted seagrass. Transplantation of mussels to improve water clarity for seagrass (Denmark) ¹²⁶ . Incidental sequential multi-habitat restoration observed after constructing and transplanting kelp to restoration units. Flat oysters (<i>Ostrea angasi</i>) then settled to the bottom of the units (Tasmania, Australia) ¹²⁷ .
Single-habitat acknowledging seascape	Restoring a single habitat type whilst being cognisant of the seascape (i.e., other habitat types) in an attempt to benefit from potential facilitative interactions. Restored or intact ecosystems can benefit.	Construction of oyster reefs near seagrass beds to assess whether restored oyster reefs benefit seagrass (Alabama, USA) ¹²⁸ . Noosa Shellfish Restoration that incorporated suitability modelling to consider the context of surrounding habitats (Queensland, Australia) ¹²⁹ .

Restoration may involve the direct manipulation of the habitat (e.g., transplanting or seed deployment, providing settlement substrate, exotic species removal), and may occur after the cessation of disturbance or environmental modification, which may allow for natural regeneration (e.g., removal of contamination, restoration of water flows).

mangrove habitat, its distance from adjacent seagrass meadows and the amount of suspended sediments flowing in from the catchment will influence the amount of sediment trapping benefit that the seagrass meadows receive from the mangroves⁹. Thus, the quantification of facilitative processes is essential to scale restoration planning. However, there is a disconnect between our understanding of the spatial scale at which processes underpinning facilitation occur, and the spatial scales at which restoration occurs. The physical, biogeochemical and biological processes supporting cross-habitat facilitation can occur across distances of at least tens to hundreds of metres¹⁰, but most examples of cross-habitat facilitation reported in the literature have been described for spatial scales less than one metre⁴. Restoration of coastal marine habitats can range from tens of square metres to thousands of hectares, depending on the objectives and type of habitat restored^{2,12}. Describing the optimal lateral distances and spatial arrangements of habitats to target in seascape restoration to help overcome this theory-practice mismatch remains a crucial knowledge gap.

With increasing understanding around the benefits of and funding for multi-habitat seascape restoration^{13,14} there is an urgent need to identify how knowledge of cross-habitat facilitation can be used to inform site selection and the spatial arrangement of restoration interventions to enhance outcomes⁵. Describing how physical, biogeochemical or biological processes are involved in cross-habitat facilitation⁴ can allow for consideration of how the presence and spacing of individual restored habitats may interact with or influence the surrounding seascape³. This information can help inform how ‘sequential multi-habitat restoration’ of one habitat can later enhance restoration success of a second habitat (Box 1). In other instances, this information may be used to overcome, offset or ameliorate compromises in restoration design caused by external constraints such as funding or permits

(Box 2). Finally, the ability to identify tradeoffs in design choices, like spacing that optimises the influence of one process over another, may help to better predict and plan for the desired outcomes of restoration projects.

Seascape restoration is a rapidly growing field¹³, and the ways in which cross-habitat facilitation might be incorporated within multi-habitat restoration approaches will vary depending on local site conditions and the restoration approach. We first provide examples of three multi-habitat restoration typologies that are used in seascape restoration (Box 1). We have included definitions of multi- and single-habitat restoration typologies in which cross-habitat facilitation can be harnessed to achieve seascape restoration. We then outline six processes that underpin seascape cross-habitat facilitation: wave attenuation, sediment dynamics, nutrient processing, water filtration, carbon dynamics and alkalinity export, and animal- and plant-mediated facilitation of other habitats. For each, we define how the process can be quantified and how it is influenced by particular parameters within various habitats. Previous studies have synthesised information on the magnitude by which habitats can regulate facilitative processes⁹, the types of facilitative interactions occurring between pairs of coastal marine habitats⁴, and, where possible, the spatial scale over which these interactions have been quantified⁴. The latter is highly biased towards studies of interactions over small spatial scales (<1 m) because empirical measurements are more feasible at this spatial scale, yet spacing between different habitats in multi-habitat restoration would, in most instances, occur over much larger scales (e.g., tens to thousands of metres). Therefore, there remains a critical knowledge gap about the strength and certainty of the lateral spatial scales over which cross-habitat facilitative processes can occur and how they can be applied to optimise seascape restoration that harnesses facilitation. Here, we seek to address this knowledge gap by

providing an overview of the known lateral spatial ranges for each of six facilitative processes. While other types of interactions (e.g., negative ones such as competition, predation or herbivory^{10,15,16}) can be just as important in influencing community structure and adjacent habitats, their impacts on restoration outcomes have received attention in other recent research^{17–19}. Finally, using a multidisciplinary and multi-organisational approach—including practitioner, academic, industry contractor and government research perspectives—we illustrate how this information can address knowledge gaps that, when filled, will allow us to better harness facilitation for effective implementation of seascape restoration at scale.

Physical, biogeochemical and biological processes that occur across habitats

This section seeks to provide information that will help inform how seascape restoration can harness cross-habitat facilitation. We first define terminology relating to six key physical, biogeochemical and ecological processes underpinning cross-habitat facilitation, and then provide examples of how they operate. While we focus on these six process categories, there are many other ways in which habitats may interact to influence each other, including negative interactions. The processes of focus were identified using a cross-disciplinary approach, which included a combination of elicitation of knowledge from experts specialised in different physical, biogeochemical or ecological areas of research, paired with a synthesis of key literature related to facilitative processes that were identified by these experts. Each subsection summarises processes that are associated with six coastal structured habitats that are created by habitat-forming plants and animals: coral reefs, shellfish reefs, macroalgae forests, seagrass beds, saltmarshes and mangrove forests (Fig. 1). Whilst discussed separately, many processes intersect and, where relevant, we identify overlaps among different processes.

We then summarise the complex interdisciplinary information on facilitative processes that is needed to guide those conducting research, making decisions and implementing seascape restoration, including known examples of facilitation and the potential spatial scales of operation. Although many facilitative processes may also benefit humans as ecosystem services (e.g., wave attenuation, water filtration, carbon trapping), here we focus on the benefit of these processes to other coastal marine habitats (facilitation). We then illustrate how this information can be used when designing seascape restoration (Box 2).

Wave attenuation

Wave attenuation is the reduction of wave energy through wave shoaling, refraction, diffraction, wave breaking and energy dissipation due to bottom friction and canopy drag²⁰. Coastal habitat-forming species modify the depth and rugosity of the seabed, helping to attenuate wave energy²¹. The attenuation of wave energy facilitates other species establishment and persistence by broadening the environmental windows of opportunity²². For example, a reef that induces wave breaking and wave attenuation, can enable the persistence of seagrass behind the reef⁸.

Key parameters that govern wave transformation include wave height, wave direction, and wavelength (the combined effect of wave period and water depth). These parameters undergo non-linear changes during wave transformation that alter the dominant wavelength spatially across the system²³. For example, swell waves shoal and break on reef-crests and headlands, generating longer infra-gravity waves that propagate into reef lagoons and along beaches²⁴. The wave height and wavelength control the movement of water at the seabed, which interacts with benthic habitats that provide flow resistance (drag) and drive sediment transport through friction.

The effectiveness of benthic habitats (“canopies”) in attenuating wave energy depends on the geometry and stiffness of the organism’s structure, such as the shape or density of the dominant species of coral, shellfish or mangrove^{25,26}. From a hydrodynamic perspective, the canopy geometry is characterised by the area presented to flow which, combined with the canopy spacing (habitat density, e.g., distance between mangrove roots) and flow velocity, reduces flow within the canopy, and dissipates wave energy

due to friction and drag²⁷. For flexible organisms (seagrass, saltmarsh, macroalgae) the geometry changes across the wave cycle as the organisms move with the flow of water. Water depth also alters the magnitude of wave reduction by benthic habitats²⁸. While the details of the wave transformation and attenuation processes are complex, non-linear and multi-scale, the reduction in wave height and changes in dominant wave frequency can be observed visually and measured quantitatively²⁸. However, modelling wave-vegetation interactions is challenging and an area of ongoing research²⁹.

The benefit of wave attenuation by one habitat for another habitat has been documented at scales ranging from metres³⁰ to kilometres^{8,31} (Fig. 1). When considering the physical parameters such as length of habitat structure or distance and area of influence, wave attenuation by one habitat may benefit other habitats over scales of tens to hundreds of kilometres away³². For example, the Great Barrier Reef provides wave attenuation to inshore habitats across these distances. The ability of benthic habitats to attenuate waves may be especially important in the future as storm events are expected to increase in intensity, since storm waves and surge place greater stress on the survival of coastal marine habitats. Coastal habitats can attenuate storm surge height³³ and may also trap and filter increased sediment or nutrients following storm conditions (see Sediment dynamics; Nutrient processing).

Sediment dynamics

Benthic sediment dynamics, including stabilisation, accretion and destabilisation, influences and are influenced by the presence and structure of coastal marine habitats. Sediment stabilisation is the trapping of particulate organic matter from the water column. This process contributes to sediment accretion, which is the accumulation and deposition of sediment particles. Sediment stabilisation also reduces the direct smothering of propagules and minimises shading caused by sediment plumes, which can detrimentally impact adult and juvenile habitat types^{34,35}. The process of sediment accretion also provides more available substrate for settlement and growth of habitats. However, excessive sedimentation can pose a risk by burying seedlings or propagules, inhibiting their growth by restricting access to light and oxygen, which can result in high mortality rates. The effects of sediment burial on seedlings are influenced by the rate and frequency of sediment deposition, as well as the resilience of the particular species³⁶.

The key parameters that govern sediment transport are current velocity, sediment settling velocity, and suspended sediment concentration³⁷. Sediment deposition and transport are driven by current velocities to reach a balance between settling and resuspension. The particle settling velocity is determined by particle shape, size and density, which are continuously changing by aggregation and disaggregation³⁸. Additionally, the bottom roughness created by marine habitats can alter friction at the seabed, ultimately affecting sediment resuspension³⁹ (see Wave attenuation).

The presence, structure and orientation of coastal marine habitat-forming species can influence sediment dynamics by reducing near-bed turbulence, promoting sediment deposition, and inhibiting the resuspension of settled mud^{40,41}. These processes reduce turbidity, thereby increasing benthic light availability and facilitating the growth of photosynthesising species like seagrass, macroalgae and coral zooxanthellae^{42,43}. The spatial scale of positive impact can be hundreds of metres to a kilometre away⁴⁴ (Fig. 1). Species-specific morphologies can affect current velocities, thereby influencing erosion and deposition of sediment and sedimentation rates. For example, there is a positive correlation between both the height of marine macrophytes and the reduction in current velocity, and the surface area of marine macrophytes and the abundance of trapped sediment particles⁴⁵. Sediment trapping from the water column reduces sediment resuspension rates and, over time, can alter the critical photic depth that determines the occurrence of marine habitats⁴⁶. Further, these habitats can also significantly impact the sediment budget of seascapes and influence the coastal processes and sediment transport in a system. Recent models (e.g., COAWST⁴⁷) have investigated the flow-vegetation interaction between habitats with flexible structures and processes regulating sediment dynamics, but further work is needed for rigid habitats. Interactions among




Fig. 1 | The strength of cross-habitat facilitative processes provided by habitat-forming species varies with the identity of the habitat type (saltmarsh, mangrove, seagrass, macroalgae, shellfish reef and coral reef) and the spacing between habitats. We illustrate the role that different habitats play in providing facilitative processes at the scales of **a)** 1–100 m and **b)** 1–100 km and **b)** 1–100 km. Arrow widths represent the relative significance of the role (major, medium or minor) and the outlines represent the level of confidence (solid white outline = higher confidence, dashed grey outline

= lower confidence) that a habitat type plays in providing a process at each scale. The absence of an arrow does not mean that the facilitative process does not occur, but rather that research has not yet demonstrated the occurrence of the process for a specific habitat type and spatial scale. The relative significance and levels of confidence for each process were elucidated from expert knowledge and supporting literature.

habitats, sediment properties, and hydrodynamic conditions can also regulate erosion, which may be more extreme after short-term episodic events like storms, which can lead to both sediment erosion and habitat loss⁴⁸.

Nutrient processing

Nutrients are essential for the growth of plants and animals, but need to be supplied in the right forms, quantities and timing. Changes in nutrient supply can have deleterious impacts on growth at the physiological level⁴⁹, and at the ecological level, will confer advantages to some taxa over others⁵⁰. Therefore, nutrient processing via habitat-forming species can help balance the appropriate nutrient provisioning to other habitats.

Many coastal ecosystems have the capacity to reduce nitrate (NO_3^-) concentrations and ameliorate the negative effects of excess nutrients on adjacent ecosystems. Mangroves, saltmarshes and seagrasses all have mildly anoxic soils that are rich in carbon (see Carbon dynamics and alkalinity export). They are located near the coast where nutrient pollution enters via rivers, groundwater, or surface runoff. These characteristics are ideal for denitrification, which is the conversion of NO_3^- to N_2 , and the permanent removal of nitrogen (N) from the water column. Additional processes for N removal are plant growth or the accumulation of N as woody biomass, soil accumulation of particulate N, and anammox or the conversion of ammonium (NH_4^+) to nitrogen gas N_2 ⁵¹.

These processes result in changes in concentrations of N and transformations among N species that could have benefits for different ecosystems within a seascape. For instance, some coastal wetlands (mangroves, saltmarsh) can transform NO_3^- to NH_4^+ through dissimilatory nitrate reduction to ammonium (DNRA), which can result in the export of NH_4^+ to waterways⁵². This process may facilitate N uptake by phytoplankton, which prefer NH_4^+ over NO_3^- , resulting in high productivity that nearby shellfish reefs can consume and convert to particulate N^{53,54}. Furthermore, the accumulation of particulate N in the soil could be remineralised back to NO_3^- and denitrified in nearby seagrasses providing a final polishing of the water column before reaching coral reefs, providing the low nutrient conditions in which they thrive. These processes have typically been documented over spatial scales of a couple metres⁵⁵ but may occur over tens or hundreds of metres, depending on the habitat area and catchment size⁵¹ (Fig. 1). Importantly, nutrients are consumed by a variety of taxa including bacteria, primary producers, phytoplankton and zooplankton⁵⁶ and as such, competition for nutrients could potentially impact cross-habitat interactions. However, competition for nutrients such as N is most likely to occur in settings where N is limited⁵⁷; therefore, understanding of location-specific parameters is needed when planning to harness cross-habitat facilitation in restoration.

The presence of benthic marine habitats can modify phosphorous and sulphide levels. Coastal wetlands can retain phosphorous within sediments⁵⁸ (see Sediment dynamics), or in dissolved forms (e.g., soluble reactive phosphorous⁵²), potentially reducing incidences of eutrophication and thereby benefitting neighbouring habitats. At the sub-metre scale, sulphide reduction by benthic bivalves benefits adjacent seagrass habitats^{59,60} (see Animal- and plant-mediated facilitation). The processes of nutrient reduction through the passive filtering of wetlands may be especially important under storm conditions, when storm surge and tidal inundation deliver higher-than-average volumes of particulate- and nutrient-rich water into wetlands⁶¹.

Water filtration

Water filtration is the process of removing suspended particles from the water column and may occur either actively or passively. Water column particulate matter may be comprised of different trophic levels of plankton, or organic and inorganic particles (see Sediment dynamics; Carbon dynamics and alkalinity export). In this section, we focus on active filtration (for passive filtration, see Nutrient processing), which in coastal ecosystems occurs by suspension feeders, such as bivalves that filter large volumes of water, ingest the edible particles, and deposit rejected particles as biodeposits on the sediment surface⁶². Furthermore, water filtration can aid in the

removal of pollutants, contaminants, and pathogens associated with suspended sediments^{63,64}. Together, these processes benefit water clarity and purity.

The water filtration capacity of a species is often assessed through an individual's filtration rate in laboratory studies where physiological factors such as size and environmental factors such as temperature, salinity, flow rate, seston concentration and particle size have been demonstrated to influence rates⁶⁵. Over reef to estuary spatial scales, filtration rates are determined by hydrodynamic factors including current speed and water residence time, and population dynamics like density or reef size⁶⁶. For shellfish, population filtration capacity and water residence time influence the rate at which particles are cleared from the water column. Restoration of shellfish reefs are expected to result in improvements to estuary water quality. For example, historical losses of oyster reefs in Chesapeake Bay resulted in declines in water quality⁶² but restored shellfish reefs can provide water-quality improvements⁶⁷.

Water filtration has important down-stream effects in estuarine systems and depends on local- and estuary-scale hydrodynamics, and the spatial position and distribution of the filtration provider (Fig. 1). In particular, the cross-habitat facilitative benefit of water filtration by shellfish improving water clarity is well-established for adjacent seagrass habitats⁶⁸. Seagrass habitats are sensitive to turbidity, which reduces seafloor light availability, thereby impacting seagrass productivity⁶⁹. The benefit of water filtration on seagrass may occur on scales as small as centimetres or up to a few kilometres, depending on the size of the estuary or embayment⁷⁰ (Fig. 1). Directionality is an important factor for cross-habitat facilitation from water filtration, with benefits generally found "down-stream". But tidal and riverine outflow processes are complex, so that "down-stream" locations may vary temporally over tidal-seasonal scales.

Carbon dynamics and alkalinity export

The marine carbon (C) cycle plays a major role in global C cycling and the growing interest in marine Carbon Dioxide Removal (mCDR) approaches to sequester excess atmospheric carbon⁷¹. The mCDR concept most related to seascape restoration is "blue carbon": the management of C stored and carbon dioxide (CO_2) removal by CO_2 fixation by vegetated marine habitats⁷². However, there is continued research on the role non-vegetated habitats may play in this process⁷³. Many aspects of the C cycle are needed for mCDR and blue carbon accounting but not all are important for cross-habitat facilitation. For clarity, we first define key components of the C cycle and then describe how some are important in cross-habitat facilitation.

Carbon sequestration by macrophytes is the uptake and storage of C via sediment trapping or accumulation of biomass, mostly as wood and roots⁷⁴. Biomass is continually transferred to sediments where physical and chemical changes in the sediment (diagenetic processing) convert part of the photosynthetically derived C to microbial C, dissolved inorganic carbon (DIC) or methane⁷⁵. This diagenetic processing influences the portion of deposited C that is sequestered in sediments as long-term C, emitted as gas or exported to surrounding waters⁷⁶. While sediment trapping sequesters C and also provides clearer water (see Sediment dynamics), the links between biomass accumulation and facilitative cross-habitat interactions are less clear⁷⁷. For example, dissolved organic C and litter export to reefs or coastal waters accounts for a large proportion of the C cycle⁷⁸; these organic C exports are likely important⁷⁷, but are highly variable in terms of the benefits to other habitat types. Further studies are needed that ascertain habitat-specific responses to C exports.

Inorganic C also plays an important role in cross-habitat facilitation. DIC is composed of four carbon species ($\text{CO}_{2\text{aq}}$, H_2CO_3 , HCO_3^- and CO_3^{2-}). Production of $\text{CO}_{2\text{aq}}$ (herein, aqueous $\text{CO}_{2\text{aq}}$ and carbonic acid H_2CO_3 ⁷⁹) from metabolic or chemical processes of habitats decreases the pH and acidifies waters, affecting adjacent habitats. Waters that are supersaturated in the partial pressure of $\text{CO}_{2\text{aq}}$ relative to the atmosphere will off-gas CO_2 , increasing pH until equilibrium is reached. Thus, the effect of acidification is greatest closest to the $\text{CO}_{2\text{aq}}$ source, decreasing over space and time. The production and export of alkalinity (including HCO_3^- and

CO_3^{-2}) increases the ability of water to buffer (reduce) acidity⁸⁰. The ability of marine habitats to generate alkalinity occurs through at least two processes: photosynthesis⁸¹ and organic matter decomposition⁸², both of which can result in increased pH. Alkalinity export is likely to be highest in estuaries with large tidal amplitudes⁸³ and after rainfall⁸⁴.

Climate change is causing acidification of the oceans⁷⁹. Alkalinity export from tidal wetlands and intertidal and submerged aquatic vegetation^{81,85} to neighbouring calcified habitats (shellfish and coral reefs) within zero to tens of metres, could support their restoration^{86,87}, especially in poorly buffered environments or those highly susceptible to acidification (Fig. 1). Although questions remain around the relative magnitude of C fluxes provided by certain habitats and how to attribute the amount of alkalinity export to a certain area of vegetation⁸⁸, we are beginning to better understand the spatial variability and scales involved. For example, C storage by vegetated habitats may vary depending on the presence of other vegetated⁸⁹ or non-vegetated⁹⁰ habitats and across different regions^{91,92}.

Animal- and plant-mediated facilitation of other habitats

Here, we describe processes attributed to the movement and/or establishment of coastal marine animals and plants that can facilitate the establishment or persistence of habitat-forming species. Many of these processes are underpinned by one of the five processes described above. However, in the context of this section, these processes are either directly provided by animal or plant movements (e.g., nutrient transport by fish, abrasion by macroalgae fronds), or result from direct plant or animal mediation of the environmental and biological conditions at the site (e.g., shading, competition reduction, herbivory). We acknowledge that in addition to the positive, facilitative interactions discussed below, there are a suite of important negative interactions that exist among plant and animal species in coastal habitats. Indeed, managing for negative species interactions has long been a foundational principle in restoration and conservation science⁹³. For instance, invasive species control has and continues to be key for restoration success¹⁹. A key research gap is to characterise the ecological contexts or spacings in which interactions switch from positive to non-existent or negative.

Cross-habitat facilitation is mediated by mobile animals transporting energy and nutrients among habitats⁹⁴ and through animals migrating across habitats and supporting key ecological functions that support ecosystem resilience in one or more of those habitats⁹⁵. In this sense, increased functional connectivity of animals between different coastal marine habitats can significantly enhance restoration outcomes if they have a positive effect on the likelihood of settlement, growth, or persistence of habitat-forming species. Across seascapes, classic examples of these processes include mangroves providing nurseries for herbivorous fish that clear algae and facilitate coral recruitment and growth (hundreds of metres to kilometres scales⁹⁶), bioturbators oxygenating sediments and processing pollutants to create suitable conditions for other species (centimetres to metre scales⁹⁷), and animal-mediated nutrient transport between different habitats via excretion (metres to hundreds of metre scales⁹⁴) (Fig. 1). However, the presence of these animals may not equate to the presence of a particular facilitative process (e.g., herbivory of a competitive algae from coral reefs⁹⁸) and necessitates context-specific consideration of whether a process(es) will occur.

Facilitation may also be mediated by sessile habitat-forming animals and plants. The mechanisms by which these organisms directly facilitate others includes competition reduction (e.g., abrasion that reduces competitor settlement within 1 m⁹⁹), or modification of environmental conditions within a few metres (e.g., shade¹⁰⁰ or temperature¹⁰¹) that allow for the establishment of another habitat (Fig. 1). However, these same processes that may be beneficial for a particular life stage and habitat type (e.g., abrasion by kelp that enables juvenile oyster settlement⁹⁹), could presumably be detrimental to other habitat life stages or types. Another important mechanism is through the process of colonisation and succession. For example, colonisation by saltmarsh vegetation can facilitate the subsequent establishment of mangroves (Box 1) by trapping propagules and altering

hydrodynamics or temperature at the site¹⁰². Alternatively, mangroves are capable of encroaching into saltmarsh (particularly as temperature increases at their poleward edge), at times leading to coastal squeeze and saltmarsh losses^{103,104}, reinforcing the need to consider both positive and negative interactions during restoration planning.

As the scale and scope of seascape restoration programmes expands, and especially when restoring multiple habitats simultaneously, the importance of these considerations likely increases. Firstly, enhancing connectivity between restoration sites and remnant habitats maximises the likelihood that species that drive facilitation will recruit to restored habitats. Secondly, maximising connectivity between new restoration sites of different habitat types helps ensure these connections can result in synergistic benefits across both new habitat types. Ultimately, the ability of animals or plants to mediate facilitation through multi-habitat connectivity depends on the species present, the ecological function by which they facilitate another habitat (e.g., grazing for competition reduction, nutrient or propagule transport, modification of environmental conditions), and the distance over which this process can effectively occur. Given the simultaneous increase in the likelihood of positive and negative interactions existing from enhancing connectivity, restoration practitioners would benefit from a sound ecological understanding of their system to account for all plausible outcomes (see ref. 19).

Minima and maxima lateral distances of cross-habitat facilitative processes

Through expert opinion of the authorship team, we characterised the likely minimum and maximum lateral distances across which each facilitative process may occur in the context of seascape restoration. The lateral minima and maxima of these processes illustrate opportunities for spacing restoration projects to harness particular processes (Fig. 2; Supplementary Table S1). For example, structure-based facilitation provided by habitat-forming species (e.g., shading, abrasion) generally occurs within distances of 0–1 m in predominately nested or co-occurring habitat patch mosaics, but larger physical processes (e.g., wave attenuation) occur from distances of tens to hundreds of metres. Through this process we identified examples of all six processes underpinning cross-habitat facilitation overlapping at distances between 1 and 100 m. At all distances considered, at least three processes overlapped (Fig. 2). Uncertainty tended to increase with increasing distance, suggesting that the practicality of harnessing facilitative processes in restoration may decrease at these larger spatial scales.

Importantly, for biologically mediated interactions, there is likely a distance-dependent shift between positive or negative interactions. Depending on context, biologically mediated interactions may be negative at close distances (<10 m; overgrazing of algae and corals by parrotfish¹⁰⁵), but positive at longer distances (>1 km; grazing of algae by parrotfish that helps corals⁹⁶). Therefore, whether the ecological processes described herein are facilitative at certain distances will be largely context-dependent, based on ecological conditions and the species present and their traits¹⁰⁶. The effects of seascape configuration on biologically mediated facilitation may depend on the ecological function resulting in facilitation. For example, in Australia, isolated coral reefs tended to have greater rates of herbivory than reefs located closer to other reefs, which tended to have higher rates of piscivory; additionally, proximity to mangrove or seagrass habitats did not have a strong influence on herbivory or piscivory in this system¹⁰⁷. There may be similar shifts between positive and negative effects of physical and biogeochemical processes at certain distances, but these did not arise during our assessment.


Interestingly, current restoration permitting practices often seek to mitigate potential negative interactions among habitats by requiring a buffer between restoration sites and existing habitats. Exact permitting requirements vary by country and jurisdiction, and local classifications of restoration actions (e.g., development or restorative), but restoration actions are generally limited 5–50 m within or adjacent to areas designated as other types of habitat. Two states in Australia (New South Wales and Queensland) require a buffer of 10–50 m around seagrass beds and at least 100 metres

Box 2 | Rules of Thumb for practitioners seeking to harness cross-habitat facilitation in seascape restoration

Here we describe key takeaway points in six “Rules of Thumb” that can help practitioners maximise cross-habitat facilitative interactions to scale-up seascape restoration. As with any restoration project, these rules assume that prior to conducting seascape restoration, practitioners will complete expected pre-planning steps including basing restoration methodologies on field trials or prior studies, identifying site-specific restoration targets, suitable habitat types to restore, and system stressors¹³⁰, including balancing the cost-effectiveness of management or restoration options under climate change¹³¹. We also provide a restoration decision flow chart that may aid the decision-making process when planning to conduct seascape restoration that seeks to harness cross-habitat facilitation (Box 2 Fig. 1). In this flow chart, we illustrate how the multi-habitat restoration typologies (Box 1) can be implemented to achieve the desired environmental conditions or configurations required for multi-habitat restoration.

- 1. Develop restoration suitability models that consider environmental, social, regulatory and facilitative processes.** Site selection and feasibility must be guided by the environmental parameters that dictate the survival of the targeted habitat-forming species to restore (e.g., water clarity, salinity, temperature and depth). Combining environmental, logistical, social and regulatory constraints at a proposed restoration space through restoration suitability models can help practitioners identify potential restoration locations, for example, in highly urbanised environments where space for restoration is very limited¹¹⁵. Simultaneously, practitioners must also determine the required size and spatial configuration of restored patches of a habitat type to achieve desired outcomes at a designated restoration site. While these conceptual or restoration suitability models can refine the areas where restoration of one or more habitat types may suit best, they typically have not focused on identifying optimal spacing that maximises cross-habitat facilitative processes for the restoration of multiple connected habitats. Therefore, future models should include information on the spatial scales of particular facilitative processes to better plan and account for overlap among habitat suitability and these processes.
- 2. Consider the order in which restoration actions are conducted to achieve desired outcomes.** During the planning process, logic models or Theory of Change activities¹³² can help practitioners work through the required steps or stages that may be required to achieve a certain restoration outcome(s). This practice aims to identify causal solutions to overcome a problem or achieve a desired goal. For example, sequential multi-habitat restoration (Box 1) is based on the underlying evidence and assumptions that restoration of habitat A will result in condition B that through facilitative processes, will enable the establishment or restoration of habitat C. Careful consideration of the assumptions being made about facilitative processes arising from restoration actions, through logic models or theory of change activities, will help ensure that desired objectives are optimised and met at each project stage or milestone.
- 3. Relate the cross-habitat facilitative processes of interest to the objectives of restoration.** Restoration projects often seek to restore degraded habitats and boost specific ecosystem services in these areas, both of which may be maximised by strategic harnessing of facilitative processes. For example, in highly urbanised environments, restoration projects may seek to prioritise habitat provision but also other co-benefits such as bioremediation, or erosion control (e.g., Project Restore¹³³). When thinking about this through a cross-habitat facilitation lens, restoration or enhancement of filter-feeding taxa can aid in bioremediation and provide clearer water for photosynthesising taxa like seagrass or macroalgae, which, in turn, can aid in erosion control. However, maximising habitat provisioning (e.g., restoring a particular area or configuration of habitat) may not be directly correlated to maximising the other co-benefits of restoration (in this example, bioremediation and erosion control). Additionally, different co-benefits of restoration may not be compatible either depending on environmental conditions or spatial constraints within the system. Therefore, prioritisation of key processes may mean that the size and spatial distribution of the restored area and habitat types is constrained to the optimum patch size and distribution to maximise the targeted facilitative processes (and ultimately, ecosystem services), rather than aiming for the largest possible restored area.
- 4. Have a local understanding of the spatial scales of processes in your seascape.** Whether processes have a facilitative, neutral or even negative effect on adjacent habitats depends on a variety of physical and environmental parameters, whose effects vary over different time and spatial scales (Fig. 2 and Box 2 Fig. 1; Supplementary Table S1). Ideally, site assessments and site data (covering many seasons and years) could help guide restoration planning decisions, but often these data may be obtained through comparisons of outcomes at other locations, published literature or through modelling approaches. It is also important to ensure that there are not any unintended, negative consequences of an action. For example, tidal reintroduction can aid in wetland, saltmarsh and mangrove restoration, but can initially lead to high sedimentation rates¹³⁴, which could inhibit the growth of adjacent habitats such as seagrasses or corals. Further, understanding the scales over which we expect processes to operate can also help mitigate unforeseen negative impacts of an action.
- 5. Optimise heterogeneity and habitat diversity within the seascape to maximise beneficial processes.** Cross-habitat facilitative processes have been documented and described across a variety of different habitat types^{4,9–11}; Supplementary Table S1). Optimising the diversity of (locally relevant) habitats in a seascape should result in the provisioning of a variety of facilitative processes. In planning and selecting for diverse habitats in seascape restoration, one must be cognisant of the appropriate species to use depending on the desired outcomes. In some instances, this may include restoration of an endemic species within its endemic ranges, using traditional planting or seeding methodologies. In other instances, it may be necessary to consider species that will help maximise the long-term restoration success by restoring a specific species that can tolerate future shifts or deterioration of conditions. Therefore, it is important to consider likely influences from changes in climate that may alter the strength of positive or negative feedbacks. Finally, it may be necessary to use modified approaches (e.g., rock substrate instead of shell for oyster restoration), or hybrid restoration approaches that are suitable to current day and future conditions.
- 6. Monitor restoration for cross-habitat-facilitation.** Following best-practice standards, restoration should be followed by a monitoring programme¹ to assess the efficacy and outcomes of restoration. These monitoring programmes should be designed so that they measure the key cross-habitat facilitative processes that were identified and targeted during the restoration planning stage. Additionally, monitoring may help inform when the restored area has grown or established sufficiently to provide the facilitative process at the scale and magnitude needed to benefit other habitats. Practitioners may not have the capacity to conduct pre- and post-restoration monitoring while also managing the successful procurement and delivery of a large-scale seascape restoration project. However, this presents a great opportunity to pair practice with research teams (e.g., universities, government agencies, citizen science groups) who can conduct monitoring to understand whether cross-habitat facilitation was achieved through seascape restoration.

(continued from previous page)



Stages of Restoration



Box 2 Fig. 1 Flow chart that describes decisions that practitioners may need to make when seeking to harness cross-habitat facilitation in seascape restoration. Adjacent habitats may refer to habitats immediately next to or in the surrounding vicinity of the focal habitat restoration area. Where each step falls within the stages of restoration (adapted from ref. 130) are indicated in colour-coordinated outlines. Harnessing facilitation in seascape restoration requires careful site selection and suitability modelling and planning and permitting, as well as careful evaluation (monitoring) of the outcomes. This flow chart represents an iterative process where decisions or questions may need to be revisited as restoration actions are implemented and monitored, or as stressors (e.g., storms, invasive species) occur in a seascape.




Fig. 2 | Approximate minima and maxima lateral distances of cross-habitat facilitative processes. The x-axis represents the log-scale lateral distance of a process in metres. Solid lines represent known distances of facilitative processes; dashed lines represent uncertainty in the distance. The horizontal grey line from 5 to 50 m

represents the common buffer that restoration sites are usually required to be placed away from other existing habitats to potentially mitigate negative effects. References in Supplementary Table S1.

around “sensitive” habitats^{108,109}. In the United States, the Florida Coastal Zone Management Act requires a 30 m buffer zone around seagrass beds in areas of coastal development¹¹⁰. However, these buffer distances are not necessarily based on an ecological understanding of all the potential interactions that can occur¹¹¹, but rather an attempt to mitigate harm. For example, long-distance negative interactions are also common and can regulate ecosystems¹⁰, and the distance over which they operate may even exceed these permitting buffers. Therefore, it is necessary to understand how both positive and negative interactions operate in seascape to move forward with scaled seascape restoration. Without this knowledge, implementing a somewhat arbitrary buffer distance may preclude the opportunity for some facilitative interactions to occur and unintentionally limit restoration outcomes.

By considering the gaps or overlaps in lateral distances of multiple processes, we can identify where there may be trade-offs in designing restoration with spacing that seeks to optimise certain processes related to project objectives. For example, a restoration project seeking to maximise both wave attenuation through the restoration of a reef system and sediment dynamics via sediment trapping through the restoration of mangroves may prioritise a spacing of 10–1000 m between these two habitats. However, at this distance range, the ability for wetlands to passively filter nutrients or aid in alkalinity export is less certain than the ability of the reef to attenuate waves, necessitating a decision in which process(es) to prioritise. Although not the focus of this manuscript, negative biotic interactions (e.g., competition, herbivory, predation) are highly context-dependent and depending

on local conditions, the spacing that influences these interactions is something that needs to be considered when designing restoration.

Incorporating cross-habitat facilitation in seascape restoration

Seascape restoration is being operationalised through multiple restoration modes (Box 1) and this review demonstrates that there are opportunities to harness cross-habitat facilitation at multiple spatial scales. Identifying new opportunities for seascape restoration that harnesses facilitation is constrained by numerous biophysical (e.g., habitat suitability for desired restoration species), logistical (e.g., contracts and site access), and social considerations (e.g., stakeholder perceptions of and support for proposed restoration projects). Therefore, practitioners working within these constraints need tools such as decision support frameworks, and conceptual and restoration suitability models that can help guide the design and implementation of seascape restoration to achieve desired objectives, notably, those related to harnessing cross-habitat facilitation (Box 2). By expanding on best-practices already implemented by practitioners when designing restoration projects, such as identifying clear targets, suitable species and local environmental conditions, the information provided in this review aims to illustrate how seascape restoration can be taken a step further to account for and harness facilitative interactions.

The spatial configuration of the seascape and habitat types therein, is a key determinant of the strength of facilitative interactions. The patch size and configuration of habitat types (including distance between patches of

the nearest neighbour as well as distance from other habitat patches in the overall seascape mosaic), varies with environmental context and in turn can influence the functioning or provisioning of facilitative processes by these habitats^{112,113}. Additionally, the magnitude of facilitation may be determined by the spatial configuration of the habitat within the broader context of the seascape. For example, we would expect mangrove facilitation of coral reefs by trapping sediment runoff to be of greater importance when mangroves are located at the mouths of rivers than when mangroves are located on an open coast¹¹⁴. The final spatial configuration of restoration projects will ultimately be influenced by a variety of environmental and social factors¹¹⁵ (Box 2). Therefore, seascape restoration planning must also consider how the eventual configuration will influence the strength of prioritised or desired facilitative interactions.

Although not a focus of this paper, the same physical, biogeochemical and biological processes of cross-habitat facilitation also underpin self-facilitation within a single habitat type⁶⁰, which in turn can influence facilitation of other habitats. Having many medium-sized seagrass beds rather than one large seagrass bed facilitates more diverse fish nurseries and overall habitat values, while also reducing erosion in the adjacent seascape¹¹⁶, which in turn can lead to a greater diversity of fish at adjacent habitats. Processes affecting habitat patch dynamics can include ontogenetic movements, hydrodynamic conditions, and the presence of grazers that influence coral settlement^{117,118}; these same processes were identified as occurring between corals and other adjacent habitat types at larger scales (Fig. 2). Therefore, it is important to understand that restoration seeking to harness cross-habitat facilitative processes will also depend on the same processes operating within a single habitat type.

Restoration remains a costly endeavour¹², requiring careful consideration of the potential successes and failures of restoration actions. Negative interactions such as competition and predation or herbivory can have negative outcomes on restoration¹⁹. For example, although saltmarshes can facilitate mangroves by trapping their seeds and modifying abiotic stressors (e.g., substrate provisioning and temperature and shading, respectively¹⁰²), competition for space between these plants at later life stages can inhibit the growth and expansion of one or both of these habitats¹⁹. Further, whether these cross-habitat interactions are positive or negative may be influenced by looming global threats such as climate change. Increasing temperatures as a result of climate change and the subsequent poleward expansion of mangrove habitat has led to the intrusion of mangroves into saltmarsh habitat at these geographic locations¹⁰⁴. Therefore, it is critical for managers and practitioners seeking to conduct seascape restoration to have a clear understanding of how their restoration sites may be influenced by both facilitative and negative cross-habitat processes, and how these processes may be altered further under a changing climate.

Restoration actions must be paired with climate change mitigation and conservation measures that seek to first protect and halt the decline of natural habitats¹²⁰. Following this, climate-smart coastal marine restoration should include clear strategies to adapt to pressures faced by climate change¹²¹. To do this, practitioners, managers and scientists must work together to implement restoration monitoring programmes that will ultimately inform required adaptation plans of these systems¹²².

Future research priorities in cross-habitat facilitation in seascapes

We have demonstrated how facilitative processes vary across lateral distances to help inform seascape restoration design. We show that different processes occur in different habitat types, suggesting that habitat heterogeneity may increase overall opportunities for cross-habitat facilitation within a seascape. Crucially, we also demonstrate how adaptive management of restoration can minimise unintended consequences of certain actions seeking to harness facilitation (Box 2). Still, there remain two key research priorities related to harnessing cross-habitat facilitation in seascapes. The first priority is to better understand the temporal (including seasonal) dependencies of cross-habitat facilitation through restoration. For example, the magnitude by which a coral reef attenuates waves depends on

its structural complexity²¹; therefore, wave attenuation by a newly restored coral reef will take time to reach the desired complexity for this process to occur. Seasonal variations in the density or structure of a habitat-forming species may also affect these processes (e.g., seagrass density and variability in wave attenuation¹²³). The second priority is to better understand the non-linearity of these processes. The magnitude of facilitation along a lateral distance over which we know it operates (Fig. 2) may be non-linear. Therefore, we need to understand the non-linearities of these processes and how they interact with each other across different spatial scales to better incorporate and plan for cross-habitat facilitation in restoration. This may need assessment on a context-specific basis, where a strong understanding of local environmental conditions (bathymetry, habitats present, and stressors) can be incorporated into a model that disentangles these non-linearities of cross-habitat facilitation. Although this review has predominantly provided examples of processes occurring across pairs of habitat types in seascapes, it is important to consider that facilitative processes may have flow-on effects (facilitation cascades¹¹) and/or interact with each other within the seascape.

Conclusion

We face an urgent need to repair nature globally as articulated by the UN Decade on Ecological Restoration and Kunming-Montreal Global Biodiversity Framework. To mitigate the combined climate and biodiversity crises, it is imperative that we scale-up restoration quickly and effectively. Facilitation is intrinsic in coastal marine environments^{6,10}. To achieve at-scale seascape restoration, cross-habitat facilitative processes need to be better understood and harnessed in restoration. While these processes are context-dependent and occur across a large range of lateral distances, we highlight where opportunities exist to achieve at-scale seascape restoration using cross-habitat facilitation and raise key questions that demand future research.

Data availability

Data is provided within the supplementary information files.

Received: 15 April 2024; Accepted: 12 November 2024;

Published online: 30 November 2024

References

- Gann, G. D. et al. International principles and standards for the practice of ecological restoration. Second edition. *Restor. Ecol.* **27**, S1–S46 (2019).
- Saunders, M. I. et al. Bright spots in coastal marine ecosystem restoration. *Curr. Biol.* **30**, R1500–R1510 (2020).
- Pittman, S. J. et al. Seascape ecology: identifying research priorities for an emerging ocean sustainability science. *Mar. Ecol. Prog. Ser.* **663**, 1–29 (2021).
- Vozzo, M. L. et al. To restore coastal marine areas, we need to work across multiple habitats simultaneously. *Proc. Natl Acad. Sci.* **120**, e2300546120 (2023).
- Gilby, B. L., Olds, A. D., Connolly, R. M., Henderson, C. J. & Schlacher, T. A. Spatial restoration ecology: placing restoration in a landscape context. *BioScience* **68**, 1007–1019 (2018).
- Silliman, B. R. et al. Facilitation shifts paradigms and can amplify coastal restoration efforts. *Proc. Natl Acad. Sci. USA* **112**, 14295–14300 (2015).
- Bruno, J. F., Stachowicz, J. J. & Bertness, M. D. Inclusion of facilitation into ecological theory. *Trends Ecol. Evol.* **8**, 119–125 (2003).
- Saunders, M. I. et al. Interdependency of tropical marine ecosystems in response to climate change. *Nat. Clim. Change* **4**, 724–729 (2014).
- Gillis, L. G. et al. Potential for landscape-scale positive interactions among tropical marine ecosystems. *Mar. Ecol. Prog. Ser.* **503**, 289–303 (2014).

10. van de Koppel, J. et al. Long-distance interactions regulate the structure and resilience of coastal ecosystems. *Ann. Rev. Mar. Sci.* **7**, 139–158 (2015).
11. Gribben, P. E. et al. in *Oceanography and Marine Biology: an Annual Review* Vol. 57 (eds Hawkins, S. J. et al.) Ch. 3, 127–168 (Taylor & Francis, 2019).
12. Bayraktarov, E. et al. The cost and feasibility of marine coastal restoration. *Ecol. Appl.* **26**, 1055–1074 (2016).
13. UNEP-WCMC et al. Progress, needs and opportunities for seascape restoration (UNEP, 2022).
14. McAfee, D. et al. Multi-habitat seascape restoration: optimising marine restoration for coastal repair and social benefit. *Front. Mar. Sci.* **9**. <https://doi.org/10.3389/fmars.2022.910467> (2022).
15. Elton, C. Competition and the structure of ecological communities. *J. Anim. Ecol.* **15**, 54–68 (1946).
16. Glasser, J. W. The role of predation in shaping and maintaining the structure of communities. *Am. Nat.* **113**, 631–641 (1979).
17. Morris, R. L. et al. Key principles for managing recovery of Kelp forests through restoration. *Bioscience* **70**, 688–698 (2020).
18. Reeves, S. E. et al. Facilitating better outcomes: how positive species interactions can improve oyster reef restoration. *Front. Mar. Sci.* **7**. <https://doi.org/10.3389/fmars.2020.00656> (2020).
19. Silliman, B. R. et al. Harnessing ecological theory to enhance ecosystem restoration. *Curr. Biol.* **34**, R418–R434 (2024).
20. Dean, R. G. & Dalrymple, R. A. *Coastal Processes with Engineering Applications* (Cambridge University Press, 2004).
21. Lowe, R. J. et al. Spectral wave dissipation over a barrier reef. *J. Geophys. Res.: Oceans* **110** (2005).
22. Denny, M. (Princeton University Press, 1988).
23. Battjes, J. A. Surf-zone dynamics. *Annu. Rev. Fluid Mech.* **20**, 257–291 (1988).
24. Holman, R. Nearshore processes. *Rev. Geophys.* **33**, 1237–1247 (1995).
25. Henderson, S. M. Motion of buoyant, flexible aquatic vegetation under waves: simple theoretical models and parameterization of wave dissipation. *Coast. Eng.* **152**, 103497 (2019).
26. Roncolato, F. et al. Ecomorphodynamics of oyster reefs and their influence on oyster reef morphology. *Geomorphology* **456**, 109213 (2024).
27. Lowe, R. J., Falter, J. L., Koseff, J. R., Monismith, S. G. & Atkinson, M. J. Spectral wave flow attenuation within submerged canopies: Implications for wave energy dissipation. *J. Geophys. Res. Oceans* **112**. <https://doi.org/10.1029/2006JC003605> (2007).
28. Horstman, E. M. et al. Wave attenuation in mangroves: a quantitative approach to field observations. *Coast. Eng.* **94**, 47–62 (2014).
29. van Rooijen, A., Lowe, R., Ghisalberti, M., McCall, R. & Hansen, J. Modelling wave attenuation through submerged vegetation canopies using a subgrid canopy flow model. *Coast. Eng.* **176**, 104153 (2022).
30. Safak, I. et al. Coupling breakwalls with oyster restoration structures enhances living shoreline performance along energetic shorelines. *Ecol. Eng.* **158**. <https://doi.org/10.1016/j.ecoleng.2020.106071> (2020).
31. Guerra-Vargas, L. A., Gillis, L. G. & Mancera-Pineda, J. E. Stronger together: do coral reefs enhance seagrass meadows “blue carbon” potential? *Front. Mar. Sci.* **7**. <https://doi.org/10.3389/fmars.2020.00628> (2020).
32. Bridges, T. S. et al. *International Guidelines on Natural and Nature-Based Features for Flood Risk Management* (U.S. Army Engineer Research and Development Center, 2021).
33. Möller, I. et al. Wave attenuation over coastal salt marshes under storm surge conditions. *Nat. Geosci.* **7**, 727–731 (2014).
34. Ricardo, G. F., Jones, R. J., Nordborg, M. & Negri, A. P. Settlement patterns of the coral *Acropora millepora* on sediment-laden surfaces. *Sci. Total Environ.* **609**, 277–288 (2017).
35. Fraser, M. W. et al. Effects of dredging on critical ecological processes for marine invertebrates, seagrasses and macroalgae, and the potential for management with environmental windows using Western Australia as a case study. *Ecol. Indic.* **78**, 229–242 (2017).
36. Cabaço, S., Santos, R. & Duarte, C. M. The impact of sediment burial and erosion on seagrasses: a review. *Estuar. Coast. Shelf Sci.* **79**, 354–366 (2008).
37. de Boer, W. F. Seagrass–sediment interactions, positive feedbacks and critical thresholds for occurrence: a review. *Hydrobiologia* **591**, 5–24 (2007).
38. Prandle, D. *Estuaries: Dynamics, Mixing, Sedimentation and Morphology* (Cambridge University Press, 2009).
39. Contti Neto, N., Pomeroy, A., Lowe, R. & Ghisalberti, M. Seagrass meadows reduce wind-wave driven sediment resuspension in a sheltered environment. *Front. Mar. Sci.* **8**. <https://doi.org/10.3389/fmars.2021.733542> (2022).
40. Reise, K. Sediment mediated species interactions in coastal waters. *J. Sea Res.* **48**, 127–141 (2002).
41. Colden, A. M., Fall, K. A., Cartwright, G. M. & Friedrichs, C. T. Sediment suspension and deposition across restored oyster reefs of varying orientation to flow: implications for restoration. *Estuaries Coasts* **39**, 1435–1448 (2016).
42. Adams, M. P. et al. Feedback between sediment and light for seagrass: where is it important? *Limnol. Oceanogr.* **61**, 1937–1955 (2016).
43. Bessell-Browne, P. et al. Impacts of turbidity on corals: the relative importance of light limitation and suspended sediments. *Mar. Pollut. Bull.* **117**, 161–170 (2017).
44. Akhand, A. et al. Lateral carbon fluxes and CO₂ evasion from a subtropical mangrove-seagrass-coral continuum. *Sci. Total Environ.* **752**. <https://doi.org/10.1016/j.scitotenv.2020.142190> (2021).
45. Gacia, E., Granata, T. C. & Duarte, C. M. An approach to measurement of particle flux and sediment retention within seagrass (*Posidonia oceanica*) meadows. *Aquat. Bot.* **65**, 255–268 (1999).
46. Carr, J., D’Odorico, P., McGlathery, K. & Wiberg, P. Stability and bistability of seagrass ecosystems in shallow coastal lagoons: role of feedbacks with sediment resuspension and light attenuation. *J. Geophys. Res. Biogeosci.* **115**. <https://doi.org/10.1029/2009JG001103> (2010).
47. Beudin, A., Kalra, T. S., Ganju, N. K. & Warner, J. C. Development of a coupled wave-flow-vegetation interaction model. *Comput. Geosci.* **100**, 76–86 (2017).
48. Dahl, M., Björk, M. & Gullström, M. Effects of seagrass overgrazing on sediment erosion and carbon sink capacity: current understanding and future priorities. *Limnol. Oceanogr. Lett.* **6**, 309–319 (2021).
49. Silbiger, N. J. et al. Nutrient pollution disrupts key ecosystem functions on coral reefs. *Proc. R. Soc. B Biol. Sci.* **285**, 20172718 (2018).
50. D’Angelo, C. & Wiedenmann, J. Impacts of nutrient enrichment on coral reefs: new perspectives and implications for coastal management and reef survival. *Curr. Opin. Environ. Sustain.* **7**, 82–93 (2014).
51. Adame, M. F. et al. A conceptual model of nitrogen dynamics for the Great Barrier Reef catchments. *Mar. Pollut. Bull.* **173**. <https://doi.org/10.1016/j.marpolbul.2021.112909> (2021).
52. Adame, M. F., Viridis, B. & Lovelock, C. E. Effect of geomorphological setting and rainfall on nutrient exchange in mangroves during tidal inundation. *Mar. Freshw. Res.* **61**, 1197–1206 (2010).
53. Collos, Y. et al. Response of coastal phytoplankton to ammonium and nitrate pulses: seasonal variations of nitrogen uptake and regeneration. *Aquat. Ecol.* **37**, 227–236 (2003).

54. Pan, K. et al. Control of phytoplankton by oysters and the consequent impact on nitrogen cycling in a Subtropical Bay. *Sci. Total Environ.* **796**. <https://doi.org/10.1016/j.scitotenv.2021.149007> (2021).
55. Mishra, A. K. & Apte, D. Ecological connectivity with mangroves influences tropical seagrass population longevity and meadow traits within an island ecosystem. *Mar. Ecol. Prog. Ser.* **644**, 47–63 (2020).
56. Philippart, C. J. M. et al. Impacts of nutrient reduction on coastal communities. *Ecosystems* **10**, 96–119 (2007).
57. Trott, L. A., McKinnon, A. D., Alongi, D. M., Davidson, A. & Burford, M. A. Carbon and nitrogen processes in a mangrove creek receiving shrimp farm effluent. *Estuar. Coast. Shelf Sci.* **59**, 197–207 (2004).
58. Xu, G., Song, J., Zhang, Y., Lv, Y. & Han, G. Enhancement of phosphorus storage capacity of sediments by coastal wetland restoration, Yellow River Delta, China. *Mar. Pollut. Bull.* **150**, 110666 (2020).
59. de Fouw, J. et al. A facultative mutualism facilitates European seagrass meadows. *Ecography* **2023**, e06636 (2023).
60. Valdez, S. R. et al. Positive ecological interactions and the success of seagrass restoration. *Front. Mar. Sci.* **7**. <https://doi.org/10.3389/fmars.2020.00091> (2020).
61. Temmerman, S. et al. Marshes and mangroves as nature-based coastal storm buffers. *Annu. Rev. Mar. Sci.* **15**, 95–118 (2023).
62. Grabowski, J. H. & Peterson, C. H. in *Ecosystem Engineers: Plants to Protists* (eds K. Cuddington, J. E. Byers, W. G. Wilson, & A. Hastings) 281–298 (Elsevier Academic Press, 2007).
63. Broszeit, S., Hattam, C. & Beaumont, N. Bioremediation of waste under ocean acidification: reviewing the role of *Mytilus edulis*. *Mar. Pollut. Bull.* **103**, 5–14 (2016).
64. Burge, C. A. et al. The use of filter-feeders to manage disease in a changing world. *Integr. Comp. Biol.* **56**, 573–587 (2016).
65. Cranford, P. J., Ward, J. E. & Shumway, S. E. in *Shellfish Aquaculture and the Environment* (ed S. E. Shumway) (John Wiley & Sons, Inc., 2011).
66. Gray, M. W. et al. Beyond residence time: quantifying factors that drive the spatially explicit filtration services of an abundant native oyster population. *Estuaries Coasts* **45**, 1343–1360 (2022).
67. Grizzle, R. E., Greene, J. K. & Coen, L. D. Seston removal by natural and constructed intertidal eastern oyster (*Crassostrea virginica*) reefs: a comparison with previous laboratory studies, and the value of in situ methods. *Estuaries Coasts* **31**, 1208–1220 (2008).
68. Newell, R. I. E. & Koch, E. W. Modeling seagrass density and distribution in response to changes in turbidity stemming from bivalve filtration and seagrass sediment stabilization. *Estuaries* **27**, 793–806 (2004).
69. Flowers, G. J. L., Needham, H. R., Bulmer, R. H., Lohrer, A. M. & Pilditch, C. A. Going under: the implications of sea-level rise and reduced light availability on intertidal primary production. *Limnol. Oceanogr.* **68**, 1301–1315 (2023).
70. Cranford, P. J. in *Goods and Services of Marine Bivalves* (eds A. C. Smaal et al.) 119–141 (Springer International Publishing, 2019).
71. NASEM & National Academies of Sciences, E., and Medicine. *A Research Strategy for Ocean-based Carbon Dioxide Removal and Sequestration* (The National Academies Press, 2021).
72. Adame, M. F. et al. All tidal wetlands are blue carbon ecosystems. *BioScience* **74**, 253–268 (2024).
73. James, K., Macreadie, P. I., Burdett, H. L., Davies, I. & Kamenos, N. A. It's time to broaden what we consider a 'blue carbon ecosystem'. *Glob. Change Biol.* **30**, e17261 (2024).
74. Duarte, C. M., Losada, I. J., Hendriks, I. E., Mazarrasa, I. & Marbà, N. The role of coastal plant communities for climate change mitigation and adaptation. *Nat. Clim. Change* **3**, 961–968 (2013).
75. Burdige, D. J. (Princeton University Press, 2007).
76. Bianchi, T. S. *Biogeochemistry of Estuaries* (Oxford University Press, 2007).
77. Hopkinson, C., Weston, N. & Cai, W.-J. in *Climate Change and Estuaries* (eds M. J. Kennish, H. W. Paerl, & J. R. Crosswell) 183–201 (CRC Press, 2023).
78. Barrón, C. & Duarte, C. M. Dissolved organic carbon pools and export from the coastal ocean. *Glob. Biogeochem. Cycles* **29**, 1725–1738 (2015).
79. Cai, W.-J. in *Climate Change and Estuaries* (eds M. J. Kennish, H. W. Paerl, & J. R. Crosswell) 171–182 (CRC Press, 2023).
80. Reithmaier, G. M. S. et al. Carbonate chemistry and carbon sequestration driven by inorganic carbon outwelling from mangroves and saltmarshes. *Nat. Commun.* **14**, 8196 (2023).
81. Su, J. et al. Chesapeake Bay acidification buffered by spatially decoupled carbonate mineral cycling. *Nat. Geosci.* **13**, 441–447 (2020).
82. Kristensen, E. & Alongi, D. M. Control by fiddler crabs (*Uca vocans*) and plant roots (*Avicennia marina*) on carbon, iron, and sulfur biogeochemistry in mangrove sediment. *Limnol. Oceanogr.* **51**, 1557–1571 (2006).
83. Taillardat, P. et al. Assessing the contribution of porewater discharge in carbon export and CO₂ evasion in a mangrove tidal creek (Can Gio, Vietnam). *J. Hydrol.* **563**, 303–318 (2018).
84. Reithmaier, G. M. S. et al. Rainfall drives rapid shifts in carbon and nutrient source-sink dynamics of an urbanised, mangrove-fringed estuary. *Estuar. Coast. Shelf Sci.* **249**, 107064 (2021).
85. Reithmaier, G. M. S. et al. Alkalinity production coupled to pyrite formation represents an unaccounted blue carbon sink. *Glob. Biogeochem. Cycles* **35**, e2020GB006785 (2021).
86. Ricart, A. M. et al. Seagrass-driven changes in carbonate chemistry enhance oyster shell growth. *Oecologia* **196**, 565–576 (2021).
87. Camp, E. F. et al. mangrove and seagrass beds provide different biogeochemical services for corals threatened by climate change. *Front. Mar. Sci.* **3** <https://doi.org/10.3389/fmars.2016.00052> (2016).
88. Adame, M. F. et al. Deconstructing the mangrove carbon cycle: gains, transformation, and losses. *Ecosphere* **15**, e4806 (2024).
89. Asplund, M. E. et al. Dynamics and fate of blue carbon in a mangrove-seagrass seascape: influence of landscape configuration and land-use change. *Landsc. Ecol.* **36**, 1489–1509 (2021).
90. McHenry, J. et al. Geographic variation in organic carbon storage by seagrass beds. *Limnol. Oceanogr.* **68**, 1256–1268 (2023).
91. Ricart, A. M. et al. High variability of Blue Carbon storage in seagrass meadows at the estuary scale. *Sci. Rep.* **10**, 5865 (2020).
92. Ward, M. A. et al. Blue carbon stocks and exchanges along the California coast. *Biogeosciences* **18**, 4717–4732 (2021).
93. Mumby, P. J. & Steneck, R. S. Coral reef management and conservation in light of rapidly evolving ecological paradigms. *Trends Ecol. Evol.* **23**, 555–563 (2008).
94. Francis, F. T. & Cote, I. M. Fish movement drives spatial and temporal patterns of nutrient provisioning on coral reef patches. *Ecosphere* **9**. <https://doi.org/10.1002/ecs2.2225> (2018).
95. Sievers, M. et al. Greater consideration of animals will enhance coastal restoration outcomes. *BioScience* **72**, 1088–1098 (2022).
96. Mumby, P. J. et al. Mangroves enhance the biomass of coral reef fish communities in the Caribbean. *Nature* **427**, 533–536 (2004).
97. Bugnot, A. B. et al. Below-ground ecosystem engineers enhance biodiversity and function in a polluted ecosystem. *J. Appl. Ecol.* **59**, 2094–2105 (2022).
98. Hoey, A. S. & Bellwood, D. R. Limited functional redundancy in a high diversity system: single species dominates key ecological process on coral reefs. *Ecosystems* **12**, 1316–1328 (2009).
99. McAfee, D., Larkin, C., Connell, S. D. & Coleman, M. Multi-species restoration accelerates recovery of extinguished oyster reefs. *J. Appl. Ecol.* **58**, 286–294 (2021).

100. Yates, K. K. et al. Diverse coral communities in mangrove habitats suggest a novel refuge from climate change. *Biogeosciences* **11**, 4321–4337 (2014).
101. Kellogg, C. A., Moyer, R. P., Jacobsen, M. & Yates, K. Identifying mangrove-coral habitats in the Florida Keys. *PeerJ* **8**, e9776 (2020).
102. McKee, K. L., Rooth, J. E. & Feller, I. C. Mangrove recruitment after forest disturbance is facilitated by herbaceous species in the caribbean. *Ecol. Appl.* **17**, 1678–1693 (2007).
103. Kelleway, J. J. et al. Review of the ecosystem service implications of mangrove encroachment into salt marshes. *Glob. Change Biol.* **23**, 3967–3983 (2017).
104. Saintilan, N., Wilson, N. C., Rogers, K., Rajkaran, A. & Krauss, K. W. Mangrove expansion and salt marsh decline at mangrove poleward limits. *Glob. Change Biol.* **20**, 147–157 (2014).
105. Venera-Ponton, D. E., Diaz-Pulido, G., McCook, L. J. & Rangel-Campo, A. Macroalgae reduce growth of juvenile corals but protect them from parrotfish damage. *Mar. Ecol. Prog. Ser.* **421**, 109–115 (2011).
106. Angelini, C., Altieri, A. H., Silliman, B. R. & Bertness, M. D. Interactions among foundation species and their consequences for community organization, biodiversity, and conservation. *BioScience* **61**, 782–789 (2011).
107. Martin, T. S. H. et al. Habitat proximity exerts opposing effects on key ecological functions. *Landsc. Ecol.* **33**, 1273–1286 (2018).
108. New South Wales Environmental Planning and Assessment Act, Australia, 1979.
109. Queensland Fisheries Act, Australia (1994).
110. Mississippi-Alabama Sea Grant Legal Program, Inventory of Shellfish Restoration Permitting and Programs in the Coastal States (Troy University, National Sea Grant Law Center, 2014).
111. Shumway, N. et al. Policy solutions to facilitate restoration in coastal marine environments. *Mar. Policy* **134**. <https://doi.org/10.1016/j.marpol.2021.104789> (2021).
112. Boström, C., Pittman, S. J., Simenstad, C. & Kneib, R. T. Seascape ecology of coastal biogenic habitats: advances, gaps, and challenges. *Mar. Ecol. Prog. Ser.* **427**, 191–217 (2011).
113. Wedding, L. M., Lepczyk, C. A., Pittman, S. J., Friedlander, A. M. & Jorgensen, S. Quantifying seascape structure: extending terrestrial spatial pattern metrics to the marine realm. *Mar. Ecol. Prog. Ser.* **427**, 219–232 (2011).
114. Adame, M. F., Neil, D., Wright, S. F. & Lovelock, C. E. Sedimentation within and among mangrove forests along a gradient of geomorphological settings. *Estuar., Coast. Shelf Sci.* **86**, 21–30 (2010).
115. Howie, A. H., Reeves, S. E., Gillies, C. L. & Bishop, M. J. Integration of social data into restoration suitability modelling for oyster reefs. *Ecol. Indic.* **158**, 111531 (2024).
116. Ferretto, G., Vergés, A., Poore, A. G. B., Glasby, T. M. & Griffin, K. J. Habitat provision and erosion are influenced by seagrass meadow complexity: a seascape perspective. *Diversity* **15**, 125 (2023).
117. Jones, G. P., Russ, G. R., Sale, P. F. & Steneck, R. S. Theme section on “Larval connectivity, resilience and the future of coral reefs”. *Coral Reefs* **28**, 303–305 (2009).
118. Hata, T. et al. Coral larvae are poor swimmers and require fine-scale reef structure to settle. *Sci. Rep.* **7**, 2249 (2017).
119. Reis, A., Rovai, A. S., Lana, P. D. C. & Barros, F. Mangrove interaction with saltmarsh varies at different life stages. *Sci. Total Environ.* **905**, 167410 (2023).
120. Cook-Patton, S. C. et al. Protect, manage and then restore lands for climate mitigation. *Nat. Clim. Change* **11**, 1027–1034 (2021).
121. Saunders, M. I. et al. A roadmap to coastal and marine ecological restoration in Australia. *Environ. Sci. Policy* **159**, 103808 (2024).
122. Rhodes, J. R. et al. Flexible conservation decisions for climate adaptation. *One Earth* **5**, 622–634 (2022).
123. Paul, M. & Amos, C. L. Spatial and seasonal variation in wave attenuation over *Zostera noltii*. *J. Geophys. Res. Oceans* **116**. <https://doi.org/10.1029/2010JC006797> (2011).
124. Milbrandt, E. C., Thompson, M., Coen, L. D., Grizzle, R. E. & Ward, K. A multiple habitat restoration strategy in a semi-enclosed Florida embayment, combining hydrologic restoration, mangrove propagule plantings and oyster substrate additions. *Ecol. Eng.* **83**, 394–404 (2015).
125. Iram, N. et al. Climate change mitigation and improvement of water quality from the restoration of a subtropical coastal wetland. *Ecol. Appl.* e2620 <https://doi.org/10.1002/eap.2620> (2022).
126. Steinfurth, R. *Applying Multiple Marine Restoration Approaches and Quantifying Ecosystem Services in Danish Estuaries* (Ph.D. thesis) (University of Southern Denmark, 2023).
127. Shelamoff, V. et al. Ecosystem engineering by a canopy-forming kelp facilitates the recruitment of native oysters. *Restor. Ecol.* **27**, 1442–1451 (2019).
128. Sharma, S. et al. Do restored oyster reefs benefit seagrasses? An experimental study in the Northern Gulf of Mexico. *Restor. Ecol.* **24**, 306–313 (2016).
129. Howie, A. H. et al. *Shellfish Reef Restoration Suitability Modelling: Technical Manual* (The Nature Conservancy, Arlington, 2023).
130. The Nature Conservancy, Australia Program, Rebuilding Australia’s Lost Shellfish Reefs: 2021 Reef Builder Annual Report (2021).
131. Saunders, M. I. et al. Simple rules can guide whether land- or ocean-based conservation will best benefit marine ecosystems. *PLoS Biol.* **15**, e2001886 (2017).
132. United Nations Development Group, *Theory of Change UNDAF Companion Guidance* (2017).
133. Project Restore, Sydney Institute of Marine Science. *Project Restore*, <https://projectrestore.sims.org.au/> (2024).
134. Oosterlee, L., Cox, T. J. S., Temmerman, S. & Meire, P. Effects of tidal re-introduction design on sedimentation rates in previously embanked tidal marshes. *Estuar., Coast. Shelf Sci.* **244**, 106428 (2020).

Acknowledgements

This work was supported by a CSIRO Julius Career Award awarded to M.I.S., M.L.V., and Z.X. were supported by CSIRO R+CERC Fellowships. M.S. was funded by Australian Research Council (ARC) Discovery Early Career Researcher Award no. DE220100079. C.D. was supported by a CSIRO Julius Career Award. Figure 1 was illustrated by Stacey McCormack (Visual Knowledge Pty Ltd). We thank Chris Brown for helpful discussions during the conceptualisation phase and for comments on an early version of this manuscript. We thank three anonymous reviewers for their helpful comments, which greatly improved the manuscript.

Author contributions

M.L.V. and M.I.S. conceptualised the manuscript. M.L.V. led the writing. M.L.V., C.A.B., M.S., M.F.A., P.B., M.B., J.R.C., C.D., B.L., F.M.B., S.R., V.R., A.S., Z.X., K.D. and M.I.S. drafted sections of the manuscript. M.L.V., C.A.B., M.S., M.F.A., P.B., M.B., J.R.C., C.D., B.L., F.M.B., S.R., V.R., A.S., Z.X., B.S., R.t.H., M.v.K. and M.I.S. edited and reviewed the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s44183-024-00095-1>.

Correspondence and requests for materials should be addressed to Maria L. Vozzo.

Reprints and permissions information is available at
<http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© Crown 2024