

## Peroxygenase-Catalysed Epoxidation of Styrene Derivatives in Neat Reaction Media

Rauch, Marine C.R.; Tieves, Florian; Paul, Caroline E.; Arends, Isabel W.C.E.; Alcalde, Miguel; Hollmann, Frank

**DOI**

[10.1002/cctc.201901142](https://doi.org/10.1002/cctc.201901142)

**Publication date**

2019

**Document Version**

Final published version

**Published in**

ChemCatChem

**Citation (APA)**

Rauch, M. C. R., Tieves, F., Paul, C. E., Arends, I. W. C. E., Alcalde, M., & Hollmann, F. (2019). Peroxygenase-Catalysed Epoxidation of Styrene Derivatives in Neat Reaction Media. *ChemCatChem*, 11(18), 4519-4523. <https://doi.org/10.1002/cctc.201901142>

**Important note**

To cite this publication, please use the final published version (if applicable). Please check the document version above.

**Copyright**

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

**Takedown policy**

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

# Peroxygenase-Catalysed Epoxidation of Styrene Derivatives in Neat Reaction Media

Marine C. R. Rauch,<sup>[a]</sup> Florian Tieves,<sup>[a]</sup> Caroline E. Paul,<sup>[a]</sup> Isabel W. C. E. Arends,<sup>[b]</sup> Miguel Alcalde,<sup>[c]</sup> and Frank Hollmann<sup>\*[a]</sup>


Biocatalytic oxyfunctionalisation reactions are traditionally conducted in aqueous media limiting their production yield. Here we report the application of a peroxygenase in neat reaction conditions reaching product concentrations of up to 360 mM.

Epoxides are important building blocks in organic synthesis. The ring opening of epoxides leads to useful  $\alpha$ - or  $\beta$ -substituted alcohols.<sup>[1]</sup> As a result, a broad range of catalytic methods for the epoxidation of C=C-double bonds have been established.<sup>[2]</sup> Compared to this variety, only few biocatalytic methods are known. The chemoenzymatic epoxidation of alkenes using lipase-borne peracids for example is receiving tremendous interest but yields racemic products.<sup>[3]</sup> Amongst the stereospecific epoxidation methods the use of flavin-dependent styrene monooxygenases<sup>[4]</sup> and P450 monooxygenases<sup>[5]</sup> are most prominent.


The latter approaches rely on reductive activation of molecular oxygen using reduced nicotinamide cofactors (NAD(P)H) as source of reducing equivalents (Scheme 1). This not only implies complicated and vulnerable electron transport chains but also, due to the exclusive water-solubility of the cofactors, largely limits these processes to aqueous reaction media.

The majority of the alkenes of interest are however hydrophobic, limiting the final product titres to the lower millimolar range. This is unacceptable from an economic and an environmental point-of-view. Current solutions focus around two-liquid-phase-system approaches (2LPS).<sup>[6]</sup>


## a) Monooxygenase-catalysed epoxidation



## b) Lipase-initiated Prilezhaev reaction



## c) Peroxygenase-catalysed epoxidation



Scheme 1. Comparison of biocatalytic epoxidation reactions.

Ideally, (bio) catalytic epoxidation reactions should occur in organic media (even neat) to enable high product concentrations. In this respect, peroxygenases represent a promising solution.<sup>[7]</sup> Peroxygenases are heme-thiolate enzymes enabling P450 monooxygenase-like oxyfunctionalisation reactions. In contrast to monooxygenases, peroxygenases do not rely on (water-soluble) redox partners but on (organic) peroxides, enabling their potential application in non-aqueous media. Pioneering works by Pu, Wang and Zhang<sup>[8]</sup> and Hofrichter<sup>[9]</sup> have established peroxygenase-catalysed epoxidation reactions using hydrogen peroxide or organic hydroperoxides as oxidants, albeit in aqueous reaction media thereby limiting the reagent concentration to the lower millimolar range.

Klibanov and co-workers reported peroxidase-reactions under non-aqueous conditions.<sup>[10]</sup> Unfortunately, these contributions have not yet found widespread attention in the biocatalysis community.

[a] M. C. R. Rauch, Dr. F. Tieves, Dr. C. E. Paul, Prof. F. Hollmann  
Department of Biotechnology  
Delft University of Technology  
Delft 2629HZ (The Netherlands)  
E-mail: f.hollmann@tudelft.nl

[b] Prof. I. W. C. E. Arends  
Faculty of Science  
Utrecht University  
Utrecht 3584CD (The Netherlands)

[c] Prof. M. Alcalde  
Department of Biocatalysis  
Institute of Catalysis, CSIC  
28049 Madrid (Spain)

Supporting information for this article is available on the WWW under <https://doi.org/10.1002/cctc.201901142>

This publication is part of a joint Special Collection with ChemBioChem on "Excellence in Biocatalysis Research". Please follow the link for more articles in the collection.

We therefore set out to establish peroxygenase-catalysed, selective oxyfunctionalisation reactions in neat reaction media. As model peroxygenase we chose an evolved recombinant peroxygenase from *Agroclybe aegerita* (*rAaeUPO*)<sup>[11]</sup> to catalyse the epoxidation of styrene and its derivatives.<sup>[9]</sup>

As oxidant, we chose *tert*-butyl hydroperoxide (<sup>t</sup>BuOOH) due to its high solubility in hydrophobic media.


To employ *rAaeUPO* in neat reaction media, we first immobilised it covalently on an epoxide-modified polyacrylic matrix (Immobead IB-COV-1). Under non-optimised conditions, 72.8% of the enzyme was immobilised (for further details see SI section 4). The remaining catalytic activity, however was only 3% (Figure S1). Further development will have to focus on optimised immobilisation procedures yielding higher activity.

Nevertheless, having the immobilised enzyme in hand, we first explored its substrate scope (Table 1). Pleasingly, all of the styrene derivatives tested were converted with satisfactory to excellent turnover numbers for the biocatalyst. In accordance with previous reports<sup>[9,12]</sup> wild-type *rAaeUPO* converted the majority of styrenes non-stereoselectively giving (near racemic) epoxides, one notable exception being *cis*- $\beta$ -methylstyrene, which was converted highly stereoselectively into (*1R,2S*)-*cis*- $\beta$ -methylstyrene oxide. It is also interesting to note that in some cases, the desired epoxide was not stable and spontaneously rearranged into the corresponding carbonyl compound (for further details see SI section 5.3).

Nevertheless, very significant product concentrations of up to 100 mM were achieved. The catalytic performance of *rAaeUPO* in terms of turnover numbers (TON = amount of product divided by the amount of enzyme, [mol  $\times$  mol<sup>-1</sup>]) was excellent.

To identify parameters influencing the productivity of the reaction, we systematically varied the biocatalyst loading and the <sup>t</sup>BuOOH feeding rate in the epoxidation of *cis*- $\beta$ -methylstyrene (Figure 1). The initial rate of the epoxidation reaction correlated directly with the dosing rate of <sup>t</sup>BuOOH. This, however did not necessarily translate in higher product titres. Most probably, increasing feed rates of the oxidant also increased the undesired oxidative inactivation of the enzyme's active site.<sup>[13]</sup> This is supported by the observation that the robustness of the reactions (i.e. the duration of product accumulation) inversely correlated with the <sup>t</sup>BuOOH feed rate (Table S2).

As mentioned above, epoxides are versatile building blocks for the synthesis of a broad range of products. Amino alcohols, for example, are common structural motifs in many pharmaceutically active ingredients.<sup>[14]</sup> We therefore envisioned a chemoenzymatic cascade reaction comprising the *rAaeUPO*-catalysed, stereoselective epoxidation of *cis*- $\beta$ -methylstyrene




**Figure 1.** Characterisation of <sup>t</sup>BuOOH feeding rate and enzyme concentration comparing the initial reaction rates (black diamonds) and final product concentrations (grey squares). General conditions: room temperature, 20 rpm in overhead rotator. Data presented are an average of duplicates and corrected from potential substrate evaporation (see the Supporting Information, Section 5.1)

followed by the chemical oxirane-opening with methyl amine yielding (pseudo)ephedrine (Scheme 2).

The epoxidation reaction was performed on a 10 mL scale with gradual <sup>t</sup>BuOOH feed (Figure 2). Although a conservative <sup>t</sup>BuOOH feed rate of 5 mMh<sup>-1</sup> was applied, inactivation of the biocatalyst represented a major challenge for the reaction, necessitating further provision of the reaction with fresh enzyme (indicated by arrows in Figure 2). It is also interesting to note that in contrast to previous experiments using *rAaeUPO* in aqueous reaction media using H<sub>2</sub>O<sub>2</sub>,<sup>[15]</sup> the peroxide utilisation efficiency was only approximately 50%. It will be interesting to further investigate this increased catalase activity of *rAaeUPO*.

From this experiment, 360 mM of (*1R,2S*)- $\beta$ -methylstyrene oxide were obtained. The turnover number of the enzyme was more than 8500. Next to the desired product, the reaction mixture also contained some benzaldehyde, originating from *rAaeUPO*-catalysed C=C-bond cleavage.<sup>[16]</sup> To avoid negative influences of this by-product, the desired product was purified chromatographically and subjected to chemical ring-opening with methylamine resulting in pseudoephedrine (58.2%), ephedrine (7%) and isoephedrine (34.8%) (Scheme S2).

Overall, with this contribution we have demonstrated that peroxygenase-catalysed epoxidations can be performed under neat reaction conditions. This opens up new possibilities for the



**Scheme 2.** Envisioned chemoenzymatic cascade to obtain (pseudo)ephedrine from *cis*- $\beta$ -methylstyrene.

**Table 1.** Substrate scope of the epoxidation of styrene derivatives with immobilised *rAaeUPO*. Data are an average of duplicates and corrected from potential substrate evaporation.<sup>[a]</sup>

1-18a  $\xrightarrow[tBuOOH]{rAaeUPO}$  1-18b  $\xrightarrow{\text{isomerisation}}$  Carbonyl product


1 R<sub>1</sub> = R<sub>2</sub> = R<sub>3</sub> = R<sub>4</sub> = H  
 2 R<sub>1</sub> = CH<sub>3</sub>, R<sub>2</sub> = R<sub>3</sub> = R<sub>4</sub> = H  
 3 R<sub>1</sub> = H, R<sub>2</sub> = CH<sub>3</sub>, R<sub>3</sub> = H, R<sub>4</sub> = H  
 4 R<sub>1</sub> = H, R<sub>2</sub> = H, R<sub>3</sub> = CH<sub>3</sub>, R<sub>4</sub> = H  
 5 R<sub>1</sub> = R<sub>2</sub> = R<sub>3</sub> = H, R<sub>4</sub> = *p*-Br  
 6 R<sub>1</sub> = R<sub>2</sub> = R<sub>3</sub> = H, R<sub>4</sub> = *m*-Br

7 R<sub>1</sub> = R<sub>2</sub> = R<sub>3</sub> = H, R<sub>4</sub> = *o*-Br  
 8 R<sub>1</sub> = R<sub>2</sub> = R<sub>3</sub> = H, R<sub>4</sub> = *p*-Cl  
 9 R<sub>1</sub> = R<sub>2</sub> = R<sub>3</sub> = H, R<sub>4</sub> = *m*-Cl  
 10 R<sub>1</sub> = R<sub>2</sub> = R<sub>3</sub> = H, R<sub>4</sub> = *o*-Cl  
 11 R<sub>1</sub> = R<sub>2</sub> = R<sub>3</sub> = H, R<sub>4</sub> = *p*-F  
 12 R<sub>1</sub> = R<sub>2</sub> = R<sub>3</sub> = H, R<sub>4</sub> = *m*-F

13 R<sub>1</sub> = R<sub>2</sub> = R<sub>3</sub> = H, R<sub>4</sub> = *p*-CH<sub>3</sub>  
 14 R<sub>1</sub> = R<sub>2</sub> = R<sub>3</sub> = H, R<sub>4</sub> = *m*-CH<sub>3</sub>  
 15 R<sub>1</sub> = CH<sub>3</sub>, R<sub>2</sub> = H, R<sub>3</sub> = H, R<sub>4</sub> = *p*-CH<sub>3</sub>  
 16 R<sub>1</sub> = R<sub>2</sub> = R<sub>3</sub> = H, R<sub>4</sub> = *p*-OCH<sub>3</sub>  
 17 1,2-dihydronaphthalene  
 18 indene

Substrate	Epoxy product [mM]	ee [%]	Carbonyl product <sup>[b]</sup> [mM(%)]	Time [h]	TON <sup>[c]</sup>
	16	12	2 (12)	42	3203
	80	30	47 (37)	86	22598
	9	35	1 (11)	86	1779
	36	> 99	2 (4)	21	6762
	59	9	21 (26)	60	14235
	24	50	102 (81)	62	22420
	10	39	90 (90)	49	17794
	8	12	22 (73)	62	5338
	16	17	12 (43)	42	4982
	15	15	22 (59)	22	6584
	11	15	25 (69)	42	6406
	3	42	1 (23)	42	712
	14	– <sup>[d]</sup>	60 (81)	62	13167
	36	28	59 (62)	97	16904
	4	6	25 (86)	49	5160
	none	–	–	–	0
	136	42	147 (52)	108	50356
	none	–	33 (100)	69	5872

[a] Reaction conditions: [rAaeUPO] = 5.62 μM, tBuOOH dosing rate = 5 mM/h, room temperature, 20 rpm in overhead rotor. [b] The concentrations of carbonyl product were calculated using the calibration curves of the epoxides. Carbonyl products are aldehyde or ketone in β position from the ring opening of the epoxides, [c] TON = [product]/[enzyme], [d] n.d. = not determined.



**Figure 2.** Time-course of the *rAaeUPO*-catalysed epoxidation of *cis*- $\beta$ -methylstyrene. Conditions: 10 mL *cis*- $\beta$ -methylstyrene, [*rAaeUPO*@IB-COV-1]<sub>total</sub> = 41.8  $\mu$ M (added in equal portions at the times indicated by the arrows), room temperature. <sup>t</sup>BuOOH was added continuously according to the feed profile shown as a dashed grey line.

preparative scale-application of this promising class of enzymes. Product concentrations of up to 360 mM have been achieved representing one of the highest product titres obtained with oxidoreductase catalysis<sup>[17]</sup> and certainly the highest product concentration with isolated enzymes.<sup>[6a,b,18]</sup>

Nevertheless, some issues remain to be solved *en route* to a truly preparatively useful system. First and foremost, more active immobilisates of *rAaeUPO* need to be found. The activity recovery of the peroxygenase needs to be improved to obtain more active catalysts.<sup>[19]</sup> We are confident that from the wealth of immobilisation methods available today,<sup>[20]</sup> a suitable method will be found in our ongoing research. Also, more enantioselective *rAaeUPO* versions are highly desirable to broaden the scope of the reaction.

## Experimental Section

**Enzyme preparation.** Recombinant expression and purification of the evolved unspecific peroxygenase mutant from *A. aegerita* in *P. pastoris* was performed following a previously described procedure.<sup>[9]</sup>

**Immobilisation protocol.** Immobeads (IB-COV-1) from ChiralVision was used to immobilise *rAaeUPO*. The beads were washed before usage and then mixed with *rAaeUPO*. Immobilisation was carried out for 5 hours using overhead rotator. After 5 hours, the immobilisation mixture was stored at 6 °C for 12 hours without stirring or shaking. After overnight incubation, the supernatant was removed, and the beads 3 times washed. The washing fractions were pooled. The peroxygenase concentration was determined *via* CO difference spectra in the supernatant and the washing fraction to calculate the amount of immobilised peroxygenase. A detailed description of the immobilisation of the enzymes is available in the Supporting Information.

**Enzymatic reaction conditions.** Reactions were performed in GC vial of 1.5 mL at room temperature. Immobilised *rAaeUPO* was first weighed in the vial according to the concentration of enzyme wanted, then pure substrate was added to the vial. Before each samples were taken, the vial was weighed in order to estimate the loss of substrate by evaporation. <sup>t</sup>BuOOH was added in the vial *via* a tube connected to a syringe pump. An overhead rotator from neoLab was mixing the reactions at 20 rpm. At intervals, aliquots were withdrawn, extracted with ethyl acetate, dried over MgSO<sub>4</sub> and analysed by chiral gas chromatography. Details of gas chromatography and temperature profiles are shown in Supporting Information.

**Chemical ring opening.** 10 mg of pure epoxide were diluted in 200  $\mu$ L of MeNH<sub>2</sub> (40% in water). The reaction was mixed during 20 hours at 60 °C. The reaction was then extracted with dichloromethane and analysed on NMR.

## Acknowledgements

Financial support by the European Research Council (ERC Consolidator Grant No. 648026) is gratefully acknowledged.

## Conflict of Interest

The authors declare no conflict of interest.

**Keywords:** Biocatalysis · Epoxidation · Peroxygenases · Neat reaction conditions · Cascade reactions


- [1] M. Pineschi, *Eur. J. Org. Chem.* **2006**, *2006*, 4979–4988.
- [2] a) E. M. McGarrigle, D. G. Gilheany, *Chem. Rev.* **2005**, *105*, 1563–1602; b) R. L. Davis, J. Stillner, T. Naicker, H. Jiang, K. A. Jørgensen, *Angew. Chem.* **2014**, *53*, 7406–7426, *Angew. Chem. Int. Ed.* **2014**, *53*, 7406–7426.
- [3] a) B. O. O. Burek, S. Bormann, F. Hollmann, J. Bloh, D. Holtmann, *Green Chem.* **2019**, *21*, 3232–3249; b) J. Dong, E. Fernández-Fueyo, F. Hollmann, C. Paul, M. Pesic, S. Schmidt, Y. Wang, S. Younes, W. Zhang, *Angew. Chem.*, **2018**, *130*, 9380–9404, *Angew. Chem. Int. Ed.* **2018**, *57*, 9238–9261.
- [4] a) E. Romero, J. R. Gómez Castellanos, G. Gadda, M. W. Fraaije, A. Mattevi, *Chem. Rev.* **2018**, *118*, 1742–1769; b) T. Heine, W. van Berkel, G. Gassner, K.-H. van Pée, D. Tischler, *Biology* **2018**, *7*, 42; c) G. Gygli, W. J. H. van Berkel, *Curr. Biotechnol.* **2015**, *4*, 100–110; d) M. M. E. Huijbers, S. Montersino, A. H. Westphal, D. Tischler, W. J. H. van Berkel, *Arch. Biochem. Biophys.* **2014**, *544*, 2–17.
- [5] a) V. B. Urlacher, M. Girhard, *Trends Biotechnol.* **2019**; b) J. B. Wang, M. T. Reetz, *Nat. Chem.* **2015**, *7*, 948–949.
- [6] a) H. Toda, R. Imae, N. Itoh, *Adv. Synth. Catal.* **2014**, *356*, 3443–3450; b) H. Toda, R. Imae, N. Itoh, *Tetrahedron: Asymmetry* **2012**, *23*, 1542–1549; c) J. B. Park, B. Buehler, T. Habicher, B. Hauer, S. Panke, B. Witholt, A. Schmid, *Biotechnol. Bioeng.* **2006**, *95*, 501–512.
- [7] a) M. Hofrichter, R. Ullrich, *Curr. Opin. Chem. Biol.* **2014**, *19*, 116–125; b) Y. Wang, D. Lan, R. Durrani, F. Hollmann, *Curr. Opin. Chem. Biol.* **2017**, *37*, 1–9.
- [8] a) L. Wang, S. Wei, X. Pan, P. Liu, X. Du, C. Zhang, L. Pu, Q. Wang, *Chem. Eur. J.* **2018**, *24*, 2741–2749; b) C. Zhang, P.-X. Liu, L.-Y. Huang, S.-P. Wei, L. Wang, S.-Y. Yang, X.-Q. Yu, L. Pu, Q. Wang, *Chem. Eur. J.* **2016**, *22*, 10969–10975.
- [9] M. Kluge, R. Ullrich, K. Scheibner, M. Hofrichter, *Green Chem.* **2012**, *14*, 440–446.
- [10] a) J. S. Dordick, M. A. Marletta, A. M. Klibanov, *Proc. Natl. Acad. Sci. USA* **1986**, *83*, 6255–6257; b) A. Zaks, A. M. Klibanov, *Science* **1984**, *224*, 1249–1251.

- [11] a) P. Molina-Espeja, S. Ma, D. M. Mate, R. Ludwig, M. Alcalde, *Enz. Microb. Technol.* **2015**, 73–74, 29–33; b) P. Molina-Espeja, E. Garcia-Ruiz, D. Gonzalez-Perez, R. Ullrich, M. Hofrichter, M. Alcalde, *Appl. Environ. Microbiol.* **2014**, 80, 3496–3507.
- [12] E. Churakova, M. Kluge, R. Ullrich, I. Arends, M. Hofrichter, F. Hollmann, *Angew. Chem. Int. Ed.* **2011**, 50, 10716–10719; *Angew. Chem.* **2011**, 123, 10904–10907.
- [13] B. Valderrama, M. Ayala, R. Vazquez-Duhalt, *Chem. Biol.* **2002**, 9, 555–565.
- [14] G. Grogan, *Curr. Opin. Chem. Biol.* **2018**, 43, 15–22.
- [15] a) F. Tieves, S. J.-P. Willot, M. M. C. H. van Schie, M. C. R. Rauch, S. H. H. Younes, W. Zhang, P. G. de Santos, J. M. Robbins, B. Bommarius, M. Alcalde, A. Bommarius, F. Hollmann, *Angew. Chem.* **2019**, 131, 7955–7959, *Angew. Chem. Int. Ed.* **2019**, 58, 7873–7877; b) Y. Ni, E. Fernández-Fueyo, A. G. Baraibar, R. Ullrich, M. Hofrichter, H. Yanase, M. Alcalde, W. J. H. van Berkel, F. Hollmann, *Angew. Chem.*, **2016**, 128, 809–812, *Angew. Chem. Int. Ed.* **2016**, 55, 798–801.
- [16] F. G. Mutti, M. Lara, M. Kroutil, W. Kroutil, *Chem. Eur. J.* **2010**, 16, 14142–14148.
- [17] a) B. Bühler, I. Bollhalder, B. Hauer, B. Witholt, A. Schmid, *Biotechnol. Bioeng.* **2003**, 82, 833–842; b) B. Bühler, I. Bollhalder, B. Hauer, B. Witholt, A. Schmid, *Biotechnol. Bioeng.* **2003**, 81, 683–694; c) A. Schmid, K. Hofstetter, H.-J. Feiten, F. Hollmann, B. Witholt, *Adv. Synth. Catal.* **2001**, 343, 732–737.
- [18] a) K. Hofstetter, J. Lutz, I. Lang, B. Witholt, A. Schmid, *Angew. Chem. Int. Ed.* **2004**, 43, 2163–2166; *Angew. Chem.* **2004**, 116, 2215–2218; b) E. Fernández-Fueyo, Y. Ni, A. Gomez Baraibar, M. Alcalde, L. M. van Langen, F. Hollmann, *J. Mol. Catal. B. Enzym.* **2016**, 134.
- [19] P. Molina-Espeja, P. Santos-Moriano, E. García-Ruiz, A. Ballesteros, F. J. Plou, M. Alcalde, *Int. J. Mol. Sci.* **2019**, 20, 1627.
- [20] U. Hanefeld, L. Gardossi, E. Magner, *Chem. Soc. Rev.* **2009**, 38, 453–468.

---

Manuscript received: June 27, 2019  
 Revised manuscript received: July 18, 2019  
 Accepted manuscript online: July 19, 2019  
 Version of record online: ■■■, ■■■■

## COMMUNICATIONS



**Very neat!** Peroxygenases catalyse epoxidation of C=C-double bonds under non-aqueous conditions.

*M. C. R. Rauch, Dr. F. Tieves, Dr. C. E. Paul, Prof. I. W. C. E. Arends, Prof. M. Alcalde, Prof. F. Hollmann\**

1 – 6

**Peroxygenase-Catalysed Epoxidation of Styrene Derivatives in Neat Reaction Media**



Open Access