

Molecular Catalysis for the Chemistry of the future a perspective

Han, Jeong Woo; Hollmann, Frank; Luque, Rafael; Song, In Kyu; Talarico, Giovanni; Tatsumi, Takashi; Yan,

DOI

10.1016/j.mcat.2022.112233

Publication date 2022

Document Version Final published version

Published in Molecular Catalysis

Citation (APA)
Han, J. W., Hollmann, F., Luque, R., Song, I. K., Talarico, G., Tatsumi, T., & Yan, N. (2022). Molecular Catalysis for the Chemistry of the future: a perspective. *Molecular Catalysis*, 522, Article 112233. https://doi.org/10.1016/j.mcat.2022.112233

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy
Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

Green Open Access added to TU Delft Institutional Repository 'You share, we take care!' - Taverne project

https://www.openaccess.nl/en/you-share-we-take-care

Otherwise as indicated in the copyright section: the publisher is the copyright holder of this work and the author uses the Dutch legislation to make this work public.

ELSEVIER

Contents lists available at ScienceDirect

Molecular Catalysis

journal homepage: www.journals.elsevier.com/molecular-catalysis





Molecular Catalysis for the Chemistry of the future: a perspective

Jeong Woo Han^a, Frank Hollmann^b, Rafael Luque^{c,*}, In Kyu Song^d, Giovanni Talarico^e, Takashi Tatsumi^f, Ning Yan^g

- a Department of Chemical Engineering, Pohang University of Science and Technology (POSTECH), Pohang, Gyeongbuk, 37673, Republic of Korea
- ^b Department of Biotechnology, Delft University of Technology, van der Maasweg 9, 2629HZ Delft, The Netherlands
- c Department of Organic Chemistry, Universidad de Córdoba, Ctra Nnal IV-A, Km 396, 14014 Córdoba, Spain
- ^d School of Chemical and Biological Engineering, Seoul National University, Seoul 151-742, Korea
- ^e Dipartimento di Scienze Chimiche, Università di Napoli Federico II, Via Cintia, 80126 Napoli, Italy
- f Laboratory for Chemistry and Life Science, Tokyo Institute of Technology, 4259 Nagatsuta, Yokohama 226-8503, Japan
- 8 National University of Singapore, Department of Chemical and Biomolecular Engineering, 4 Engineering Drive 4, 117585 Singapore

The chemistry of the future will have to find solutions for the societal challenges of tomorrow (and today) and undoubtedly, catalysis will play a central role. Molecular catalysis for the chemistry of the future will openly face significant challenges for present and future generations including: 1) scarcity of resources (food, water, fossil resources etc.) combined with 2) an ever increasing degradation of the environment, 3) a "new life" for plastics and particularly plastic waste that is not recycled, and 4) the development of a circular economy based on the use of renewable energies limiting or ideally reversing CO₂ emissions.

All types of catalysis are profoundly linked with each other at the molecular level. However, they are mostly studied by different communities at present with little scientific exchange between them. It is important to develop unifying concepts of catalysis (thermal-, bio-, photo- and electro-catalytic pathways) by using state-of-the-art techniques from different subdisciplines. In this manner, new understandings will be generated on important design parameters in catalysis including solvent and pH effects, synergies between metals in alloy catalysts, synergy between metals and ligands and support in heterogeneous catalysts. The vision is to achieve fundamental understanding of catalytic processes culminating in a great unifying theory of catalysis.

To elucidate the catalytic mechanism at the atomic and molecular level and predict the catalytic activity and stability before trial-and-error based experiments to work out sophisticated catalyst designs, computational approaches are also essential. Over the last two decades, the computer-aided design of catalysts has been very successful owing to the prosperous establishment of computational methodologies accompanied with tremendous progress of computing power. Many examples can be found, in which computational screening has played a major role in experimentally finding a new class of molecular catalysts and

computational analyses could be successfully explained for the underlying mechanism of catalysts' behaviours. However, there are still many challenges to overcome for the use of these approaches, and continuous efforts on this research field are further necessary.

As editors of *Molecular Catalysis*, we feel in an advantageous position to directly follow (and impact) various academic trends and industrial realities to deal with such problems administrating different contributions submitted to our journal. This privileged position allowed us to visualize a number of hot areas (Figure 1) currently under development for a more sustainable future:

- a) Establishing new chemistries for safe-by-design/benign-by-design products, not only comprises synthesis methods avoiding immediate hazards for the environment such as emissions and wastes generated as well as safety issues of the production process. More importantly, it will also yield safer products. Safer in the sense that they will exhibit only the desired properties in human use and further on if exposed to the environment, i.e. bio-degradable by design. [1]
- b) Highly selective catalytic transformations. Reducing the amount of wastes formed in chemical transformations is key to reducing the environmental impact of the chemical industry. This comprises, but is not limited to: more selective transformations (avoiding undesired side products), avoidance of protection group chemistry and reduction of the energy demand of chemical transformations (reducing energy-linked CO₂ emissions). Biocatalysis will play a central role here. [2] When it comes to selectivity, enzymatic methods are unparalleled. Further exploring this catalytic wealth and designing reaction concepts compatible with

E-mail address: q62alsor@uco.es (R. Luque).

^{*} Corresponding author: Professor Rafael Luque, Quimica Orgánica, Universidad de Cordoba, Campus de Rabanales, Edificio Marie Curie (C-3) Ctra Nna..., E14014 Córdoba, Spain




Figure 1. Impact of Molecular Catalysis in the chemistries of the future

the current chemical industry will represent one of the challenges for future White Biotechnology.

- c) A more general and widespread attention to Metal Organic Frameworks (MOFs) as practical catalysts not only for the synthesis of specialty and fine chemicals but also for other relevant environmental-related applications in photocatalysis, the design of bionanoMOF hybrids and related topics. [3].
- d) Stereoselective catalysis of monomers obtained by renewable resources and biomass-derived. An interesting example is the stereoselective ring opening polymerization (ROP) of lactide. The polylactides (PLAs) found important industrial applications and their mechanical and thermal properties are strongly depending on the microstructures. [4].
- e) CO₂ utilization as a C1-feedstock by incorporating into suitable products (e.g. stereoselective CO₂/epoxide copolymerization) and/or CO₂ catalytic transformation into biosourced organic carbonates. Both aspects are combined with the primary task to develop selective, fast and efficient reduction processes of CO₂ into valuable products in a sustainable manner (e.g. carbon monoxide, formic acid, methanol, methane and so on). [5]
- f) The development of chemical recycling to monomers (CRM) where the final polymers are transformed back into monomers and the monomers are reconsidered for repolymerization without loss in properties. This process, although at an early stage, creates an ideal circular polymer economy pushing the necessary economic and environmental synergies for a circular economy. [6]. Efficient catalytic synthesis of monomers building up biodegradable polymers, ideally from renewable resources in a carbon neutral manner, is also a key target.
- g) Modification of polyolefins with polar functionalities. The controlled incorporation of polar monomers into the polyolefins backbone chain strongly enhances the polymer properties. The copolymerization of α -olefins with polar monomers catalyzed by late transition metals achieved significant breakthroughs in the academia although no commercial process currently exists for such reactions due to the lack of high-performance heterogeneous catalysts. [7]
- h) The design of single-atom based systems for advanced catalytic prospects, potentially applicable to industrial premises [8].

- i) Living cells as chemical factories with unprecedented flexibility with respect to the source of energy and material input [9]. Photoautroph organisms for example can build up complex molecules from CO₂ using visible light as energy source. To fully explore this potential, combined efforts of metabolic-, protein-and reactor engineering will be needed. Similarly, electrotroph organisms principally can utilise electrical energy to build complex products from simple carbon sources.
- j) Organocatalysis, following up the steps of the recently granted 2021 Nobel Prize will play a fundamental role, particularly enantioselective organocatalysis in the advance synthesis of drugs and pharmaceuticals (e.g. greener enantioselective halogenations) for a more sustainable future. [10].
- k) Electro- and photo-catalysis for the catalytic synthesis of the molecules of the future, with the potential to be extended to selective functionalisations (e.g. electro/photochemical upgrading), synthesis of pharmaceuticals and relevant high added value chemicals, [11, 12].
- Auxiliary promotion methods including mechanochemistry, microwaves, ultrasound, electric and magnetic fields, and electric potentials, have all shown promise in tuning the adsorbate—active size interactions, thereby increasing the reactivity. These strategies should be widely tested in promoting various types of catalytic reactions, in particular those that are notoriously difficult to occur under mild conditions. [13, 14].
- m) *The combination of Flow Chemistry and Catalysis* as a valuable tool for more scalable catalytic applications able to bring us one step closer to industrial premises [15].

The main question is which of the above lines will be predominant as the way forward in Molecular Catalysis for the Chemistry of the future? There is not an unique answer and we believe that the near future will see a flourishing demand of all identified research topics. Certainly, chemistry papers will show even more a combination of experimental and theoretical aspects with the recent input of machine learning techniques for fast catalyst screening. The combination of these key identified topics/areas (and various others) hint how chemists are preparing themselves for the challenges to come of a still undisclosed and uncertain future. At *Molecular Catalysis*, we will certainly welcome contributions providing alternatives and solutions to some of the highlighted future challenges that we hope to be enjoying for the journal in the years to come.

CREDIT AUTHOR Statement

All authors contributed to writing, revising and editing of the manuscript including conceptualization and overall shown ideas/topics.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper

References

- J.F.M. van der Waals, A. Falk, P. Fantke, P. Filippousi, R.C.H. Flipphi, D. Mottet, X. Trier (2019). Safe-by-design for materials and chemicals: Towards an innovation programme in Horizon Europe.
- [2] U. Hanefeld, F. Hollmann, C.E. Paul, Chem. Soc. Rev. 51 (2022) 594–627.
- [3] K. Shen, L. Zhang, X. Chen, L. Liu, D. Zhang, Y. Han, J. Chen, J. Long, R. Luque, Y. Li, B. Chen, Science 359 (2018) 206–210.
- [4] X. Zhang, M. Fevre, G.O. Jones, R.M. Waymouth, Chem. Rev. 118 (2018), 839885.
- [5] A. Akhundi, A. Badiei, G.M. Ziarani, A. Habibi-Yangjeh, M.J. Muñoz-Batista, R. Luque, Mol. Catal. 488 (2020), 110902.
- [6] Y. Jing, Y. Wang, S. Furukawa, J. Xia, C. Sun, M.J. Hülsey, H. Wang, Y. Guo, X. Liu, N. Yan, Angew. Chem. Int. Ed. 60 (2021) 5527–5535.
- [7] S.L.J. Luckham, K. Nozaki, Acc. Chem. Res. 54 (2021) 344-355.

- [8] R. Ciriminna, M. Gharemani, B. Karimi, M. Pagliaro, R. Luque, Current Op. Green Sust. Chem. 25 (2020), 100358.
- [9] M. Eisenstein, Nature 531 (2016) 401-403.
- [10] A. Carlone, L. Bernardi, P. McCormack, T. Warr, S. Oruganti, C.J. Cobley, Org. Process Res. Dev. 25 (2021) 2795-2805.
- [11] C. Zhu, N.W.J. Ang, T.H. Meyer, Y. Qiu, L. Ackermann, ACS Cent. Sci. 7 (2021) 415–431.
- [12] S. Song, J. Qu, P. Han, M.J. Hülsey, G. Zhang, Y. Wang, S. Wang, D. Chen, J. Lu, N. Yan, Nat. Commun. 11 (2020) 4899.
- [13] K. Kubota, Y. Pang, A. Miura, H. Ito, Science 366 (2019) 1500–1504.
- [14] M.J. Hülsey, C.W. Lim, N. Yan, Chem. Sci. 11 (2020) 1456–1468.
 [15] G. Laudadio, Y. Deng, K. Van der Wal, D. Ravelli, M. Nuño, M. Fagnoni, D. Guthrie, Y. Sun, T. Noel, Science 369 (2020) 92-96.