

## Interlinkages between human agency, water use efficiency and sustainable food production

Lyu, Haoyang; Dong, Zengchuan; Pande, Saket

**DOI**

[10.1016/j.jhydrol.2019.124524](https://doi.org/10.1016/j.jhydrol.2019.124524)

**Publication date**

2020

**Document Version**

Accepted author manuscript

**Published in**

Journal of Hydrology

**Citation (APA)**

Lyu, H., Dong, Z., & Pande, S. (2020). Interlinkages between human agency, water use efficiency and sustainable food production. *Journal of Hydrology*, 582, Article 124524.  
<https://doi.org/10.1016/j.jhydrol.2019.124524>

**Important note**

To cite this publication, please use the final published version (if applicable).  
Please check the document version above.

**Copyright**

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

**Takedown policy**

Please contact us and provide details if you believe this document breaches copyrights.  
We will remove access to the work immediately and investigate your claim.

1 **Interlinkages between human agency, water use efficiency and sustainable food**  
2 **production**

3

4 Haoyang Lyu<sup>(1,2)</sup>; Zengchuan Dong<sup>(2)</sup> and Saket Pande<sup>(1)</sup>

5 (1) Delft University of Technology, Delft, Netherlands; (2) Hohai University, Nanjing, China

6

7

8 **Abstract:**

9 Efficient use of water and nutrients in crop production are critical for sustainable  
10 water and crop production systems. Understanding the role of humans in ensuring  
11 water and nutrient use efficiency is therefore an important ingredient of sustainable  
12 development. Crop production functions are often defined either as functions of water  
13 and nutrient deficiency or are based on economic production theory that  
14 conceptualizes production as a result of economic activities that take in inputs such as  
15 water, capital and labor and produce crop biomass as output. This paper fills a gap by  
16 consistently treating water and nutrient use and human agency in crop production,  
17 thus providing a better understanding of the role humans play in crop production.  
18 Uptake of water and nutrients are two dominant biophysical processes of crop growth  
19 while human agency, including irrigation machine power, land-preparing machine  
20 power and human labor force, determine limits of water and nutrient resources that  
21 are accessible to crops. Two crops, i.e., winter wheat and rice, which account for the  
22 majority of food crop production are considered in a rapidly developing region of the  
23 world, Jiangsu Province, China, that is witnessing the phenomenon of rural to urban  
24 migration. Its production is modeled in two steps. First water and nutrient efficiencies,  
25 defined as the ratios of observed uptake to quantities applied, are modeled as  
26 functions of labor and machine power (representing human agency). In the second  
27 step, crop yields are modeled as functions of water and nutrient efficiencies multiplied  
28 by amounts of water and fertilizers applied. As a result, crop production is predicted  
29 by first simulating water and nutrient uptake efficiencies and then determining yield

30 as a function of water and nutrients that are actually taken up by crops. Results show  
31 that modeled relationship between water use efficiency and human agency explains 68%  
32 of observed variance for wheat and 49% for rice. The modeled relationship between  
33 nutrient use efficiency and human agency explains 49% of the variance for wheat and  
34 56% for rice. The modeled relationships between yields and actual uptakes in the  
35 second step explain even higher percentages of observed the variance: 73% for wheat  
36 and 84% for rice. Leave-one-out cross validation of yield predictions shows that  
37 relative errors are on average within 5% of the observed yields, reinforcing the  
38 robustness of the estimated relationship and of conceptualizing crop production as a  
39 composite function of bio-physical mechanism and human agency. Interpretations  
40 based on the model reveal that after 2005, mechanization gradually led to less labor  
41 being used relative to machinery to achieve same levels of water use efficiency. Labor  
42 and irrigation equipment, on the other hand, were found to be complimentary inputs  
43 to water use efficiency. While the results suggest interventions targeting machinery  
44 are most instrumental in increasing wheat productivity, they may exasperate rural –  
45 urban migration. Policy strategies for alleviating rural-urban migration while ensuring  
46 regional food security can nonetheless be devised where appropriate data are  
47 available.

48

49

50 **1. Introduction**

51 Changing climate and growing population in the Anthropocene (Vörösmarty et al.,  
52 2013; Savenije, H. H. G et al., 2014) are amplifying the tension between water supply  
53 and demand across the planet (Vörösmarty et al., 2000; Arnell and Lloyd-Hughes,  
54 2014; Flörke et al., 2018; Duan et al., 2019; Brown et al., 2019; Di Baldassarre et al.,  
55 2019). Increased average temperatures and variability in rainfall are making water  
56 infrastructures for urban and agriculture water supply obsolete as they have often been  
57 designed for a stationary climate (Milly et al., 2008; Wagener et al., 2010). Such  
58 changes are coupled with rising population in emerging, mostly agrarian, economies  
59 such as China and India that rely on agricultural water (Parry, 2019). Such changes  
60 are major threats for sustainable development as it renders societies water insecure,  
61 food insecure and at the same time stripping rural communities of livelihood  
62 opportunities (Novoa et al., 2019).

63

64 Often human ingenuity is assumed to be able to overcome water and food challenges  
65 posed by changing climate, by conquering climate determinism of human fate and  
66 engender human prosperity through technological innovation in spite of climatic  
67 adversity (Kreibich et al., 2017; Kendall and Spang, 2019). Technological innovations  
68 are assumed to not only mitigate water insecurity but also adapt to it by internalizing  
69 climate change in new water infrastructure and technological designs (Fletcher et al.,  
70 2019; Levin-Koopman et al., 2019; Allen et al., 2019). As part of the solutions offered,  
71 it is assumed that human agency makes the use of water and other related inputs in

72 food production more efficient, thereby releasing pressures of increasing water  
73 scarcity and sustaining food production and human wellbeing (Sivapalan et al., 2014;  
74 Konar et al., 2016).

75

76 Human well-being comprising of food and economic security in water insecure rural  
77 areas depends on crop production that efficiently uses water and nutrients  
78 (Haines-Young & Potschin, 2010; Herrero et al., 2012). This includes the production  
79 of feed for livestock production, thereby making it the foundation of agricultural  
80 systems in general and linking it to phenomena such as migration, agrarian crisis  
81 across the globe and even dispersal of agrarian societies (Afifi et al., 2014; Elshafei et  
82 al., 2014; Pande et al., 2014; FAO et al., 2018). Agricultural systems in Jiangsu  
83 Province, China are one such example, where crop production is a major consumer of  
84 water, accounting for nearly 73.5% of total water consumption in Jiangsu Province,  
85 China (BSJ, 2018). It is also a major human activity, employing 5.82 million people in  
86 Jiangsu province in 2017 (BSJ, 2018). Crop production critically couples wellbeing of  
87 human with water and nutrient cycles and has been shown to be linked with rural to  
88 urban migration in Jiangsu Province, China (Lyu et al., 2019).

89

90 Crop production is influenced both by hydro-climatic variability and the interventions  
91 of humans in terms of provisioning of irrigation and labor. There are therefore several  
92 conceptualizations of interactions between human agency and the environment  
93 (Sivapalan et al., 2015) in how crops are produced, often reflecting the disciplines

94 from which such models have originated. Water proxies such as transpiration, nutrient  
95 proxies such as fertilizer use and their joint-effect effects have been incorporated in  
96 multivariate linear regressions to estimate crop yield–input relationships in  
97 agricultural sciences community (Insam et al., 1991; Heaton et al., 2004). Meanwhile  
98 biophysical models such as CROPWAT (Smith, 1992), Aquacrop (Steduto et al., 2009;  
99 Raes et al., 2009; Hsiao et al., 2009), WOFOST (WOrld FOod STudies) (de Wit et al.,  
100 2018; Lecerf et al., 2018; Ceglar et al., 2019), APSIM (The Agricultural Production  
101 Systems sIMulator) (Holzworth et al., 2014; Gaydon et al., 2017), and statistical  
102 models such as by Sheldrick et al. (2003) explicitly explain the underlying  
103 mechanisms. Similarly, Hatirli et al. (2006) focus on nonlinear water, temperature and  
104 nutrients constraints on biomass production (Ferrero et al., 2018; Hoffman et al.,  
105 2018). These models emphasize the critical role of water and other nutrients in crop  
106 yields and incorporate the role of humans as multipliers that scale optimal yields to  
107 reflect less than optimal efforts of humans. For example, less than optimal crop yield  
108 is often linked to water deficit via linear function, i.e.,  $1 - \frac{Y_a}{Y_x} = K_y(1 - \frac{ET_a}{ET_x})$ , where  
109 the crop yield response factor  $K_y$  changes with crop characteristics (FAO, 2012; Liu  
110 et al. 2002).

111

112 On the other side of the spectrum are conceptualizations of production models based  
113 on economic theory, which emphasize less on biophysical constraints but more on  
114 human agency based on the principles of economics. Models derived based on  
115 Cobb-Douglas production functions (Cobb and Douglas, 1928) have often been

116 applied, such as by Goldsmith et al. (2004), which consider water as an input  
117 alongside other inputs such as machinery and labor in the production of a crop as an  
118 economic good. Other forms of production functions have also been used in this  
119 context, see e.g. McCarl (1982). Others examples include linear programming models  
120 of agricultural production (Howitt, 1995, Pattanayak and Sills 2001), multi-crop  
121 micro-econometric models to interpret farmers' production acreage choice (Femenia et  
122 al., 2018), complex integrated economic-hydrologic models to model the interactions  
123 between water allocation, farmer input choice, agricultural productivity and water  
124 demand (Rosegrant et al. 2000, Roobavannan et al. 2017a) and system dynamics  
125 based socio-hydrological models to understand the interlinkages between water  
126 availability, labor demand and migration (Roobavannan et al. 2017b).

127

128 While such models have proved powerful in simulating yields and modeling labor  
129 employment in the agriculture sector as a function of water availability and other  
130 inputs, it remains a challenge to consistently estimate both yield and labor demand  
131 from the same function. Bio-physical models represent labor as a scaling factor on  
132 potential yields, while economic theory-based models often include water as one of  
133 the inputs into an economic activity while deemphasizing the biophysical role played  
134 by water in biomass production.

135

136 The present paper fills this gap by focusing not only on the bio-physical relationships  
137 of crop yield with water and nutrient inputs, but also by considering how humans, e.g.



138 through irrigation and land-preparation, which would influence the efficiencies of  
139 water and nutrients uptake. The paper acknowledges that understanding the critical  
140 role played by human agency in efficient use of water and nutrients for crop  
141 production is key to facilitating a sustainable future, especially in fast developing  
142 parts of the world. Jiangsu Province in China is one such region, a typical example of  
143 rapidly urbanizing region with a significant flow of economic migrants from rural to  
144 urban areas. Jiangsu is a producer of crops such as rice and wheat, which occupy  
145 almost 60% of the total planted area. Though agriculture production is closely linked  
146 to water availability and is influenced by climatic factors, several government  
147 initiatives have produced rapid development and industrialization of agriculture in  
148 Jiangsu Province. At the same time, it is undergoing an industrial revolution. The  
149 proportion of agriculture output is being gradually reduced by modern secondary and  
150 tertiary industries, affecting income sources of rural families and exasperating rural to  
151 urban migration. Understanding the interlinkages between water security, water and  
152 nutrient efficiency and food production would therefore enable policy makers to  
153 devise and implement appropriate hydrological or economic instruments to address  
154 the migration phenomenon in the province.

155

156 The paper is organized as follows. Section 2 introduces the methodology of  
157 incorporating both bio-physical mechanisms and human agency into a single  
158 crop-production modeling framework and the study area. Modeling results including  
159 the calibrated parameters of the crop production model, together with the results of

160 cross validation, are then shown in section 3. Section 4 then discusses how  
161 substitutions between labor and machinery have changed over time in Jiangsu China,  
162 how it matches with patterns estimated based on independent data and what it means  
163 in terms of rapid mechanization of agriculture in China. Section 5 gives the  
164 conclusions of the study.

165

## 166 2. Methodology

167

168 The light reactions of photosynthesis absorb energy from the sun that is then used by  
169 the dark reactions to convert nutrients into crop biomass (Foyer, 1984; Leegood et al.,  
170 2000; Ke, 2001; Allakhverdiev, 2015). Crop greenness resulting from energy  
171 absorption by light reactions is therefore an important indicator of crop biomass  
172 accumulation and can be measured by reflectance-based vegetation indexes (VI).  
173 Such indices have been widely used as indicators of crop yields (Quarmby et al., 1993;  
174 Ren et al., 2008; Mkhabela et al., 2011; Kogan et al., 2013; Sharma et al., 2015).  
175 Given that transpiration, carbon & nitrogen fixation, and phosphorus consumption  
176 occur in leaves (Foyer, 1984), reflectance measurements have also been utilized to  
177 assess crop water and nutrient status (Sembiring, 1998; Albayrak, 2008; Caturegli et  
178 al., 2016).

179

180 Human agencies, representing labor force and machinery utilization, on the other  
181 hand, contribute to crop production by enabling crops to access water (Allen et al.,  
182 1998) and nutrient resources. Irrigation devices such as pumps and drip-irrigation  
183 systems (Brouwer et al., 1988) help conveying and concentrating water in the root  
184 zone whereas land-preparation machinery, such as tractors create appropriate growing  
185 space for seeds to get access to nutrients (Klein and Zaid Date, 2002). Such agencies  
186 are crucial for improving the efficiencies of water and nutrients use for variety of  
187 crops (Bhiuyan et al. 1992; Erkossa et al., 2005; Johnston and Bruulsema, 2014; Ma


188 et al., 2014).

189

190 Given that human agency supplements water and nutrients in order to efficiently  
191 produce biomass from photosynthesis, we conceptualize crop production as a  
192 (composite) function of human agency induced use efficiency of water and nutrients  
193 and resulting biomass production.

194

## 195 2.1 Conceptual Model Structure



196

197

198 Fig. 1 Crop production conceptualized as a composite function of biophysical  
199 mechanisms and human agency. Human agency influences uptake efficiencies, which  
200 then influence biomass production for given levels of water and nutrient resources.

201


202 Figure 1 illustrates the overall methodology described in detail in Section 2.3. A crop

203 production function is conceptualized as a composite function of biomass production  
204 and efficiency with which water and nutrients are used up. Labor, irrigation and  
205 land-preparation machinery are considered as factors that impact the efficiency of  
206 water and nutrient use by crops. Human agency therefore does not directly contribute  
207 to crop biomass accumulation but determines the amounts of accessible water and  
208 nutrient resources for crops.

209

## 210 2.2 Research Area and Data

211



212

213 Fig. 2 Study area: Jiangsu Province of China. Also shown are the locations of  
214 agro-meteorological stations at which Normalized Difference Vegetation Index

215 (NDVI) and water-crop-related data were used.

216

217 Figure 2 shows the study area. Crop production is modelled in Jiangsu Province,

218 China. Jiangsu Province is in the central area of the south-east coast of China. Being a

219 part of the Yangzi River Delta, Jiangsu Province rapidly developing, together with one  
220 of the highest population densities in the country. The main climate pattern of Jiangsu  
221 Province is subtropical monsoon, with annual precipitation around 1000 mm/year.  
222 Dominated by plain terrain, Jiangsu Province has the highest water surface proportion  
223 among all the administrative regions in China, taking advantage of abundant surface  
224 water resources. The total planted area under food crops in Jiangsu Province reached  
225 to about 5.41 million hectares in 2017. Wheat and rice have the highest two  
226 proportions of plant area, which are 28.69% and 29.94%, respectively. Jiangsu  
227 Province has been undergoing a rapid process of agricultural mechanization, i.e. more  
228 and more machines are being used to replace human labor. The total power of  
229 agricultural mechanics reached to approx. 50 million Kw in 2017, which is nearly 6  
230 times of the value in 1978 (approx. 8.6 million Kw) (Bureau of Statistics of Jiangsu.,  
231 2018). As a result, it is also witnessing rural to urban migration and urgently seeks  
232 solutions that increase water and food security while balancing it with employment in  
233 rural areas.

234

235 Crop growth information, including crop type and growing status, were obtained from  
236 nine agro-meteorological monitoring stations across Jiangsu Province (shown in  
237 Figure 1). Two major types of food crops, i.e., winter wheat (growing season: starts  
238 from October of previous year, 8 months in total) and rice (growing season: starts  
239 from May of current year, 5 months in total) were selected as modeling objects. Of the  
240 nine stations, six stations located in Fengxian, Ganyu, Xuyi, Huaiyin, Yangzhou and

241 Kunshan provided crop growth information for winter wheat; three stations, including  
 242 Ganyu, Dantu and Gaochun, provided information for rice. Time series of  
 243 precipitation, rootzone moisture, transpiration, and provincial crop yield per area, are  
 244 also used. The data sources are listed in Table 2.1.

245

<i>Data categories</i>	<i>Variables (symbol)</i>	<i>Unit</i>	<i>Period</i>	<i>Spatial Resolution</i>	<i>Temporal Resolution</i>	<i>Data source</i>
Hydro-climatic	Precipitation (P)	mm	2000 -2017	0.5*0.5 °	Derived from monthly data. Growing-season-accu- mulated value for each year.	CRU (CRU, 1901-2017; Harris et al., 2014)
	Transpiration (T)			0.25*0.25 °		GLDAS Noah Land Surface Model L4 monthly
	Rootzone Moisture (S <sub>w</sub> )			0.25*0.25 °		0.25*0.25 ° V2.1 (Rodell et al., 2004)
Crop Information	NDVI (g)	-	2000 -2017	30 meters	Derived from 8-day data. Growing-season-maxi- mum value for each year.	Landsat 7 NDVI (imported from Google Earth Engine: 'LANDSAT/LE07/C01/T1_8 DAY_NDVI', Gorelick et al., 2017)
	Crop type & Growing season		1991 -2010	Station-level	Yearly	National Meteorological Information Center of China (2006)

	Provincial crop yield (Y)	kg/ha	2001-2017	Provincial	Yearly	Statistical Yearbook of Jiangsu (BSJ, 2018)
Human Agencies	Labor force in crop cultivation ( $L_C$ )	Capita/1000ha	2001-2017	Provincial	Yearly	Statistical Yearbook of Jiangsu (BSJ, 2018)
	Irrigation machinery ( $M_I$ )	Kw/1000ha				
	Land-preparing machinery ( $M_L$ )	1000ha				
	Fertilizer use (F)	Ton/1000ha				

246 Table 2.1 Description of the data sets used. The overlapping period from 2001-2017 was used for  
247 regression and related analysis.

248

249

### 250 2.3 Model Set-up, calibration and validation

251

252 Let crop yield  $Y$  be represented by a function  $G(.,.)$  of actual water,  $x_W$ , and nutrient,  
253  $x_N$ , uptakes. Then  $Y = G(x_W, x_N)$ . However, actual amounts of uptakes are often less  
254 than total amount of water available  $W_T$  in the form of rainfall  $R$ , rootzone moisture  
255  $S_W$  and nutrients available after fertilizer amount  $F$  has been applied. The actual  
256 amount of water and nutrient uptakes relative to their available supply defines





279 Burke, 2010; Cai et al., 2014; Álvarez et al., 2017) are performed in two steps. This  
280 also synthesizes observations at different locations and provides general relationships  
281 *across stations*.

282

283 *Fixed Effect Estimation of the model in two stages*

284 **Step 1:** In order to understand regional water and nutrient use efficiencies across  
285 locations, panel regression is performed across stations to estimate  $\eta_W$  and  $\eta_N$  as  
286 functions of human activities,  $H$ . We use the ratios  $\eta_W = x_W/W_T$  and  $\eta_N = x_N/F$   
287 (efficiencies of water and nutrient uptakes respectively) as dependent variables and  
288 use inputs,  $H$ , such as machineries linked to labor and irrigation as independent  
289 variables to estimate the following equations for stations  $i = 1, \dots, S$ :

290

$$291 \quad \eta_W^i = \Lambda H^i + \delta^i + \epsilon_W$$

$$292 \quad \eta_N^i = \Theta H^i + \theta^i + \epsilon_N$$

293 (2a, b)

294 Here,  $H^i$  represent station-specific human activities but its effect on efficiencies,  
295  $(\Lambda, \Theta)$ , are general across all the stations. Hence,  $(\Lambda, \Theta)$  are independent of station  $i$ .  
296 Fixed station-specific effects are quantified by  $(\delta^i, \theta^i)$ , and  $(\epsilon_W, \epsilon_N)$  represent the  
297 residuals accounting for variance of efficiencies not explained by  $H$ . The estimation of  
298 effects is based on linear regression of equations in Equation 2a,b and implemented  
299 by using Álvarez et al. (2017).

300

301 **Step 2:** Panel regressions are again employed to estimate crop yields as functions of  
 302 observed water and nutrients uptakes, independent of the stations. We assume that  
 303  $G(x_W, x_N) = kx_W^\alpha x_N^\beta$  (Kouka et al., 1994; Gowariker et al., 2009; Xin et al., 2016;  
 304 Li et al., 2016). This is done by estimating the following equation in log-space,  
 305 accounting for station specific fixed effects.

$$306 \quad \log(Y^i) = \alpha \log(x_W^i) + \beta \log(x_N^i) + \log(k) + \pi^i + \epsilon_Y$$

307 (3)

308 Here  $(x_W^i, x_N^i)$  are ‘observed’ water and nutrient inputs and  $Y^i$  corresponding  
 309 observed yields. Note that the effects  $(\alpha, \beta, k)$  are independent of the stations while  
 310  $\pi^i$  is station specific fixed effect and  $\epsilon_Y$  represents variance of  $Y$  not explain by the  
 311 independent variables. The *station independent* prediction of yields are obtained by  
 312 removing fixed effects  $Y = kx_W^\alpha x_N^\beta$ . The regression is performed using Álvarez et  
 313 al. (2017).

314

### 315 *Model based prediction*

316 When *predicting* yields, predictions of station specific water,  $\hat{x}_W^i$ , and nutrient  
 317 uptakes,  $\hat{x}_N^i$ , are obtained as products of station-specific predictions of water and  
 318 nutrient use efficiencies with location specific water and nutrient availability  
 319 respectively.

320

321 That is,  $\hat{x}_W^i = \hat{\eta}_W^i W_T^i$  and  $\hat{x}_N^i = \hat{\eta}_N^i F^i$  with  $\hat{\eta}_W^i = \Lambda H^i + \delta^i$  and  $\hat{\eta}_N^i = \Lambda H^i + \theta^i$   
 322 being predictions of first stage regressions.

323

### 324 *2.3.1 Proxies for water and nutrient uptakes*

325

326 Transpiration,  $T$ , is chosen as the proxy for water uptake by plants. Since it is harder  
327 to detect nutrient uptake directly, proxy for nutrient uptake is estimated based on  
328 Normalized Difference Vegetation Index (NDVI), (Landsat 7, 2001-2017). NDVI  
329 reflects the joint effect of water and nutrient uptakes on plant greenness (Quarmby et  
330 al., 1993; Ren et al., 2008; Mkhabela et al., 2011; Kogan et al., 2013; Sharma et al.,  
331 2015). Therefore, the effect of water uptake on NDVI is first filtered out and the  
332 remaining variance of NDVI is then assumed to approximate the uptake of nutrients.

333

334 In order to filter out the effect of water uptake from NDVI, a fixed effect regression  
335 across the stations, similar to fixed effect regressions described above, is conducted in  
336 log space with NDVI as dependent variable,  $g$ , and water uptake  $x_W$ , as represented  
337 by transpiration  $T$ , as the independent variable. This regression provides  $\hat{g}$  (an  
338 estimate of  $g$ ), which is the part of greenness that is explained only by water uptake.  
339 The difference between  $g$  and  $\hat{g}$  in log space, i.e. residuals, then provides the part of  
340 greenness that is only a function of nutrients taken up by crops. Such residuals are  
341 then taken as proxy of nutrient uptake  $N$ , i.e.

$$342 \quad \log(N) = \log(g) - \log(\hat{g})$$

343

(4)

344 The yearly maximum value of NDVI during the growing season is chosen to represent

345 the maximum level of crop greenness because peak NDVI is most sensitive to the  
346 levels of water and nutrient uptakes (Gamon et al., 1995). We assume that higher peak  
347 NDVI also implies that the crop has undergone lower water and nutrient stress during  
348 other critical growth stages. The growing season considered for winter wheat was  
349 from 1<sup>st</sup> October of previous year to 1<sup>st</sup> June of next year (8 months), while the  
350 growing period of rice was set as 1<sup>st</sup> May to 1<sup>st</sup> of October (5 months).

351

### 352 2.3.2 Water and nutrient use efficiency

353

354 Water and nutrient use efficiencies are defined as the ratio of transpiration  $T$  and  
355 nutrient proxy  $N$  to total available water,  $W_T$ , and nutrient resources respectively.  
356 Total available water resources,  $W_T$ , is defined as the sum of root zone moisture  $S_W$   
357 at the beginning of crop growing season and precipitation  $P$  during crop growing  
358 season. Nutrient availability,  $F$ , is represented by the total amount of fertilizer applied  
359 per unit area – assuming that yield response to increased amounts of residual soil  
360 nutrients are much less than to freshly applied fertilizer (Prihar et al., 1985). The  
361 *observed* water use efficiency  $\eta_W$  and nutrient use efficiency  $\eta_N$  are then calculated  
362 as follows and used to calibrate its predictive equations (equations 2a, b).

$$363 \quad \eta_W = \frac{T}{P + S_W}$$

364 (5a)

$$365 \quad \eta_N = \frac{N}{F}$$

366 (5b)

367 Human factors such as labor used in crop production  $L_C$ , irrigation machinery power  
368  $M_I$  and land-preparing machinery power  $M_L$  per unit area are considered in the set  
369 of independent variables  $H$  (see equations 2a, b). All combinations of joint and  
370 individual effects (such as  $L_C M_I M_L$ ,  $L_C M_I$ ,  $M_I M_L$ ,  $L_C M_I$ ,  $L_C$ ,  $M_I$  and  $M_L$ ) were first  
371 regressed and only those effects that were statistically significant were selected in the  
372 final model.

373

### 374 *2.3.3 Validation*

375

376 Leave-one-out cross validation was implemented to test the robustness of estimated  
377 crop production for each crop. For each station, data was available for 17 years  
378 (2001-2017, see Table 1). In each round of validation, 16 out of 17 years for each  
379 station were chosen to train the model, while the remaining year was used to validate  
380 the estimated model. This was repeated 17 times, each time with a unique year left out  
381 for validation. Boxplots of relative errors show the distribution of relative errors in  
382 leave one out cross validation. The calculation of relative errors is defined as:

$$383 \quad RE = (\hat{y} - y)/x$$

384 (5)

385 where  $\hat{y}$  represents the predicted yield for a crop using all except one year of data,  
386 whereas  $y$  represents the observed yield.

387

## 388 **3. Model interpretation: substitution between labor and machinery in winter**

389 **wheat cultivation**

390

391 The proposed crop production function is a composite function of crop yield and  
392 efficiency with which water and nutrients are taken up, as facilitated by human  
393 agency.

394

395 One can therefore interrogate such a model to understand how tradeoffs between  
396 different components of human agencies have evolved over time. The water use  
397 efficiency of winter wheat, as shown in Table 3.2, is supported by labor and irrigation  
398 machinery (pumps) and land-preparation machinery (tractors and supporting tools).  
399 Thus, winter wheat serves as an interesting example to investigate how different  
400 elements have substituted one another and shed light on the mechanization of  
401 agriculture in Jiangsu. Here we show that such estimations based on the composite  
402 production function are consistent with observed data.

403

404 Here, by substitution of one factor by another we mean how much of one factor can  
405 be substituted by one unit of another factor such that water use efficiency remains the  
406 same. This requires, for example, the estimation of  $\frac{dL_C}{dM_I}$  such that  $d(\eta_W) = 0$  (so  
407 that the level of water use efficiency remains the same).

$$408 \quad d(\eta_W) = \frac{\partial \eta_W}{\partial L_C} dL_C + \frac{\partial \eta_W}{\partial M_I} dM_I + \frac{\partial \eta_W}{\partial M_L} dM_L = 0$$

409 (4.1)

410 To obtain  $\frac{dL_C}{dM_I}$ , we divide both sides of Eq. 4.1 by  $dM_I$ :

411 
$$\frac{\partial \eta_W}{\partial L_C} \frac{dL_C}{dM_I} + \frac{\partial \eta_W}{\partial M_I} + \frac{\partial \eta_W}{\partial M_L} \frac{dM_L}{dM_I} = 0$$

412 (4.2)

413 Obtaining  $\frac{dM_L}{dM_I}$  by using data from the statistical yearbooks of the province (BSJ,  
 414 2001~2018),  $\frac{dL_C}{dM_I}$  can be calculated as

415 
$$\frac{dL_C}{dM_I} = -\frac{\partial L_C}{\partial \eta_W} \left( \frac{\partial \eta_W}{\partial M_I} + \frac{\partial \eta_W}{\partial M_L} \frac{dM_L}{dM_I} \right)$$

416 (4.3a)

417 Note here that partial derivatives can be estimated from the regressed equations in

418 2.3.2. Similarly,  $\frac{dL_C}{dM_L}$  can be calculated as:

419 
$$\frac{dL_C}{dM_L} = -\frac{\partial L_C}{\partial \eta_W} \left( \frac{\partial \eta_W}{\partial M_L} + \frac{\partial \eta_W}{\partial M_I} \frac{dM_I}{dM_L} \right)$$

420 (4.3b)

421 The calculated  $\frac{dL_C}{dM_I}$  and  $\frac{dL_C}{dM_L}$  are shown in figure 4.1a, b.

422



423 **4. Model Results**

424 **4.1 Yield-uptake relationship**

425 Table 3.1 gives the coefficients of proxies correspond to the effects (i.e.,  $\alpha$  and  $\beta$  in  
 426 equation 3) of water and nutrient respectively. It reports that estimated effects for both  
 427 the crops were significant.

428

<i>Crops</i>	<i>X</i>	<i>Coefficients</i>	<i>Std. Error</i>
<i>Winter wheat</i>	$(x_W)^\alpha$	0.53	0.06
	$(x_N)^\beta$	0.12	0.03
	<i>Overall</i>	<i>R-squared</i>	0.73
<i>Rice</i>	$(x_W)^\alpha$	0.17	0.02
	$(x_N)^\beta$	0.04	0.01
	<i>Overall</i>	<i>R-squared</i>	0.84

429 Table 3.1 Yield-uptake fixed effect estimation of  
 430  $\alpha$  and  
 431  $\beta$  for the two crops. All effects are significant with  $p < 0.01$ .

432

433 Figure 3a,b show observed yields in comparison with the modeled yields for rice and  
 434 wheat. Modeled yields for various stations are obtained by incorporating station  
 435 specific fixed effects (from equation 3) for stations  $i = 1, \dots, S$  with parameters given  
 436 in Table 3.1 (fixed effects  $k^i = ke^{\pi^i}$  not shown).




Fig. 3a, b Modeled and observed crop yields based on estimated effects shown in

Table 3.1.

443 **4.2 Water Use Efficiency**

444 Table 3.2 shows that several elements of  $H$  were found to be statistically significant in  
 445 explaining water use efficiency of winter wheat. In case of rice, only irrigation  
 446 machinery  $M_I$  demonstrated significant effect (regression coefficient) on water use  
 447 efficiency, which reflects that water access is most important for its water uptake.

448

<i>Crops</i>	<i>H</i>	<i>Coefficients</i>	<i>Std. Error</i>
<b>Winter wheat</b>	$L_C$	$8.57e-3$	$2.76e-3$
	$M_I$	$5.60e-3$	$2.00e-3$
	$M_L$	$2.94e-3$	$0.75e-3$
	$L_C * M_I$	$-9.92e-6$	$0.34e-5$
	$L_C * M_L$	$-3.77e-6$	$1.05e-6$
	$M_I * M_L$	$-3.41e-6$	$0.92e-6$
	$L_C * M_I * M_L$	$4.45e-9$	$1.29e-9$
	<i>Overall</i>	<i>R-squared</i>	<i>0.68</i>
<b>Rice</b>	$M_I$	$3.01e-4$	$0.77e-4$
	<i>Overall</i>	<i>R-squared</i>	<i>0.49</i>


449 Table 3.2 Fixed effect estimation of water use efficiency for the two crops. All effects  
 450 are significant at  $p < 0.01$ .

451

452 Again, Figure 4a, b show ‘observed’ (see equation 5a how water use efficiency, i.e.,  
 453 WUE, has been defined) WUE in comparison with the modeled WUE for rice and

454 wheat (from equation 2a).

455



456

457 Fig. 4a, b Modeled and observed crop water use efficiency (WUE) across the stations.

458

459 **4.3 Nutrient Use Efficiency**

460 Table 3.3 shows that land-preparing machinery is the only significant factor for the  
 461 nutrient use efficiency of winter wheat. This indicates that better-prepared farmland is  
 462 the only significant factor that facilitated better nutrient access for winter wheat. On  
 463 the other hand, the major contributing factor to rice nutrient use efficiency is labor  
 464 power, together with the joint-effect factor of crop labor and land-preparation  
 465 machinery.


466

<i>Crops</i>	<i>X</i>	<i>Coefficients</i>	<i>Std. Error</i>
<i>Winter Wheat</i>	$M_L$	$1.26e-7$	$0.23e-7$
	<i>Overall</i>	<i>R-squared</i>	$0.49$
<i>Rice</i>	$L_C$	$2.35e-8$	$0.58e-8$
	$L_C * M_L$	$5.35e-12$	$1.25e-12$
	<i>Overall</i>	<i>R-squared</i>	$0.56$

467 Table 3.3 Fixed effect estimation of nutrient use efficiency. All effects are significant  
 468 at  $p < 0.01$ .

469

470



471

472 Fig. 5a, b Modeled vs observed crop nutrient use efficiency (NUE) across stations for

473 two crop types.


474

475

476 **4.4 Leave-one-out cross validation**

477

478 According to Fig 6a, b, for all crops at all stations, the median values (shown by red  
479 line at the center of the box plots) of relative errors are within  $\pm 0.05$  of observed  
480 values (as indicated by the red shadows). This indicates that the proposed model is  
481 robust in modeling winter wheat and rice production within Jiangsu Province.



482

483

484 Fig. 6a, b: Leave-one-out cross validation across the stations and two crops.

485 Distribution of relative errors (equation 5) are shown with zero error shown by the  
486 grey dashed line, and  $\pm 0.05$  error shown by the red shadows


487

488 **4.5 Substitution between labor and machinery in winter wheat cultivation**

489

490 Fig. 7a, b indicate that during years before 2005, more labor,  $L_C$ , was used relative to  
491 machinery  $M_L$ . After 2005, mechanization gradually led to less labor being used  
492 relative to machinery while achieving same level of water use efficiency. The  
493 derivative between  $L_C$  and  $M_L$ , i.e.,  $\frac{dL_C}{dM_L}$ , however fluctuated around 0, indicating  
494 that they are complimentary and do not tend to substitute one another. The close  
495 resemblance of substitution effects estimated based on regressed relationships,  
496 together with those estimated based on statistical year books (indicated as data in  
497 Figure 6), further suggests that the proposed production function is capable of  
498 providing robust interpretation of how one input has been, or can be, substituted with  
499 another without affecting water use efficiency.

500



501

502

Fig.7a, b Rates of substitution between labor and machinery

503

504

505



## 506 **5. Discussion and Conclusion**

507

508 This paper conceptualized crop production as a composite function of bio-physical  
509 mechanisms and human-agency. While the former links water and nutrient uptakes to  
510 crop biomass production, the latter influences the efficiencies with which water and  
511 nutrients are taken up.

512

513 The model was calibrated using hydro-climatic and agricultural statistics from 2001 to  
514 2017 for two main food crops in Jiangsu province, i.e., winter wheat and rice, using  
515 panel regressions across agro-meteorological monitoring stations (six for winter  
516 wheat, three for rice). The median performance of the composite function was found  
517 to be within 5% of the observed based on leave one out cross validation.

518

519 The fixed effect regressions were used to filter out station specific effects of human  
520 agency on use efficiencies and of efficiencies on crop production. This data-driven  
521 approach was key to commensurate, to certain extent, different scales of the data sets  
522 used and to obtain a generic relationship that is devoid of any station specific effects.

523 The yield data used was at provincial level, transpiration and soil moisture was at  $0.25^\circ$   
524  $\times 0.25^\circ$  scale based on GLDAS reanalysis data, and NDVI and human agency data  
525 was station specific. However, transpiration and soil moisture data used is at much  
526 coarse resolution compared to NDVI, which means that, for example, transpiration  
527 would give an aggregate for both (irrigated) crops and native vegetation and other

528 land surfaces. It is assumed that higher peak NDVI also implies that the crop has  
529 undergone lower water and nutrient stress during other critical growth stages. Further,  
530 irrigation has been ignored when calculating water use efficiency. Results therefore  
531 demonstrate a proof of concept at best, which can be made more reliable with higher  
532 resolution data sets.

533

534 The data driven approach treated crop production as a composite function of water  
535 and nutrient use efficiency and human agency. This approach fills a gap in our  
536 coupled human-water system understanding of crop production, which either has been  
537 focused on bio-physical mechanisms or based on economic production theory. The  
538 proposed method demonstrated its novelty by not only modeling the bio-physical  
539 relationships of crop yield with water and nutrient inputs, but also considering how  
540 humans, e.g. through irrigation and land-preparation, influence the efficiencies of  
541 water and nutrients uptake. Crop and labor machinery were found to be important for  
542 nutrient use efficiency. Irrigation machinery was most important for water use  
543 efficiency of rice production. However, all aspects of human agency were important  
544 for water use efficiency of winter wheat production.

545

546 These results are intuitive, suggesting the labor and land preparation machinery are  
547 key to crop production in the region in general. While the variance explained by  
548 human agency in explaining nutrient use efficiency was similar across the two crops,  
549 human agency appeared to explain water use efficiency of winter wheat a lot better

550 than rice. This indicates that WUE of rice is less sensitive to human agency and  
551 perhaps more dependent on water scarcity. On the other hand water and nutrient use  
552 efficiency and therefore crop productivity of winter wheat was sensitive to various  
553 aspects of human agency such as labor and land preparation machinery. The  
554 differences in the effects between the two crops indicate that rice production is a  
555 water intensive crop and its yield exclusively depends on how well the crop is  
556 irrigated. Even though rice cultivation is labor intensive, the role of human agency in  
557 various stages of the crop growth appears to be less complicated. In contrast, winter  
558 wheat, often grown in autumn, relies on a complex interplay of water and nutrient  
559 availability that is facilitated by human agency during its growing period.

560

561 The substitution analysis revealed that more labor was used relative to machinery in  
562 winter wheat production before 2005. Post 2005, mechanization gradually led to less  
563 labor being used relative to machinery while achieving similar level of water use  
564 efficiency. Labor ( $L_C$ ) and irrigation machinery ( $M_I$ ) were found to be complimentary  
565 to water use efficiency of winter wheat production. Therefore, interventions targeting  
566 machinery are most instrumental in increasing wheat productivity.

567

568 Lyu et al. (2019) have recently found that under-employment in rural areas of Jiangsu  
569 Province has been fueling the rural to urban migration. Given the gains in efficiency  
570 that mechanization produces and the observed transition to mechanization, any sound  
571 policy aimed at alleviating under-employment and hence migration should target

572 more skilled employment in the non-agricultural sectors of rural areas. This will  
573 ensure rural employment, sustainable rural communities (Li, 2010) as well as regional  
574 food security.

575

576 The methodology is transferrable to other regions as well. This is because the data  
577 sets used are regional agricultural statistics on crop yields and open access  
578 hydrological data, such as reanalysis data for transpiration and soil moisture (Rodell  
579 et al., 2004) and high resolution LANDSAT7 based NDVI data (Gorelick et al., 2017).

580 Policy strategies for alleviating migration while ensuring regional food security  
581 therefore can be devised based on crop production simulations, as shown in this paper,  
582 in regions where agricultural statistics data are available. This can be done by  
583 analyzing the implications of crop production simulations that ensure food security on  
584 rural employment under future climate and socio-economic scenarios. As Lyu et al.  
585 (2019) have found, rural under-employment is a major driver of rural-urban migration.

586 Target regions could be fast developing regions such as Maghreb region of Africa and  
587 South Asia that are witnessing massive flux of rural to urban economic migrants. Yet,  
588 given that the dataset that the approach relies on is either reanalysis or at different  
589 scales, such policy designs will need to be handled with caution and be validated  
590 based on field campaigns where possible.

591

592 **References**

- 593 Afifi, T., Liwenga, E., & Kwezi, L. (2014). Rainfall-induced crop failure, food  
594 insecurity and out-migration in Same-Kilimanjaro, Tanzania. *Climate and*  
595 *Development*, 6(1), 53-60. doi: <https://doi.org/10.1080/17565529.2013.826128>
- 596 Albayrak, S. (2008). Use of reflectance measurements for the detection of N, P, K,  
597 ADF and NDF contents in sainfoin pasture. *Sensors*, 8(11), 7275-7286.  
598 doi: <https://doi.org/10.3390/s8117275>
- 599 Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). *FAO Irrigation and*  
600 *drainage paper No. 56. Rome: Food and Agriculture Organization of the*  
601 *United Nations*, 56(97), e156.
- 602 Allen, T. R., Crawford, T., Montz, B., Whitehead, J., Lovelace, S., Hanks, A. D., ... &  
603 Kearney, G. D. (2019). Linking Water Infrastructure, Public Health, and Sea  
604 Level Rise: Integrated Assessment of Flood Resilience in Coastal Cities.  
605 *Public Works Management & Policy*, 24(1), 110-139. doi:  
606 <https://doi.org/10.1177/1087724X18798380>
- 607 Álvarez, I., Barbero, J., & Zofio, J. (2017). A Panel Data Toolbox for  
608 MATLAB. *Journal of Statistical Software*, 76(6), 1 - 27.  
609 doi: <http://dx.doi.org/10.18637/jss.v076.i06>
- 610 Arias-Jiménez, E. (2002). *FAO Plant Production and protection paper No. 156 Rev. 1:*  
611 *Date palm cultivation*, chapter VI by Klein and Zaid Date, 2002. Retrieved  
612 from: <http://www.fao.org/3/y4360e/y4360e00.htm>
- 613 Arnell, N. W., & Lloyd-Hughes, B. (2014). The global-scale impacts of climate

614 change on water resources and flooding under new climate and  
615 socio-economic scenarios. *Climatic Change*, 122(1-2), 127-140. doi:  
616 10.1007/s10584-013-0948-4

617 Bureau of Statistics of Jiangsu. (2018). *Statistical Yearbook of Jiangsu: China*  
618 Statistics Press. Retrieved from  
619 <http://tj.jiangsu.gov.cn/col/col70123/index.html> (Chinese version, access for  
620 free.)

621 Brown, T. C., Mahat, V., & Ramirez, J. A. (2019). Adaptation to future water  
622 shortages in the United States caused by population growth and climate  
623 change. *Earth's Future*, 7(3), 219-234. doi:  
624 <https://doi.org/10.1029/2018EF001091>

625 Brouwer, C., Prins, K., Kay, M., & Heibloem, M. (1988). *Irrigation water*  
626 *management: irrigation methods. Training manual*, 9.

627 Cai, R., Yu, D., & Oppenheimer, M. (2014). Estimating the Spatially Varying  
628 Responses of Corn Yields to Weather Variations using Geographically  
629 Weighted Panel Regression. *Journal of Agricultural and Resource*  
630 *Economics*, 39(1835-2016-149505), 230-252.

631 Ceglar, A., Van der Wijngaart, R., De Wit, A., Lecerf, R., Boogaard, H., Seguni, L., ...  
632 & Baruth, B. (2019). Improving WOFOST model to simulate winter wheat  
633 phenology in Europe: Evaluation and effects on yield. *Agricultural*  
634 *Systems*, 168, 168-180. doi: <https://doi.org/10.1016/j.agsy.2018.05.002>

635 CHE, S.-g., ZHAO, B.-q., LI, Y.-t., Liang, Y., Lin, Z.-a., HU, S.-w., & Bing, S. (2016).

636 Nutrient uptake requirements with increasing grain yield for rice in China.  
637 *Journal of integrative agriculture*, 15(4), 907-917.  
638 doi: [https://doi.org/10.1016/S2095-3119\(15\)61143-1](https://doi.org/10.1016/S2095-3119(15)61143-1)

639 Cobb, C. W., & Douglas, P. H. (1928). A Theory of Production. *American Economic*  
640 *Review*, 18(1), 139-165.

641 CRU TS4.02: Climatic Research Unit (CRU) Time-Series (TS) version 4.02 of  
642 high-resolution gridded data of month-by-month variation in climate (Jan.  
643 1901- Dec. 2017).  
644 Retrieved from <https://crudata.uea.ac.uk/cru/data/hrg/>

645 de Wit, A., Boogaard, H., Fumagalli, D., Janssen, S., Knapen, R., van Kraalingen, D.,  
646 Supit, I., van der Wijngaart, R., & van Diepen, K. (2018). 25 years of the  
647 WOFOST cropping systems model. *Agricultural systems*. doi:  
648 <https://doi.org/10.1016/j.agsy.2018.06.018>

649 Di Baldassarre, G., Sivapalan, M., Rusca, M., Cudennec, C., Garcia, M., Kreibich,  
650 H., ... & Sanderson, M. R. (2019). Sociohydrology: Scientific challenges in  
651 addressing the sustainable development goals. *Water Resources Research*. doi:  
652 <https://doi.org/10.1029/2018WR023901>

653 Duan, K., Caldwell, P. V., Sun, G., McNulty, S. G., Zhang, Y., Shuster, E., ... &  
654 Bolstad, P. V. (2019). Understanding the role of regional water connectivity in  
655 mitigating climate change impacts on surface water supply stress in the United  
656 States. *Journal of hydrology*, 570, 80-95. doi:  
657 <https://doi.org/10.1016/j.jhydrol.2019.01.011>

658 Elshafei, Y., Sivapalan, M., Tonts, M., & Hipsey, M. R. (2014). A prototype  
659 framework for models of socio-hydrology: identification of key feedback  
660 loops and parameterisation approach. *Hydrology and Earth System  
661 Sciences*, 18(6), 2141-2166. doi: <https://doi.org/10.5194/hess-18-2141-2014>

662 Erkossa, T., Stahr, K., & Gaiser, T. (2005). Effect of different methods of land  
663 preparation on runoff, soil and nutrient losses from a Vertisol in the Ethiopian  
664 highlands. *Soil Use and Management*, 21(2), 253-259.  
665 doi: <https://doi.org/10.1111/j.1475-2743.2005.tb00132.x>

666 FAO. (2012). Crop yield response to water. *FAO Irrigation and Drainage Paper,  
667 Paper 66. ISSN 0254-5284.*

668 FAO IFAD IOM WFP. (2018). The Linkages between Migration, Agriculture, Food  
669 Security and Rural Development. Rome. 80pp.  
670 (<http://www.fao.org/3/CA0922EN/CA0922EN.pdf>). Licence: CC BY-NC-SA  
671 3.0 IGO

672 Femenia, F., Carpentier, A., & Koutchade, O. P. (2018). Dealing with corner solutions  
673 in multi-crop micro-econometric models: an endogenous regime approach  
674 with regime fixed costs. Post-Print hal-01879042, *HAL*. Retrieved from  
675 <https://ideas.repec.org/p/hal/journal/hal-01879042.html>

676 Ferrero, R., Lima, M., & Gonzalez-Andujar, J. L. (2018). Crop production structure  
677 and stability under climate change in South America. *Annals of applied  
678 biology*, 172(1), 65-73. doi: <https://doi.org/10.1111/aab.12402>

679 Fletcher, S., Lickley, M., & Strzepek, K. (2019). Learning about climate change



680 uncertainty enables flexible water infrastructure planning. *Nature*  
681 *communications*, 10(1), 1782. doi:  
682 <https://doi.org/10.1038/s41467-019-09677-x>

683 Flörke, M., Schneider, C., & McDonald, R. I. (2018). Water competition between  
684 cities and agriculture driven by climate change and urban growth. *Nature*  
685 *Sustainability*, 1(1), 51. doi: <https://doi.org/10.1038/s41893-017-0006-8>

686 Foyer, C. (1984). *Photosynthesis (Cell biology, v. 1)*. New York: Wiley.

687 Gamon, J. A., Field, C. B., Goulden, M. L., Griffin, K. L., Hartley, A. E., Joel, G., ...  
688 & Valentini, R. (1995). Relationships between NDVI, canopy structure, and  
689 photosynthesis in three Californian vegetation types. *Ecological*  
690 *Applications*, 5(1), 28-41.

691 Gaydon, D. S., Wang, E., Poulton, P. L., Ahmad, B., Ahmed, F., Akhter, S., ... &  
692 Choudhury, B. U. (2017). Evaluation of the APSIM model in cropping  
693 systems of Asia. *Field Crops Research*, 204, 52-75. doi:  
694 <https://doi.org/10.1016/j.fcr.2016.12.015>

695 Goldsmith, P. D., Gunjal, K., & Ndarishikanye, B. (2004). Rural–urban migration and  
696 agricultural productivity: the case of Senegal. *Agricultural economics*, 31(1),  
697 33-45. doi: <https://doi.org/10.1111/j.1574-0862.2004.tb00220.x>

698 Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R.  
699 (2017). Google Earth Engine: Planetary-scale geospatial analysis for  
700 everyone. *Remote Sensing of Environment*, 202, 18-27.  
701 doi: <https://doi.org/10.1016/j.rse.2017.06.031>

702 Gowariker, V., Krishnamurthy, V. N., Gowariker, S., Dhanorkar, M., & Paranjape, K.  
703 (2009). *The fertilizer encyclopedia*. John Wiley & Sons.

704 Haines-Young, R., & Potschin, M. (2010). The links between biodiversity, ecosystem  
705 services and human well-being. *Ecosystem Ecology: a new synthesis, 1*,  
706 110-139.

707 Harris, I. P. D. J., Jones, P. D., Osborn, T. J., & Lister, D. H. (2014). Updated high-  
708 resolution grids of monthly climatic observations—the CRU TS3. 10  
709 Dataset. *International journal of climatology*, 34(3), 623-642.  
710 doi: <https://doi.org/10.1002/joc.3711>

711 Hatirli, S. A., Ozkan, B., & Fert, C. (2006). Energy inputs and crop yield relationship  
712 in greenhouse tomato production. *Renewable Energy*, 31(4), 427-438.  
713 doi: <https://doi.org/10.1016/j.renene.2005.04.007>

714 Heaton, E., Voigt, T., & Long, S. P. (2004). A quantitative review comparing the  
715 yields of two candidate C4 perennial biomass crops in relation to nitrogen,  
716 temperature and water. *Biomass and bioenergy*, 27(1), 21-30.  
717 doi: <https://doi.org/10.1016/j.biombioe.2003.10.005>

718 Herrero, M. T., Thornton, P. K., Notenbaert, A. M. O., Msangi, S., Wood, S., Kruska,  
719 R. L., Dixon, J.A., Bossio, D.A., Steeg, J.V.D., Freeman, H.A., & Li, X.  
720 (2012). Drivers of change in crop–livestock systems and their potential  
721 impacts on agro-ecosystems services and human wellbeing to 2030: A study  
722 commissioned by the CGIAR Systemwide Livestock Programme. Retrieved  
723 from:

724 [https://cgspace.cgiar.org/bitstream/handle/10568/3020/SLPdriversstudyfinaldr](https://cgspace.cgiar.org/bitstream/handle/10568/3020/SLPdriversstudyfinaldraft.pdf?sequence=4)  
725 [aft.pdf?sequence=4](https://cgspace.cgiar.org/bitstream/handle/10568/3020/SLPdriversstudyfinaldraft.pdf?sequence=4)

726 Hoffman, A. L., Kemanian, A. R., & Forest, C. E. (2018). Analysis of climate signals  
727 in the crop yield record of sub-Saharan Africa. *Global change biology*, 24(1),  
728 143-157. doi: <https://doi.org/10.1111/gcb.13901>

729 Holzworth, D. P., Huth, N. I., deVoil, P. G., Zurcher, E. J., Herrmann, N. I., McLean,  
730 G., ... & Moore, A. D. (2014). APSIM—evolution towards a new generation of  
731 agricultural systems simulation. *Environmental Modelling & Software*, 62,  
732 327-350. doi: <https://doi.org/10.1016/j.envsoft.2014.07.009>

733 Howitt, R. E. (1995). A calibration method for agricultural economic production  
734 models. *Journal of agricultural economics*, 46(2), 147-159.  
735 doi: <https://doi.org/10.1111/j.1477-9552.1995.tb00762.x>

736 Hsiao, T. C., Heng, L., Steduto, P., Rojas-Lara, B., Raes, D., & Fereres, E. (2009).  
737 AquaCrop—the FAO crop model to simulate yield response to water: III.  
738 Parameterization and testing for maize. *Agronomy Journal*, 101(3), 448-459.  
739 doi: [doi:10.2134/agronj2008.0218s](https://doi.org/10.2134/agronj2008.0218s)

740 Insam, H., Mitchell, C. C., & Dormaar, J. F. (1991). Relationship of soil microbial  
741 biomass and activity with fertilization practice and crop yield of three  
742 ultisols. *Soil Biology and Biochemistry*, 23(5), 459-464.  
743 doi: [https://doi.org/10.1016/0038-0717\(91\)90010-H](https://doi.org/10.1016/0038-0717(91)90010-H)

744 Johnston, A., & Bruulsema, T. (2014). 4R nutrient stewardship for improved nutrient  
745 use efficiency. *Procedia Engineering*, 83, 365-370.

746 doi: <https://doi.org/10.1016/j.proeng.2014.09.029>

747 Ke, B. (2001). *Photosynthesis photobiochemistry and photobiophysics (Vol. 10)*:  
748 Springer Science & Business Media.

749 Kendall, A., & Spang, E. S. (2019). The role of industrial ecology in food and  
750 agriculture's adaptation to climate change. *Journal of Industrial Ecology*. doi:  
751 <https://doi.org/10.1111/jiec.12851>

752 Kogan, F., Kussul, N., Adamenko, T., Skakun, S., Kravchenko, O., Kryvobok, O., . . .  
753 Lavrenyuk, A. (2013). Winter wheat yield forecasting in Ukraine based on  
754 Earth observation, meteorological data and biophysical models. *International*  
755 *Journal of Applied Earth Observation and Geoinformation*, 23, 192-203.  
756 doi: <https://doi.org/10.1016/j.jag.2013.01.002>

757 Konar, M., Evans, T. P., Levy, M., Scott, C. A., Troy, T. J., Vörösmarty, C. J., &  
758 Sivapalan, M. (2016). Water resources sustainability in a globalizing world:  
759 Who uses the water? *Hydrological Processes*, 30(18), 3330–3336.  
760 <https://doi.org/10.1002/hyp.10843>

761 Kouka, P. J., Jolly, C. M., & Henao, J. (1994). Agricultural response functions for  
762 limited resource farmers in Sub-Saharan Africa. *Fertilizer research*, 40(2),  
763 135-141.

764 Kreibich, H., di Baldassarre, G., Vorogushyn, S., Aerts, J. C. J. H., Apel, H., Aronica,  
765 G. T., Arnbjerg - Nielsen, K., Bouwer, L. M., Bubeck, P., Caloiero, T., Chinh,  
766 D. T., Cortès, M., Gain, A. K., Giampá, V., Kuhlicke, C., Kundzewicz, Z. W.,  
767 Llasat, M. C., Mård, J., Matczak, P., Mazzoleni, M., Molinari, D., Dung, N. V.,

768 Petrucci, O., Schröter, K., Slager, K., Thieken, A. H., Ward, P. J., & Merz, B.  
769 (2017). Adaptation to flood risk: Results of international paired flood event  
770 studies. *Earth's Future*, 5(10), 953–965.  
771 <https://doi.org/10.1002/2017EF000606>

772 Lecerf, R., Ceglar, A., López-Lozano, R., Van Der Velde, M., & Baruth, B. (2019).  
773 Assessing the information in crop model and meteorological indicators to  
774 forecast crop yield over Europe. *Agricultural Systems*, 168, 191-202.  
775 doi: <https://doi.org/10.1016/j.agsy.2018.03.002>

776 Leegood, R. C., Sharkey, T. D., & Von Caemmerer, S. (2006). *Photosynthesis:  
777 physiology and metabolism (Vol. 9)*: Springer Science & Business Media.

778 Lemaire, G., Salette, J., Sigogne, M., & Terrasson, J.-P. (1984). Relation entre  
779 dynamique de croissance et dynamique de prélèvement d'azote pour un  
780 peuplement de graminées fourragères. I.--Etude de l'effet du milieu.  
781 *Agronomie*, 4(5), 423-430.

782 Levin-Koopman, J., Kuik, O., Tol, R., Van Der Vat, M., Hunink, J., & Brouwer, R.  
783 (2019). Distributing water between competing users in the Netherlands  
784 (Economy-Wide Modeling of Water at Regional and Global Scales by  
785 Springer; Presented at the 22nd Annual Conference on Global Economic  
786 Analysis, Warsaw, Poland). Purdue University, West Lafayette, IN: Global  
787 Trade Analysis Project (GTAP). Retrieved from  
788 [https://www.gtap.agecon.purdue.edu/resources/res\\_display.asp?RecordID=584](https://www.gtap.agecon.purdue.edu/resources/res_display.asp?RecordID=584)

790 Li, J. (2010). The Path Choice of Jiangsu's Agricultural Transformation and  
791 Upgrading. *Jiangsu Rural Economy* (305), 15-19.

792 Li, X., Zhang, X., Niu, J., Tong, L., Kang, S., Du, T., Li, S., & Ding, R. (2016).  
793 Irrigation water productivity is more influenced by agronomic practice factors  
794 than by climatic factors in Hexi Corridor, Northwest China. *Scientific  
795 reports*, 6, 37971. doi: <https://doi.org/10.1038/srep37971>

796 Liu, W., Hunsaker, D., Li, Y., Xie, X., & Wall, G. (2002). Interrelations of yield,  
797 evapotranspiration, and water use efficiency from marginal analysis of water  
798 production functions. *Agricultural Water Management*, 56(2), 143-151.  
799 doi: [https://doi.org/10.1016/S0378-3774\(02\)00011-2](https://doi.org/10.1016/S0378-3774(02)00011-2)

800 Lobell, D. B., & Burke, M. B. (2010). On the use of statistical models to predict crop  
801 yield responses to climate change. *Agricultural and forest  
802 meteorology*, 150(11), 1443-1452.  
803 doi: <https://doi.org/10.1016/j.agrformet.2010.07.008>

804 Ma, L., Feng, S., Reidsma, P., Qu, F., & Heerink, N. (2014). Identifying entry points  
805 to improve fertilizer use efficiency in Taihu Basin, China. *Land Use Policy*, 37,  
806 52-59. doi: <https://doi.org/10.1016/j.landusepol.2013.01.008>

807 McCarl, B. A. (1982). Cropping activities in agricultural sector models: a  
808 methodological proposal. *American Journal of Agricultural Economics*, 64(4),  
809 768-772.

810 Milly, P. C. D., Betancourt, J., Falkenmark, M., Hirsch, R. M., Kundzewicz, Z. W.,  
811 Lettenmaier, D. P., & Stouffer, R. J. (2008). Stationarity is dead: Whither

812 water management? Science, 319, 573–574.  
813 <https://doi.org/10.1126/science.1151915>

814 Mkhabela, M., Bullock, P., Raj, S., Wang, S., & Yang, Y. (2011). Crop yield  
815 forecasting on the Canadian Prairies using MODIS NDVI data. *Agricultural*  
816 *and Forest Meteorology*, 151(3), 385-393.  
817 doi: <https://doi.org/10.1016/j.agrformet.2010.11.012>

818 National Meteorological Information Center of China, *Crop growth and development*  
819 *and farmland soil moisture data set in China*. 2006.

820 Novoa, V., Ahumada-Rudolph, R., Rojas, O., Munizaga, J., Sáez, K., & Arumí, J. L.  
821 (2019). Sustainability assessment of the agricultural water footprint in the  
822 Cachapoal River basin, Chile. *Ecological indicators*, 98, 19-28. doi:  
823 <https://doi.org/10.1016/j.ecolind.2018.10.048>

824 Pande, S., Ertsen, M., & Sivapalan, M. (2014). Endogenous technological and  
825 population change under increasing water scarcity. *Hydrology and Earth*  
826 *System Sciences*, 18(8), 3239-3258.  
827 doi: <https://doi.org/10.5194/hess-18-3239-2014>

828 Parry, M. L. (2019). Climate change and world agriculture (Chapter 1: The Sensitivity  
829 of Agriculture to Climate). Routledge.

830 Pattanayak, S. K., & Sills, E. O. (2001). Do tropical forests provide natural insurance?  
831 The microeconomics of non-timber forest product collection in the Brazilian  
832 Amazon. *Land economics*, 77(4), 595-612. doi: 10.2307/3146943

833 Prihar, S., Gajri, P., & Arora, V. (1985). Nitrogen fertilization of wheat under limited

834 water supplies. *Fertilizer research*, 8(1), 1-8.  
835 doi: <https://doi.org/10.1007/BF01048902>

836 Raes, D., Steduto, P., Hsiao, T. C., & Fereres, E. (2009). AquaCrop—the FAO crop  
837 model to simulate yield response to water: II. Main algorithms and software  
838 description. *Agronomy Journal*, 101(3), 438-447.  
839 doi: [doi:10.2134/agronj2008.0140s](https://doi.org/10.2134/agronj2008.0140s)

840 Ren, J., Chen, Z., Zhou, Q., & Tang, H. (2008). Regional yield estimation for winter  
841 wheat with MODIS-NDVI data in Shandong, China. *International Journal of*  
842 *Applied Earth Observation and Geoinformation*, 10(4), 403-413.  
843 doi: <https://doi.org/10.1016/j.jag.2007.11.003>

844 Rodell, M., P.R. Houser, U. Jambor, J. Gottschalck, K. Mitchell, C.-J. Meng, K.  
845 Arsenault, B. Cosgrove, J. Radakovich, M. Bosilovich, J.K. Entin, J.P. Walker,  
846 D. Lohmann, and D. Toll, The Global Land Data Assimilation System, *Bull.*  
847 *Amer. Meteor. Soc.*, 85(3), 381-394, 2004.  
848 doi: <https://doi.org/10.1175/BAMS-85-3-381>

849 Roobavannan, M., Kandasamy, J., Pande, S., Vigneswaran, S., & Sivapalan, M.  
850 (2017a). Role of sectoral transformation in the evolution of water management  
851 norms in agricultural catchments: A sociohydrologic modeling analysis. *Water*  
852 *Resources Research*, 53, 8344–8365. <https://doi.org/10.1002/2017WR020671>

853 Roobavannan, M., Kandasamy, J., Pande, S., Vigneswaran, S., & Sivapalan, M.  
854 (2017b). Allocating environmental water and impact on basin unemployment:  
855 Role of a diversified economy. *Ecological Economics*, 136, 178-188. doi:



856 <https://doi.org/10.1016/j.ecolecon.2017.02.006>

857 Rosegrant, M. W., Ringler, C., McKinney, D. C., Cai, X., Keller, A., & Donoso, G.  
858 (2000). Integrated economic - hydrologic water modeling at the basin scale:  
859 The Maipo River basin. *Agricultural economics*, 24(1), 33-46.  
860 doi: <https://doi.org/10.1111/j.1574-0862.2000.tb00091.x>

861 Savenije, H. H. G., Hoekstra, A. Y., & van der Zaag, P. (2014). Evolving water  
862 science in the Anthropocene. *Hydrology and Earth System Sciences*, 18(1),  
863 319 - 332. <https://doi.org/10.5194/hess - 18 - 319 - 2014>

864 Sembiring, H., Raun, W., Johnson, G., Stone, M., Solie, J., & Phillips, S. (1998).  
865 Detection of nitrogen and phosphorus nutrient status in bermudagrass using  
866 spectral radiance. *Journal of plant nutrition*, 21(6), 1189-1206.  
867 doi: <https://doi.org/10.1080/01904169809365477>

868 Sivapalan, M., & Blöschl, G. (2015). Time scale interactions and the coevolution of  
869 humans and water. *Water Resources Research*, 51, 6988–7022.  
870 <https://doi.org/10.1002/2015WR017896>

871 Sivapalan, M., Konar, M., Srinivasan, V., Chhatre, A., Wutich, A., Scott, C. A.,  
872 Wescoat, J. L., & Rodriguez - Iturbe, I. (2014). Sociohydrology: Use -  
873 inspired water sustainability science for the Anthropocene. *Earth's Future*, 2,  
874 225 - 230. <https://doi.org/10.1002/2013EF000164>

875 Smith, M. (1992). *CROPWAT: A computer program for irrigation planning and*  
876 *management* (No. 46). Food & Agriculture Org.

877 Steduto, P., Hsiao, T. C., Raes, D., & Fereres, E. (2009). AquaCrop—The FAO crop

878 model to simulate yield response to water: I. Concepts and underlying  
879 principles. *Agronomy Journal*, 101(3), 426-437.  
880 doi: oi:10.2134/agronj2008.0139s

881 Vörösmarty, C. J., Green, P., Salisbury, J., & Lammers, R. B. (2000). Global water  
882 resources: vulnerability from climate change and population growth. *science*,  
883 289(5477), 284-288. doi: 10.1126/science.289.5477.284

884 Vörösmarty, C. J., Pahl - Wostl, C., Bunn, S. E., & Lawford, R. (2013). Global water,  
885 the Anthropocene and the transformation of a science. *Current Opinion in*  
886 *Environmental Sustainability*, 5(6), 539 - 550.  
887 <https://doi.org/10.1016/j.cosust.2013.10.005>

888 Wagener, T., Sivapalan, M., Troch, P. A., McGlynn, B. L., Harman, C. J., Gupta, H. V.,  
889 Kumar, P., Rao, P. S. C., Basu, N. B., & Wilson, J. S. (2010). The future of  
890 hydrology: An evolving science for a changing world. *Water Resources*  
891 *Research*, 46, W05301. <https://doi.org/10.1029/2009WR008906>

892 Xiao, X., Boles, S., Frohling, S., Salas, W., Moore Iii, B., Li, C., . . . Zhao, R. (2002).  
893 Landscape-scale characterization of cropland in China using Vegetation and  
894 Landsat TM images. *International Journal of Remote Sensing*, 23(18),  
895 3579-3594. doi: <https://doi.org/10.1080/01431160110106069>

896 Xin, H., Peiling, Y., Shumei, R., Yunkai, L., Guangyu, J., & Lianhao, L. (2016).  
897 Quantitative response of oil sunflower yield to evapotranspiration and soil  
898 salinity with saline water irrigation. *International Journal of Agricultural and*  
899 *Biological Engineering*, 9(2), 63-73. doi: 10.3965/j.ijabe.20160902.1683

