TU Delft

Delft University of Technology

Citation (APA)
Dai, S., Ma, Q., Zhu, S., Liu, M., Andersen, T., Fei, Z., Goldflam, M. D., Wagner, M., Watanabe, K., Taniguchi, T., Thiemens, M., Keilmann, F., Janssen, G. C. A. M., Jarillo-Herrero, P., Fogler, M., & Basov, D. N. (2015). *Tunable polaritons from plasmon-phonon coupling in hyperbolic media*. Abstract from APS March meeting 2015, San Antonia, TX, United States. http://meetings.aps.org/link/BAPS.2015.MAR.F2.6

Important note
To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright
In case the licence states “Dutch Copyright Act (Article 25fa)”, this publication was made available Green Open Access via the TU Delft Institutional Repository pursuant to Dutch Copyright Act (Article 25fa, the Taverne amendment). This provision does not affect copyright ownership.
Unless copyright is transferred by contract or statute, it remains with the copyright holder.

Sharing and reuse
Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy
Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

*This work is downloaded from Delft University of Technology.*

Abstract Submitted
for the MAR15 Meeting of
The American Physical Society

**Tunable polaritons from plasmon-phonon coupling in hyperbolic media** SIYUAN DAI, Univ of California - San Diego, QIONG MA, Massachusetts Institute of Technology, SHOU-EN ZHU, Delft University of Technology, Netherlands, MENGKUN LIU, University of California, San Diego, TROND ANDERSEN, Massachusetts Institute of Technology, ZHE FEI, MICHAEL GOLDFLAM, MARTIN WAGNER, Univ of California - San Diego, KENJI WATANABE, TAKASHI TANIGUCHI, National Institute for Materials Science, Japan, MARK THIEMENS, Univ of California - San Diego, FRITZ KEILMANN, Ludwig-Maximilians-Universität and Center for Nanoscience, G.C.A.M. JANSSEN, Delft University of Technology, Netherlands, PABLO JARILLO-HERRERO, Massachusetts Institute of Technology, MICHAEL FOGLER, D.N. BASOV, Univ of California - San Diego — Using infrared nano-imaging and nano-spectroscopy, we report on the tunable hyperbolic response in heterostructures comprised of a monolayer graphene deposited on hexagonal boron nitride (G-hBN). Electrostatic gating of the top graphene layer allows for modification of the wavelength and intensity of hyperbolic phonon polaritons in bulk hBN. When compared with the pristine hBN polaritons, the graphene modified ones exhibit a longer wavelength in the Type II hyperbolic region and shorter wavelength in the Type I region. Because of this modification, we achieve a 90% increase of the propagation length for Type II polaritons in hBN. The physical origin of the modification is attributed to plasmon-phonon coupling in the G-hBN heterostructure. Our work provides a comprehensive study of plasmon-phonon coupling in a hyperbolic medium with the exploration of graphene's potential for modification of collective modes in other materials.

Siyuan Dai
Univ of California - San Diego

Date submitted: 12 Nov 2014 Electronic form version 1.4