

## Quantifying Within-Individual Elbow Load Variability in Youth Elite Baseball Pitchers and Its Role in Overuse Injuries

van Trigt, B.; Bouman, F.F.; Leenen, Antonius J. R. ; Hoozemans, Marco J.M.; van der Helm, F.C.T.; Veeger, H.E.J.

**DOI**

[10.3390/app12136549](https://doi.org/10.3390/app12136549)

**Publication date**

2022

**Document Version**

Final published version

**Published in**

Applied Sciences

**Citation (APA)**

van Trigt, B., Bouman, F. F., Leenen, A. J. R., Hoozemans, M. J. M., van der Helm, F. C. T., & Veeger, H. E. J. (2022). Quantifying Within-Individual Elbow Load Variability in Youth Elite Baseball Pitchers and Its Role in Overuse Injuries. *Applied Sciences*, 12(13), Article 6549. <https://doi.org/10.3390/app12136549>

**Important note**

To cite this publication, please use the final published version (if applicable). Please check the document version above.

**Copyright**

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

**Takedown policy**

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

## Article

# Quantifying Within-Individual Elbow Load Variability in Youth Elite Baseball Pitchers and Its Role in Overuse Injuries

Bart Van Trigt <sup>1,\*</sup>, Foskien F. Bouman <sup>1</sup>, Antonius J. R. Leenen <sup>2</sup>, Marco J. M. Hoozemans <sup>2</sup>, Frans C. T. Van der Helm <sup>1</sup> and Dirkjan Veeger <sup>1</sup>

<sup>1</sup> Department of Biomechanical Engineering, Delft University of Technology, 2628 CD Delft, The Netherlands; foskienbouman@gmail.com (F.F.B.); f.c.t.vanderhelm@tudelft.nl (F.C.T.V.d.H.); h.e.j.veeger@tudelft.nl (D.V.)

<sup>2</sup> Department of Human Movement Sciences, Faculty of Behavioural and Movement Sciences, Vrije Universiteit Amsterdam, Amsterdam Movement Sciences, 1081 HV Amsterdam, The Netherlands; a.j.r.leenen@vu.nl (A.J.R.L.); m.j.m.hoozemans@vu.nl (M.J.M.H.)

\* Correspondence: b.vantrigt@tudelft.nl

**Abstract:** Medial elbow overuse injuries are rising in baseball. The external valgus torque magnitude is a possible risk factor for medial elbow injuries. The magnitude on its own cannot explain why one pitcher sustains an injury and another does not. Therefore, the aim of this study is to describe the within-individual external valgus torque variability and to determine whether the within-individual external valgus torque variability can be described by a Gaussian distribution. Eleven youth elite baseball pitchers threw twenty-five fastball pitches. Body kinematics were measured with VICON motion capture at 400 Hz. Elbow valgus torques of the total 270 pitches were calculated with a custom-made inverse dynamic model in Python. Visual inspection and the Shapiro–Wilk test were performed to test for the within-individual elbow valgus torque normality. The results showed that within-individual valgus torque variability was present in pitchers and differed among pitchers. Furthermore, it was shown that the within-individual valgus torque variability was normally distributed in nine out of eleven subjects. In conclusion, the presence of and differences in within-individual elbow load variability among baseball pitchers can be useful variables as they might be related to overuse elbow injuries.

**Keywords:** fastball; biomechanics; valgus torque; kinetics; ulnar collateral ligament (UCL); injury prevention

**Citation:** van Trigt, B.; Bouman, F.F.; Leenen, A.J.R.; Hoozemans, M.J.M.; van der Helm, F.C.T.; Veeger, D. Quantifying Within-Individual Elbow Load Variability in Youth Elite Baseball Pitchers and Its Role in Overuse Injuries. *Appl. Sci.* **2022**, *12*, 6549. <https://doi.org/10.3390/app12136549>

Academic Editor: Enrique Navarro

Received: 25 May 2022

Accepted: 23 June 2022

Published: 28 June 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

In baseball, faster and more accurate pitches increase the chances of beating the batter and thus winning the game. Professional pitchers can throw a baseball, accurately, at speeds over 100 mph. Fast pitches—often at maximal effort—produce high loads on the body, especially on the throwing arm. The pitchers' body parts are exposed to these high loads repetitively in practice and competition. The combination of these high loads and the repetitive nature of the pitch motion induces upper extremity injuries [1]. The most common upper extremity injury is the medial Ulnar Collateral Ligament (UCL) sprain or tear. In Major League Baseball, the highest level of baseball competition in the USA, 25% of the professional pitchers sustain a UCL injury during their career [2]. In addition, UCL reconstructions—a treatment modality for UCL tears—are reported with a yearly incidence of 4.4% in collegiate pitchers [3]. The number of UCL reconstructions has increased substantially at all levels of play over recent years [4]. It is therefore important to prevent athletes from injuries such as UCL injury.

In a systematic review, Bullock et al. [5] investigated biomechanical factors that are associated with elbow injuries. They concluded that elbow valgus torque and early trunk

rotation are positively related to UCL injuries. In addition, in a professional group of pitchers, it was found that maximum ball speed is associated with UCL injuries [6]. Thus, reducing elbow valgus torque while maintaining ball speed—for example, by the use of a “proper” kinetic chain [7,8]—would be the best way of reducing UCL injuries. While the positive association between valgus torque and ball speed at group level is weak, it is considerably stronger on the individual level [9]. Reducing elbow valgus torque without loss of performance will, however, be complex at the least.

As ball speed and elbow load (i.e., elbow valgus torque) are positively associated with UCL injuries, one could argue that recreational pitchers, who throw slower and with less elbow load, should report fewer UCL injuries compared with professional pitchers. However, this lower level of play also shows a high prevalence of UCL injuries [3]. In addition, within a homogenous group such as professional baseball pitchers, the average individual level of ball speed or elbow load cannot explain why 25% sustain a UCL injury and others do not, assuming that the individual magnitude is similar in this homogenous group of professionals. Most of the aforementioned studies analyzed a single pitch [10] or the average of the fastest three–five pitches [11,12], neglecting to consider the within-individual variability in ball speed and elbow load. Assuming that reaching a critical (peak) value of valgus torque is a major determinant in UCL injury, this magnitude will depend on the average torque level as well as the within-individual variability of this torque [13]. Information on both parameters is therefore necessary.

A simple explanatory computational model illustrates that a higher within-individual (elbow) load variability increases (elbow) injury risk [13]. We modeled the load as a Gaussian distribution with an average, standard deviation, and different numbers of pitches as inputs. In this model, the average and standard deviation are interpreted as the load magnitude and within-individual load variability, respectively. Practice shows that individual pitchers throw baseballs at various speeds. It is, therefore, also likely that the various individual pitches (within for instance a game, practice, or season) are associated with a variation in (peak) elbow load. However, the characteristics of the distributions of the within-individual variability of the elbow load are unknown. If, for instance, it can be established that these distributions of individual variability can be described as Gaussian distributions, individual values of means and standard deviations of elbow load can be used to model each individual pitcher’s risk of suffering an elbow injury.

To date, it is still not possible to measure elbow load, and more specifically, UCL load, during pitching. However, inverse dynamic models can estimate the external valgus torque as the best proxy for UCL load [5,14], which can thus be used as a parameter for the study of within-individual variability in elbow load.

Therefore, the aim of this study was to describe the within-individual external valgus torque variability of full effort fastball pitches of elite youth baseball pitchers and determine whether the within-individual external valgus torque variability can be described by a Gaussian distribution.

## 2. Materials and Methods

### 2.1. Participants

Data were collected from eleven male Dutch national (AAA) pitchers, at a mean age of 17.4 years (SD 2.2, range 15–23). Mean body mass was 80.6 kg (SD 11.7, range 68.2–107.0), mean body height was 186.7 cm (SD 6.3, range 177.0–194.0), and mean ball speed was 76.6 mph (SD 3.19, range 67–83). These elite youth pitchers are the best pitchers of their age group in the Netherlands. Nine pitchers threw with their right hand, while two pitchers threw with their left. All participants were healthy and experienced no pain, soreness, or range-of-motion restrictions. Before involvement in the study, participants were informed of the procedure and study aims and informed consent forms were signed. This research was conducted in accordance with the Declaration of Helsinki and the local ethics

committee of the Faculty of Behavioural and Movement Sciences at the Vrije Universiteit Amsterdam approved the measurement protocol (reference number: VCWE-2019-033).

## 2.2. Procedure

The measurements were performed at the campus indoor facility of the Royal Dutch Football Association. The pitchers wore sneakers, athletic stretch shorts, catching gloves, and no shirts. Forty-three reflective markers (10 mm diameter) were attached with double-sided tape directly on the skin at bony landmarks. The pitchers were given an unlimited amount of time to perform their warm-up, including running, stretching, and a specific throwing session. Next, they threw several pitches off the mound to become familiar with the research setup. Once the pitchers were ready, they were instructed to throw fastballs as fast and as accurately as possible. The time between each pitch was not controlled but regulated by the pitcher himself, like in a normal game. To investigate the within-individual load variability, the pitchers threw 25 fastballs. The pitchers threw from a pitching mound towards a square strike zone (height 0.64 m; width 0.38 m) 0.55 m above the ground at a regular game distance of 18.44 m.

## 2.3. Data Acquisition

Marker positions were captured using a Vicon eight-camera motion capture system (model V5; Vicon Motion Systems Ltd., Yarnton, UK) sampled at 400 Hz and stored on a local computer. The ball speed of each pitch was captured from a position next to the strike zone with a radar gun (Stalker Radar, Plano, TX, USA) pointing in the direction of the pitcher.

## 2.4. Data Analysis

Three-dimensional marker position data were withdrawn in  $x$ ,  $y$ , and  $z$  coordinates from the Vicon system, and all the calculations were performed in Python [15]. All data were analyzed between foot contact and ball release, which includes the critical moment of peak external valgus torque. Foot contact was defined as the moment that the forward velocity of the toe was smaller than 0.3 m/s. Ball release was defined as the moment that the wrist exceeded the position of the elbow in the forward direction. The data were interpolated with a 3rd order cubic spline polynomial and filtered with a 4th order Butterworth filter with a cut-off frequency of 12.5 Hz. If a marker flew off before ball release or if a marker could not correctly be reconstructed, the corresponding pitch was not included. Each trial was visually checked for errors and mistakes. Additionally, participant 11 was not allowed, due to pitching restrictions in his training schedule, to throw more than 20 full-effort pitches. Altogether, 250 of the total 270 pitches performed were used for analysis.

## 2.5. Inverse Dynamics

For the hand and forearm, an anatomical local coordinate system was made according to the ISB recommendations [16]. The following bony landmarks on the throwing arm were used: third proximal interphalangeal, ulna processes styloid, radius processes styloid, lateral humeral epicondyle, and medial humeral epicondyle. Positions of the centers of mass and the moments of inertia were estimated according to Zatsiorsky (2002) [17] and De Leva et al. (1996) [18]. The baseball was modeled with a mass of 0.145 kg attached to the hand, where the ball mass linearly reduced by 10% over the last ten samples (0.025 s) before ball release. The elbow joint torque was expressed in the anatomical local coordinate system of the elbow, located in the middle of the lateral and medial humeral epicondyle. Using this anatomical coordinate system with the segment data and with the scaling factors of De Leva et al. (1996) and Zatsiorsky et al. (1990), the kinetics of the segments were calculated [17–19]. Elbow joint torques were calculated using the top-down inverse dynamic analysis based on the Newton–Euler equation of motions, starting in the

hand of the throwing arm. The external elbow valgus torque was calculated as a time series for each single pitch between foot contact and ball release. Subsequently, the peak external valgus torque was obtained from this time series.

### 2.6. Statistical Analysis

The peak external valgus torque values of the series of pitches of each participant were visually checked for normality with Q–Q plots and tested with the Shapiro–Wilk test. A ridgeline density plot was made for the distribution of the external valgus torque for each participant. The ridgeline density plot included the original data of the external valgus torque. The measures of central tendency and dispersion were calculated. If the data were normally distributed, the standard deviation was used to indicate within-individual elbow valgus torque variability. In addition, a normal distribution was calculated from the external valgus torque mean and standard deviation for each participant and added to the figure with the individual density plots. All statistical analyses were performed in Rstudio (version 2022.2.0.443) [20].

## 3. Results


### *Within-Individual Valgus Torque Magnitude and Variability*

Figure 1 presents the density plots and normal distributions of the (peak) external elbow valgus torque values for the series of 17–25 fastball pitches for each of the eleven participants. The descriptives and normality test results are shown in Table 1. The Shapiro–Wilk test revealed that the external valgus torque scores in nine of the eleven participants were normally distributed. The scores of participants 6 and 9 did not show a normal distribution. The scores of participant 6 were positively skewed and the scores of participant 9 were negatively skewed. The skewness of both participants is visualized in Figure 1. The Q–Q plots of the eleven participants are attached in the supplementary materials.

Visual inspection of Figure 1 and the measures of central tendency and dispersion in Table 1 show differences between pitchers in within-individual external elbow valgus torque variability and the actual location of the distribution (i.e., overall elbow load magnitude).

**Table 1.** Descriptives of the set of scores for each participant for their series of (peak) external elbow valgus torque. The two columns on the right show the results of the Shapiro–Wilk test. SD = standard deviation; IQR = inter quartile range; \* = significant.

Participant	Analyzed Pitches	Mean	Median	SD	IQR	Skewness	Kurtosis	Shapiro–Wilk Test	
								Sig.	
1	24	58.91	59.86	4.40	6.55	−0.07	−1.27	0.93	0.11
2	25	53.40	53.61	2.07	3.04	−0.73	0.15	0.95	0.27
3	25	36.35	36.58	2.34	3.25	0.17	−0.26	0.98	0.83
4	23	43.68	43.55	1.34	2.00	0.20	−0.68	0.98	0.94
5	24	47.07	47.21	2.82	4.23	0.39	−0.52	0.97	0.68
6	19	59.68	58.79	4.08	4.15	1.11	0.57	0.88	0.02 *
7	23	48.67	48.96	2.55	4.65	−0.51	−0.41	0.96	0.39
8	23	64.54	65.10	3.25	3.33	−1.01	1.65	0.94	0.16
9	23	51.53	52.69	2.05	3.22	−0.76	−0.71	0.87	0.01 *
10	24	68.51	68.28	3.25	2.98	0.08	0.71	0.97	0.61
11	17	52.77	53.31	1.96	1.97	−1.21	1.44	0.90	0.07



**Figure 1.** Density plots and matching normal distributions of the within-individual variability in external elbow valgus torque. The participants are shown on the y-axis and the external elbow valgus torque on the x-axis. Each color represents a participant. The dots show the original data of each external valgus torque of a single pitch. The height of the dots is randomly jittered in the density plot. The three vertical solid lines inside the density plots show the 25th, 50th, and 75th percentile of the distribution, respectively. The thin line inside the density plot shows the modeled normal distribution based on the distribution's mean and standard deviation.


#### 4. Discussion

The aim of this study was to quantify the within-individual elbow valgus torque variability when pitching fastballs in elite youth baseball pitchers. The results showed that within-individual elbow valgus torque variability is present in pitchers and differed among pitchers. In the majority of these pitchers, the variability can be described as a Gaussian distribution.

Pitching at full effort produces slightly different elbow valgus torques between fastball pitches within a pitcher. As expected, and in line with other studies, the magnitude (average) of the elbow valgus torque was different between pitchers, showing the between-pitcher variability [8,11,21]. The group average elbow valgus torque in this study population was 53.19 Nm (SD 9.34; range: 36.35–68.5). Similar valgus torques were reported in other studies with comparable ball speeds and study population [8,11]. In these studies, the valgus torque magnitude differences between pitchers has been regularly reported, but the within-individual valgus torque variability has mostly been neglected. In our study, we show that this within-individual valgus torque variability is present in pitchers and differs between pitchers ranging from 1.34 Nm to 4.40 Nm (Figure 1). All pitchers were instructed to throw 25 full effort fastballs, and some pitchers were able to throw with less valgus torque variability compared with others. It is known that within-individual variability is smaller compared with between-individual variability [21]. The magnitude (average) external valgus torque between pitchers can be explained mainly by personal characteristics (body length, mass), motivation, well-being, fatigue, and technique (kinetic chain, energy flow), whereas the within-individual valgus torque

variability can mainly be explained by motivation, technique, and fatigue. So, in addition to the magnitude (average) valgus torque, the within-individual valgus torque variability is also different between pitchers.

A higher valgus torque is considered to lead to an increased UCL injury risk, according to previous research [8,10,22]. Within a homogenous group of professional pitchers, 25% sustained a UCL injury [2]. The valgus torque magnitude on its own cannot completely explain this percentage, as all professional pitchers show high valgus torques. To explain this prevalence, the within-individual load variability and the injury threshold are both important metrics to include [13]. The injury threshold in this model is the maximal load the UCL can sustain before it strains or tears. The demonstrated within-individual valgus torque variability and the fact that this differs between subjects is a step towards validating the injury model of van Trigt et al. (2020). However, it is suggested that pitchers who throw with a higher magnitude and within-individual variability of valgus torque might have—but do not necessarily have—a higher chance of sustaining an injury, because the injury threshold is also of importance [13] (Figure 2). It is important to mention that we did not investigate the relationship between valgus torque variability and injuries.



**Figure 2.** Schematic overview of within-individual elbow load variability (width of the normal distributions) and injury thresholds (red vertical lines). All four quadrants have the same load magnitude (average), shown as the vertical dashed lines. The horizontal axis of the quadrant shows the load variability, and the vertical axis shows the load capacity. The state on the left top panel shows the lowest chance, the right top and left bottom have the same chance, and the right bottom has the highest chance of sustaining an injury. Be aware that a shift in the load magnitude is also possible, but not represented here.


In the model of van Trigt et al. (2020), a load distribution is formed with a load average and standard deviation. The results showed that the elbow load distribution in nine out of eleven pitchers could be described as a Gaussian distribution. The standard deviation of this Gaussian distribution can be used as a parameter to quantify the within-individual load variability. This study shows that, based on 25 thrown pitches, an individual load distribution can be built with the within-individual load variability and the load magnitude (average), which can be used for injury assessment.

Longitudinal studies have shown that fatigue influences injury risk [23,24]. Thus, it is too simplistic to assume that the load magnitude, load variability, and injury threshold are fixed parameters over time and are not influenced by, for example, fatigue. It was found that the development of fatigue over time increases the external valgus torque magnitude [25], indicating a shift of the load distribution toward the injury threshold.



However, it is unknown whether fatigue also influences the within-individual load variability on a micro-level (within a session) or macro-level (between sessions). Executing a study that investigates fatigue in relation to load variability and the effects on an estimated injury threshold based on our model could provide us with new insights.

In terms of practical implications, the presence of within-individual load variability raises the question of whether this load variability is also related to performance. We, therefore, performed a post hoc analysis and analyzed how the pitchers with load variability were distributed in a graph displaying the relationship between ball speed and external valgus torque (Figure 3). Each dot in Figure 3 represents a single pitch and each color represents a pitcher, corresponding with the colors in Figure 1. On a group level, a weak relationship between external valgus torque and ball speed is present. When individual characteristics are included, this results in a much stronger relationship, which is in agreement with the study of Slowik et al. (2019) [9]. The heights of the ellipses, indicating the within-individual load variability, are randomly scattered in this graph. Linear regression showed that, unlike the valgus torque magnitude, the within-individual valgus torque variability is unrelated to the pitcher average ( $p = 0.309$ ,  $R^2 = 0.1146$ ) or maximal ( $p = 0.195$ ,  $R^2 = 0.1791$ ) ball speed. Thus, based on these preliminary results, it seems that within-individual elbow load variability is not related to performance in this group of pitchers. In practical terms, this means that a pitcher should strive to minimize their elbow load variability as it might reduce the chance of sustaining an injury without influencing performance. It is known that a “proper” technique is related to the elbow valgus torque [7], and thus it might also be related to the load variability. For injury prevention, it should be investigated whether a pitcher is able to minimize their elbow load variability consciously.



**Figure 3.** The relationship between ball speed and valgus torque. Every dot is a single pitch of a participant. Each color represents a participant, corresponding with the colors in Figure 1. The ellipses show the within-individual external valgus torque variability and the ball speed variability. The height of the ellipse is twice the standard deviation of individual scores of the external valgus



torque, and the width of the ellipse is twice the ball speed standard deviation. The center of the ellipse is the mean of the scores of the external elbow valgus torque and ball speed of a single participant.

The study population was a homogenous group of youth elite baseball pitchers (age 15–22). It is known that the magnitude of external elbow valgus torque increases with the level of play [11]. The group average valgus torque variability did not show significant differences between levels of play according to Fleisig et al. (2009) [26], indicating that our results showing differences between pitchers in within-individual load variability can be extended to other levels of play. A large difference between our study and the study of Fleisig et al. (2009) is that they only included the 5 fastest pitches in their analysis to describe within-individual variability, which is less compared to the 25 analyzed pitches in our study. Five pitches are too few to define whether the valgus torque is normally distributed. Therefore, it is unclear whether the finding that a load distribution can be described with a Gaussian distribution can be extended to other levels of play.

A limitation of this study is that the external valgus torque magnitude and variability are assumed to be representative of the magnitude and variability of the UCL load. The literature shows that, in addition to the UCL, elbow muscles and joint articulation can also counteract the external valgus torque [27]. How the magnitude and variability of the valgus torque load are distributed over these anatomical structures is yet unknown. Thus, how much of the valgus torque magnitude and variability can be translated to the UCL load and variability should be investigated in the future. In conclusion, as it is known that the external valgus torque is related to UCL load and UCL injuries [8,14], we conclude that UCL load variability is present within baseball pitchers. However, the exact values of magnitude and variability depend on other elbow stabilizers and the individual pitcher.

## 5. Conclusions

In conclusion, this study showed that within-individual elbow load variability is present in youth elite baseball pitchers and that the within-individual load variability differs between pitchers. It is possible to describe the elbow load distribution with a Gaussian distribution, in which the standard deviation describes the within-individual load variability. Therefore, in addition to the load magnitude, the within-individual load variability seems important in injury assessment and possibly in injury prevention. To make it relevant to injury prevention, future research should investigate whether load variability changes over time and whether it can be minimized by pitchers.

**Supplementary Materials:** The following supporting information can be downloaded at: [www.mdpi.com/article/10.3390/app12136549/s1](http://www.mdpi.com/article/10.3390/app12136549/s1), Figure S1: Q–Q plots external valgus torque.

**Author Contributions:** Conceptualization, B.V.T. and D.V.; methodology, B.V.T. and M.J.M.H.; software, B.V.T. and A.J.R.L.; formal analysis, B.V.T. and F.F.B.; investigation, B.V.T. and A.J.R.L.; resources, M.J.M.H.; data curation, B.V.T. and A.J.R.L.; writing—original draft preparation, B.V.T. and F.F.B.; writing—review and editing, B.V.T., M.J.M.H., A.J.R.L. and D.V.; visualization, B.V.T. and F.F.B.; supervision, D.V., F.C.T.V.d.H. and M.J.M.H.; project administration, D.V.; funding acquisition, D.V., M.J.M.H. and F.C.T.V.d.H. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was supported by the NWO Domain Applied and Engineering Sciences (AES) under project number [R/003635]. This NWO-funded project, named Breaking the High Load—Bad Coordination Multiplier in Overhead Sports Injuries part of the Citius Altius Sanius perspective program (Project 7), is a cooperative effort between the Royal Dutch Baseball and Softball Federation (KNBSB), Royal Dutch Tennis Federation (KNLTB), Vrije Universiteit Amsterdam, Delft University of Technology, Milé Fysiotherapy, PitchPerfect and PLUX.

**Institutional Review Board Statement:** The study was conducted in accordance with the Declaration of Helsinki and approved by the Institutional Review Board (or Ethics Committee) of Vrije Universiteit Amsterdam (protocol code VCWE-2019-033 and date 1 April 2019).

**Informed Consent Statement:** Informed consent was obtained from all subjects involved in the study.

**Data Availability Statement:** The data of this study can be found at doi:10.4121/19868722.

**Acknowledgments:** The authors would like to thankfully acknowledge the Dutch Royal Football Association (KNVB) for facilitating the research program by giving access to their research facilities. We also would like to thank Yvette Leeftink and Lars Welten for helping during the experiments. Thanks to all the pitchers that participated in this study.

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; or in the writing of the manuscript, or in the decision to publish the results.

## References

- Fortenbaugh, D.; Fleisig, G.S.; Andrews, J.R. Baseball Pitching Biomechanics in Relation to Injury Risk and Performance. *Sports Health* **2009**, *1*, 314–320. <https://doi.org/10.1177/1941738109338546>.
- Conte, S.A.; Fleisig, G.S.; Dines, J.S.; Wilk, K.E.; Aune, K.T.; Patterson-flynn, N.; ElAttrache, N. Prevalence of Ulnar Collateral Ligament Surgery in Professional Baseball Players. *Am. J. Sports Med.* **2015**, *43*, 1764–1769. <https://doi.org/10.1177/0363546515580792>.
- Rothermich, M.A.; Conte, S.A.; Aune, K.T.; Fleisig, G.S.; Jr, E.L.C.; Dugas, J.R.; Cain, E.L.; Dugas, J.R. Incidence of Elbow Ulnar Collateral Ligament Surgery in Collegiate Baseball Players. *Orthop. J. Sports Med.* **2018**, *6*, 2325967118764657. <https://doi.org/10.1177/2325967118764657>.
- Zaremski, J.L.; Vincent, K.R.; Vincent, H.K. Elbow Ulnar Collateral Ligament: Injury, Treatment Options, and Recovery in Overhead Throwing Athletes. *Curr. Sports Med. Rep.* **2019**, *18*, 338–345.
- Bullock, G.S.; Menon, G.; Nicholson, K.; Butler, R.J.; Arden, N.K.; Filbay, S.R. Baseball Pitching Biomechanics in Relation to Pain, Injury, and Surgery: A Systematic Review. *J. Sci. Med. Sport* **2021**, *24*, 13–20. <https://doi.org/https://doi.org/10.1016/j.jsams.2020.06.015>.
- Bushnell, B.D.; Anz, A.W.; Noonan, T.J.; Torry, M.R.; Hawkins, R.J.; Carolina, S. Association of Maximum Pitch Velocity and Elbow Injury in Professional Baseball Pitchers. *Am. J. Sports Med.* **2010**, *38*, 728–732. <https://doi.org/10.1177/0363546509350067>.
- Urbini, M.A.; Fleisig, G.S.; Abebe, A.; Andrews, J.R. Associations between Timing in the Baseball Pitch and Shoulder Kinetics, Elbow Kinetics, and Ball Speed. *Am. J. Sports Med.* **2013**, *41*, 336–342. <https://doi.org/10.1177/0363546512467952>.
- Aguinaldo, A.L.; Chambers, H. Correlation of Throwing Mechanics with Elbow Valgus Load in Adult Baseball Pitchers. *Am. J. Sports Med.* **2009**, *37*, 2043–2048. <https://doi.org/10.1177/0363546509336721>.
- Slowik, J.S.; Aune, K.T.; Diffendaffer, A.Z.; Cain, E.L.; Dugas, J.R.; Fleisig, G.S. Fastball Velocity and Elbow-Varus Torque in Professional Baseball Pitchers. *J. Athl. Train.* **2019**, *54*, 296–301.
- Sabick, M.B.; Torry, M.R.; Lawton, R.L.; Hawkins, R.J.; Frics, C. Valgus Torque in Youth Baseball Pitchers: A Biomechanical Study. *J. Shoulder Elb. Surg.* **2004**, *13*, 349–355. <https://doi.org/10.1016/j.jse.2004.01.013>.
- Fleisig, G.S.; Barrentine, S.W.; Zheng, N.; Escamilla, R.F.; Andrews, J.R. Kinematic and Kinetic Comparison of Baseball Pitching among Various Levels of Development. *J. Biomech.* **1999**, *32*, 1371–1375. [https://doi.org/10.1016/S0021-9290\(99\)00127-X](https://doi.org/10.1016/S0021-9290(99)00127-X).
- Aguinaldo, A.; Escamilla, R. Segmental Power Analysis of Sequential Body Motion and Elbow Valgus Loading During Baseball Pitching: Comparison Between Professional and High School Baseball Players. *Orthop. J. Sports Med.* **2019**, *7*, 232596711982792. <https://doi.org/10.1177/232596711982792>.
- van Trigt, B.; Leenen, A.J.R.; Hoozemans, M.J.M.; van der Helm, F.C.T.; Veeger, H.E.J.; van Trigt, B.; Leenen, T.; Hoozemans, M.J.M.; Helm, F. van der; Veeger, D. Are UCL Injuries a Matter of Bad Luck?: The Role of Variability and Fatigue Quantified. *Proceedings* **2020**, *49*, 107. <https://doi.org/10.3390/proceedings2020049107>.
- Anz, A.W.; Bushnell, B.D.; Griffin, L.P.; Noonan, T.J.; Torry, M.R.; Richard, J.; Hawkins, R.J.; Carolina, S. Correlation of Torque and Elbow Injury in Professional Baseball Pitchers. *Am. J. Sports Med.* **2010**, *38*, 1368–1374. <https://doi.org/10.1177/0363546510363402>.
- Van Rossum, G.; Drake, F.L. *Python 3 Reference Manual*; CreateSpace: Scotts Valley, CA, USA, 2009.
- Wu, G.; Van Der Helm, F.C.T.; Veeger, H.E.J.; Makhssous, M.; Van Roy, P.; Anglin, C.; Nagels, J.; Karduna, A.R.; McQuade, K.; Wang, X.; et al. ISB Recommendation on Definitions of Joint Coordinate Systems of Various Joints for the Reporting of Human Joint Motion-Part II: Shoulder, Elbow, Wrist and Hand. *J. Biomech.* **2005**, *38*, 981–992. <https://doi.org/10.1016/j.jbiomech.2004.05.042>.
- Zatsiorsky, V.M. *Kinetics of Human Motion*; Human Kinetics: Champaign, IL, USA, 2002.
- de Leva, P. Adjustments to Zatsiorsky-Seluyanov's Segment Inertia Parameters. *J. Biomech.* **1996**, *29*, 1223–1230.
- Zatsiorsky, V.M.; Seluyanov, V.N.; Chugunova, L.G. Methods of Determining Mass-Inertial Characteristics of Human Body Segments. In *Contemporary Problems of Biomechanics*; CRC Press: Boca Raton, FL, USA, 1990; pp. 272–291.
- RStudio Team. *RStudio: Integrated Development for R*; RStudio: Boston, MA, USA, 2020.
- Escamilla, R.F.; Barrentine, S.W.; Fleisig, G.S.; Zheng, N.; Takada, Y.; Kingsley, D.; Andrews, J.R. Pitching Biomechanics as a Pitcher Approaches Muscular Fatigue during a Simulated Baseball Game. *Am. J. Sports Med.* **2007**, *35*, 23–33.

22. Wilk, K.E.; Macrina, L.C.; Fleisig, G.S.; Aune, K.T.; Porterfield, R.A.; Harker, P.; Evans, T.J.; Andrews, J.R. Deficits in Glenohumeral Passive Range of Motion Increase Risk of Elbow Injury in Professional Baseball Pitchers A Prospective Study. *Am. J. Sports Med.* **2014**, *42*, 2075–2081. <https://doi.org/10.1177/0363546514538391>.
23. Olsen, S.J.; Fleisig, G.S.; Dun, S.; Loftice, J.; Andrews, J.R. Risk Factors for Shoulder and Elbow Injuries in Adolescent Baseball Pitchers. *Am. J. Sports Med.* **2006**, *34*, 905–912.
24. Yang, J.; Mann, B.J.; Guettler, J.H.; Dugas, J.R.; Irrgang, J.J.; Fleisig, G.S.; Albright, J.P. Risk-Prone Pitching Activities and Injuries in Youth Baseball: Findings from a National Sample. *Am. J. Sports Med.* **2014**, *42*, 1456–1463. <https://doi.org/10.1177/0363546514524699>.
25. Okoroha, K.R.; Meldau, J.E.; Lizzio, V.A.; Meta, F.; Stephens, J.P.; Moutzouros, V.; Makhni, E.C. Effect of Fatigue on Medial Elbow Torque in Baseball Pitchers: A Simulated Game Analysis. *Am. J. Sports Med.* **2018**, *46*, 2509–2513. <https://doi.org/10.1177/0363546518782451>.
26. Fleisig, G.; Chu, Y.; Weber, A.; Andrews, J. Variability in Baseball Pitching Biomechanics among Various Levels of Competition. *Sports Biomech.* **2009**, *8*, 10–21. <https://doi.org/10.1080/14763140802629958>.
27. Van Trigt, B.; Vliegen, L.; Leenen, T.; Veeger, D. The Ulnar Collateral Ligament Loading Paradox between In-Vitro and in-Vivo Studies on Baseball Pitching (Narrative Review). *Int. Biomech.* **2021**, *8*, 19–29. <https://doi.org/10.1080/23335432.2021.1916405>.