

PFA and ω_1 -free compact spaces

Hart, K.P.; Dow, Alan

DOI

[10.1007/s10474-021-01197-9](https://doi.org/10.1007/s10474-021-01197-9)

Publication date

2022

Document Version

Final published version

Published in

Acta Mathematica Hungarica

Citation (APA)

Hart, K. P., & Dow, A. (2022). PFA and ω_1 -free compact spaces. *Acta Mathematica Hungarica*, 166(1), 57-64. <https://doi.org/10.1007/s10474-021-01197-9>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Green Open Access added to TU Delft Institutional Repository

'You share, we take care!' - Taverne project

<https://www.openaccess.nl/en/you-share-we-take-care>

Otherwise as indicated in the copyright section: the publisher is the copyright holder of this work and the author uses the Dutch legislation to make this work public.



PFA AND ω_1 -FREE COMPACT SPACES

A. DOW¹ and K. P. HART^{2,*}

¹Department of Mathematics, UNC-Charlotte, 9201 University City Blvd.,
Charlotte, NC 28223-0001, USA
e-mail: adow@uncc.edu

²Faculty EEMCS, TU Delft, Postbus 5031, 2600 GA Delft, the Netherlands
e-mail: k.p.hart@tudelft.nl

(Received May 10, 2021; revised October 9, 2021; accepted October 25, 2021)

Abstract. The Proper Forcing Axiom implies that compact Hausdorff spaces are either first-countable or contain a converging ω_1 -sequence.

Introduction

This paper continues the investigation from [5] into a conjectured dichotomy for compact Hausdorff spaces: every such space is either first-countable or it has a non-trivial converging ω_1 -sequence. Whether this dichotomy holds was asked by Juhász and a consistent counterexample to this conjecture was given by him, Koszmider, and Soukup in [8].

The main result in [5] states that the dichotomy holds in a large class of ccc forcing extensions of models in which the Continuum Hypothesis holds. This result does not encompass all ccc extensions and one can rightfully ask what the status of the dichotomy is under Martin's Axiom.

In Section 2 we point out that a space constructed by the first author in [4] witnesses that the combination $\text{MA} + \mathfrak{c} = \aleph_2$ is not strong enough to imply the dichotomy.

Our main result appears in Section 3: the Proper Forcing Axiom implies the dichotomy. This then also implies that the dichotomy is consistent with and independent of $\text{MA} + \mathfrak{c} = \aleph_2$. It also improves upon the first author's result in [3, Theorem 5.1], which implies that, in turn, PFA implies that a compact space without convergent ω_1 -sequences must be Fréchet–Urysohn.

* Corresponding author.

Key words and phrases: PFA, compact, convergence, ω_1 -sequence, first-countable.

Mathematics Subject Classification: primary 54D30, secondary 03E35, 03E50, 03E57, 54A20, 54A35, 54D20.



1. Some preliminaries

Here we collect the notions that will be used throughout the paper. Others will be defined when needed.

Converging ω_1 -sequences. The central notion is that of a converging ω_1 -sequence. An ω_1 -sequence $\langle x_\alpha : \alpha \in \omega_1 \rangle$ *converges* to a point x if for every neighbourhood U of x there is an α such that $\{x_\beta : \beta \geq \alpha\} \subseteq U$. Unless stated otherwise our converging sequences are assumed to be non-trivial, that is, injective. A space that contains no non-trivial converging ω_1 -sequences will be called *ω_1 -free*.

In all cases where we construct converging ω_1 -sequences we use the regularity of the space to show convergence: an ω_1 -sequence $\langle x_\alpha : \alpha \in \omega_1 \rangle$ converges to a point x if and only if for every neighbourhood U of x there is an α such that $\{x_\beta : \beta \geq \alpha\} \subseteq \text{cl}U$.

We should record here that the dichotomy is indeed a dichotomy. If a point has a countable local base then it is not the limit of a non-trivial ω_1 -sequence: that sequence would have to be constant on a tail.

Free sequences. At one point we shall need to know that a compact ω_1 -free space has countable tightness, which for the purposes of this paper is best defined by a characterization: there is no free ω_1 -sequence.

An ω_1 -sequence $\langle x_\alpha : \alpha \in \omega_1 \rangle$ is *free* if for every α the sets $\{x_\beta : \beta < \alpha\}$ and $\{x_\beta : \beta \geq \alpha\}$ have disjoint closures.

2. MA is not enough

In this section we show that the conjunction of MA and the equality $\mathfrak{c} = \aleph_2$ is not strong enough to give a positive answer to Juhász' question. For this we recall the definition of initial ω_1 -compactness: every open cover of cardinality at most \aleph_1 has a finite subcover.

In [4, Corollary 5.5] one finds a model of $\text{MA} + \mathfrak{c} = \aleph_2$ in which there is a compact space X that is of countable tightness, and has a proper dense subspace Y that is first-countable and initially ω_1 -compact. As Y is a fortiori countably compact the space X is not first-countable at all points that are not in Y . In addition every compact subset of $X \setminus Y$ is finite. We shall show that X is ω_1 -free.

Assume $\langle x_\alpha : \alpha \in \omega_1 \rangle$ is a converging ω_1 -sequence with limit x . Since, as noted above, a point of first-countability cannot be the limit of a non-trivial ω_1 -sequence we must have $x \in X \setminus Y$.

Since x is the only complete accumulation point of the set $\{x_\alpha : \alpha \in \omega_1\}$ the initial ω_1 -compactness of Y implies that all but countably many x_α are in $X \setminus Y$.

Thus we assume the x_α are all in $X \setminus Y$. For each limit ordinal δ we choose a cofinal subset C_δ of order type ω and an accumulation point y_δ of $\{x_\alpha : \alpha \in C_\delta\}$; by the assumption on $X \setminus Y$ we can take y_δ in Y .

There is a point z in Y with the property that for every neighbourhood U of z the set $\{\delta : y_\delta \in U\}$ is uncountable. But then so is $\{\alpha : x_\alpha \in U\}$ for every such neighbourhood. But then $z = x$ and we have our contradiction.

3. The main result

In this section we prove our main result.

THEOREM 1 (PFA). *Every compact Hausdorff space either contains a non-trivial converging ω_1 -sequence or is first-countable.*

We shall prove the theorem in a number of steps. We assume PFA and we let X be a compact Hausdorff space that is non first-countable and also ω_1 -free, and we intend to reach a contradiction by the end of this section.

LEMMA 2. *Our space X has countable tightness.*

PROOF. This is established by Juhász and Szentmiklóssy in [9]: a compact space of uncountable tightness contains (even) a free ω_1 -sequence that converges. \square

LEMMA 3. *We may assume, without loss of generality, that X is separable.*

PROOF. This was established by the present authors in [5]: a compact ω_1 -free space that is not first-countable contains a closed separable subspace that is not first-countable. \square

Henceforth we add separability to the assumptions on our space X .

LEMMA 4. *Our space X has cardinality at most \mathfrak{c} .*

PROOF. In [1] Balogh proved that PFA implies all compact Hausdorff spaces of countable tightness are in fact sequential.

Now apply the elementary fact that the sequential closure of a countable set has cardinality at most \mathfrak{c} . \square

Our next step is to investigate what happens to our space X when we force with the partial order $\text{Fn}(\omega_1, 2, \aleph_1)$: the set of countable partial functions from ω_1 to 2 (notation as in [10]).

The reason for this is that the main steps towards the contradiction involve proper partial orders that are built in two stages, the first of which is the aforementioned partial order $\text{Fn}(\omega_1, 2, \aleph_1)$; this partial order is proper and forces CH to hold.

LEMMA 5. *Upon forcing with $\text{Fn}(\omega_1, 2, \aleph_1)$ our space X remains compact.*

PROOF. By “our space X remains compact” we mean that the set X , with the topology generated by the ground-model open sets, is a compact space.

To begin: our space X does remain countably compact. This is so because every countably infinite subset of X belongs to the ground model and hence has an accumulation point there. Because the old sets form a base for the new topology that point remains an accumulation point.

Now, if X is no longer compact in this extension then [2, Proposition 5.8] applies and there is a proper partial order \mathbb{P} that introduces an embedding e of the ordinal space ω_1 into X . We shall specify \aleph_1 many dense sets in the partial order $\text{Fn}(\omega_1, 2, \aleph_1) * \dot{\mathbb{P}}$ to determine a map $f: \omega_1 \rightarrow X$, in our ground model, that is a free sequence in X .

First we take, for each α , the set $E_\alpha = \{p : (\exists x \in X)(p \Vdash \dot{e}(\alpha) = x)\}$. An application of PFA to this family yields a map $f: \omega_1 \rightarrow X$.

Next we note that in the extension the set $e[\alpha + 1]$ is compact and does not meet the closure of $e[[\alpha + 1, \omega_1]]$. Therefore there is an open set U_α in the ground model topology with the property that $e[\alpha + 1] \subseteq U_\alpha$ and $\text{cl}U_\alpha \cap e[[\alpha + 1, \omega_1]] = \emptyset$. As above we get dense sets D_α that determine the sequence $\langle U_\alpha : \alpha \in \omega_1 \rangle$.

Now apply PFA to the union of the families to get the desired free sequence $f: \omega_1 \rightarrow X$.

This contradicts the countable tightness of X and establishes the lemma.

□

If we combine this with Lemma 4 then we know that in the extension by $\text{Fn}(\omega_1, 2, \aleph_1)$ our space X has become a compact space of cardinality and weight at most \aleph_1 . This will be used in the next step.

From [5] we quote (and modify) the notion ultra-Fréchet, a useful technical property that will be used in our proof.

DEFINITION 6. A space Z will be said to be *ultra-Fréchet* if it has countable tightness and for each countable subset D of Z and each free ultrafilter \mathcal{U} on D that converges in Z there is a countable subfamily \mathcal{U}' of \mathcal{U} with the property that every infinite pseudointersection of \mathcal{U}' converges.

This is a weakening of the original definition, which did not require that the ultrafilter \mathcal{U} be convergent and thus would imply that the space is countably compact. In the present context of compact spaces there is no difference but the notion may be useful outside this class.

We also remark that all pseudointersections of the countable subfamily converge to the same point (the union of two pseudointersections is again a pseudointersection), to wit the limit of the ultrafilter.

LEMMA 7. *Our space X is ultra-Fréchet, also upon forcing with*

$$\text{Fn}(\omega_1, 2, \aleph_1).$$

PROOF. We argue by contradiction and assume that in the extension there are a countable set D and an ultrafilter \mathcal{U} on D where the property fails; that is: every countable subfamily of \mathcal{U} has an infinite pseudointersection that does not converge.

This will also cover the ground model case: because the forcing adds no new countable sets a ‘bad’ pair $\langle D, \mathcal{U} \rangle$ from the ground model is still ‘bad’ in the extension.

We note that by sequentiality of X every countably infinite subset of X contains an infinite subset that converges: if the set is closed then it is compact metrizable and hence contains a convergent sequence, and if it is not closed then sequentiality gives us a sequence in the set that converges to a point outside the set.

We now construct, recursively, an almost disjoint family $\{a_\alpha : \alpha \in \omega_1\}$ of infinite subsets of D that converge, say a_α converges to y_α , where $y_\alpha \neq x$. Note that none of the a_α s can belong to \mathcal{U} : the countable, indeed singleton, family $\{a_\alpha\}$ would contradict our assumption. We also choose, for each α , a neighbourhood O_α of x such that $y_\alpha \notin \text{cl } O_\alpha$.

We enumerate \mathcal{U} as $\langle U_\alpha : \alpha \in \omega_1 \rangle$.

At stage α we let c_α be an infinite pseudointersection of $\{U_\beta : \beta < \alpha\} \cup \{D \cap O_\beta : \beta < \alpha\}$ that does not converge.

As noted above c_α has an infinite subset that converges; if every infinite convergent subset of c_α were to converge to x then by sequentiality the sequential closure of c_α would be just $c_\alpha \cup \{x\}$, and c_α would converge after all. So there is an infinite convergent subset a_α of c_α that converges to some point y_α distinct from x .

The family $\{a_\alpha : \alpha \in \omega_1\}$ and the ultrafilter \mathcal{U} satisfy the conditions of [6, Lemma 1.8]. Therefore there is a ccc partial order \mathbb{Q} that introduces an uncountable subset I of ω_1 such that $\{a_\alpha : \alpha \in I\}$ is a Luzin family. This means that for every $\alpha \in I$ and every natural number n the set $F(\alpha, n) = \{\beta \in I \cap \alpha : a_\beta \cap a_\alpha \subseteq n\}$ is finite.

It takes only \aleph_1 many dense sets in the partial order $\text{Fn}(\omega_1, 2, \aleph_1) * \dot{\mathbb{Q}}$ to determine such an uncountable set I , the function $\alpha \mapsto a_\alpha \cup \{y_\alpha\}$ and the finite sets $F(\alpha, n)$. Therefore an application of PFA yields the existence of such sets in the ground model, and we may as well assume $I = \omega_1$.

The sequence $\langle y_\alpha : \alpha \in \omega_1 \rangle$ converges, in the ground model. Indeed, let $K = \bigcap_\alpha \text{cl}\{y_\beta : \beta \geq \alpha\}$. By compactness K is non-empty of course and by the Luzin property of the family $\{a_\alpha : \alpha \in \omega_1\}$ it consists of just one point and $\langle y_\alpha : \alpha \in \omega_1 \rangle$ converges to that point, which contradicts our standing assumption on X .

To see this, let $x \in K$ and let O be an arbitrary neighbourhood of x . The set $A = \{\alpha : y_\alpha \in O\}$ is uncountable, and for each $\alpha \in A$ there is an $m_\alpha \in \omega$ such that $a_\alpha \setminus O \subseteq m_\alpha$. We may assume that there is a single m such that $m_\alpha = m$ for all $\alpha \in A$.

We claim that $\{\beta : y_\beta \notin \text{cl}O\}$ is countable; by regularity of X this suffices to prove that $\langle y_\alpha : \alpha \in \omega_1 \rangle$ converges to x .

If the set is uncountable then, by the same argument as above, we find an uncountable set B and an $n \geq m$ such that $a_\beta \cap O \subseteq n$ for all $\beta \in B$. But now take $\alpha \in A$ such that $\alpha \cap B$ is infinite. Then $a_\beta \cap a_\alpha \subseteq n$ for all $\beta \in \alpha \cap B$, contradicting the Luzin property of our family. \square

We can now take the last step towards our contradiction. To recapitulate: we assume our space X is compact, separable, not first-countable and ω_1 -free. From this we deduced that X remains compact after forcing with the partial order $\text{Fn}(\omega_1, 2, \aleph_1)$, and that it is ultra-Fréchet before and after this forcing. In addition, its separability and non-first-countability are preserved by $\text{Fn}(\omega_1, 2, \aleph_1)$ too.

LEMMA 8. *The assumptions that X is compact, separable, not first-countable and ultra-Fréchet, before and after forcing with $\text{Fn}(\omega_1, 2, \aleph_1)$, imply that X has a converging ω_1 -sequence.*

PROOF. We work in the extension by $\text{Fn}(\omega_1, 2, \aleph_1)$ and take a sequence $\langle M_\alpha : \alpha \in \omega_1 \rangle$ of countable elementary substructures of $H(\kappa)$ for a suitable large cardinal κ such that

- X belongs to M_0
- $M_\alpha = \bigcup_{\beta < \alpha} M_\beta$ if α is a limit
- $\langle M_\beta : \beta \leq \alpha \rangle \in M_{\alpha+1}$ and $M_\alpha \prec M_{\alpha+1}$ for all α .

By elementarity there are $z \in M_0 \cap X$ and $D \in M_0$ such that X is not first-countable at z and D is countable and dense in X . We fix such z and D .

Elementarity also implies that $M_\alpha^\omega \in M_{\alpha+1}$ for all α and this, combined with the Continuum Hypothesis, implies that every countable subset of $M = \bigcup_{\alpha < \omega_1} M_\alpha$ is a member of M . In particular $\mathcal{P}(D) \subseteq M$.

At this point we could refer to [5, Proof of Proposition 2.8], which produces sequences $\langle a_\alpha : \alpha \in \omega_1 \rangle$ and $\langle b_\alpha : \alpha \in \omega_1 \rangle$ of infinite subsets of D and a sequence $\langle y_\alpha : \alpha \in \omega_1 \rangle$ of points in $X \setminus \{z\}$ with the following properties:

- (1) a_α converges to z
- (2) b_α converges to y_α
- (3) $a_\alpha \cap b_\alpha = \emptyset$
- (4) $\{A : a_\alpha \cup b_\alpha \subseteq^* A\}$ is an ultrafilter in $M_\alpha \cap \mathcal{P}(D)$
- (5) $a_\alpha, b_\alpha,$ and y_α belong to $M_{\alpha+1}$.

However, for the benefit of the reader we will give a somewhat more streamlined construction than the one referred to above.

Fix an $\alpha \in \omega_1$ and let y_α be a point in the intersection

$$\bigcap \{U : U \in M_\alpha, z \in U, U \text{ is open in } X\}$$

that is distinct from z . Such a point exists because $\{z\}$ is not a G_δ -set in X . Moreover, by elementarity we can assume $y_\alpha \in M_{\alpha+1}$.

Next observe that $y_\alpha \notin \text{cl} A$ whenever $A \in M_\alpha \cap \mathcal{P}(D)$ and $z \notin \text{cl} A$. There is an ultrafilter \mathcal{U}_1 on the Boolean algebra $M_\alpha \cap \mathcal{P}(D)$ that extends the filter dual to $\{A : z \notin \text{cl} A\}$ and is such that $y_\alpha \in \bigcap \{\text{cl} U : U \in \mathcal{U}_1\}$; again we can assume $\mathcal{U}_1 \in M_{\alpha+1}$. By the contrapositive of the first sentence of this paragraph we have $z \in \bigcap \{\text{cl} U : U \in \mathcal{U}_1\}$. Next we take ultrafilters \mathcal{U}_2 and \mathcal{U}_3 on D that extend \mathcal{U}_1 and converge to z and y_α respectively, yet again we take these ultrafilters in $M_{\alpha+1}$.

Now apply the ultra-Fréchet property to \mathcal{U}_2 and y_α , and to \mathcal{U}_3 and z to find countable subfamilies \mathcal{U}'_2 and \mathcal{U}'_3 as in the definition, where we may assume that both families contain \mathcal{U}_1 . These choices can be made in $M_{\alpha+1}$ and we can then take pseudointersections a_α and b_α of \mathcal{U}'_2 and \mathcal{U}'_3 respectively that belong to $M_{\alpha+1}$. By the remark after Definition 6 a_α converges to z and b_α converges to y_α , and we can assume $a_\alpha \cap b_\alpha = \emptyset$.

All five conditions are met: we took care of (1), (2), (3), and (5) explicitly; (4) holds because a_α and b_α are both pseudointersections of \mathcal{U}_1 .

Conditions (4) and (5) imply that the sequences $\langle a_\alpha : \alpha \in \omega_1 \rangle$ and $\langle b_\alpha : \alpha \in \omega_1 \rangle$ are *not* σ -separated in a strong way: whenever \mathcal{A} is a countable family of subsets of D there is an α such that for every $A \in \mathcal{A}$ either $A \cap (a_\alpha \cup b_\alpha)$ is finite or $a_\alpha \cup b_\alpha \subseteq^* A$.

We may then apply a result of Todorčević, see [7, p. 145], that implies there is a proper partial order \mathbb{T} that produces an uncountable subset J of ω_1 such that $(a_\alpha \cap b_\beta) \cup (a_\beta \cap b_\alpha)$ is non-empty whenever $\alpha, \beta \in J$ are distinct.

As before we need \aleph_1 many dense sets in the proper partial order $\text{Fn}(\omega_1, 2, \aleph_1) * \mathbb{T}$ to determine the sequences $\langle a_\alpha : \alpha \in \omega_1 \rangle$, $\langle b_\alpha : \alpha \in \omega_1 \rangle$, and $\langle y_\alpha : \alpha \in \omega_1 \rangle$ as well as the uncountable set J . We apply PFA and obtain items with properties (1), (2) and (3) in our list above as well as the uncountable set J from Todorčević's result, which we may assume to be ω_1 itself.

We claim that $\langle y_\alpha : \alpha \in \omega_1 \rangle$ converges to z , which is the final contradiction that we seek. Let O be a neighbourhood of z . Because every a_α converges to z we know that $a_\alpha \setminus O$ is always finite. We claim that $b_\alpha \cap O$ is finite for only countably many α , and hence that there is an α such that $y_\beta \in \text{cl} O$ for $\beta \geq \alpha$.

Indeed, assume $b_\alpha \cap O$ is finite for uncountably many α and fix two finite sets F and G such that the set A of those α for which $F = a_\alpha \setminus O$ and $G = b_\alpha \cap O$ is uncountable. Now let $\alpha, \beta \in A$ be distinct. Then

$$\begin{aligned} a_\alpha &= F \cup (a_\alpha \cap O) = (a_\beta \setminus O) \cup (a_\alpha \cap O), \\ b_\beta &= G \cup (b_\beta \setminus O) = (b_\alpha \cap O) \cup (b_\beta \setminus O), \end{aligned}$$

which implies $a_\alpha \cap b_\beta = \emptyset$. Likewise $a_\beta \cap b_\alpha = \emptyset$ and we are done. \square

References

- [1] Zoltán T. Balogh, On compact Hausdorff spaces of countable tightness, *Proc. Amer. Math. Soc.*, **105** (1989), 755–764.
- [2] Alan Dow, An introduction to applications of elementary submodels to topology, *Topology Proc.*, **13** (1988), 17–72.
- [3] Alan Dow, Set-theoretic update on topology, in: *Recent Progress in General Topology. III*, Atlantis Press (Paris, 2014), pp. 329–357.
- [4] Alan Dow, Generalized side-conditions and Moore–Mrówka, *Topology Appl.*, **197** (2016), 75–101.
- [5] Alan Dow and Klaas Pieter Hart, Reflecting Lindelöf and converging ω_1 -sequences, *Fund. Math.*, **224** (2014), 205–218.
- [6] Alan Dow and Saharon Shelah, Martin’s axiom and separated mad families, *Rend. Circ. Mat. Palermo (2)*, **61** (2012), 107–115.
- [7] Ilijas Farah, Analytic quotients: theory of liftings for quotients over analytic ideals on the integers, *Mem. Amer. Math. Soc.*, **148** (2000), 177 pp.
- [8] István Juhász, Piotr Koszmider, and Lajos Soukup, A first countable, initially ω_1 -compact but non-compact space, *Topology Appl.*, **156** (2009), 1863–1879.
- [9] István Juhász and Zoltán Szentmiklóssy, Convergent free sequences in compact spaces, *Proc. Amer. Math. Soc.*, **116** (1992), 1153–1160.
- [10] Kenneth Kunen, *Set Theory. An Introduction to Independence Proofs*, Studies in Logic and the Foundations of Mathematics, vol. 102, North-Holland Publishing Co. (Amsterdam, 1980).