

A note on the minimum degree of minimal Ramsey graphs

Bishnoi, Anurag; Lesgourgues, Thomas

Publication date 2025 **Document Version** Final published version

Published in Australasian Journal of Combinatorics

Citation (APA)

Bishnoi, A., & Lesgourgues, T. (2025). A note on the minimum degree of minimal Ramsey graphs. Australasian Journal of Combinatorics, 92(1), 65-69.

Important note To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

A note on the minimum degree of minimal Ramsey graphs

Anurag Bishnoi*

Delft Institute of Applied Mathematics TU Delft, Netherlands A.Bishnoi@tudelft.nl

THOMAS LESGOURGUES[†]

University of Waterloo Waterloo, ON Canada tlesgourgues@uwaterloo.ca

Abstract

In this note, we briefly rectify oversights in the works of several authors on $s_r(K_k)$, the Ramsey parameter introduced by Burr, Erdős and Lovász in 1976, which is defined as the smallest minimum degree of a graph Gsuch that any *r*-colouring of the edges of G contains a monochromatic K_k , whereas no proper subgraph of G has this property. We show that $s_r(K_{k+1}) = O(k^3r^3\ln^3 k)$, improving the best known bounds when $k \ge 8$ and $k^2 \le r \le O(k^4/\ln^6 k)$.

1 Introduction

A graph G is called r-Ramsey for another graph H, denoted by $G \to (H)_r$, if every r-colouring of the edges of G contains a monochromatic copy of H. Observe that if $G \to (H)_r$, then every graph containing G as a subgraph is also r-Ramsey for H. Some very interesting questions arise when we study graphs G which are minimal with respect to $G \to (H)_r$, that is, $G \to (H)_r$ but there is no proper subgraph G' of G such that $G' \to (H)_r$. We call such graphs r-Ramsey minimal for H, and we denote the set of all r-Ramsey minimal graphs for H by $\mathcal{M}_r(H)$. It follows from the

 $^{^{\}ast}\,$ Research supported by a Discovery Early Career Award of the Australian Research Council (No. DE190100666).

[†] Research supported by an Australian Government Research Training Program Scholarship and the School of Mathematics and Statistics, UNSW.

classical result of Ramsey [12] that $\mathcal{M}_r(H)$ is non-empty for any choices of graph H and positive integer r.

Many questions on $\mathcal{M}_r(H)$ have been explored; for example, the Ramsey number $R_r(H)$ denotes the smallest number of vertices of any graph in $\mathcal{M}_r(H)$ and the size Ramsey number $\hat{R}_r(H)$ denotes the smallest number of edges. We refer the reader to [2, 4, 10, 13] for various results on Ramsey minimal problems. In this paper, we will be interested in the smallest minimum degree of an r-Ramsey minimal graph, defined by

$$s_r(H) \coloneqq \min_{G \in \mathcal{M}_r(H)} \delta(G),$$

for a finite graph H and positive integer r, where $\delta(G)$ denotes the minimum degree of G. Trivially, we have $s_r(H) \leq R_r(H) - 1$, since the complete graph on $R_r(H)$ vertices is r-Ramsey for H and is $(R_r(H) - 1)$ -regular (taking minimal Ramsey subgraphs of this graph cannot increase the minimum degree). This parameter was introduced by Burr, Erdős and Lovász [3] in 1976. They were able to show the rather surprising exact result, $s_2(K_{k+1}) = k^2$, where K_{k+1} is the complete graph on k + 1 vertices, which is far away from the trivial exponential bound of $s_2(K_{k+1}) \leq R_2(k+1) - 1$.

While no precise values are known for $s_r(K_{k+1})$ for r > 2, Fox, Grinshpun, Liebenau, Person, and Szabó [6] showed that $s_r(K_{k+1})$ is quadratic in r, up to a polylogarithmic factor, when the size of the clique is fixed. Formally, they showed that for all $k \ge 2$ there exist constants $c_k, C_k > 0$ such that for all $r \ge 3$, we have

$$c_k r^2 \frac{\ln r}{\ln \ln r} \leqslant s_r(K_{k+1}) \leqslant C_k r^2 (\ln r)^{8k^2}.$$
 (1.1)

When k = 2, Guo and Warnke [7] settled the exact polylogarithmic factor, following earlier work in [6]. The constant in the upper bound of (1.1) is rather large $(C_k \sim k^2 2^{8k^2})$, and in particular not polynomial in k. To remedy this, Fox, Grinshpun, Liebenau, Person, and Szabó [6] also proved an upper bound which is polynomial in both k and r and is applicable for small values of r and k.

Theorem 1.1 (Fox, Grinshpun, Liebenau, Person, Szabó). For all $k \ge 2$, $r \ge 3$, $s_r(K_{k+1}) \le 8k^6r^3$.

In the other regime, when the number of colours is fixed, Hàn, Rödl, and Szabó [8] showed that $s_r(K_{k+1})$ is quadratic in the clique size k, up to a polylogarithmic factor. They showed that there exists a constant k_0 such that for every $k > k_0$ and $r < k^2$, we have $s_r(K_{k+1}) \leq 80^3 (r \ln r)^3 (k \ln k)^2$. Combined with (1.1), this result implies the existence of a large absolute constant C and a polynomial upper bound for $s_r(K_{k+1})$.

Theorem 1.2 (Hàn, Rödl, Szabó). There exists an absolute constant C such that for every $k \ge 2$ and $r < k^2$,

$$s_r(K_{k+1}) \leq C(r \ln r)^3 (k \ln k)^2.$$

Finally, using a group theoretic model of generalised quadrangles introduced by Kantor in 1980 [9], Bamberg and the authors [1] proved another polynomial bound,

reducing the dependency in r, and improving on Theorem 1.1 for any k, r and on Theorem 1.2 when $r > k^6$.

Theorem 1.3 (Bamberg, Bishnoi, Lesgourgues). There exists an absolute constant C such that for all $k \ge 2$, $r \ge 3$, $s_r(K_{k+1}) \le Ck^5r^{5/2}$.

These theorems all use the equivalence between $s_r(K_k)$ and another extremal function, called the *r*-colour *k*-clique packing number [6]. Theorems 1.1 and 1.3 further use some 'triangle-free' point-line geometries, for which, under certain conditions on their parameters, any packing of these geometries implies an upper bound on the *r*-colour *k*-clique packing number. This argumentation, initially developed by Dudek and Rödl [5] and then by Fox et al. in [6], has been slightly optimized by Bamberg et al. in [1, Lemma 3.1], allowing for the use of more general geometries and optimising the choice of some parameters. We then rectify the oversight of [1, 6], showing that, by using the optimized argumentation in [1] and the finite geometric construction of Fox et al. in [6], we immediately obtain the following upper bound, improving on the best known bounds for $k \ge 8$ and r in the range $k^2 \le r \le O(k^4/\ln^6 k)$.

Theorem 1.4. For all $k \ge 2$, $r \ge 3$, $s_r(K_{k+1}) \le (8kr \ln k)^3$.

Table 1 contains a summary of the bounds presented above, explaining which theorem gives the best known upper bound for $s_r(K_{k+1})$, depending on the range of r as a function of k.

Range for r	$r < k^2$	$k^2 \leqslant r \leqslant O(k^4 / \ln^6 k)$	$r = \Omega(k^4 / \ln^6 k)$
Upper bound	$C(r\ln r)^3(k\ln k)^2$	$(8kr\ln k)^3$	$Ck^5r^{5/2}$
Source	Theorem 1.2 [8]	Theorem 1.4	Theorem 1.3 [1]

Table 1: Upper bounds for $s_r(K_{k+1})$.

2 Packing partial linear spaces

A partial linear space is an incidence structure of points \mathcal{P} and lines \mathcal{L} , with an incidence relation such that there is at most one line through every pair of distinct points. If every line is incident with exactly s + 1 points and every point is incident with exactly t + 1 lines, then the partial linear space has order (s, t). If there are no three distinct lines pairwise meeting each other in three distinct points, then the partial linear space is *triangle-free. Generalised quadrangles* are standard examples of triangle-free partial linear spaces, with the additional property that for every non-incident point-line pair x, ℓ there exists a unique point x' incident to ℓ such that x and x' are collinear (see the book by Payne and Thas [11] for a standard reference on finite generalised quadrangles).

The next lemma can be found in [1, Lemma 3.1]. Its proof follows a methodology initially developed by Dudek and Rödl [5], using the *r*-colour *k*-clique packing number developed in [6].

Lemma 2.1 (Bamberg, Bishnoi, Lesgourgues). Let r, k, s, t be positive integers. Say there exists a family $(\mathcal{I}_i)_{i=1}^r$ of triangle-free partial linear spaces of order (s, t), on the same point set \mathcal{P} and with pairwise disjoint line sets $\mathcal{L}_1, \ldots, \mathcal{L}_r$, such that the point-line geometry $(\mathcal{P}, \bigcup_{i=1}^r \mathcal{L}_i)$ is also a partial linear space. If $s \ge 3rk \ln k$ and $t \ge 3k(1 + \ln r)$, then $s_r(K_{k+1}) \le |\mathcal{P}|$.

While Theorem 1.1 from [6] was the motivation behind the general Lemma 2.1 and its use in conjunction with the new group-based construction in [1], the authors did not check then the impact of their improved Lemma 2.1 directly on the motivating construction of [6]. This note aims at rectifying this oversight. The following lemma is a reformulation in the language of (triangle-free) partial linear space of the construction that can be found in [6, Proof of Lemma 4.4]. Theorem 1.4 is then a direct consequence of Lemmas 2.1 and 2.2.

Lemma 2.2. Let q be any prime power. There exists a family $(\mathcal{I}_i)_{i=1}^{q-1}$ of trianglefree partial linear spaces of order (q-1, q-2), on the same point set \mathcal{P} of size q^3 and with pairwise disjoint line-sets $\mathcal{L}_1, \ldots, \mathcal{L}_{q-1}$, such that the point-line geometry $(\mathcal{P}, \bigcup_{i=1}^{q-1} \mathcal{L}_i)$ is also a partial linear space.

We include a short proof of Theorem 1.4 for completeness. We note again that this is a replica of the proof of Fox et al. [6], with an optimal choice of q allowed by the work of Bamberg et al [1] presented in Lemma 2.1. The proof of [6] (using their own construction) works word by word if the prime q is chosen to be at least $Ckr \ln k$, for some constant C, instead of k^2r as done [6].

Proof of Theorem 1.4. Let $k \ge 2$, $r \ge 3$, and let q be the smallest prime such that $q \ge 4kr \ln k$. By Bertrand's postulate, $q \le 8kr \ln k$. By Lemma 2.2, there exists a family of r < q triangle-free partial linear spaces of order (q - 1, q - 2), on the same point set \mathcal{P} and pairwise disjoint line-sets $\mathcal{L}_1, \ldots, \mathcal{L}_r$, such that the point-line geometry $(\mathcal{P}, \bigcup_{i=1}^r \mathcal{L}_i)$ is also a partial linear space. Note that with $k \ge 2$ and $r \ge 3$, we have $q - 1 \ge 3rk \ln k$ and $q - 2 \ge 3k(1 + \ln r)$. By Lemma 2.1, $s_r(K_{k+1}) \le |\mathcal{P}|$, and then $|\mathcal{P}| = q^3$ yields the desired bound.

A careful review of the arguments in [1, Lemma 3.1 and 5.2] would allow a small optimisation on the multiplicative constant of this corollary. However, in light of the conjectured quadratic upper bound [1, Conjecture 5.2], we did not push this further.

References

- J. Bamberg, A. Bishnoi and T. Lesgourgues, The minimum degree of minimal Ramsey graphs for cliques, *Bull. London Math. Soc.* 54 (2022), 1827–1838.
- [2] S. A. Burr, P. Erdős, R. J. Faudree, C. C. Rousseau and R. H. Schelp, Ramseyminimal graphs for star-forests, *Discrete Math.* 33(3) (1981), 227–237.

- [3] S. A. Burr, P. Erdős and L. Lovász, On graphs of Ramsey type, Ars Combin. 1(1) (1976), 167–190.
- [4] S. A. Burr, J. Nešetřil and V. Rödl, On the use of senders in generalized Ramsey theory for graphs, *Discrete Math.* 54(1) (1985), 1–13.
- [5] A. Dudek and V. Rödl, On K_s -free subgraphs in K_{s+k} -free graphs and vertex Folkman numbers, *Combinatorica* **31**(1) (2011), 39–53.
- [6] J. Fox, A. Grinshpun, A. Liebenau, Y. Person and T. Szabó, On the minimum degree of minimal Ramsey graphs for multiple colours, J. Combin. Theory Ser. B 120 (2016), 64–82.
- [7] H. Guo and L. Warnke, Packing nearly optimal Ramsey R(3,t) graphs, *Combinatorica* **40**(1) (2020), 63–103.
- [8] H. Hàn, V. Rödl and T. Szabó, Vertex Folkman numbers and the minimum degree of minimal Ramsey graphs, SIAM J. Discrete Math. 32(2) (2018), 826– 838.
- [9] W. M. Kantor, Generalized polygons, SCABs and GABs, In: Buildings and the geometry of diagrams (Como, 1984), Lecture Notes in Math. Vol. 1181 (1986), 79–158.
- [10] T. Łuczak, On Ramsey minimal graphs, *Elec. J. Combin.* 1 (1994), #R4.
- [11] S. E. Payne and J. A. Thas, *Finite generalized quadrangles*, EMS Series of Lectures in Mathematics, European Mathematical Society (EMS), Zürich, second ed., 2009.
- [12] F. P. Ramsey, On a Problem of Formal Logic, Proc. London Math. Soc. 30(4) (1929), 264–286.
- [13] V. Rödl and M. Siggers, On Ramsey minimal graphs, SIAM J. Discrete Math. 22(2) (2008), 467–488.

(Received 14 Apr 2024; revised 23 Nov 2024, 31 Mar 2025)