

Document Version

Final published version

Licence

CC BY

Citation (APA)

Pagliari, F., Spadea, M.-F., Montay-Gruel, P., Puspitasari-Kokko, A., Seco, J., Tirinato, L., Accardo, A., De Angelis, F., & Gentile, F. (2025). Nano-Topography Enhanced Topological-Cell-Analysis in Radiation-Therapy. *Advanced Healthcare Materials*, 14(12), Article 2405187. <https://doi.org/10.1002/adhm.202405187>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

In case the licence states "Dutch Copyright Act (Article 25fa)", this publication was made available Green Open Access via the TU Delft Institutional Repository pursuant to Dutch Copyright Act (Article 25fa, the Taverne amendment). This provision does not affect copyright ownership.
Unless copyright is transferred by contract or statute, it remains with the copyright holder.

Sharing and reuse

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Nano-Topography Enhanced Topological-Cell-Analysis in Radiation-Therapy

Francesca Pagliari, Maria-Francesca Spadea, Pierre Montay-Gruel, Anggraeini Puspitasari-Kokko, Joao Seco, Luca Tirinato, Angelo Accardo,* Francesco De Angelis,* and Francesco Gentile*

Radiotherapy (RT) is a cancer treatment technique that involves exposing cells to ionizing radiation, including X-rays, electrons, or protons. RT offers promise to treat cancer, however, some inherent limitations can hamper its performance. Radio-resistance, whether innate or acquired, refers to the ability of tumor cells to withstand treatment, making it a key factor in RT failure. This perspective hypothesizes that nanoscale surface topography can impact on the topology of cancer cells network under radiation, and that this understanding can possibly advance the assessment of cell radio-resistance in RT applications. An experimental plan is proposed to test this hypothesis, using cancer cells exposed to various RT forms. By examining the influence of 2D surface and 3D scaffold nanoscale architecture on cancer cells, this approach diverges from traditional methodologies, such as clonogenic assays, offering a novel viewpoint that integrates fields such as tissue engineering, artificial intelligence, and nanotechnology. The hypotheses at the base of this perspective not only may advance cancer treatment but also offers insights into the broader field of structural biology. Nanotechnology and label-free Raman phenotyping of biological samples are lenses through which scientists can possibly better elucidate the structure-function relationship in biological systems.

1. Introduction

This perspective proposes to explore the combined effects of surface topography, scaffold geometry, and RT treatment on the topology of cancer cell networks, examining how these responses are modulated by the intrinsic radiation resistance (RR) of cells (Figure 1). The work is structured as follows.

In the *introduction*, we first briefly describe how RT works, and suggest that surface-enhanced topological cell analysis can be a powerful tool for RT assessment (1.1). Then, we discuss the notion that nano-topography can efficiently guide cell behavior, and the spatial arrangement of cells, i.e., structure (1.2). Subsequently, we recall that, in biological systems, structure and function are correlated (1.3) providing examples of how, by measuring the internal architecture of a system one can have an indication of how that system works (1.4). In the last section of the introduction,


F. Pagliari, J. Seco
Division of BioMedical Physics in Radiation Oncology
German Cancer Research Center
69120 Heidelberg, Germany
M.-F. Spadea
Institute of Biomedical Engineering
Karlsruhe Institute of Technology (KIT)
76131 Karlsruhe, Germany
P. Montay-Gruel
Radiation Oncology Department
Iridium Network
Antwerp 2610, Belgium

P. Montay-Gruel
Antwerp Research in Radiation Oncology (AreRO)
Center for Oncological Research (CORE)
University of Antwerp
Antwerp 2020, Belgium
A. Puspitasari-Kokko
Research and Development - HollandPTC
Delft University of Technology
Delft, The Netherlands
J. Seco
Department of Physics and Astronomy
Heidelberg University
Im Neuenheimer Feld 227, 69120 Heidelberg, Germany
L. Tirinato
Department of Medical and Surgical Sciences
University Magna Graecia of Catanzaro
Catanzaro 88100, Italy
A. Accardo
Department of Precision and Microsystems Engineering
Faculty of Mechanical Engineering
Delft University of Technology
Mekelweg 2, Delft 2628 CD, The Netherlands
E-mail: a.accardo@tudelft.nl

The ORCID identification number(s) for the author(s) of this article can be found under <https://doi.org/10.1002/adhm.202405187>

© 2025 The Author(s). Advanced Healthcare Materials published by Wiley-VCH GmbH. This is an open access article under the terms of the [Creative Commons Attribution](#) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

DOI: 10.1002/adhm.202405187



Introduction	Radiation Therapy (RT) Nanoscale details influence cell behaviour Structure-function relationship Experimental plan
1	
Review	Conventional assays in RT Topological cell analysis in RT Nanotopography as a regulator of cell behaviour 3D scaffolds as in vitro-models of RT studies Nanomedicine-based strategies for overcoming cell RT
2	
the Idea	Nano-topography enhanced topological cell analysis
3	
4-Plan	An experimental plan of implementation
Conclusions	The impact of the method in RT and biomedical sciences
5	

Figure 1. Schematic representation of the proposed approach detailed in the perspective. Cancer cells are cultured in biomimetic scaffolds a) and exposed to radiation b). The scaffold's surface nano-topography c) and overall architecture d) can be precisely engineered using nanotechnology. The interplay between surface nano-topography, scaffold geometry, and radiation exposure leads to the organization of cells into specific architectures e). A topological analysis of these cellular networks provides insights into their biological behavior and potential resistance to radiotherapy. The inset f) summarizes the structure of this work, which includes i) an introduction, ii) a review section, iii) a discussion of the proposed approach—nano-topography-enhanced topological cell analysis, iv) an experimental implementation plan, and v) a conclusion.

we outline a possible experimental plan necessary to develop these ideas (1.5).

Thereafter, we *review* the existing body of literature related to RT, either pertaining to conventional methods of analysis, such as clonogenic assays (2.1), or to a recently introduced technique that leverages cell-network topology (2.2). Then, we critically review reports on nano-topography as a regulator of cell behavior (2.3) and comment on the use of 3D scaffolds as an in-vitro-model for RT studies (2.4). The final section of the review focuses on analyzing current nanomedicine strategies to overcome cell RR (2.5).

In the next section, we introduce the object of this perspective, i.e., *nano-topography enhanced topological cell analysis*, and substantiate the claim that this new field of study can improve

over the existing methods of RT assessment, in terms of precision, sensitivity, accuracy, and cost-effectiveness (3). In doing so, we describe possible outcomes related to the research, including technology, scientific and social outcomes. A possible *experimental plan of implementation* of this idea is briefly outlined in a separate study design section (4). In the *conclusions*, we discuss the potential impact of the concepts outlined in this perspective on cancer care and the broader field of biomedical sciences (5).

1.1. Radiation Therapy - RT

RT is a cancer treatment approach that involves exposing cancerous tissues and organs to ionizing radiation, such as X-rays, electrons, or protons. This radiation damages the genetic material and molecular structure of cells, disrupting their functions, including proliferation and growth, altering their cellular interactions, and ultimately leading to their death.^[1,2] In modern clinical practice, around half of all cancer patients undergo RT, either as a standalone treatment or in combination with chemotherapy and surgery.^[3] Although radiotherapy holds great potential for cancer treatment, certain inherent limitations can reduce its effectiveness. RR, whether innate or acquired, enables tumor cells to withstand treatment through various mechanisms. These include enhanced DNA damage repair, autophagy, adaptation to hypoxia, and the presence of cancer stem cells, all of which are linked

F. De Angelis
Plasmon nano-technologies
Italian Institute of Technology
Genova 16163, Italy
E-mail: francesco.deangelis@iit.it

F. Gentile
Nanotechnology Research Center
Department of Experimental and Clinical Medicine
University of Magna Graecia of Catanzaro
Catanzaro 88100, Italy
E-mail: francesco.gentile@unicz.it

to alterations in the tumor microenvironment.^[4,5] RR is a key contributor to the failure of RT. Developing and strengthening techniques and methods to evaluate cell sensitivity to internal or external radiation fields is thus critically important. *The aim of this perspective is to propose a method for assessing the RR of cancer cells and to outline a study design for its implementation.* In particular, the research is designed to explore the influence of the nanoscale architecture of 2D surfaces and 3D scaffolds on the *topology* of cancer cell networks under radiation. The idea behind this perspective stems from the observation that external radiation fields alter cellular DNA, influencing the collective behavior of cellular systems. Here, we hypothesize that nano-topography can amplify these effects. This approach leverages detailed knowledge of cell anatomy and spatial organization within colonies to assess biological characteristics, using topological measures instead of traditional clonogenic assays for evaluating RT treatment effectiveness.

1.2. Nano-Topography Drives Cell Assembly

The formation of multicellular structures is governed by adhesion forces, which regulate interactions both between individual cells and between cells and their surrounding environment. This interaction determines the evolution of cell systems and guides them toward ordered structures.^[6–8] Nano-topography, in particular, has been identified as a critical factor in directing cell fate at the bio-interface,^[9–12] making it a significant area of interest in tissue engineering, biosensors, the analysis of neurodegenerative disorders. Studies^[13–15] have demonstrated that cells cultured on surfaces with nanoscale motifs tend to arrange into clusters with high clustering and short average path lengths. *Thanks to this peculiar architecture, similar structures exhibit enhanced computational capabilities compared to cells randomly distributed on a surface without any special layout.* Thus, in the context of this perspective, nano-topography plays a major role because it represents a factor – a parameter – through which one can possibly control the shape of biological aggregates and, notably, their function.

The organization of cells into networks with specific topologies enhances overall network performance—much like how an organism's functionality depends more on the collective cooperation of its elements than on their individual properties. In the brain, complex functions such as memory, language, and thought emerge from the continuous exchange of signals (action potentials). The dynamic positioning of signal sources and drains influences the distances and paths these signals must travel, directly impacting their efficiency. Similar to the traveling salesman problem, system performance is highly sensitive to component layout, a principle studied in topology and network science. This fundamental relationship explains why computational power is tied to network topology. While the correlation between structure and function in biological systems has long been recognized, a seminal study in the early 1980s^[16] first demonstrated how neural network computation depends on collective behavior—a discovery that earned a Nobel Prize in physics 45 years later. More recently, researchers have investigated how the cell-material interface regulates information flow in biological networks, using both experimental^[14,15,17] and numerical^[18] approaches.

1.3. In Biological Systems, Structure and Function are Correlated

The way in which cells of a system are arranged in the space, and the way in which they are interconnected, govern the functionality and overall behavior of biological systems.^[16,19,20] The peculiar characteristics of similar systems rest more on the cooperation of numerous cells at different organizational levels, and less on the attributes of individual cells.^[21–23] (This is, notably, the definition of complex systems.^[24–27]) Thus, studying and understanding the functions of biological systems inevitably requires a deep comprehension of their structure. This includes examining their architecture, form, the connections between individual components, and how these components communicate. To efficiently do this, it is convenient representing systems as networks.^[28,29] In simple terms, a network (or a graph) is a set of objects that are connected together. The objects of a network, in turn, are regarded as nodes: the connections between the nodes are called edges or links. Networks science is the discipline that studies networks. In the context of networks science, notably, correlation implies causation: objects of a network that are connected do also interact (**Figure 2**). This also suggests that biological systems can be modeled as networks, where the nodes represent system elements and the links describe the interactions between them. One key advantage of networks is that they can be quantified. Networks' *topology* defines the way a network is shaped and how its internal components are associated. There exist quantitative metrics that precisely measure topology, and make networks a universal tool confidently used in different areas across biology, ecology, and social science.^[28,29]

1.4. Examples of Structure-Function Relationships in Biological Systems and Vision

One notable example of a biological system where structure and function are manifestly correlated is the brain: a tightly interweaving network of neural cells. In this system, the topological structure of the network is responsible for increased computational ability, low energy consumption, and non-deterministic functions like language and memory. The complexity of human behavior is driven less by the specialization of individual nerve cells and more by the formation of intricate networks composed of a large number of these cells.^[18,26,38,39] Theories like the integrated information theory propose that consciousness itself emerges from the way information is processed within a system, indicating a direct correlation between interconnectedness and higher cognitive functions.^[40] Understanding how individual cells interact to form complex systems is paramount for fields like tissue engineering and regenerative medicine, as it offers insights into the formation of in-vitro models, the analysis of neuro-degenerative diseases, the design of efficient prosthesis implants, and neuromorphic engineering. The exploration of biological systems through the lens of their internal architecture, topology, and spatial layout reveals a fascinating interplay between structure and function. The influence of 3D scaffold architecture on cell network topology opens up new avenues in understanding how biological systems self-organize and function. This insight not only enhances our comprehension of natural biological processes but also guides the development of synthetic




Figure 2. A graphical representation of networks science. The study of networks, known as network science, has gained prominence across various disciplines as a tool for analyzing complex systems: including social, biological, and technological systems.^[26,28,30–32] In particular, biological systems can be efficiently described as networks a). We can use different parameters to describe networks, including the network degree, the clustering coefficient b), the characteristic path length c), and the small world coefficient d). The network degree (k) represents the average number of links per node in a graph. The clustering coefficient (cc) measures graph connectivity, ranging from 0 to 1. It quantifies the proportion of existing links among the nodes surrounding a given element compared to the total possible connections among them, averaged across the entire network. The characteristic path length (cpl) refers to the average shortest path between network elements. Among these parameters, the small-world coefficient (SW) is particularly significant. It is derived from the combination of cc and cpl , representing the balance between network connectivity (cc) and proximity (cpl) relative to a random graph of the same size.^[33] – SW is non-dimensional. Small-world networks with $SW > 1$ are topologically biased to enhance local connectivity,^[28,31,32] which in turn optimizes system's performance.^[14,18,34–37] The study of small-world networks has recently gained momentum because many real systems, phenomena and processes can be modelled after this network topology.




Figure 3. Interdisciplinary approach. Various disciplines have to be integrated to enhance our understanding of cancer RT.

systems and therapeutic interventions. The implications of these findings are vast, extending from tissue engineering to the intricate workings of the human brain, showing the profound impact of architecture and topology on the functionality of biological systems.

It is important to note that both the geometry of 2D surfaces and the architecture of 3D scaffolds play a crucial role in shaping cell network topology. A number of examples, conveniently reported in a separate Supporting Information 1, demonstrate the effectiveness of bidimensional surfaces in regulating of cell-assembly. Nanoscale 2D surfaces serve as the crucial link between traditional 2D cell culture models and more advanced 3D templates. They provide the necessary complexity for cell colony growth while enhancing physiological relevance, drug response accuracy, and predictive power—hallmarks of 3D cell models.^[41]

1.5. Outline of the Work

This article postulates that cancer-cell response to radiation can be modulated by an engineered micro-environment. Moreover, we suggest that the topology of a biological system subjected to external radiation fields can be measured using the methods of networks science, and that such a topological measure can be indicative of the system's state. Throughout the text, stemming from a critical review of the existing body of literature on cancer cell RT, nano-topography, and the development of 3D scaffolds for in vitro-studies of biological problems, we provide evidence of this hypothesis. In addition, we sketch a possible experimental plan that can be used to test the hypothesis of nano-topography enhanced topological-cell-analysis in RT. Such a plan involves exposing cancer cells to various RT modalities, in both conventional and ultra-high dose rate settings. The architecture of the resulting cell networks, conveniently analyzed using Network Science and Artificial Intelligence, can possibly enable the construction of predictive models to forecast patient responses to radiation based on ex vivo radio-sensitivity (RS) tests. Additionally, Raman spectroscopy and electrophysiological monitoring can be designed to examine the correlation between the topology of cell networks and the phenotypic traits expressed by the cells. By investigating the correlation between the topology of cell networks and the phenotypic traits expressed by the cells, this research aims to establish a link between structure

and function of a biological system. This study-design represents a pioneering approach, integrating nanotechnology, cell biology, radiation therapy, AI, and network science, to advance understanding of cancer treatment and response mechanisms (Figure 3).

2. Literature Review

This section presents a thorough review of the existing body of literature related to i) RT assessment, ii) topological cell analysis, iii) nano-topography driven cell assembly, iv) 3D scaffolds for in vitro-models study of RT, and v) nanomedicine strategies in RT.

2.1. Consolidated Methods of Analysis of RT

State-of-the-art techniques for assessing cell RR and RS include cell proliferation assays,^[42] the neutral comet assay,^[43] gamma-H2AX immunofluorescence staining,^[44] and the clonogenic (colony formation) assay.^[45] Among these, the clonogenic assay is widely utilized in radiation biology, as it establishes a correlation between cell RS, survival, and clinical responses to RT in both tumor cell lines and patient-derived tumor cells.^[46] This in-vitro cell survival assay evaluates the capacity of a single cell to develop into a colony. In this method, cancer cells are seeded onto a substrate, supplemented with a cell culture medium, exposed to ionizing radiation, and observed over time. The plating efficiency and surviving fraction values obtained from this assay provide key insights into a cancer cell's ability to withstand radiation. Recognized as the gold standard for studying cancer cell responses to radiation, the clonogenic assay, however, is influenced by several factors, including cell type, doubling time, cell density, cellular interactions, and nutrient availability. In radiobiology studies using the clonogenic assay, each colony is traditionally treated as a *black box*, meaning the individual characteristics of cancer cells within each surviving colony are often overlooked. Colonies are typically considered as single entities, without accounting for intrinsic cellular variability. While conventional clonogenic assays primarily measure the number of surviving colonies post-treatment, the technique has advanced over time, incorporating greater complexity. Recently, several research groups^[47,48] have proposed using the morphological characteristics of cells as an additional parameter to assess their RR or RS.




Figure 4. Comparative assessment of RR techniques: clonogenic assay versus topological cell analysis. Schematic representation of cells exposed to RT and their post-exposure spatial organization, highlighting the importance of networks science for describing the impact of radiation on cell clustering and network formation a). Traditional clonogenic assays assess RR by counting the number of surviving colonies, without considering their internal spatial structure b). Topological cell analysis quantitatively characterizes cell-network architecture using metrics such as clustering coefficient, characteristic path length, and small-world coefficient. These network parameters provide a higher-resolution measure of cellular response to RT by capturing the connectivity and organization of surviving cell populations. By integrating nano-engineered surfaces, topological analysis can enhance sensitivity in detecting subtle alterations in cell behavior post-irradiation, offering a novel approach for evaluating RR and tailoring treatment strategies c).

2.2. Topological Cell Analysis of Radiation-Therapy: A Paradigm Shift

In a recent study,^[49] researchers have investigated the relationship between the topological properties of lung cancer networks in 2D cultures and their response to ionizing radiation. In experiments in which the dose was varied up to 8 Gy, the study showed that the topological measures of cancer lung cell-graphs also varied significantly. In the continuation of the research,^[50] the same team demonstrated that radiation in the 0 – 6 Gy range influences the topology of cancer cell lines different from cancer, including epithelial neuroglioma cells, PC3 bone metastasis of prostate cancer and urinary bladder cancer cells (a more detailed analysis of the correlation between external radiation dose and cancer cell network topology for different cell lines is provided in Supporting Information 2). Current clonogenic protocols^[45] aim to assess cell RS and RR by analyzing the number of cell colonies that form on a 2D substrate after irradiation.

However, the number of colonies in a sample is an estimate of how cells behave in response to radiotherapy. Methods devised in references^[49,50] are an attempt to surpass the limitations found in conventional clonogenic assays.

In these cited works, *topology* is utilized to explore the behavior of cancer cells exposed to radiation. The topological struc-

ture of a large number of cells within a colony is determined by the spatial arrangement of their nuclei, reflecting how cells are interconnected and organized. Compared to simple cell counting, topology provides a more comprehensive, detailed, and quantitative representation of the system (Figure 4). Cancer cell graphs emerge from the interactions among all elements within a colony, encapsulating the maximum amount of information about the cellular system. As a result, network variables offer a robust and reliable means of describing how external factors, such as radiation, influence the system.

This concept can be promptly applied to clinical practice. Tumor tissue obtained through solid biopsy can be dissociated, and the resulting cancer cells cultured on a support. As these cell colonies develop, they can be exposed to progressively higher radiation doses and analyzed using the methods outlined in this study. The small-world coefficient values derived from these samples will then be compared to those of non-cancerous cells from the same tissue or organ. This comparison will help determine whether the patient's cells are radio-sensitive or radio-resistant and to what extent—critical factors in personalizing the most effective treatment strategy for the patient.

Thus, *topological cell analysis* offers promises to overcome the limitations of clonogenic assays. To place this work in context and substantiate the claims of the perspective, in a separate




Figure 5. Influence of surface nano-topography and scaffold geometry on cancer cell network topology. Engineered nano-topographical surfaces and 3D scaffolds introduce controlled variations in roughness, porosity, and surface architecture, modulating cell adhesion and clustering. The biological mechanisms at the basis of nano-topography driven cell behavior are understood, and recapitulated in a separate supporting information a). Nano-scale surfaces and scaffolds guide the self-assembly of cells into interconnected networks, influencing their spatial organization and functional properties b). Quantification of topological features—such as node degree, clustering coefficient, and small-world properties—demonstrates that nano-engineered substrates are effective driving network formation. This understanding can be possibly used in the assessment of RT by topological cell analysis: surface nano-topography can amplify differences in cancer cell network topology post-irradiation. This amplification effect enhances in turn the resolution of RR assessment, facilitating the identification of radio-sensitive and radio-resistant phenotypes with greater accuracy.

Supporting Information 3 it is thoroughly described how clonogenic assays work, especially pertaining to tissue culture plastic models –, i.e., still the gold standard for determining the RS of cells. This section examines the critical aspects of clonogenic assays, including their robustness and inter-assay precision. Additionally, it presents a quantitative comparison of the sensitivity of clonogenic assays versus topological cell analysis, based on reported cases in the literature. The findings suggest that topological cell analysis, along with its future advancements, offers significant improvements over conventional clonogenic assays.

The image of cells in inset (a) is adapted under the terms of the CC-BY Creative Commons Attribution 4.0 International license (<https://creativecommons.org/licenses/by/4.0/>):^[49] Copyright 2022, published by Nature Portfolio. The clonogenic assay image in insets (b) is adapted under the terms of the CC-BY Creative Commons Attribution 4.0 International license (<https://creativecommons.org/licenses/by/4.0/>):^[49] Copyright 2020, published by Nature Portfolio. The measured values of small-world coefficient as a function of external dose for the T24, H460, PC3 and H4 cell lines shown in inset (c), are adapted under the terms of the CC-BY Creative Commons Attribution 4.0 International license (<https://creativecommons.org/licenses/by/4.0/>):^[50] Copyright 2023, published by Springer.

2.3. Nano-Topographical Control of Cell Assembly

The development of cellular systems on a surface is governed by the interplay between cell-substrate interaction forces, cell-cell adhesion forces, and their competition. Cell adhesion to a substrate is influenced by a series of chemical and mechanical signals that begin at the bio-interface and travel through the cell membrane to the cytoskeleton.^[51] Cell-adhesion molecules (CAMs), which operate at a characteristic low nanometer scale, effectively mediate these signals.^[52] Consequently, cell adhesion and clustering can be adjusted by designing surface topography at the nanoscale.^[11,53] By employing nanofabrication techniques one can create surfaces with precisely controlled nano-topography, where roughness and *fractal dimension* are modulated independently. Fractal dimension measures surface complexity, with higher values indicating the presence of details at increasingly smaller scales.^[54] Among the several different techniques that exist to produce nanoscale surfaces, the simplest ones imply wet-etching techniques^[14,54,55] and electro-chemical etching.^[13,56] By fine-tuning the parameters of these fabrication processes, one can produce surfaces with specific morphological characteristics that influence cell behavior,^[11,51–53,57] (Figure 5) potentially enhancing the effects of RT on cells. Previous research has shown that surfaces with intermediate roughness (20 – 40 nm) and high fractal dimension (> 2.4) enhance

adhesion and clustering of various cell types, including human lung carcinoma,^[54] neuroblastoma,^[13,58] and breast cancer cells.^[56] These findings have been supported by biomechanical models^[7] that attribute the collective behavior of cells to the combined influence of cell-surface and cell–cell interactions.

2.4. 3D Scaffolds for In Vitro-Models Study of RT

2D models of cells may sometimes fail to accurately reproduce the functions of tissues and organs. While conventional clonogenic assays are based themselves on 2D set-ups, to examine more faithfully the response of cancer cells to radiation one has to shift to 3D cell systems. Compared to 2D cultures, 3D cell models offer greater physiological relevance, more accurately mimicking tissue microenvironments, cell-to-cell interactions, and biological processes. 3D in vitro cancer constructs have emerged as effective tools to gain insights into cancer biology, striking a balance between simplistic 2D cultures and more intricate cancer xenografts in immunocompromised animals or syngeneic models. Recently, different 3D models have been applying to the field of radio-oncology, showing good correlations between experimental models and actual patient's responses to RT.^[59–61]

In particular, 3D scaffold-based constructs provide structural support to cells and enable the investigation of cellular behavior under well-defined culture conditions, approaching a “quasi in vivo” level. These controlled systems foster improved reproducibility and facilitate the development of more precise, scalable, high-throughput tools, thereby producing more consistent and reliable data and saving time and costs. In 3D systems, cells preserve their spatial interactions, which are vital in determining their morphology, polarity and consequently their functions and fate. A number of studies has shown dose-response curves, which consistently demonstrate heightened cell RR in a 3D environment compared to a 2D setting.^[62,63]

In scaffold-based 3D models of cell cultures, cells are typically inserted in a 3D preformed structure with overall size spanning the millimeter size, but with details as small as some microns (system's resolution) and roughness in the nanometer range. Similar systems provide adequate support for cell growth enabling, among other things, the formation of cell–cell junctions, cell proliferation, the expression of specific genes and proteins: in similar systems, cells interact actively with the scaffold at the cell-material interface. The scaffold surface can be conveniently modified to stimulate cell clustering and boost cellular response to radiation. Scaffolds can be designed using computer aided designed software, and fabricated by two-photon polymerization following well-assessed methods reviewed, for examples, in references.^[64] The surface-characteristics of the scaffold can be conveniently tuned acting on the parameters of the process (hatching and slicing) to obtain the desired value of surface roughness at the nanoscale.

Images of the scaffold and 3D distributions of cells are adapted under the terms of the CC-BY Creative Commons Attribution 4.0 International license (<https://creativecommons.org/licenses/by/4.0>):^[49] Copyright 2024, published by Wiley.

Images of the nano-rough surface and of the 2D clustered distribution of cells are adapted under the terms of the CC-BY Creative Commons Attribution 4.0 International license ([\[creativecommons.org/licenses/by/4.0\]\(https://creativecommons.org/licenses/by/4.0\)\):^{\[49\]} Copyright 2017, published by Nature Portfolio.](https://</p></div><div data-bbox=)

The first mesoporous silicon surface image is adapted under the terms of the CC-BY Creative Commons Attribution 4.0 International license (<https://creativecommons.org/licenses/by/4.0>):^[49] Copyright 2020, The Authors, published by Nature MDPI. The second mesoporous silicon surface image is adapted with permission^[58] Copyright 2012, published by the American Chemical Society.

2.5. Nanomedicine-Based Strategies for Overcoming Cell RR

While this perspective focuses on developing a method to assess cellular RR to radiation, it is useful to briefly review the current nanotechnology and nanomedicine therapeutic approaches aimed at overcoming RR to provide context for the work. Nanomedicine has emerged as a promising approach to overcoming RR in cancer therapy by leveraging nano-scale drug delivery systems (NDDS) and nanomaterials as radiosensitizers.^[65–67] These platforms enhance the therapeutic efficacy of radiotherapy, facilitating precise tumor targeting, and minimizing damage to surrounding healthy tissues. Various nano-radiosensitizers, including gold, silver, platinum, gadolinium, and titanium-based nanoparticles, as well as liposomes, polymeric nanocarriers, and carbon-based nanomaterials, have been developed to enhance radiation-induced damage at multiple biological levels.^[68–70] These materials exhibit excellent biocompatibility and, at the nano-scale, can improve radiotherapy responses through their unique physicochemical, electric, and optical properties.^[71] The primary biological mechanisms of nanomaterial-induced radio-sensitization involve alterations in the cell cycle, inhibition of DNA repair, and induction of mitochondrial dysfunction.^[68] Functionalized nanocarriers incorporating microRNAs (miRNAs) and long non-coding RNAs (lncRNAs) further enhance radio-sensitization by targeting key genetic and epigenetic pathways responsible for tumor resistance. Additionally, metal nanoparticles loaded with chemotherapeutic agents enable combinatorial effects, amplifying therapeutic outcomes through enhanced targeting, biodistribution, and controlled drug release. Hybrid nanoplatforms integrating radiosensitizers, immune modulators, and chemotherapeutic drugs show promise in overcoming RR in cancers such as breast, lung, and glioblastoma. The combination of nanotechnology with molecular targeting has led to personalized and precision treatment strategies, optimizing the therapeutic index of radiotherapy while reducing systemic toxicity. Future advancements in NDDS are expected to focus on multifunctional nanoparticles that integrate imaging, targeting, and therapeutic capabilities, allowing real-time monitoring and adaptive treatment strategies. With continued research and clinical translation, nano-based radiosensitizers hold great potential for improving patient outcomes, reducing treatment-related side effects, and addressing the longstanding challenge of RR in oncology.

3. Nano-Topography Enhanced Topological Analysis of Cells

To improve sensitivity, enhance precision and optimize performance of topological cell analysis, we propose to integrate

networks analysis of cancer-cell graphs and nanotechnology. We aim at boosting the topological response of cells to radiation using devices modified at the nanoscale (Figure 5). Cell-surface interactions and radiation effects can combine and direct cells to cluster into networks. Cells that are sensitive or resistant to radiation will form significantly different networks – this difference is made even larger by the underlying nanoscale geometry of the substrate. Thus, cells with low but still-not null responsiveness to radiation (otherwise invisible to conventional analysis) can be detected thanks to the amplification effect of the substrate.

The core idea of this perspective is that both the nanoscale geometry of a surface or scaffold and radiation can influence cell clustering, and their combined effects may amplify cellular responses. As a result, this synergy can significantly enhance the sensitivity, precision, and accuracy of the method, similarly to the increased Raman signal observed in Surface-Enhanced Raman Scattering (SERS) applications. While this perspective primarily draws from existing literature on nano-topography-guided cell behavior, radiation therapy, and topological cell analysis—most of which are phenomenological in nature—it remains crucial to further investigate the fundamental biological mechanisms underlying these effects. Understanding these mechanisms could reveal a common link between different drivers of cell assembly, such as nano-topography and radiation. If elucidated, this connection could serve as a guiding principle for engineering nanoscaffolds with enhanced efficacy for RT applications. Although further research is needed to fully explore these concepts, in a separate Supporting Information 4, we propose possible biological mechanisms underlying this work. Specifically, we first review key findings on cell-material interactions at the nanoscale from a cell biology perspective, emphasizing recent evidence that nano-topography can influence cell phenotype. This observation suggests that such interactions can be analyzed using epigenetics and chromatin evolution in cell nuclei. Next, we examine the biological aspects of cell-radiation interactions in RT applications, highlighting how chromatin structure affects DNA sensitivity to radiation. We then propose that chromatin serves as the crucial link between radiation therapy and surface nano-roughness, demonstrating how these factors can jointly drive cell behavior and clustering. Lastly, we further elaborate on the biological significance of key networks metrics, such as the small-world coefficient.

Further to this end, much of the claims of this perspective assume that nano-topography enhances the formation of cellular networks. In fact, this hypothesis has long been proven. The work we are submitting is part of ongoing research effort aimed at engineering bio-interfaces and developing mathematical models^[13–15,17,72–75] to elucidate how nano-topography promotes cellular networking and, in turn, influences system performance. While many of these studies are discussed throughout this perspective, we provide a comprehensive overview in a separate Supporting Information 5, offering additional context, further demonstrating the feasibility and significance of this approach, and adding perspective to this work.

Thus, nano-topography enhanced topological-cell-analysis can achieve accurate analysis of how cells respond to radiation therapy.

In turn, precise quantification of the sensitiveness of cells can help to choose the right dose and the right time of the treat-




Figure 6. A nanoscale device is the main project outcome.

ment, increasing life expectancy of cancer patient and reducing the costs of follow-up. Both these aspects can positively impact on the efficiency of national health services. To investigate cell-radiation interaction, it is recommended to use both 2D and 3D models. Conventional and time modulated (i.e., FLASH RT) administration of radiation can be used to optimize the method even further and find the best treatment conditions of cancer. Moreover, different RT techniques (i.e., X-ray, electron, proton RT) may help to demonstrate repeatability, reliability and accuracy of the method under several different working conditions. *While the use of nano-topography to guide cell behavior is very well assessed, the integration of nanopatterned scaffolds and topological cell analysis does not have any precedent in the literature for the evaluation of the effects of radiotherapy – and offers ways to measure the performance of radiotherapy with unprecedented accuracy.*

The outcomes of a similar integration are manifold, and can be reviewed as follows.

3.1. Technology Outcome

One of the main objectives of this perspective is suggesting ways for evaluating the RR and RS of cancer cells. Thus, the first product of a similar suggested approach is a nanoscale device (Figure 6). If used as a substrate for cells extracted from a patient with cancer, the device will guide cells into structures with some level of complexity. Cells will be then inspected by conventional fluorescence microscopy, and resulting images examined employing the methods developed using Networks Science and AI. Convenient algorithms will give an estimate of the RS and RR of cells with elevated sensitivity and specificity, high accuracy, and low limit of detection. The RS and RR of cancer cells – assessed through the method and the device – will be used to tailor the right dose and the right time of the treatment with RT.

3.2. Scientific Outcome

Alongside with the development of the device, problems that are still open will be possibly addressed. The first is elucidating the exact nature of the relationship between tissue topology,

tissue physiology, and scaffold topology. While the effects of nano-topography on cell clustering have been occasionally examined, still a systematic study on how the topology of a scaffold is correlated to the characteristics of growing cells and networks is lacking. The study herein proposed will examine how the topology, of the scaffold and of the cells, correlates to the functions of cells cultured on them. The correspondence – *equivalence* – between structure and function is a paradigm shift, in that it allows to estimate the effects of the geometry of a system on the performance of that system. Moreover, since Raman Phenotyping will be used to characterize cells, it will also be dissected whether there is a correlation between the topology of cell networks and the phenotypic distribution in those networks. This will allow establishing a correlation between scaffold topology, cell topology, and the spatial distribution of phenotypes: indicating which phenotypes are expressed by cells in response to a specific combination of parameters. Many groups worldwide are trying to develop 3D, self-organizing, in-vitro cell-culture organ models. These 3D structures harbor key features of their native organs and can be used to investigate and understand biological mechanisms and diseases, including cancer. Against this background, we are setting a new powerful paradigm. The perception that the topological characteristics of a system, cell's phenotype, and the function of that system are tightly interwoven – can offer guidance to design 3D cell models and organoids with unprecedented efficiency.

3.3. Social outcome

This perspective centers on enhancing cancer treatment efficiency, particularly in radiotherapy. Cancer, a leading cause of morbidity and death globally, incurs substantial financial costs. RT, a crucial treatment modality, involves high expenses. This method promises to personalize treatment, potentially increasing survival rates, reducing treatment costs, and improving healthcare resource utilization. It offers a cost-effective alternative to traditional methods, potentially revolutionizing cancer care by enabling more targeted and efficient treatment strategies. While this concept is here immediately applied to RT, it can be easily translated to other fields of biomedical research. *As for an example, the technology of prosthetic implants and medical implants.* The still-high rejection rate of implanted devices from the human body is cause of concern of the medical industry, especially because there are no consistently effective strategies to prevent or suppress it. An effectively designed implant, guided by the principle of structure-function equivalence, can reduce the risk of rejection and enhance performance. Further to this end, a similar principle can contribute to reduce the chance of postoperative infection, simply choosing values of geometric variables of the implant that can possibly impair the growth and proliferation of bacterial films.

4. Study Design and Implementation

Here we outline an experimental plan for the development and optimization of surface-enhanced topological cell analysis. The plan is here just sketched, its exact implementation does depend on the resources, the instrumentation, and the facilities of the research groups involved in the research.

4.1. Fabrication

2D surfaces and 3D scaffolds with controlled geometric characteristics and nano-topography are designed, fabricated and characterized. These platforms enable interaction of cancer cells with engineered environments.

4.2. Biological Preparation

Human cancer cells, including those derived from melanoma, breast, and lung cancers, provide a foundation for this research. Using conventional biological assays, one can evaluate cellular RR and RS, and build a biological database that is necessary for the rest of the study. Moreover, simple adhesion and growth of cells can be examined on 2D support and 3D devices. This approach ensures the identification of optimal conditions to enhance cellular response to radiotherapy. Selected devices will be incubated with the cells identified for this study.

4.3. RT Experiments

By exposing selected cell-support systems to various RT modalities—including conventional X-ray RT, ultra-high dose rate FLASH electron RT, and FLASH proton RT—one can generate invaluable data on how these factors shape cellular response as a function of the supporting devices. Fluorescence microscopy can be used to acquire cellular networks on the devices and dissect the nature of cell-cell interactions.

4.4. Topological Cell Analysis

Morphological and topological characteristics of cancer-cell networks will be then examined using computational tools. Using topological data determined from networks-science analysis, one will dissect whether and to what extent cancer traits, scaffold geometry, radiation dose, and network topology are associated. This relationship promises to unravel previously unrecognized patterns in cellular behavior.

4.5. Data Analysis

Once the topological characteristics of the cancer-cell networks are determined, researchers will use statistical techniques to examine whether treatment with radiations is a special-cause of variation of networks topology, and find specific cause-effect (radiation-topology) laws.

4.6. Artificial Intelligence

Artificial intelligence (AI), encompassing both deep learning (DL) and machine learning (ML), will serve as a powerful ally, identifying intricate relationships within extensive imaging datasets. These insights will be instrumental in predicting RR and RS across cancer cell types with increased accuracy compared to simple analytical techniques. A more detailed description on how AI can be practically implemented to examine cell response to radiation, and its integration with networks science, is reported in a separate Supporting Information 6.

4.7. Raman Phenotyping

At the same point, Raman spectroscopy will complement this effort by providing multifactorial characterization of 2D and 3D cultures, adding another layer of depth to our understanding of cellular phenotypes and their environmental interactions.

4.8. General Model of Cell Response

Finally, basing on data, a general model of correlation between scaffold-topography, cell-topology, and phenotypes spatial distribution will be built.

Thus, a side effect of this study-design is understanding how the topology of cell networks and the phenotypic traits expressed by the cells correlate: it aims to establish a direct link between the physical and geometric properties of the scaffolds, the organizational patterns of cells, and their phenotypic responses. By establishing a *direct link between the physical and geometric properties of scaffolds and the functional responses of cells*, this study not only advances cancer research but also introduces a transformative framework for the design of engineered cell models and targeted therapeutic strategies.

One key ingredient in the study is *networks science*. This study proposal extends the existing knowledge on how cancer cell-networks behave in response to RT^[49,50] and sets a novel, powerful paradigm in radiotherapy, adding a very significant improvement with clinical implications.

5. Conclusion

Current RR and RS assessment of human cancer cells primarily relies on estimating the survival fraction of cells on a surface, a method with limited sensitivity, precision, and accuracy. In this perspective, we propose a multifactorial approach that considers colony shape, internal architecture, and biochemical properties. A key novelty of this approach is the integration of previously overlooked factors, such as topology, into RT assessment. Additionally, we introduce nano-topography as an amplifying agent, enhancing the visibility of otherwise imponderable differences between radiosensitive and radioresistant cells by leveraging nanoscale materials. The introduction of materials science and nanotechnology in the field of RT, along with alternative methods of analysis based on networks science and topology, can realistically improve RT treatment of patients and, as a collateral effect, reduce costs.

Beyond RT, our approach aims to establish a broader correlation between structure and function in biological systems—a concept with far-reaching biomedical implications. In systems like the nervous system or neuromorphic computing, function is inherently tied to structure. By measuring one, we can predict the other. The direct utilization of this principle, *from structure to function*, is useful in diagnostics, i.e., for estimating the effects of radiotherapy, the progression of cancer, the progression of a neurodegenerative disease. The indirect utilization of this principle, *from function to structure*, can be exploited in tissue engineering, reverse engineering, and related fields.

This technology has the potential to impact a wide range of fields, including cancer diagnostics and therapy, tissue repair and

regeneration, neurodegenerative disease diagnosis, and neuromorphic engineering.

In the context of cancer therapy, it is essential to further elaborate on this method. The nanotechnology that we have devised is designed to assess the RR and RS of cells by analyzing their collective behaviour upon radiation exposure. This approach enables physicians to evaluate a patient's condition more accurately and design personalized treatment strategies accordingly. However, the method can be improved even further. The nanoscale material used to boost cell clustering can be, among many other possibilities, mesoporous silicon. Mesoporous silicon has the remarkable property of working as a nanoscale drug delivery system. Consequently, the method can be advanced into a theranostic system, simultaneously facilitating both diagnosis and treatment. The same platform that directs and analyzes cell assembly could also be employed to deliver radiosensitizers to targeted cells. This method could be used to study the effects of therapeutic agents on radio-resistant cells, enabling a more sophisticated and personalized diagnostic approach for each patient. Potentially improving over the existing nanoscale drug delivery systems already employed to overcome RR and previously reviewed.

Supporting Information

Supporting Information is available from the Wiley Online Library or from the author.

Acknowledgements

This study was partially funded by the Italian Association for Cancer Research (AIRC) under the grant number AIRC IG 2021 ID 25656.

Open access publishing facilitated by Universita degli Studi Magna Graecia di Catanzaro, as part of the Wiley - CRUI-CARE agreement.

Conflict of Interest

The authors declare no conflict of interest.

Keywords

AI, biomaterials, nano-topography, networks science, radiation-therapy, Raman phenotyping, scaffolds, topology

Received: December 23, 2024
Revised: March 11, 2025
Published online: March 22, 2025

- [1] J. Sia, R. Szmyd, E. Hau, H. E. Gee, *Front. Cell Dev. Biol.* **2020**, *8*, 41.
- [2] P. Liu, R. Song, G. L. Elison, W. Peng, M. Acar, *Nat. Commun.* **2017**, *8*, 680.
- [3] M. K. Thompson, P. Poortmans, A. J. Chalmers, C. Faivre-Finn, E. Hall, R. A. Huddart, Y. Lievens, D. Sebag-Montefiore, C. E. Coles, *Br. J. Cancer*. **2018**, *119*, 389.
- [4] L. Tang, F. Wei, Y. Wu, Y. He, L. Shi, F. Xiong, Z. Gong, C. Guo, X. Li, H. Deng, *J. Exp. Clin. Cancer Res.* **2018**, *37*, 87.
- [5] L. Tirinato, M. G. Marafioti, F. Pagliari, J. Jansen, I. Aversa, R. Hanley, C. Nisticò, D. Garcia-Calderón, G. Genard, J. F. Guerreiro, *Elife*. **2021**, *10*, 72943.

- [6] R. Ananthakrishnan, A. Ehrlicher, *Int. J. Biol. Sci.* **2007**, *3*, 303.
- [7] F. Gentile, *J. Biomech.* **2021**, *115*, 110134.
- [8] C. Heisenberg, Y. Bellaïche, *Cell* **2013**, *153*, 948.
- [9] C. Cimmino, L. Rossano, P. A. Netti, M. Ventre, *Front. Bioeng. Biotechnol.* **2018**, *6*, 190.
- [10] H. Lou, W. Zhao, X. Li, L. Duan, A. Powers, M. Akamatsu, F. Santoro, A. F. McGuire, Y. Cui, D. G. Drubin, *Proc. Natl. Acad. Sci. USA* **2019**, *116*, 23143.
- [11] M. M. Stevens, J. H. George, *Science* **2005**, *310*, 1135.
- [12] M. Ventre, C. F. Natale, C. Rianna, P. A. Netti, *J. R. Soc., Interface* **2014**, *11*, 20140687.
- [13] G. Marinaro, R. La Rocca, A. Toma, M. Barberio, L. Cancedda, E. Di Fabrizio, P. Decuzzi, F. Gentile, *Integr. Biol.* **2015**, *7*, 184.
- [14] V. Onesto, L. Cancedda, M. L. Coluccio, M. Nanni, M. Pesce, N. Malara, M. Cesarelli, E. Di Fabrizio, F. Amato, F. Gentile, *Sci. Rep.* **2017**, *7*, 9841.
- [15] V. Onesto, M. Villani, R. Narducci, N. Malara, A. Imbrogno, M. Allione, N. Costa, N. Coppedè, A. Zappettini, C. V. Cannistraci, *Sci. Rep.* **2019**, *9*, 4021.
- [16] J. J. Hopfield, *Proc. Natl. Acad. Sci. USA* **1982**, *79*, 2554.
- [17] G. Marinaro, L. Bruno, N. Pirillo, M. L. Coluccio, M. Nanni, N. Malara, E. Battista, G. Bruno, F. De Angelis, L. Cancedda, *Commun. Biol.* **2024**, *7*, 617.
- [18] F. Aprile, V. Onesto, F. Gentile, *NPJ Syst. Biol. Appl.* **2022**, *8*, 4.
- [19] M. El-Gaby, H. M. Reeve, V. Lopes-dos-Santos, N. Campo-Urriza, P. V. Perestenko, A. Morley, L. A. Strickland, I. P. Lukács, O. Paulsen, D. Dupret, *Nat. Neurosci.* **2021**, *24*, 694.
- [20] L. Ma, J. M. Hyman, A. J. Lindsay, A. G. Phillips, J. K. Seamans, *Nat. Neurosci.* **2014**, *17*, 1100.
- [21] U. S. Bhalla, R. Iyengar, *Science* **1999**, *283*, 381.
- [22] W. R. MacLellan, Y. Wang, A. J. Lusis, *Nat. Rev. Cardiol.* **2012**, *9*, 172.
- [23] L. Rizik, L. Daniai, M. Habib, R. Weiss, R. Daniel, *Nat. Commun.* **2022**, *13*, 5602.
- [24] J. M. Ottino, *Nature* **2004**, *427*, 399.
- [25] Y. Liu, A. Barabási, *Rev. Mod. Phys.* **2016**, *88*, 035006.
- [26] S. H. Strogatz, *Nature* **2001**, *410*, 268.
- [27] J. Kwapień, S. Drożdż, *Phys. Rep.* **2012**, *515*, 115.
- [28] M. Pósfai, A. Barabási, *Netw. Sci.* **2016**.
- [29] J. Wang, Y. Zhang, C. Xu, J. Li, J. Sun, J. Xie, L. Feng, T. Zhou, Y. Hu, *Nat. Commun.* **2024**, *15*, 2849.
- [30] A. Barabási, *Nat. Phys.* **2012**, *8*, 14.
- [31] D. J. Watts, S. H. Strogatz, *Nature* **1998**, *393*, 440.
- [32] D. J. Watts, *Small Worlds: The Dynamics of Networks between Order and Randomness*, Princeton University Press, New Jersey United States, **2003**, p. 1.
- [33] M. D. Humphries, K. Gurney, *PLoS One* **2008**, *3*, e0002051.
- [34] A. Abend, C. Steele, S. Schmidt, R. Frank, H. Jahnke, M. Zink, *Int. J. Mol. Sci.* **2020**, *21*, 6249.
- [35] D. S. Bassett, E. D. Bullmore, *Neuroscientist* **2006**, *12*, 512.
- [36] L. F. Lago-Fernández, R. Huerta, F. Corbacho, J. A. Sigüenza, *Phys. Rev. Lett.* **2000**, *84*, 2758.
- [37] V. Latora, M. Marchiori, *Phys. Rev. Lett.* **2001**, *87*, 198701.
- [38] M. Ballerini, N. Cabibbo, R. Candelier, A. Cavagna, E. Cisbani, I. Giardina, V. Lecomte, A. Orlandi, G. Parisi, A. Procaccini, *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 1232.
- [39] J. P. Crutchfield, *Nat. Phys.* **2012**, *8*, 17.
- [40] G. Tononi, G. M. Edelman, *Science* **1998**, *282*, 1846.
- [41] C. Jensen, Y. Teng, *Front. Mol. Biosci.* **2020**, *7*, 33.
- [42] G. A. Romar, T. S. Kupper, S. J. Divito, *J. Invest. Dermatol.* **2016**, *136*, e1.
- [43] E. Boutet-Robinet, D. Trouche, Y. Canitrot, *Bio-protocol* **2013**, *3*.
- [44] L. G. Mariotti, G. Pirovano, K. I. Savage, M. Ghita, A. Ottolenghi, K. M. Prise, G. Schettino, *PLoS One* **2013**, *8*, e79541.
- [45] N. A. Franken, H. M. Rodermond, J. Stap, J. Haveman, C. Van Bree, *Nat. Protoc.* **2006**, *1*, 2315.
- [46] A. L. Dunne, M. E. Price, C. Mothersill, S. R. McKeown, T. Robson, D. G. Hirst, *Br. J. Cancer.* **2003**, *89*, 2277.
- [47] N. R. Sham, N. Hasan, N. A. H. Hasani, M. K. Karim, M. J. Ibahim, *J. Phys.* **2020**, *1497*, 012032.
- [48] M. Yasser, R. Shaikh, M. K. Chilakapati, T. Teni, *PLoS One* **2014**, *9*, e97777.
- [49] L. Tirinato, V. Onesto, D. Garcia-Calderon, F. Pagliari, M. Spadea, J. Seco, F. Gentile, *Sci. Rep.* **2022**, *12*, 12980.
- [50] L. Tirinato, V. Onesto, D. Garcia-Calderon, F. Pagliari, M. Spadea, J. Seco, F. Gentile, *Ann. Biomed. Eng.* **2023**, *51*, 1859.
- [51] B. Geiger, J. P. Spatz, A. D. Bershadsky, *Nat. Rev. Mol. Cell Biol.* **2009**, *10*, 21.
- [52] P. Kanchanawong, G. Shtengel, A. M. Pasapera, E. B. Ramko, M. W. Davidson, H. F. Hess, C. M. Waterman, *Nature* **2010**, *468*, 580.
- [53] P. Decuzzi, M. Ferrari, *Biomaterials* **2010**, *31*, 173.
- [54] F. Gentile, L. Tirinato, E. Battista, F. Causa, C. Liberale, E. M. di Fabrizio, P. Decuzzi, *Biomaterials* **2010**, *31*, 7205.
- [55] F. Gentile, R. Medda, L. Cheng, E. Battista, P. E. Scopelliti, P. Milani, E. A. Cavalcanti-Adam, P. Decuzzi, *Sci. Rep.* **2013**, *3*, 1461.
- [56] M. L. Coluccio, V. Onesto, G. Marinaro, M. Dell'Apa, S. De Vitis, A. Imbrogno, L. Tirinato, G. Perozziello, E. Di Fabrizio, P. Candeloro, *Pharmaceutics* **2020**, *12*, 481.
- [57] B. Geiger, A. Bershadsky, R. Pankov, K. M. Yamada, *Nat. Rev. Mol. Cell Biol.* **2001**, *2*, 793.
- [58] F. Gentile, R. La Rocca, G. Marinaro, A. Nicastrì, A. Toma, F. Paonessa, G. Cojoc, C. Liberale, F. Benfenati, E. Di Fabrizio, *ACS Appl. Mater. Interfaces* **2012**, *4*, 2903.
- [59] Q. Akolawala, M. Rovituso, H. H. Versteeg, A. M. Rondon, A. Accardo, *ACS Appl. Mater. Interfaces* **2022**, *14*, 20778.
- [60] Q. Akolawala, F. Keuning, M. Rovituso, W. van Burik, E. van der Wal, H. H. Versteeg, A. M. Rondon, A. Accardo, *Adv. Healthcare Mater.* **2024**, *13*, 2302988.
- [61] K. Ganesh, C. Wu, K. P. O'Rourke, B. C. Szeglin, Y. Zheng, C. G. Sauvè, M. Adileh, I. Wasserman, M. R. Marco, A. S. Kim, *Nat. Med.* **2019**, *25*, 1607.
- [62] A. Acheva, M. Ghita, G. Patel, K. M. Prise, G. Schettino, *PLoS One* **2014**, *9*, e86092.
- [63] G. Xue, Z. Ren, P. W. Grabham, Y. Chen, J. Zhu, Y. Du, D. Pan, X. Li, B. Hu, *J. Radiat. Res.* **2015**, *56*, 656.
- [64] N. Barin, H. E. Balcioglu, I. de Heer, M. de Wit, M. L. Lamfers, M. E. van Royen, P. J. French, A. Accardo, *Small* **2022**, *18*, 2204485.
- [65] C. R. Arnold, J. Mangesius, I. Portnaia, U. Ganswindt, H. A. Wolff, *Front. Oncol.* **2024**, *14*, 1379986.
- [66] X. Song, Z. Sun, L. Li, L. Zhou, S. Yuan, *Front. Oncol.* **2023**, *13*, 1088878.
- [67] D. J. J. Poon, L. M. Tay, D. Ho, M. L. K. Chua, E. K. Chow, E. L. L. Yeo, *Cancer Lett.* **2021**, *511*, 56.
- [68] Y. Liu, P. Zhang, F. Li, X. Jin, J. Li, W. Chen, Q. Li, *Theranostics* **2018**, *8*, 1824.
- [69] J. F. Hainfeld, F. A. Dilmanian, D. N. Slatkin, H. M. Smilowitz, *J. Pharm. Pharmacol.* **2008**, *60*, 977.
- [70] S. Pinel, N. Thomas, C. Boura, M. Barberi-Heyob, *Adv. Drug Delivery Rev.* **2019**, *138*, 344.
- [71] G. Le Duc, I. Miladi, C. Alric, P. Mowat, E. Bräuer-Krisch, A. Bouchet, E. Khalil, C. Billotey, M. Janier, F. Lux, *ACS Nano* **2011**, *5*, 9566.
- [72] F. Aprile, V. Onesto, F. Gentile, *NPJ Syst. Biol. Appl.* **2022**, *8*, 4.
- [73] V. Onesto, R. Narducci, F. Amato, L. Cancedda, F. Gentile, *Integr. Biol.* **2018**, *10*, 121.
- [74] T. Limongi, F. Cesca, F. Gentile, R. Marotta, R. Ruffilli, A. Barberis, M. Dal Maschio, E. M. Petrini, S. Santoriello, F. Benfenati, *Small* **2013**, *9*, 402.
- [75] V. Onesto, A. Accardo, C. Vieu, F. Gentile, *Neural Regen. Res.* **2020**, *15*, 759.




Francesca Pagliari is a Senior Research Fellow at the German Cancer Research Center (DKFZ), specializing in oxidative stress, cancer cell radio resistance, and lipid metabolism. With a PhD in *Materials for Environment and Energy* (2011), her interdisciplinary expertise bridges stem cell biology, nanomedicine, and radiation oncology. Francesca has led pivotal research projects, supervised students, and co-authored over 30 publications, including studies on lipid droplets' role in cancer therapy. She holds a patent on cancer cell migration prevention and actively reviews for high-impact journals. Her innovative work continues to advance understanding of cancer biology and regenerative medicine.



Maria Francesca Spadea currently holds the chair of Modelling and Simulation for Medical Engineering at Karlsruhe Institute of Technology (Karlsruhe, Germany) where she co-leads the Institute of Biomedical Engineering. Her main research interests are the implementation and the validation of computational methodologies in the biomedical field for supporting diagnosis and therapy. Research lines can be summarized as following: optical and surface localization for patient positioning in radiotherapy and surgery; medical image processing and registration; 4D image reconstruction; image-to-image translation; radiomics analysis for patient stratifications; Medical Data Science.




Pierre Montay-Gruel is a radiobiologist and research group leader at the Iridium Network in Antwerp, Belgium. He also serves as a visiting professor at the University of Antwerp's Centre for Oncological Research. His research is centered on ultra-high dose rate FLASH radiation therapy and innovative approaches in radiation therapy. Montay-Gruel's team develops both in vitro and in vivo models to investigate the responses of normal tissue and tumors to ultra-high dose rate FLASH-RT, exploring molecular mechanisms and functional endpoints. Their ultimate goal is to unravel the mechanisms underlying the FLASH effect and translate FLASH-RT into clinical practice.



I – Anggraeni Puspitasari Kokko (she/her), born on February 10, 1985 – am a medical doctor and radiation biologist specializing in radiation neurobiology and cancer research. I hold an MD from Maranatha Christian University, Indonesia (2011), and a PhD in medicine from Gunma University, Japan (2016). I completed my postdoctoral research in Biophysics at GSI, Germany (2019–2024). Since February 2024, I have been working as a Radiation Biologist at HollandPTC and a research Collaborator at Gunma University. My research explores radiation effects on the central nervous system, cancer metastasis, FLASH irradiation, and hibernation for radiation protection, including collaborations with JAXA and ESA.



Joao Seco graduated with a PhD from the University of London, at the Institute of Cancer Research (ICR) and Royal Marsden Hospital in London, UK. He then went on to become an Assistant Professor of Radiation Oncology at Harvard Medical School in Boston, working at the Massachusetts General Hospital (MGH). He then returned to Europe to work at the German Cancer Research Center, DKFZ in Heidelberg, heading up a new group dedicated to ion beam research and with the focus on 1) novel imaging technologies to reduce Bragg peak positioning errors in patients and 2) on investigating the mechanism of radiation triggered DNA damage via reactive oxygen species. He is also presently the Chair of Medical Physics at the Department of Physics and Astronomy, Heidelberg University and is a member of the EFOMP Scientific Committee, representing the DGMP, German Society for Medical Physics.



Luca Tirinato is Associate Professor at the Department of Medical and Surgical Sciences at the University of Magna Graecia of Catanzaro. He has several years of experience in top international research centers such as IIT, KAUST and DKFZ. His scientific interests are focused on the study and understanding of lipid alterations and accumulation (e.g., lipid droplets) in metabolic and neoplastic diseases. He is also interested in understanding the molecular mechanisms underlying the formation and secretion of small extracellular vesicles. Particular attention is given to the use of label-free imaging techniques (e.g., Raman spectroscopy) to study the above biological systems.



Angelo Accardo is Associate Professor at the Faculty of Mechanical Engineering, Department of Precision and Microsystems Engineering of Delft University of Technology, co-founder of the Delft Young Academy, and Steering Committee Member of the International Micro and Nanoengineering (MNE) Conference. His main research interest is to develop versatile multi-scale (nano-micro-meso) fabrication paradigms to manufacture 3D engineered cell microenvironments. By exploiting high-resolution light-assisted 3D additive manufacturing techniques (two-photon polymerization, stereolithography, micro-digital light projection), he creates engineered polymeric/hydrogel microenvironments for fundamental mechanobiology, in vitro disease/treatment modeling and tissue engineering applications involving brain (e.g., microglia, iPSCs-derived/primary neurons), cancer (e.g., glioblastoma), or bone (e.g., preosteoblasts) cells.



Francesco De Angelis earned his MD in Physics from the University of Rome in 2000 and a PhD in Physics of Matter in 2004. He currently heads the Plasmon Nanotechnologies Unit at the Italian Institute of Technology. He has authored over 180 publications, primarily in the fields of nanotechnology and biomedical physics. His expertise lies in designing and fabricating nanostructures for applications in optics, electronics, and mechanics, with a particular focus on molecular sensing in living tissues or biological samples. His team is composed of physicists, chemists, biotechnologists, pharmacologists and engineers collaborating in a very multi-disciplinary group that aims to control biological systems at the nanoscale.



Francesco Gentile (FG) does research in biomedical nanotechnology. He uses mathematical modelling and nanotechnology to engineer solutions to biomedical problems. FG earned his MS in Mechanical Engineering at the University of Calabria in 2003, and a PhD in Biomedical Engineering at the University Magna Graecia (UMG) of Catanzaro in 2008. FG has been a Visiting Scholar at the Ohio State University (2004), the Synchrotron Light Laboratory in Trieste (2005), the University of Texas at Houston (2007). From 2010 to 2013, FG worked as a post doc, and then as a Scientist, at the iit in Genoa, Italy. In 2014, FG was appointed associate professor of biomedical engineering at the University Federico II of Naples. From 2020, FG works at UMG.