

How lichen's 3D architecture temporally and spatially regulates the passage of water, gases and food

ten Veldhuis, Marie-claire; Dismukes, Charles; Ananyev, Gennady

Publication date

2018

Document Version

Final published version

Published in

Geophysical Research Abstracts (online)

Citation (APA)

ten Veldhuis, M., Dismukes, C., & Ananyev, G. (2018). How lichen's 3D architecture temporally and spatially regulates the passage of water, gases and food. *Geophysical Research Abstracts (online)*, 20, Article EGU2018-16048.

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.



How lichen's 3D architecture temporally and spatially regulates the passage of water, gases and food

Marie-Claire ten Veldhuis (1), Charles Dismukes (2,3), Gennady Ananyev (2,3)

(1) Delft University of Technology, Watermanagement Department, Delft, Netherlands (j.a.e.tenveldhuis@tudelft.nl), (2) Rutgers University, the State University of New Jersey, Department of CCB, (3) Rutgers University, the State University of New Jersey, Waksman Institute of Microbiology

Lichens have developed unique strategies to control water, oxygen and photosynthate (sugar product) fluxes between the algal photobiont and its fungal host that provide a model for gas-liquid separation and transport. We measured in parallel the kinetics of dehydration (water content), light trapping by charge separation and oxygen evolution (chlorophyll fluorescence emission) from full hydration to the dehydrated state. Kinetics show two distinct temporal stages in lichen dehydration: rapid loss of free, extracellular water, followed by slower loss of intracellular water associated with a discontinuous (rapid) decrease in charge separation. The former shows that water saturation does not prevent lichens from being photosynthetically active and allows gas transport (CO_2 and O_2) to occur during water logging. The latter observation indicates that dehydration is used to cooperatively turn off photosynthesis. This is hypothesized to trigger transport of algal-derived sugars to the fungal host. Confocal microscopy of lichens shows that the 3D architecture has distinct zones of physical space for gas and water transport between alga and fungus. Algal cells are encased in a hydrophobic shell that limits water accessibility and channels CO_2 and O_2 gas respectively to the photobiont and fungus (for respiration). Algae are connected to a single hydrophilic fungal filament that delivers water and recovers algal products (sugar & O_2), synchronized to the daily solar cycle. This bipolar chemical asymmetry enables such counter transport. Hence, the daily hydration/dehydration cycle serves as a pump.